

## Differential *in vitro* inhibition of polyphenoloxidase from a wild edible mushroom *Lactarius salmonicolor*

NURCAN DEDEOGLU & OZEN OZENSOY GULER

Department of Chemistry, Balikesir University Science & Art Faculty, Cagis-Kampus, 10100 Balikesir, Turkey

(Received 11 February 2008)

### Abstract

The polyphenol oxidase (LsPPO) from a wild edible mushroom *Lactarius salmonicolor* was purified using a Sepharose 4B-L-tyrosine-*p*-amino benzoic acid affinity column. At the optimum pH and temperature, the  $K_M$  and  $V_{Max}$  values of LsPPO towards catechol, 4-methylcatechol and pyrogallol were determined as 0.025 M & 0.748 EU/mL,  $1.809 \times 10^{-3}$  M & 0.723 EU/mL and  $9.465 \times 10^{-3}$  M & 0.722 EU/mL, respectively.

Optimum pH and temperature values of LsPPO for the three substrates above ranged between the pH 4.5–11.0 and 5–50°C. Enzyme activity decreased due to heat denaturation with increasing temperature. Effects of a variety of classical PPO inhibitors were investigated upon the activity of LsPPO using catechol as the substrate.  $IC_{50}$  values for glutathione, *p*-aminobenzenesulfonamide, L-cysteine, L-tyrosine, oxalic acid,  $\beta$ -mercaptoethanol and syringic acid were determined as  $9.1 \times 10^{-4}$ ,  $2.3 \times 10^{-4}$  M,  $1.5 \times 10^{-4}$  M,  $3.8 \times 10^{-7}$  M,  $1.2 \times 10^{-4}$  M,  $4.9 \times 10^{-4}$  M, and  $4 \times 10^{-4}$  M respectively. Thus L-tyrosine was by far the most effective inhibitor. Interestingly, sulfosalicylic acid behaved as an activator of LsPPO in this study.

**Keywords:** *Lactarius salmonicolor*, affinity chromatography, inhibition, enzymatic browning, polyphenoloxidase

### Introduction

The collection and consumption of wild edible fungi has traditionally been important to the livelihoods of many people in Northern Eurasia, and is still important, particularly in developing countries. It is a healthy food that can make a useful dietary contribution, being a good source of digestible proteins with low but balanced lipid content, and possessing useful amounts of phosphorus, potassium, selenium, zinc, magnesium, copper and B vitamins [1]. Some dried mushrooms and concentrated extracts are also used for their medicinal properties and as dietary supplements. Thus there is also a strong and increasing commercial interest in fungi, with demand often outstripping local supply [2]. Unfortunately mushrooms are easily prone to browning when they are subjected to forces that can disrupt cellular

integrity, such as vibration, rough handling, and ageing [3,4].

Tyrosinases are widely distributed among animals, plants and fungi [5,6]. They are responsible for many biologically essential functions, such as pigmentation, sclerotization, primary immune response and host defense but in mushrooms they are responsible for the browning reactions. Browning reactions are major causes of quality loss during harvesting, post-harvest handling/storage, and processing of fruits, plants and vegetables in food industry [7]. The enzymatic browning causes deterioration of sensory and nutritional quality and affects appearance and organoleptic properties, inactivation of PPO is desirable for preservation of foods [8]. Several methods such as the addition of antioxidants and the exclusion of oxygen as well as thermal processing have been used to inhibit enzymatic browning. For inactivation of PPO,

Correspondence: Dr Ozen Ozensoy Guler, Department of Chemistry, Balikesir University Science & Art Faculty, Cagis/Kampus, 10100 Balikesir, Turkey. Tel: 90 266 612 1263/1112. Fax: 90 266 612 1215. E-mail: ozensoy@balikesir.edu.tr

thermal processing has limits like loss of sensory and nutritional quality of food products. Therefore, high-pressure treatment has been considered as an alternative [9,10].

Enzymatic browning via PPO causes deterioration of sensory and nutritional quality and affects appearance and organoleptic properties, thus inactivation of PPO is desirable in food preservation [8]. Several methods such as the addition of antioxidants and the exclusion of oxygen, as well as thermal processing, have been used to inhibit enzymatic browning. However, thermal processing has limits dictated by consequent loss of sensory and nutritional quality of food products. Therefore, high-pressure treatment has been considered as an alternative process of PPO inactivation [9,10].

Here we have purified PPO from the edible wild mushroom, *Lactarius salmonicolor*, by a single-step affinity procedure, and investigated various enzymic characteristics, including its substrate specificity, kinetics, pH and temperature optima, temperature inactivation, and chemical inhibition.

## Materials and methods

Edible mushrooms (*Lactarius salmonicolor*) used in this study were harvested in the middle of November from a forest near Balıkesir in Turkey. The extract of mushroom was prepared as quickly as possible and stored deep-frozen at  $-80^{\circ}\text{C}$  until used. All chemicals used in this study were the best grade available. Enzyme assays were measured with the aid of a Biotek automated recording spectrophotometer. Sepharose 4B,L-tyrosine-p-amino benzoic acid affinity gel which used in this study was synthesized at Balıkesir University, Research Center of Applied Sciences (BURCAS/Balıkesir, Turkey) in Biology section laboratory [7].

### Purification of PPO

All purification steps were carried out at  $4^{\circ}\text{C}$ . The extraction procedure was adopted from Wesche-Ebeling & Montgomery [11]. Firstly, the *Lactarius salmonicolor* mushrooms were washed with distilled water three times. Secondly, to prepare the crude extract, 50 g of mushrooms were cut quickly into thin slices and homogenized in a Waring blender for 2 min using 100 ml of 0.1 M phosphate buffer, pH:7.3 containing 5% poly(ethylene glycol) and 10 mM ascorbic acid. After filtration of the homogenate through a muslin, the filtrate was centrifuged at  $15\,000 \times g$  for 30 min, and the supernatant was collected. A crude protein precipitate was made by adding  $(\text{NH}_4)_2\text{SO}_4$  to 80% saturation. The resulting precipitate was suspended in a minimum volume of 5 mM phosphate buffer and then dialyzed against 5 the same buffer overnight. The enzyme solution was then

applied to the Sepharose 4B-tyrosine-p-amino benzoic acid affinity column [7], pre equilibrated with 5 mM phosphate buffer, pH 5.0. The affinity gel was extensively washed with the same buffer before the *Lactarius salmonicolor* PPO (LsPPO) was eluted with 1 M NaCl, 5 mM phosphate, pH 7,0.

### LsPPO activity

Enzyme activity was determined, using different mono- or di-phenolic compounds, by measuring the increase in absorbance at 494 nm for 4-methylcatechol and 500 nm for all other substrates [12] respectively, in a Biotek automated recording spectrophotometer. Enzyme activity was calculated from the linear portion of the curve. One unit of PPO activity was defined as the amount of enzyme that causes an increase in absorbance of  $0.001 \text{ units min}^{-1}$  for 1 ml of enzyme at  $25^{\circ}\text{C}$  [7].

### Protein determination

Protein was determined by the method of Bradford [13] using bovine serum albumin as a standard. In chromatography, protein was expressed as absorbance at 280 nm.

### Polyacrylamide gel electrophoresis

Polyacrylamide gel slab electrophoresis of purified enzyme was carried out according to the method of Laemmli [14].

### Enzyme kinetics and substrate specificity

LsPPO activity was assayed using pyrogallol, catechol and 4-methyl catechol as substrates. The rate of reaction was measured as the increase in absorbance at the absorption maxima of the corresponding quinone product for each substrate. One unit of enzyme activity was defined as the amount of enzyme causing a change of 0.001 in absorbance per minute., LsPPO activities were measured with the substrates at varying concentrations (2.0, 4.0, 6.0, 8.0, 10.0, 12.0, and 15.0 mM) under optimum conditions of pH, ionic strength, and temperature. For each substrate, the Michaelis-Menten constant  $K_M$  and maximum velocity  $V_{\text{max}}$  values were calculated from a plot of  $1/V$  against  $1/[S]$  by the method of Lineweaver and Burk (Figure 2).

### Effect of pH

LsPPO activity as a function of pH was determined using catechol as substrate (0.1 M stock concentration). The buffers used were 0.1 M acetate (pH 4.5–6.0) and 0.1 M phosphate (pH 6.0–9.5)




Figure 1. Purification of LsPPO by elution from the affinity gel.

### Effect of temperature

LsPPO activity, as a function of temperature, was determined under standard assay conditions, using temperatures from 20 to 80°C with pyrogallol and catechol as substrates. The desired temperatures were provided by using an ice bath for temperatures under 20°C and a Tempette Junior TE-85 temperature controller attached to the cell-holder of the spectrophotometer for temperatures above 20°C. The reaction mixtures, containing all the reagents except enzyme, were incubated for 5 min at various temperatures, as indicated above. After the purified enzyme extract was added to the incubated reaction mixture, the activity of LsPPO was determined spectrophotometrically. The final reaction mixture contained 0.6 mL of substrate (0.02 M final concentration), 2.3 mL of 0.1 M buffer solution, and 0.1 mL of enzyme solution. Each assay was repeated twice using the same stock of enzyme extract.

### Inhibition of LsPPO activity

An aliquot of each inhibitor at various final concentrations was added to the standard reaction solution immediately before the addition of enzyme extract. The concentration of inhibitor (L-cysteine, p-aminobenzenesulfonamide, glutathion and sulfosalicylic acid) giving 50% inhibition ( $IC_{50}$ ) was determined from a plot of residual activity against inhibitor concentration, with 10 mM catechol as substrate. The control was activity without inhibitor.

## Results and discussion

### Extraction and purification of LsPPO

In the past, purification of PPO from different sources has used methods such as Triton X-100 extraction, ammonium sulfate precipitation, dialysis, affinity chromatography, Sephadex G-200 gel filtration chromatography, and Phenyl-Sepharose hydrophobic

chromatography [7,15,16]. In the present study LsPPO was purified in one step from a crude 80%  $(NH_4)_2SO_4$  precipitate by affinity chromatography on a Sepharose 4B-L-tyrosine-p-amino benzoic acid affinity column (Figure 1). This affinity chromatography achieved a 26.6-fold purification, as shown in Table I, This is higher than the 17.2, 10.8, 9.0, 4.9 and 6.5 fold purifications variously described for guava [17], yali pear [18] and yam [19]. Previously, our group had obtained 31.5-fold purification for wild pear (*Pyrus elaeagnifolia*) PPO [20] and a 74-fold purification for mulberry (*Morus Alba L.*) PPO [7] using the same affinity column.

PPO is widely distributed in all plants. The molecular weight of PPO varies, however, between species [21]. In the present work, the purified LsPPO migrated as a band of approximately 65 kDa (data not shown), upon SDS-polyacrylamide gel electrophoresis. This is the same molecular weight as the PPO isolated from Chinese cabbage [22], mulberry fruit [7] and wild pear [20].

### Optimum pH and substrate specificity

The pH profile of LsPPO activity was determined between pH 4.5 and 11.0. As shown in Figure 3, the optimum pH for maximum LsPPO activity was very dependent on the nature of the substrate used for the assay. Optimum pH values were 6, 7.5 and 7.5 for 4-methyl catechol, catechol, and pyrogallol, respectively. These values were different from those of raspberry pH 8.0 [23], Allium sp. pH 7.5 [24], and mulberry fruit pH 7.0 [7] using catechol as substrate. Different optimum pH values for PPO obtained from different sources are reported in the literature. The optimum pH values are 5.5 for strawberry [11], 6.0 for DeChaunac grape [25], 7.0 for Amasya apple [26], 6.0 for *Pyrus elaeagnifolia* (PePPO) [20]. The optimum pH of mulberry PPO (MPPO) [7] was similar to that of sago log (*Metroxylon sagu*) pH 7.5 [27] using pyrogallol as substrate.




Figure 2.  $V_{MAX}$  and  $K_M$  values of LsPPO with (a) catechol; (b) 4-methyl catechol; (c) pyrogallol substrates.

#### Optimum temperature & thermal inactivation

To investigate the temperature dependence of LsPPO, catechol was used as a substrate. A temperature range of 5–50°C was investigated and activity monitored after treatment of the enzyme for between 5 and 40 min at each temperature (Figure 3a). It can be seen that at the shortest treatment time (5 minutes) the highest activity was seen at 5°C. Even this short period

at higher temperatures led to reduced activity. Whereas activity was reasonably maintained after 40 min of incubation at temperatures of  $\leq 30^\circ\text{C}$ , significant inactivation became apparent at 40°C, and more so at 50°C. In the literature, the maximum activity of various PPO species, using catechol as substrate, has been reported as being at 22°C for potato [28], 40°C for Chinese cabbage [22], 12°C for *Ferula* sp. [29], and 25°C for artichoke [30].

Table I. Purification of polyphenol oxidase from *Lactaris salmonicolor*.

Steps	Volume (mL)	Activity (U/mL)	Total Protein (mg)	Protein (mg/mL)	Total Activity (U)	Specific Activity (U/mg)	Purification fold	Overall Yield (%)
Crude Extract	87	6425	120.1	1.38	558975	4654.2	–	100
Ammonium sulphate precipitation	60	4975	16.2	0.27	298500	18425.9	3.96	34.2
Dialysis	10.8	9450	18.8	1.74	102060	5428.7	1.17	18.3
Affinity chromatography	3	13000	0.315	0.105	39000	123809.5	26.6	7

### Enzyme inhibition

A considerable number of inhibitors of PPO-induced browning of fruits and vegetables have been identified. Reducing agents have been widely used, but these may have adverse health effects and can also react with other components in the food system [31]. Another important group of browning inhibitors is comprised of compounds that are structurally analogous to the

phenolic substrates of PPO, the inhibitory capabilities of which will depend on the enzyme source and the substrate used [32]. Here we investigated the effects of some classical PPO inhibitors, namely glutathione, L-cysteine, *p*-aminobenzenesulfonamide, sulfosalicylic acid and L-tyrosine, on the activity of LsPPO using catechol as the substrate.  $IC_{50}$  values of  $9.1 \times 10^{-4}$  M,  $2.3 \times 10^{-4}$  M,  $1.5 \times 10^{-4}$  M,  $3.8 \times 10^{-7}$  M,  $1.2 \times 10^{-4}$  M,  $4.9 \times 10^{-4}$  M, and




Figure 3. (a) The effect of temperature on the purified LsPPO activity; (b) Effect of pH on the activity of LsPPO with different substrates.



Table II. Effects of inhibitors on LsPPO.

Inhibitor	Type of inhibition	Ki (M)	IC <sub>50</sub> (M)
L-cysteine	Uncompetitive	$1.45 \times 10^{-5}$	$1.5 \times 10^{-4}$
L-tyrosine	Uncompetitive	$1.75 \times 10^{-3}$	$3.8 \times 10^{-7}$
Glutathione	Mixed type(Competitive + non competitive)	K1: $0.067 \times 10^{-5}$ < K2: $1.6 \times 10^{-5}$	$9.1 \times 10^{-4}$
p-aminobenzenesulfonamide	Competitive	$2.67 \times 10^{-5}$	$2.3 \times 10^{-4}$
Oxalic acid	Uncompetitive	$7.43 \times 10^{-8}$	$1.2 \times 10^{-4}$
β-mercapto-ethanol	Uncompetitive	$0.31 \times 10^{-5}$	$4.9 \times 10^{-4}$
Syringic acid	Uncompetitive	$2.14 \times 10^{-5}$	$4 \times 10^{-4}$

$4 \times 10^{-4}$  M were obtained for glutathione, p-aminobenzenesulfonamide, L-cysteine, L-tyrosine, oxalic acid, β-mercaptoethanol and syringic acid, respectively. Depending on kinetic analysis, mixed-type inhibition (glutathione), competitive inhibition (p-aminobenzenesulfonamide) and uncompetitive inhibition (L-cysteine, L-tyrosine, oxalic acid and syringic acid) were all seen in this study (Table II). Arslan et. al reported glutathione and L-cysteine as displaying competitive inhibition of mulberry PPO [7] but L-cysteine behaved as an uncompetitive inhibitor for LsPPO. Similarly, sulfosalicylic acid was an uncompetitive inhibitor in that study, but for LsPPO this compound was an activator.

## Conclusions

There are so many kinds of harvested edible mushroom species in the world. Several mushrooms are especially tasty and many are rich on nutrients and some of them are also toxic and dangerous for the human health as well. Mushrooms are also easily preserved, and historically have provided additional nutrition over winter. In some parts of Eurasia, especially in Russia and Nordic countries, mushrooms are an important part of the diet. Around six percent of edible species also have medicinal properties. This contribution to human welfare is difficult to assess and has received little attention. The medicinal properties of mycorrhizal fungi have not been well investigated [33]. Edible fungi already play an important role in the lives of many people and more benefits could be achieved for many years. Mushrooms are considered to be a good source of digestible proteins, and while the lipid content is low, the main classes of lipid compounds are represented including phospholipids, sterols, sterol esters, mono-, di- and triglycerides as well as free fatty acids [1].

The principal enzyme responsible for the browning reactions is a binuclear copper containing enzyme, polyphenol oxidase (PPO; E.C 1.14.18.1), which uses molecular oxygen to catalyze the o-hydroxylation of monophenols to o-diphenols and their further oxidation lead to react with endogenous amino acids and proteins to form complex brown pigments and fort this the colour of mushrooms after processing

is strongly influenced by the activity of polyphenoloxidase. The browning of mushrooms might also be caused by the action of bacteria and mold on the mushroom. *Pseudomonas tolaasii* is regarded as a normal constituent of the microflora of the mushroom bed which could produce a metabolite toxic compounds to mushrooms under certain conditions. The infection appears as a brown injury on mushrooms tissues [34,35]. According to this, wild edible fungi provide a source of food and income benefits to people, for this purpose preventing the food quality of this type of mushroom attempts are being made to investigate the purification polyphenol oxidase activity and its some kinetic properties on *Lactarius salmonicolor* which is commonly used as a food source in Turkey in the middle of November in Balikesir.

LsPPO has the same apparent mass as most other reported PPO species. Its pH optimum is near neutral (pH 6.0–7.5), and like many other PPO enzymes it is, to some extent, substrate-dependent. Comparing catechol, 4-methyl catechol and pyrogallol the  $K_M$  and  $V_{Max}$  values of LsPPO towards these three substrates were determined as 0.025 M & 0.748 EU/mL,  $1.809 \times 10^{-3}$  M & 0.723 EU/mL, and  $9.465 \times 10^{-3}$  M & 0.722 EU/mL, respectively. Its pH optimum is unusually low and shows evidence of sensitivity to thermal denaturation with increasing temperature (Figure 3a, 3b). Inhibition studies indicate a particularly marked inhibitory potential of L-tyrosine (Table II). Further characterisation of purified fungal PPO enzymes, like LsPPO, may lead to the identification of the most suitable strategies for inhibition of PPO-mediated browning reactions and loss of quality in commercial mushroom processing.

## Acknowledgements

The authors would like to thank the group researchers Murat Sayın, Aysegul Sahin and Semra Isık for their technical laboratory support, and also Balikesir University, Research Center of Applied Sciences (BURCAS / Balikesir, Turkey) for providing the research facilities. We also thank Dr. Malcolm Lyon (University of Manchester. Paterson Institute for Cancer Research) for his invaluable contribution to this paper.

**Declaration of interest:** The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

## References

- [1] Smith JE, Rowan NJ, Sullivan R. Medicinal mushrooms: a rapidly developing area of biotechnology for cancer therapy and other bioactivities (review). *Biotechnol Lett* 2002;22:1839–1845.
- [2] Boa ER. How do local people make use of wild edible fungi? Personal narratives from Malawi. In: Hall IR, Wang Y, Zambonelli A, Danell E, editors. *Edible ectomycorrhizal mushrooms and their cultivation*. Proceedings of the second international conference on edible ectomycorrhizal mushrooms. July 2001, Christchurch. CD-ROM. Christchurch: New Zealand Institute for Crop and Food Research Limited; 2002.
- [3] Guthrie BD. Studies on the control of bacteria deterioration of fresh, Washed mushrooms (*Agaricus bisporus/brunescens*); Master Thesis, Pennsylvania State University; 1984.
- [4] Morte Pérez-Gilabert A, Honrubia M, García-Carmona F. Partial purification, characterization, and histochemical localization of fully latent desert truffle (*terfezia clavaryi* chatin) polyphenol oxidase. *J Agric Food Chem* 2001;49:1922–1927.
- [5] Van Gelder CWG, Flurkey WH, Wichers HJ. Sequence and structural features of plant and fungal tyrosinases. *Phytochemistry* 1997;45:1309–1323.
- [6] Whitaker JR. In: Wong DWS, editor. *Food enzymes, structure and mechanisms*. New York: Chapman and Hall; 1995. p 271–307.
- [7] Arslan O, Erzenin M, Sinan S, Ozensoy O. Purification of mulberry (*Morus alba* L.) polyphenol oxidase by affinity chromatography and investigation of its kinetic and electrophoretic properties. *Food Chem* 2004;88(3):479–484.
- [8] Hendrickx M, Ludikhuyze Van den Broeck I, Weemaes C. Effects of high pressure on enzymes related to food quality. *Trends Food Sci Technol* 1998;9:197–203.
- [9] Asaka M, Hayashi R. Activation of polyphenol oxidase in pear fruit by high pressure treatment. *Agric Biol Chem* 1991;5(9):2439–2440.
- [10] Knorr D. Effects of high hydrostatic pressure processes on food safety and quality. *Food Technol* 1993;47:156–161.
- [11] Wesche-Ebeling P, Montgomery MW. Strawberry polyphenol oxidase: extraction and partial characterization. *J Food Sci* 1990;55:1320–1325.
- [12] Espin JC, Morales M, Varon R, Tudela J, Garcia-Canovas F. A continuous spectrophotometric method for determining the monophenolase and diphenolase activities of apple polyphenol oxidase. *Anal Biochem* 1995;43:2807–2812.
- [13] Bradford M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976;72:248.
- [14] Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970;227:680–685.
- [15] Jiang YM. Purification and some properties of polyphenol oxidase of longan fruit. *Food Chem* 1999;66:75–79.
- [16] Weemaes CA, Ludikhuyze LR, Broeck IV, Hendrickx ME, Tobback PP. Activity, Electrophoretic Characteristics And Heat Inactivation Of Polyphenoloxidases From Apples, Avocados, Grapes, Pears And Plums. *Lebensmittel-Wissenschaft und-Technologie* 1998;31:44–49.
- [17] Augustin MA, Ghazali HM, Hashim H. Polyphenoloxidase from guava (*Psidium guajava* L.). *J Agric Food Chem* 1985;36:1259–1265.
- [18] Zhou H, Feng X. Polyphenol oxidase from yali pear (*Pyrus bretschneideri*). *J Sci Food Agric* 1991;57:307–313.
- [19] Martin FW, Ruberte RM. The polyphenol of *Dioscorea alata* (yam) tubers associated with oxidative browning (varieties). *J Agric Food Chem* 1976;24(1):67–70.
- [20] Arslan O, Sinan S, Yerlitürk FU, Gencer N, Özensoy Ö. Characterization of polyphenol oxidase from wild pear (*Pyrus elaeagnifolia*). *J Food Biochem* (Article in press vol 33, issue 2 in 2009) reviewed 2007.
- [21] Cheng T-M, Huang P-C, Pan J-P, Lin K-Y, Mao S-J. Gel electrophoresis of polyphenol oxidase with instant identification by *in situ* blotting. *J Chromatog B* 2007;849(1–2):331–336.
- [22] Nagai T, Suzuki N. Partial purification of polyphenol oxidase from Chinese cabbage *Brassica rapa* L. *J Agric Food Chem* 2001;49:3922–3926.
- [23] Gonzalez EM, deAncos B, Cano MP. Partial characterization of polyphenoloxidase activity in raspberry fruits. *J Agric Food Chem* 1999;47:4068–4072.
- [24] Arslan O, Temur A, Tozlu I. Polyphenol oxidase from *Allium* sp. *J Agric Food Chem* 1997;45:2861.
- [25] Lee CY, Smith NL, Pennesi AP. Polyphenol oxidase from DeChaunac grapes. *J Sci Food Agric* 1983;34:987–991.
- [26] Oktay M, Kufrevioglu I, Kocacaliskan I, Sakiroglu H. Polyphenol oxidase from Amasya apple. *J Food Sci* 1995;60:495–499.
- [27] Vomas-Vigyazo L. Polyphenoloxidase and peroxidase in fruits and vegetables. *CRC Critical Reviews Food Sci Nutr* 1981;14:44.
- [28] Matheis G, Belitz H-D. Studies on enzymic browning of potatoes (*Solanum tuberosum*). III. Kinetics of potato phenoloxidase (EC 1.14.18.1 monophenol, dihydroxyphenylalanine: Oxygenoxidoreductase). *Z Lebensm Unters Forsch* 1977;163:191.
- [29] Erat M, Şakiroglu H, Kufrevioglu OI. Purification and characterization of polyphenol oxidase from *Ferula* sp. *Food Chem* 2006;95:503–508.
- [30] Aydemir T. Partial purification and characterization of polyphenol oxidase from artichoke (*Cynara scolymus* L.) heads. *Food Chem* 2004;87:59–67.
- [31] McEvily AJ, Iyengar R, Otwell WS. Inhibition of enzymatic browning in foods and beverages. *Crit Rev Food Sci Nutr* 1992;32:253–273.
- [32] Mayer AM, Harel E. Polyphenol oxidase in plants. *Phytochemistry* 1979;18:193–215.
- [33] Reshetnikov SV, Wasser SP, Tan KK. Higher basidiomycota as a source of Antitumour and immunostimulating polysaccharides. A review. *Inter J Med Mushrooms* 2001;3:361–394.
- [34] Paine SG. Studies in bacteriosis II. A brown blotch disease of cultivated mushrooms. *Ann Appl Biol* 1919;5:206–219.
- [35] Royse DJ, Wuest PJ. Mushroom brown blotch. Effects of chlorinated water on disease intensity and bacterial populations in casing soil and pilei. *Phytopathology* 1980;70(9):902–905.