

I-43

THE REACTION OF URANIUM HEXAFLUORIDE WITH SILVER FLUORIDE IN ANHYDROUS HYDROGEN FLUORIDE AND THE CHEMICAL PROPERTIES OF THE PRODUCT Ag_2UF_8

John G. Malm

Chemistry Division, Argonne National Laboratory, Argonne, IL 60439 (U.S.A.)

UF_6 reacts with AgF dissolved in anhydrous hydrogen fluoride to precipitate Ag_2UF_8 . Ag_2UF_8 has some unexpected properties: On reaction with water it produces O_2 and reduced uranium. No adequate explanation could be found of why UF_6 and AgF combined in this manner should produce a powerful oxidant. Raman spectra and chemical properties of the solid products are given.

I-44

FLUORINATION OF URANIUM RESIDUES IN ISOTOPE ENRICHMENT PLANTS

W. Bacher, W. Bier, A. Maner

Institut für Kernverfahrenstechnik, Kernforschungszentrum Karlsruhe, D-7500 Karlsruhe 1 (F.R.G.)

E. Jacob

M.A.N. – Neue Technologie, D-800 Munich 50 (F.R.G.)

One possible method to remove solid deposits of uranium hexafluoride decomposition products (UF_5 , $\text{UO}_2\text{F}_2 \cdot \text{H}_2\text{O}$) which may be formed during long-time operation of uranium enrichment plants, is the application of a gaseous mixture of CBrF_3 and F_2 at a total pressure well below atmospheric pressure.

The feasibility of this procedure has been demonstrated in technical components of the "Separation Nozzle"-process for the enrichment of U-235 which is under development at the Nuclear Research Center of Karlsruhe, w.-Germany.¹⁾

The mixture of CBrF_3 and F_2 is not stable thermodynamically but shows slow decomposition which is catalyzed by metallic surfaces. The reaction $\text{CBrF}_3 + 3\text{F}_2 \rightarrow \text{CF}_4 + \text{BrF}_5$ produces BrF_5 , but the intermediate formation of BrF_3 seems reasonable. Since BrF_3 is a well known agent for the fluorination of uranium fluorides and oxifluorides into UF_6 , but is hardly applicable because of its low vapour pressure, the system CBrF_3/F_2 offers the possibility to decontaminate isotope enrichment facilities from uranium residues.

¹ E.W. Becker, W. Bier, P. Bley, W. Ehrfeld, K. Schubert, D. Seidel (paper presented to the AIChE 1982 winter meeting, Orlando, Fla., Feb. 28 - March 3, 1982, also published as report KFK 3310 (1982))