


0-61

ADDITIONS TO HEXAFLUOROBUT-2-YNE


D. Bielefeldt, R. D. Chambers, C. G. P. Jones and M. J. Silvester*

University of Durham, Durham City, DH1 3LE (U.K.)

Earlier in our laboratory^[1], we have shown that hexafluorobut-2-yne (1) reacts with sulphur, in an aprotic solvent, to give F-tetramethylthiophene and the reaction was formulated as a nucleophilic cyclisation process. Here we will describe attempts to obtain cyclised products via the generalised process shown below and using a variety of systems.



D.M.S.O. forms an adduct (2), rather than undergo cyclisation, and the



properties of (2) will be discussed.

Other additions to (1) will be described.

1 R.D. Chambers and D.B. Speight, unpublished results.


0-62

NEW CONVENIENT AND EFFICIENT ROUTES TO LONG-CHAIN (PERFLUORO-ALKYL) ALKYNES


Patrice Moreau*, Patrick Calas, Raad Albadri and Auguste Commeyras

Laboratoire de Chimie Organique, ERA No. 555, Université des Sciences et Techniques du Languedoc, Place E. Bataillon, 34060 Montpellier Cédex (France)

Perfluorinated alkynes are potential intermediates for the introduction of perfluoroalkyl chains in organic compounds. We report here new convenient and efficient preparations of various long-chain (perfluoroalkyl) acetylenes. (Perfluoroalkyl) alkynes $R_F-C\equiv C-H$ ($R_F=C_8F_{17}$, C_6F_{13} , C_4F_9) are obtained from perfluoroalkyl iodides in excellent yields (80-90%) by means of a simple electrochemical technique (electrocatalysis at a carbon fibre cathode), according to :



The corresponding 1-substituted alkynes $R_F-C\equiv C-R_H$ ($R_H=C_2H_5$, C_4H_9 , C_8H_{17} , C_6H_5 , $O-CH_2-C_6H_4$, $O-CH_3-O-C_6H_4$) are obtained in good yields (70%), in a one-pot reaction, from the thermal decomposition of perfluoroalkyl grignard reagents in the presence of excess alkyl or aryl magnesium halides, according to :



The syntheses are described and mechanisms of the reactions are discussed.