

Received January 18 1989 accepted July 2 1989

NEW POLYFLUOROALKOXYLSULFONYL FLUORIDES (II)

LI-FO CHEN, JAVID MOHTASHAM and GARY L GARD

Department of Chemistry, Portland State University, Portland, Oregon
97207-0751 (U S A)

SUMMARY

The reaction of $\overline{\text{CF}_2\text{CF}_2\text{OSO}_2}$ with haloalkanes (RX, X = I, Br) in the presence of metal fluorides, MF (M = Cs⁺, Ag⁺) was studied as a means for preparing novel reactive polyfluoroalkoxysulfonyl fluorides. The following compounds have been prepared and characterized: $\text{ROCF}_2\text{CF}_2\text{SO}_2\text{F}$ where R = $\overline{\text{OCH}_2\text{CHCH}_2}$, $(\text{CH}_3)_3\text{SiCH}_2$, $(\text{CF}_3)_2\text{CFCH}_2\text{CH}_2$, $\text{HC}\equiv\text{CCH}_2$, $\text{BrCH}=\text{CHCH}_2$ (cis and trans), $\text{CH}_2=\text{CBrCH}_2$, $(\text{CH}_3\text{CH}_2\text{O})_2\text{P}(\text{O})\text{CH}_2\text{CH}_2$, $\text{CH}_3\text{CH}_2\text{OPF}(\text{O})\text{CH}_2\text{CH}_2$. Infrared, mass and nmr spectra are presented in order to support the assigned structures.






INTRODUCTION

Fluorocarbon sulfonyl fluorides (RSO_2F) continue to be subjects of considerable interest because of their utilization as ion-exchange resins, surface active agents and strong sulfonic acids [1-4]. Previously, we have reported on a method involving the use of metal fluorides, $\overline{\text{CF}_2\text{CF}_2\text{OSO}_2}$, and RX (X = Br, I) in preparing fluorocarbon ether containing sulfonyl fluorides [5]. In this paper we wish to report our success in extending this method to include

fluorocarbon derivatives that not only contain the sulfonyl fluoride grouping but other functional groups such as $\overline{\text{OCH}_2\text{CHCH}_2}$, $(\text{CH}_3)_3\text{SiCH}_2$, $\text{HC}=\text{CCH}_2$, $(\text{CF}_3)_3\text{CFCH}_2\text{CH}_2$, $\text{BrCH}=\text{CHCH}_2$, $\text{CH}_2=\text{CBrCH}_2$, $(\text{CH}_3\text{CH}_2\text{O})_2\text{P}(\text{O})\text{CH}_2\text{CH}_2$, and $\text{CH}_3\text{CH}_2\text{OPF}(\text{O})\text{CH}_2\text{CH}_2$

RESULTS AND DISCUSSION

The reactions of $\overline{\text{CF}_2\text{CF}_2\text{OSO}_2}$ with haloalkanes were carried out according to the following equations



In general, the reactions involved two stages, in the first stage the rearrangement of the sultone occurs (eqn 1) followed by formation of the alkoxide anion (eqn 2) In the second stage, a nucleophilic substitution of RX by $\text{^-OCF}_2\text{CF}_2\text{SO}_2\text{F}$ occurred

We found that silver fluoride was a more efficient reagent for both the catalytic rearrangement of the sultone and the stabilization of $\text{^-OCF}_2\text{CF}_2\text{SO}_2\text{F}$ anion We found that using AgF required milder conditions (20-40°C) with better overall yields It was also found that when 1,3 or 2,3-dibromopropene was used, only products attributable to allylic displacement were obtained, no vinyl substituted product was observed For a multi-functional compound such as diethyl-2-bromoethyl phosphonate, a mixture of

$(\text{CH}_3\text{CH}_2\text{O})_2\text{P}(\text{O})\text{CH}_2\text{CH}_2\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$ and $(\text{CH}_3\text{CH}_2\text{O})\text{PF}(\text{O})\text{CH}_2\text{CH}_2\text{OCF}_2\text{CF}_2\text{SO}_2$ was formed. It was possible to convert the monofluoro derivative the diethoxy product by reaction with absolute ethanol. With most of the reactions mentioned above, a by-product, $\text{CH}_3\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$, was found. This result suggests the possibility that the solvent, diglyme, participated in the reaction and formed CH_3Br . Further reaction of CH_3Br with $^-\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$ anion produced the $\text{CH}_3\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$.

It is worthwhile to note that bromomethyl trimethylsilane reacted to give, as expected, the product $(\text{CH}_3)_3\text{SiCH}_2\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$. This means that the bulky substituent $(\text{CH}_3)_3\text{Si}$, at the reaction carbon center, did not prevent the nucleophilic attack of the $^-\text{OCF}_2\text{CF}_2\text{SO}_2$ ion. This indicates that a $\text{S}_{\text{N}}1$ mechanism may be involved in this reaction.

The infrared spectra of all new sulfonyl fluoride compounds have several common features. The characteristic SO_2 asym, SO_2 sym and S-F stretching frequencies are found in 1465-1455, 1244-1210, 803-792 cm^{-1} regions, respectively. These assignments agree with the results obtained previously [3-7]. The strong carbon fluorine absorption bands found at 1109-1244 cm^{-1} can be correlated with the CF_2 group [8]. The C-H absorption bands are located in the 2903-3016 cm^{-1} region. The characteristic C-H absorption of $\text{CH}=\text{C}$ is found at 3311(s) cm^{-1} region. The double bond and triple bond vibrational bands for $\text{BrCH}=\text{CHCH}_2\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$ and $\text{HC}=\text{CCH}_2\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$ appeared at 1631 and 2137 cm^{-1} , respectively.

In the $(\text{CI})^+$ or $(\text{EI})^+$ mass spectra, the $(\text{M}+1)$ or molecular ion peak for most of the compounds was found. For all compounds, a cracking pattern was found that was supportive of the assigned structure.

TABLE I

^{19}F NMR Data of $\text{ROCF}_2\text{CF}_2\text{SO}_2\text{F}$ Chemical Shift (ppm) and Coupling Constant (Hz)

R	OCF_2	CF_2	SO_2F
 $(\text{CH}_3)_3\text{SiCH}_2$	-86 0 (d-t)	-113 7 (d-t)	43 0 (t-t)
	$J_{\text{OCF}_2-\text{SO}_2\text{F}}=4$ 9	$J_{\text{OCF}_2-\text{CF}_2}=3$ 7	$J_{\text{CF}_2-\text{SO}_2\text{F}}=4$ 4
	-87 7 (d-t)	-112 7 (d-t)	42 6 (t-t)
	$J_{\text{OCF}_2-\text{SO}_2\text{F}}=5$ 8	$J_{\text{OCF}_2-\text{CF}_2}=4$ 7	$J_{\text{CF}_2-\text{SO}_2\text{F}}=5$ 4
$(\text{CF}_3)_2\text{CFCH}_2\text{CH}_2$	-87 5 (d-t)	-113 8 (d-t)	42 5 (t-t)
	$J_{\text{OCF}_2-\text{SO}_2\text{F}}=6$ 1	$J_{\text{OCF}_2-\text{CF}_2}=4$ 2	$J_{\text{CF}_2-\text{SO}_2\text{F}}=5$ 2
	$(\text{CF}_3)_2 = -78$ 6 (d)	$F = -187$ 2 (t-hept)	
	$J_{\text{CF}_3-\text{CF}}=6$ 6	$J_{\text{CF}-\text{CH}_2}=19$ 8	
HC C- CH_2	-85 0 (d-t)	-112 0 (d-t)	44 2 (t-t)
	$J_{\text{OCF}_2-\text{SO}_2\text{F}}=5$ 6	$J_{\text{OCF}_2-\text{CF}_2}=3$ 9	$J_{\text{CF}_2-\text{SO}_2\text{F}}=4$ 6

BrCH=CH₂CH₂

-84 2 (d-t) -113 0 (d-t) 43 8 (overlap)
J_{(trans)OCF₂-SO₂F} = 6 6 J_{(trans)OCF₂-CF₂} = 4 9 J_{(trans)CF₂-SO₂F} = 5 0
J_{(cis)OCF₂-SO₂F} = 5 2 J_{(cis)OCF₂-CF₂} = 4 9 J_{(cis)CF₂-SO₂F} = 5 0
-85 2 (d-t) -112 7 (d-t) 44 2 (t-t)
J_{OCF₂-SO₂F} = 6 4 J_{OCF₂-CF₂} = 4 7 J_{CF₂-SO₂F} = 5 1
-84 5 (d-t) -111 5 (d-t) 44 4 (t-t)
J_{OCF₂-SO₂F} = 5 2 J_{OCF₂-CF₂} = 3 7 J_{CF₂-SO₂F} = 4 2
-84 9 (d-t) -111 5 (d-t) 44 2 (t-t) PF 60 7
J_{OCF₂-SO₂F} = 5 2 J_{OCF₂-CF₂} = 3 7 J_{CF₂-SO₂F} = 4 2
J_{PF} = 1075



CH₂=CBrCH₂

(CH₃CH₂O)₂P(O)CH₂CH₂

(CH₃CH₂O)PF(O)CH₂CH₂

TABLE II

 ^1H NMR Data of $\text{ROCF}_2\text{CF}_2\text{SO}_2\text{F}$

R	Chemical Shift (ppm)	Coupling Constant (Hz)
	$\text{H}_a = 3.35$ (m) $\text{H}_b = 2.94$ (d-d) $\text{H}_c = 2.77$ (d-d) $\text{H}_d = 4.50$ (d-d) $\text{H}_e = 4.05$ (d-d)	$J_{ac} = 5.0$ $J_{ab} = 5.8$ $J_{bc} = 4.5$ $J_{de} = 11.7$
$(\text{CH}_3)_3\text{SiCH}_2$	$(\text{CH}_3)_3 = 0.32$ (s) $\text{CH}_2 = 3.98$ (s)	
$(\text{CF}_3)_2\text{CFCH}_2\text{CH}_2$	$\text{CFCH}_2 = 2.80$ (d-t) $\text{CH}_2\text{O} = 4.62$ (t)	$J_{\text{CF}-\text{CH}_2} = 19.8$ $J_{\text{CH}_2-\text{CH}_2} = 7.2$
$\text{HC}=\text{C}-\text{CH}_2$	$\text{CH} = 3.00$ (t) $\text{CH}_2 = 4.95$ (d)	$J_{\text{CH}-\text{CH}_2} = 2.5$
$\text{BrCH}=\text{CHCH}_2$	$\text{CH}=\text{CH} = 6.90$ (m) $\text{CH}_2(\text{trans}) = 5.25$ (d) $\text{CH}_2(\text{cis}) = 5.02$ (d)	$J_{(\text{trans})\text{CH}-\text{CH}_2} = 5.04$ $J_{(\text{cis})\text{CH}-\text{CH}_2} = 5.58$
	$\text{H}_a = 6.32$ (s) $\text{H}_b = 6.05$ (s) $\text{CH}_2 = 4.97$ (s)	
$(\text{CH}_3\text{CH}_2\text{O})_2\text{P}(\text{O})\underline{\text{CH}_2}\text{CH}_2$	$\text{CH}_3 = 1.75$ (t) $\text{O}\underline{\text{CH}_2}\text{CH}_3 = 4.37$ (d-q) $\text{CH}_2\text{OCF}_2 = 4.63$ (t) $\text{PCH}_2 = 2.55$ (d-t)	$J_{\text{CH}_3-\text{CH}_2\text{O}} = 7.38$ $J_{\text{P}(\text{O})-\text{OCH}_2} = 7.38$ $J_{\underline{\text{CH}_2}-\text{CH}_2} = 7.56$ $J_{\text{P}(\text{O})-\underline{\text{CH}_2}} = 19.80$
$(\text{CH}_3\text{CH}_2\text{O})\text{PF}(\text{O})\underline{\text{CH}_2}\text{CH}_2$	$\text{CH}_3 = 1.78$ (t) $\text{PCH}_2 = 2.90$ (d-m) $\text{CH}_2\text{O} = 4.30 \approx 5.05$ (m)	$J_{\text{CH}_3-\text{CH}_2\text{O}} = 5.4$ $J_{\text{P}-\underline{\text{CH}_2}} = 18.0$ $J_{\text{P}(\text{O})-\text{OCH}_2} = \text{overlap}$

The structures of all new products were determined from their respective ^1H and ^{19}F nmr spectra, with most compounds, first order couplings were found. The ^{19}F nmr chemical shifts and coupling constants are reported in Table I. It is found that consistent chemical shift values for similar groupings are maintained for all compounds. For OCF_2 , CF_2 , SO_2F , the chemical shift values range from -84.2 to -87.7, -112.0 to -113.8, 42.5 to 44.0 ppm, respectively. The ^1H nmr chemical shifts and coupling constants are listed in Table II.

For $\text{BrCH}=\text{CHCH}_2\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$, both ^1H and ^{19}F nmr data confirm the presence of isomers (trans 60%, cis 40%). In the ^1H nmr spectra, the two protons situated at a vinyl position exhibited an overlap pattern in the region of 6.9 ppm. However, the CH_2 group, adjacent to the vinyl group, appeared at the following chemical shift values δ 5.02 (cis) and δ 5.25 (trans) ppm.

The reaction of diethyl-2-bromoethylphosphonate with $\text{AgOCF}_2\text{CF}_2\text{SO}_2\text{F}$ gave a mixture, which from the ^{19}F nmr spectrum was found to consist of $(\text{CH}_3\text{CH}_2\text{O})\text{PF}(\text{O})\text{CH}_2\text{CH}_2\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$ (45%) and the expected product $(\text{CH}_3\text{CH}_2\text{O})_2\text{P}(\text{O})\text{CH}_2\text{CH}_2\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$ (55%). The coupling constant of the P-F bond in $\text{CH}_3\text{CH}_2\text{OPF}(\text{O})\text{CH}_2\text{CH}_2\text{O}(\text{CF}_2)_2\text{SO}_2\text{F}$ was 1075 Hz and the fluorine chemical shift appeared at -60.7 ppm, the chemical shift for OCF_2 was 0.5 ppm higher than that found for the non-P-F containing product. The ^{19}F chemical shift and the coupling constant for the PF group are in agreement with values reported for compounds with a similar molecular structure [9]. When treated with absolute alcohol, the P-F bond at -60.7 ppm disappeared and only the pure product, $(\text{CH}_3\text{CH}_2\text{O})_2\text{P}(\text{O})(\text{CH}_2)_2(\text{CF}_2)_2\text{SO}_2\text{F}$, was present.

EXPERIMENTAL

The sulfone $\overline{\text{CF}_2\text{CF}_2\text{OSO}_2}$ was prepared according to the literature method [10]. Cesium fluoride and silver fluoride were dried under vacuum before use. All other chemicals were obtained from commercial sources and used as received.

General Procedure Gases were manipulated in a conventional Pyrex vacuum apparatus equipped with a Heise-Bourdon tube gauge and televac thermocouple gauge. Infrared spectra were obtained by using a Pyrex-glass cell with KBr windows or as liquids between KBr disks on a Nicolet 20DX spectrometer. The nmr spectra were recorded with a Varian model EM-390 spectrometer operating at 90.0 MHz for proton and 84.67 MHz for the fluorine resonance. TMS and F-11 were used as external standards. In some cases, compounds were purified via gas chromatography using an Aerograph autoprep (model A-700) gas chromatograph. The mass spectra were taken on a VG-7070 HS mass spectrometer with an ionization potential of 70 eV. Perfluoro-kerosene was used as an internal standard.

Elemental analyses were determined by Beller Microanalytical Laboratory in Göttingen, Federal Republic of Germany.

Preparation of $\overline{\text{OCH}_2\text{CHCH}_2\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}}$

Into a 125 mL Pyrex-glass vessel equipped with a Kontes Teflon valve and a Teflon stirring bar were added 5.0 g (39.0 mmol) of dried silver fluoride and 10.0 mL of diglyme, 9.50 g (52.7 mmol) of $\overline{\text{CF}_2\text{CF}_2\text{OSO}_2}$ was vacuum transferred into the reaction vessel cooled to -196 °C. The reaction was warmed to room temperature and stirred at

35°C for 15 h. During this time a homogenous brown solution was formed. The vessel was again cooled to -196°C and 4.9 g (35.7 mmol) of epibromohydrin was added. The reaction was warmed to room temperature and heated (35°C) for 3 d. The volatile material (5.09 g), recovered at room temperature in a trap cooled to -196°C, was found to be $\text{FC(O)CF}_2\text{SO}_2\text{F}$. A second fraction (15.0 g) was collected by heating the reaction vessel under vacuum. This fraction was washed four times with water and gave 5.6 g of an oil which was dried with MgSO_4 and distilled to give 4.0 g (15.6 mmol) product, in 44% yield, b.p. 92-93°C/40 mm.

The infrared spectrum had the following bands (cm^{-1})
 3016(vw), 1457(vs), 1364(w), 1335(m), 1242(s), 1205(s), 1142(s),
 1115(s), 1038(w), 1000(m), 948(w), 907(w), 800(vs), 763(w), 655(m),
 614(s), 551(w)

In the (EI)⁺ mass spectrum, the molecular ion was observed at 256 [(M)⁺, 156], other main fragment ions were detected at 226 [(M-CH₂O)⁺, 534], 213 [(M-C₂H₃O)⁺, 2114], 149 [(C₅F₃O₂)⁺, 2488], 145 [(CF₂=CFSO₂)⁺, 766], 133 [(CF₂SO₂F)⁺, 474], 127 [(C₅H₃O₂S)⁺, 1758], 119 [(C₃FO₂S)⁺, 1181], 100 [(C₂F₄)⁺, 3370], 99 [(C₄O₂F)⁺, 2825], 97 [(C₂F₃O)⁺, 6309], 95 [(CFSO₂)⁺, 677], 93 [(C₃H₃F₂O)⁺, 4756], 83 [(SO₂F)⁺, 276], 73 [(OCH₂CHCH₂O)⁺, 866], 71 [(C₃H₃O₂)⁺, 296], 69 [(C₃HO₂)⁺, 222], 67 [(SOF)⁺, 100], 65 [(C₄OH)⁺, 544], 63 [(CFS)⁺, 602]

Anal. Calcd for C₅H₅F₅O₄S C, 23.43, H, 1.95, F, 37.1, S, 12.50
 Found C, 23.61, H, 1.94, F, 37.5, S, 12.30

Preparation of $(\text{CH}_3)_3\text{SiCH}_2\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$

In a similar procedure as previously described, 3.70 g (29.1 mmol) of silver fluoride in 50 mL of diglyme with 6.5 g (36.1 mmol) of $\overline{\text{CF}_2\text{CF}_2\text{OSO}_2}$ was heated at 36°C for 26 h to give a homogeneous solution of $\text{AgOCF}_2\text{CF}_2\text{SO}_2\text{F}$ and diglyme. At -196°C, 4.1 g (24.5 mmol) of $(\text{CH}_3)_3\text{SiCH}_2\text{Br}$ was added by vacuum transfer to the reaction, and then maintained at 37°C for 4 d. Only 0.2 g of a low boiling material was recovered. The volatile material at room temperature was transferred, washed with water, dried and distilled to give 2.2 g (7.6 mmol) of the product in 33% yield (b.p. 125-127°C) and 0.30 g of a minor product, $\text{CH}_3\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$ in 5.7% yield, b.p. 83-85°C.

The infrared spectrum had the following bands (cm^{-1}): 2964(w), 1459(s), 1344(s), 1293(w), 1257(s), 1238(s), 1206(s), 1137(s), 1109(s), 994(m), 856(vs), 819(w), 792(m), 755(w), 709(w), 649(w), 608(s), 539(w).

In the $(\text{CI})^+$ mass spectrum, no molecular ion was observed. Main fragmentations were found at: 271 $[(\text{M}-\text{CH}_3)^+, 0.37]$, 187 $[(\text{C}_5\text{H}_7\text{F}_4\text{OSi})^+, 0.45]$, 183 $[(\text{CF}_2\text{CF}_2\text{SO}_2\text{F})^+, 0.56]$, 145 $[(\text{CF}_2=\text{CFSO}_2)^+, 1.13]$, 137 $[(\text{C}_4\text{H}_7\text{F}_2\text{OSi})^+, 1.87]$, 135 $[(\text{C}_4\text{H}_5\text{F}_2\text{OSi})^+, 1.87]$, 134 $[(\text{C}_4\text{H}_4\text{F}_2\text{OSi})^+, 0.88]$, 133 $[(\text{CF}_2\text{SO}_2\text{F})^+, 0.81]$, 123 $[(\text{C}_3\text{H}_5\text{F}_2\text{OSi})^+, 1.39]$, 120 $[(\text{C}_3\text{H}_2\text{F}_2\text{OSi})^+, 1.18]$, 119 $[(\text{C}_3\text{HF}_2\text{OSi})^+, 1.66]$, 107 $[(\text{C}_2\text{HF}_2\text{OSi})^+, 1.67]$, 100 $[(\text{C}_2\text{F}_4)^+, 8.50]$, 97 $[(\text{C}_2\text{F}_3\text{O})^+, 7.86]$, 95 $[(\text{CFSO}_2)^+, 2.18]$, 92 $[(\text{C}_3\text{H}_2\text{F}_2\text{O})^+, 3.65]$, 91 $[(\text{C}_3\text{HF}_2\text{O})^+, 27.15]$, 89 $[(\text{C}_2\text{H}_2\text{FOSi})^+, 4.47]$, 88 $[(\text{C}_2\text{HFOSi})^+, 9.31]$, 87 $[(\text{CH}_3)_3\text{SiCH}_2^+, 100]$, 83 $[(\text{SO}_2\text{F})^+, 2.13]$, 81 $[(\text{C}_2\text{F}_3)^+, 2.65]$, 79 $[(\text{C}_2\text{HF}_2\text{O})^+, 6.37]$, 73 $[(\text{CH}_3)_3\text{Si}^+, 71.70]$, 67 $[\text{SOF}]^+, 4.51]$, 65 $[(\text{C}_4\text{OH})^+, 2.83]$, 63 $[(\text{CFS})^+, 2.31]$.

Anal Calcd for $C_6H_{11}F_5O_3S_1S$ C, 25.17, H, 3.85, F, 33.2, S, 11.19
Found C, 25.41, H, 3.88, F, 33.5, S, 11.39

Preparation of $(CF_3)_2CFCH_2CH_2OCF_2CF_2SO_2F$

In a similar procedure as previously described, 3.50 g (27.5 mmol) of dried silver fluoride, 5.0 mL of diglyme and 6.80 g (37.8 mmol) of $CF_2CF_2OSO_2$ were reacted at room temperature (24 h) to give a homogenous clear solution of $AgOCF_2CF_2SO_2F$ and diglyme. The mixture was cooled and 8.0 g (24.7 mmol) of $(CF_3)_2CFCH_2CH_2I$ was added. The reaction mixture was warmed and maintained at room temperature for 3 d. The products were transferred under vacuum and washed with water. 8.10 g of an oily material was collected, dried over P_4O_{10} and distilled to give 5.1 g (12.9 mmol) of product in 52.1% yield, b.p. 150–152°C.

The infrared spectrum had the following bands (cm^{-1})
 2945(vw), 2853(vw), 1462(vs), 1335(m), 1314(m), 1237(vs), 1166(s), 1145(s), 1117(s), 1068(m), 1005(m), 934(w), 801(vs), 723(m), 653(v), 612(s), 541(w).

In the $(CI)^+$ mass spectrum, no molecular ion was observed. Main fragment ions were found at 377 $[(M-F)^+]$, 0.77], 357 $[(M-HF_2)^+]$, 3.72], 337 $[(M-H_2F_3)^+]$, 2.67], 213 $[(CF_3)_2CFCH_2CH_2O^+]$, 3.51], 197 $[(CF_3)_2CFCH_2CH_2^+]$, 4.75], 183 $[(CF_3)_2CFCH_2^+]$, 5.09], 181 $[(C_4F_7)^+]$, 2.78], 178 $[(C_5H_4F_6)^+]$, 1.24], 177 $[(C_5H_3F_6)^+]$, 1.00], 175 $[(C_5HF_6)^+]$, 4.14], 161 $[(C_2F_3O_3S)^+]$, 7.05], 157 $[(C_3F_3O_2S)^+]$, 8.01], 155 $[(C_5H_3F_4O)^+]$, 5.45], 149 $[(C_6H_4F_3O)^+]$, 6.79], 145 $[(CF_2=CFSO_2)^+]$, 3.19], 133 $(CF_2SO_2F)^+$, 5.36], 127 $[(C_5H_3O_2S)^+]$, 2.49], 119

[(C₃FO₂S)⁺, 11 15], 115 [(C₄H₃O₂S)⁺, 28 86], 113 [(C₄HO₂S)⁺, 10 48],
 109 [(C₅HOS)⁺, 5 89], 100 [(C₂F₄)⁺, 25 65], 97 [(C₂F₃O)⁺, 17 11], 95
 [(CFSO₂)⁺, 11 81], 93 [(C₃H₃F₂O)⁺, 2 30], 87 [(C₄H₄FO)⁺, 9 32], 83
 [(SO₂F)⁺, 1 03], 81 [(CF₂=CF)⁺, 15 6], 79 [(C₂HF₂O)⁺, 19 51], 77
 [(C₅HO)⁺, 11 26], 67 [(SOF)⁺, 37 20], 65 [(C₄OH)⁺, 12 76]

Anal Calcd for C₇H₄F₁₂O₃S C, 21 12, H, 1 01, F, 57 6, S,
 8 08 Found C, 21 36, H, 1 12, F, 58 0, S, 7 90

Preparation of HC=C-CH₂OCF₂CF₂SO₂F

Into the same reaction vessel previously described, 6 70 g
 (52 7 mmol) of dried silver fluoride, 10 0 mL diglyme and 18 0 g
 (100 mmol) of $\overline{\text{CF}_2\text{CF}_2\text{OSO}_2}$ were added at -196°C A clear, slightly
 blue solution of AgOCF₂CF₂SO₂F was obtained after heating at 30-35 C
 for 2 h At -196 C, 8 10 g of a solution containing toluene (20%)
 and HC=CCH₂Br (80% and 54 6 mmol) was added into the reaction
 vessel

The reaction mixture was maintained at 5-8°C for 2 h, and at
 18-20 C for another 16 h All volatile materials (27 60 g) were
 distilled to give 4 0 g of FC(O)CF₂SO₂F, b p 32-34 C, 6 20 g of
 CH₃OCF₂CF₂SO₂F, 55% yield, b p 78-87°C, and 3 5 g of a mixture
 The G C indicated that the mixture contained 50% HC=CCH₂OCF₂CF₂SO₂F
 (14% yield) and 40% toluene The pure HC=CCH₂OCF₂CF₂SO₂F can be
 separated from G C (SE-30, 3M, 150°C), b p 107°C

The infrared spectrum had the following bands (cm⁻¹) 3311(s),
 2966(vw), 2903(vw), 2137(m), 1459(vs), 1383(w), 1334(m), 1280(m),
 1242(s), 1204(s), 1144(s), 1123(s), 1036(m), 1004(s), 917(w),
 803(vs), 766(m), 695(w), 657(m), 608(s)

In the (CI)⁺ mass spectrum, the (M+1)⁺ ion was observed at 239 [(M+1)⁺, 19 14], other main fragment ions were detected at 237 [(M-1)⁺, 2 24], 155 [(C₅H₃OF₄)⁺, 5 93], 154 [(C₅H₂OF₄)⁺, 4 54], 153 [(C₅HOF₄)⁺, 2 81], 127 [(C₅H₃O₂S)⁺, 60 19], 126 [(C₅H₂O₂S)⁺, 24 23], 125 [(C₅HO₂S)⁺, 13 51], 123 [(C₄H₂OF₃)⁺, 8 20], 119 [(C₃FO₂S)⁺, 16 58], 105 [(C₄H₃OF₂)⁺, 5 96], 103 [(C₄HF₂O)⁺, 7 77], 100 [(C₂F₄)⁺, 81 34], 97 [(C₂F₃O)⁺, 6 43], 95 [(CFSO₂)⁺, 3.84], 94 [(C₂F₂S)⁺, 8 86], 91 [(C₃HOF₂)⁺, 3 26], 90 [(C₃OF₂)⁺, 3 28], 83 [(SO₂F)⁺, 3 22], 79 [(C₂HOF₂)⁺, 38 79], 78 [(C₅H₂O)⁺, 2 13], 77 [(C₅HO)⁺, 65 62], 75 [(C₂FS)⁺, 10 22], 71 [(C₃OF)⁺, 22 53], 67 [(SOF)⁺, 43 43], 65 [(C₄OH)⁺, 22 53], 63 [(CFS)⁺, 5 84], 59 [(C₂OF)⁺, 100], 55 [(C₃H₃O)⁺, 32 81], 54 [(C₃H₂O)⁺, 29 60], 53 [(C₃HO)⁺, 21 09]

Anal. Calcd for C₅H₃O₃F₅S C, 25.21, H, 1.26, F, 39.9, S, 13.44
Found C, 25.30, H, 1.29, F, 39.8, S, 13.48

Preparation of BrCH=CHCH₂OCF₂CF₂SO₂F

(A) AgF Method Into a similar reaction vessel previously described, 4.20 g (33.0 mmol) of dried silver fluoride, 7.0 mL of diglyme and 7.70 g (42.7 mmol) of $\overline{\text{CF}_2\text{CF}_2\text{OSO}_2}$ were added. The reaction mixture was stirred at room temperature for 5 h, followed by the addition of 6.00 g (30.0 mmol) of BrCH=CHCH₂Br. The mixture was maintained at 0°C for 1.5 h and at 35–37°C for 24 h. The mixture was filtered in order to remove the AgBr. The filtrate was washed three times with water to give 7.0 g of a crude product which was dried over P₄O₁₀ and distilled to give 0.40 g (6.2% yield) of CH₃OCF₂CF₂SO₂F, b p 45–46°C/143 mm, and 5.2 g (16.3 mmol) of product (trans 60%, cis 40%) in 54% yield, b p 119–122°C/143 mm

The infrared spectrum had the following bands (cm^{-1})
 3100(vw), 2966(vw), 2903(vw), 1631(m), 1455(vs), 1335(m), 1300(w),
 1244(s), 1202(s), 1138(s), 1117(s), 1033(w), 998(m), 934(m),
 801(vs), 725(w), 681(w), 653(m), 612(s), 548(w)

In the (EI)⁺ mass spectrum, the molecular ion was observed at
 320 [(H⁸¹BrC=CHCH₂OCF₂CF₂SO₂F), 2 47] and 318
 [(H⁷⁹BrC=CHCH₂OCF₂CF₂SO₂F), 3 88] The other main fragments were
 found at 239 [(M-HBr)⁺, 100 0], 155 [(C₄H₂F₃OS)⁺, 6 05], 134
 [(C₅HOF₃)⁺, 5 98], 133 [(C₅OF₃)⁺, 4 12], 123 [(C₄H₂OF₃)⁺, 8 33], 121
 [(⁸¹BrCH=CHCH₂)⁺, 55 28], 119 [(⁷⁹BrCH=CHCH₂)⁺, 51 31], 100
 [(C₂F₄)⁺, 20 54], 97 [(C₂F₃O)⁺, 4 57], 95 [(CF₂SO₂)⁺, 5 06], 93
 [(C₃H₃F₂O)⁺, 4 89], 67 [(SOF)⁺, 65 28], 59 [(C₂FO)⁺, 81 77], 55
 [(C₃H₂O)⁺, 24 52]

Anal Calcd for C₅H₄F₅O₃BrS C, 18 81, H, 1 25, F, 29 8, S,
 10 03, Br, 25 08 Found C, 18 97, H, 1 30, F, 29 8, S, 9 88, Br,
 24 85

(B) CsF Method In a similar procedure as reported above,
 4 80 g (31 5 mmol) of dried CsF, 7 0 mL diglyme, 6 80 g (37 7 mmol)
 of $\overline{\text{CF}_2\text{CF}_2\text{OSO}_2}$ were stirred at room temperature for 4 h, followed by
 reaction with 5 00 g (25 0 mmol) of BrCH=CHCH₂Br at 100-105°C for
 48 h The liquid product was washed with water in order to remove
 any solvent The crude product was dried over MgSO₄ and distilled
 to give 1 90 g product (24% yield) with a b p 76-78 C/20 mm

Preparation of CH₂=CBrCH₂OCF₂CF₂SO₂F

(A) AgF Method In a similar procedure as previously
 described, 3 90 g (30 7 mmol) of dried AgF, 6 0 mL of diglyme and

7.30 g (40.5 mmol) of $\overline{\text{CF}_2\text{CF}_2\text{OSO}_2}$ were mixed together. The mixture was stirred at room temperature for 5 h, and then 6.0 g (30.0 mmol) of $\text{CH}_2=\text{CBrCH}_2\text{Br}$ was added. The reaction mixture was stirred at 0 °C for 2 h, and at room temperature for 16 h. The AgBr was removed by filtration and the liquid filtrate was washed, dried over P_4O_{10} and distilled to give 3.00 g (9.4 mmol) of the olefinic product (33.1% yield) with a b.p. 107–109 °C/130 mm and 1.6 g of $\text{CH}_3\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$, 26% yield.

The infrared spectrum had the following bands (cm^{-1})
 2985(vw), 1655(m), 1640(m), 1465(s), 1405(w), 1375(vw), 1337(m),
 1300(s), 1210(s), 1145(s), 1125(s), 1040(w), 1005(m), 965(w),
 915(m), 821–795(vs), 655(m), 610(s), 546(w)

In the $(\text{CI})^+$ mass spectrum, the molecular ion was observed at 320 [$(\text{H}_2\text{C}=\text{C}^{81}\text{BrCH}_2\text{OCF}_2\text{CF}_2\text{SO}_2\text{F})$, 7.95] and 318 [$(\text{H}_2\text{C}=\text{C}^{79}\text{BrCH}_2\text{OCF}_2\text{CF}_2\text{SO}_2\text{F})$, 7.90]. The other main fragments were found at 239 [$(\text{M}-\text{Br})^+$, 12.89], 238 [$(\text{M}-\text{HBr})^+$, 11.73], 155 [$(\text{C}_4\text{H}_2\text{F}_3\text{OS})^+$, 5.01], 137 [$(\text{CH}_2=\text{C}^{81}\text{BrCH}_2\text{O})^+$, 1.34], 135 [$(\text{CH}_2=\text{C}^{79}\text{BrCH}_2\text{O})^+$, 1.17], 134 [$(\text{C}_5\text{HOF}_3)^+$, 1.52], 133 [$(\text{C}_5\text{OF}_3)^+$, 1.29], 121 [$(\text{CH}_2=\text{C}^{81}\text{BrCH}_2)^+$, 0.9472], 119 [$(\text{CH}_2=\text{C}^{79}\text{BrCH}_2)^+$, 1.000], 100 [$(\text{C}_2\text{F}_4)^+$, 1.282], 97 [$(\text{C}_2\text{F}_3\text{O})^+$, 0.470], 94 [$(\text{CFSO}_2)^+$, 2.69], 93 [$(\text{C}_3\text{H}_3\text{F}_2\text{O})^+$, 2.43], 79 [$(\text{C}_2\text{HF}_2\text{O})^+$, 0.657], 67 [$(\text{SOF})^+$, 3.257], 65 [$(\text{C}_4\text{HO})^+$, 1.543], 59 [$(\text{C}_2\text{FO})^+$, 0.3092], 55 [$(\text{CH}_2\text{CCH}_2\text{O})^+$, 1.086].

Anal. Calcd for $\text{C}_5\text{H}_5\text{F}_5\text{O}_3\text{BrS}$ C, 18.81, H, 1.25, F, 29.78, S, 10.03, Br, 25.08. Found C, 18.87, H, 1.22, F, 30.3, S, 10.03, Br, 24.97.

(B) CsF Method. In a similar way as reported above, 4.80 g (31.5 mmol) of dried cesium fluoride, 7.0 mL diglyme and 5.40 g

(30.0 mmol) of $\overline{\text{CF}_2\text{CF}_2\text{OSO}_2}$ were added in order to prepare the cesium salt, $\text{CsOCF}_2\text{CF}_2\text{SO}_2\text{F}$. The cesium salt was then reacted with 6.00 g (30.0 mmol) of $\text{CH}_2=\text{CBrCH}_2\text{Br}$ at 90-97°C for 24 h. The CsBr which formed was filtered and the filtrate was washed with water, dried over P_4O_{10} and distilled to give 1.15 g product in 12% yield.

Preparation of $(\text{CH}_3\text{CH}_2\text{O})_2\text{P}(\text{O})\text{CH}_2\text{CH}_2\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$ and $(\text{CH}_3\text{CH}_2\text{O})\text{PF}(\text{O})\text{CH}_2\text{CH}_2\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$

Into the reaction vessel previously described were added 4.10 g (32.3 mmol) of dried silver fluoride, 5.0 mL diglyme and 7.70 g (42.7 mmol) of $\overline{\text{CF}_2\text{CF}_2\text{OSO}_2}$ at -196°C. The mixture was stirred at room temperature for 16 h and 4.50 g (18.3 mmol) of diethyl-2-bromoethyl-phosphonate was then added at low temperature (-90 to -100°C). The reaction mixture was warmed and maintained at 35°C for 4 d. The volatile $\text{FC}(\text{O})\text{CF}_2\text{SO}_2\text{F}$ (2.50 g) was recovered. The AgBr precipitate was filtered in a dry atmosphere and the AgBr was washed twice with anhydrous diethyl ether. The washings were combined with the filtrate and then distilled to give a mixture of $(\text{CH}_3\text{CH}_2\text{O})_2\text{P}(\text{O})(\text{CH}_2)_2(\text{CF}_2)_2\text{SO}_2\text{F}$ (55%) and $(\text{CH}_3\text{CH}_2\text{O})\text{PF}(\text{O})(\text{CH}_2)_2\text{O}(\text{CF}_2)_2\text{SO}_2\text{F}$ (45%). Subsequently, this mixture was reacted with an excess of absolute alcohol at room temperature for 14 h and distilled to give 3.30 g of $(\text{CH}_3\text{CH}_2\text{O})_2\text{P}(\text{O})(\text{CH}_2)_2\text{O}(\text{CF}_2)_2\text{SO}_2\text{F}$, b.p. 127-129°C/0.7 mm in 54% yield.

Anal. Calcd for $\text{C}_8\text{H}_{14}\text{F}_5\text{O}_6\text{PS}$ C, 26.37, H, 3.85, F, 26.1, S, 8.79. Found C, 26.39, H, 3.75, F, 27.1, S, 8.96.

The infrared spectrum of $(\text{CH}_3\text{CH}_2\text{O})_2\text{P}(\text{O})\text{CH}_2\text{CH}_2\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$ had the following bands (cm^{-1}) 2987(w), 2924(w), 1455(s), 1398(w), 1335(m), 1244(s), 1195(m), 1138(m), 1110(m), 1033(s), 998(m), 955(w), 864(w), 800(vs), 737(w), 611(m)

In the $(\text{CI})^+$ mass spectrum of $(\text{CH}_3\text{CH}_2\text{O})_2\text{P}(\text{O})\text{CH}_2\text{CH}_2\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$, the $(\text{M}+1)^+$ peak was observed at 365 $[(\text{M}+1)^+, 47\ 80]$, the other main fragments were found at 311 $[(\text{M}-\text{CH}_3\text{F}_2)^+, 43\ 49]$, 291 $[(\text{M}-\text{CH}_4\text{F}_3)^+, 18\ 57]$, 227 $[(\text{CH}_2\text{CH}_2\text{OCF}_2\text{CF}_2\text{SO}_2\text{F})^+, 6\ 26]$, 225 $[(\text{C}_2\text{H}_2\text{COCF}_2\text{CF}_2\text{SO}_2\text{F})^+, 10\ 35]$, 205 $[(\text{C}_4\text{HF}_4\text{O}_3\text{S})^+, 9\ 46]$, 181 $[(\text{CH}_3\text{CH}_2\text{O})_2\text{P}(\text{O})\text{CH}_2\text{CH}_2\text{O}^+, 4\ 31]$, 165 $[(\text{CH}_3\text{CH}_2\text{O})_2\text{P}(\text{O})\text{CH}_2\text{CH}_2^+, 25\ 05]$, 151 $[(\text{CH}_3\text{CH}_2\text{O})_2\text{P}(\text{O})\text{CH}_2^+, 2\ 14]$, 137 $[(\text{CH}_3\text{CH}_2\text{O})_2\text{P}(\text{O})^+, 25\ 05]$, 123 $[(\text{C}_4\text{H}_2\text{OF}_3)^+, 9\ 32]$, 113 $[(\text{C}_4\text{HO}_2\text{S})^+, 16\ 13]$, 112 $[(\text{C}_4\text{O}_2\text{S})^+, 5\ 04]$, 97 $[(\text{C}_2\text{F}_3\text{O})^+, 5\ 10]$, 93 $[(\text{C}_3\text{H}_3\text{F}_2\text{O})^+, 18\ 75]$, 91 $[(\text{C}_2\text{H}_4\text{O}_2\text{P})^+, 12\ 56]$, 83 $[(\text{SO}_2\text{F})^+, 4\ 00]$, 81 $[(\text{C}_2\text{F}_3)^+, 15\ 79]$, 79 $[(\text{C}_2\text{HF}_2\text{O})^+, 5\ 18]$, 77 $[(\text{CH}_2\text{O}_2\text{P})^+, 7\ 57]$, 67 $[(\text{SOF})^+, 12\ 90]$, 65 $[(\text{C}_4\text{OH})^+, 18\ 33]$

It was very difficult to obtain a pure sample of $\text{CH}_3\text{CH}_2\text{OPF}(\text{O})\text{CH}_2\text{CH}_2\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$. However, it was possible to assign some characteristic fragments and the $(\text{M}+1)^+$ ion from mass spectrum patterns of a mixture of $(\text{CH}_3\text{CH}_2\text{O})_2\text{P}(\text{O})\text{CH}_2\text{CH}_2\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$ and $\text{CH}_3\text{CH}_2\text{OPF}(\text{O})\text{CH}_2\text{CH}_2\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$. The $(\text{M}+1)^+$ ion of $\text{CH}_3\text{CH}_2\text{OPF}(\text{O})\text{CH}_2\text{CH}_2\text{OCF}_2\text{CF}_2\text{SO}_2\text{F}$ appeared at 339 $[(\text{M}+1), 14\ 37]$, other main fragment ions were 337 $[(\text{M}-1)^+, 36\ 22]$, 323 $[(\text{M}-\text{CH}_3)^+, 4\ 99]$, 309 $[(\text{M}-\text{C}_2\text{H}_5)^+, 19\ 74]$, 139 $[(\text{CH}_3\text{CH}_2\text{OPF}(\text{O})\text{CH}_2\text{CH}_2)^+, 9\ 99]$, 138 $[(\text{CH}_3\text{CH}_2\text{OPF}(\text{O})\text{CH}=\text{CH}_2)^+, 27\ 58]$, 125 $[(\text{CH}_3\text{CH}_2\text{OPF}(\text{O})\text{CH}_2)^+, 16\ 99]$, 109 $[(\text{CH}_2=\text{CHOP}(\text{O})\text{F})^+, 38\ 78]$

ACKNOWLEDGEMENT

We express our appreciation to the U S Department of Energy, Grant No DE-FG21-88MC25142, for support of this work Dr Gary Knerr (University of Idaho) obtained the mass spectra

REFERENCES

- 1 G A Olah, P S Iyer and P Surya, *Synthesis*, (1986) 513
- 2 C Bunyagldj, H Plotrowska and M H Aldridge, *J Chem Eng Data*, 26 (1981) 344
- 3 J M Canich, M M Ludvig, G L Gard and J M Shreeve, *Inorg Chem* , 23 (1984) 4403
- 4 R J Terjeson, J Mohtasham and G L Gard, *Inorg Chem* , 27 (1988) 2916
- 5 L F Chen, J Mohtasham and G L Gard, *J Fluorine Chem* , 46 (1990) 21
- 6 T Gramstad and R N Haszeldine, *J Chem Soc* , (1956) 173
- 7 N S Ham, A N Hambly and R H Laby, *Aust J Chem* , 13 (1960) 443
- 8 J K Brown and K J Morgan, *Adv Fluorine Chem* , 4 (1965) 256
- 9 G S Reddy and R Schmutzler, *Z Naturforsch* , 256 (1970) 1199
- 10 D C England, M A Dietrich and R V Lindsey, *J Am Chem Soc* , 82 (1960) 6181