

Received: October 4, 1989; accepted January 20, 1990

PRELIMINARY NOTE

A Novel Nucleophilic Polyfluoroalkylating Agent $R_fI-P(NEt_2)_3$

WEI-YUAN HUANG*, CHANG-MING HU, YA-BO HE and HUI-FEN ZHOU


Shanghai Institute of Organic Chemistry, Academia Sinica,
Shanghai 200032 (China)**SUMMARY**

In the presence of tris(diethylamino)phosphine polyfluoroalkyl iodides reacted with various aroyl chlorides to give the corresponding aryl polyfluoroalkyl ketones.

Nigumi et al. reported that treatment of perfluoroalkyl iodides with an electron rich phosphine, i.e. tris(dialkylamino)phosphine ($P(NR_2)_3$), gave dialkylaminoperfluoroalkylphosphines ($(R_2N)_2PR_f$) [1]. Recently, Ruppert and Bürger et al. reported that, in the presence of tris(dialkylamino)phosphine, trifluoromethyl bromide reacted with electropositive elements E, e.g. R_3Si- , R_2B- , to give CF_3E [2,3].

We now report $R_fI-P(NEt_2)_3$ as a novel nucleophilic polyfluoroalkylating agent for aroyl chlorides.

In the presence of tris(diethylamino)phosphine, polyfluoroalkyl iodides reacted with aroyl chlorides in refluxing methylene chloride to give aryl polyfluoroalkyl ketones in moderate yield. The results were summarized in the Table.



TABLE


The reaction of polyfluoroalkyl iodides with tris(diethylamino) phosphine and aroyl chlorides^a

ArCOCl	R _f I	t(h)	Yield(%) ^b
p-ClC ₆ H ₄ COCl	CF ₃ (CF ₂) ₅ I	6	41
2,4-Cl ₂ C ₆ H ₃ COCl	CF ₃ (CF ₂) ₇ I	6	36
p-ClC ₆ H ₄ COCl	Cl(CF ₂) ₄ I	3.5	42
C ₆ H ₅ COCl	"	6	43
2,4-Cl ₂ C ₆ H ₃ COCl	"	4	43
p-ClC ₆ H ₄ COCl	Cl(CF ₂) ₆ I	5	41
p-BrC ₆ H ₄ COCl	"	7	40
o-ClC ₆ H ₄ COCl	"	6.5	32
C ₆ H ₅ COCl	"	6	41
2,4-Cl ₂ C ₆ H ₃ COCl	"	6.5	40

^a All reactions were carried out at 45°C in methylene chloride.

^b Isolated yields based on the starting ArCOCl.

The first step of such reaction is considered as a nucleophilic attack of phosphine on the positive iodine atom. In the presence of electrophiles, i.e. aroyl chloride, further nucleophilic attack of the anionic polyfluoroalkyl group on the carbonyl group leads to the formation of a ketone rather than diethylaminopolyfluoroalkyl phosphine[1].



Typical procedure: To a mixture of 3g(8.3 mmol) of ω -chlorooctafluorobutyl iodide and 525mg(3 mmol) of p-chlorobenzoyl chloride in dry CH_2Cl_2 (10 ml), was added 885mg(3 mmol) of tris-(diethylamino)phosphine within 5 min. The mixture was then refluxed at 45°C for 3.5h. After usual workup, 470 mg of p-chlorophenyl ω -chlorooctafluorobutyl ketone was isolated by column chromatography in a 42% yield. Anal. for $\text{C}_{11}\text{H}_4\text{Cl}_2\text{F}_8\text{O}$:

Calcd, C,35.23, H,1.07, Cl,18.91, F, 40.53; Found, C,34.86, H,0.91, Cl,18.72, F,41.16. IR(film) 1710, 1590, 1490, 1400, 1200-1150, 820 cm^{-1} ; $^1\text{H-NMR}(\text{CCl}_4)$: 7.26(d, J=9Hz, 2H), 7.81(d, J=9Hz, 2H); $^{19}\text{F-NMR}(\text{CCl}_4)$: -8.2(s,2F), +36.6(S, 2F), +42.8(S, 2F), +44.0(s,2F); MSm/z: 375(M^+), 377(M^++2), 311, 140, 111, 75. Elemental analyses and spectra of other products were all in accordance with their structure.

Further work on this nucleophilic polyfluoroalkylating agent is in progress.

- 1 Y. O. El Nigumi and H. J. Emeleus, J. Inorg. Nucl. Chem., 32 (1970) 3211.
- 2 I. Ruppert, K. Schlich and W. Volbach, Tetrahedron Lett., 25 (1984) 2195.
- 3 H. Burger, M. Grunwald and G. Pawelke, J. Fluorine Chem., 31 (1986) 89.