

Synthesis of pentafluorobenzoyloxy mono- to tetra-fluoroacetophenones

Linxiao Xu and Roger W. Giese*

Department of Pharmaceutical Sciences in the Bouve College of Pharmacy and Health Professions, Chemistry Department and Barnett Institute of Chemical Analysis, Northeastern University, Boston, MA 02115 (USA)

(Received December 30, 1992; accepted March 13, 1993)

Abstract

Pentafluorobenzoyloxyacetophenones are formed by reacting 2',3',4',5',6'-pentafluoroacetophenone and 2',3',4',5'-tetrafluoroacetophenone with pentafluorobenzyl alcohol (PFBA) in refluxing toluene under phase-transfer conditions. This provided seven new such acetophenones. Pentafluorobenzoyloxy-substituted mono- and difluoroacetophenones were synthesized by reacting *p*-hydroxyacetophenone with pentafluorobenzyl bromide.

Introduction

Studies are being undertaken in our laboratories on electrophoric derivatives of acetophenone as molecular labels to replace radioisotopes in some analytical procedures. An electrophore is a compound that readily combines with an electron in the gas phase, and hence can be measured by gas chromatography using electron capture detection (GC-ECD) or a related technique. Previously, we have described in detail the use of electrophores as labels [1], including the synthesis of *p*-(pentafluorobenzoyloxy)acetophenone, and have detected electrophore-labeled DNA and albumin as a combined sample by GC-ECD methods [2]. Establishing a diversity of electrophoric derivatives of acetophenone is of interest since a key advantage of electrophores as labels is the potential for measuring many of them as a mixture by GC-ECD methods. In particular, polyfluoro derivatives of acetophenone are attractive because fluorine atoms promote volatility and electron capture. Here we report some reactions which help to build up the diversity of volatile pentafluorobenzoyloxy-substituted mono- to tetra-fluoroacetophenones.

Results and discussion


Recently, we have found that the pentafluorobenzyl group is lost when 2,6-difluorophenyl pentafluorobenzyl ether is subjected to Friedel-Crafts acetylation conditions [3]. Hence, such reaction conditions are not

attractive for the formation of pentafluorobenzoyloxyacetophenones from the corresponding pentafluorobenzyl aromatic ethers. Both *o*- and *p*-fluorophenyl acetate are known to undergo the Fries rearrangement smoothly in the presence of Lewis acid catalysts [4]. However, no examples have been reported of the Fries rearrangement of difluoroaryl esters. As shown below, we have prepared 3'-fluoro- and 3',5'-difluoro-4'-hydroxyacetophenone (**3** and **4**) under Fries rearrangement conditions. The former compound forms more readily, providing a 65% yield of **3** after 3 h and a 52% yield of **4** after 5 h at 140–150 °C. By reacting pentafluorobenzyl bromide (PFB-Br) with **3** and **4** in the presence of potassium carbonate in refluxing acetone, we obtained 3'-fluoro- and 3',5'-difluoro-4'-pentafluorobenzoyloxyacetophenones (**5** and **6**) in high yield (91% and 88%, respectively).


It is well known that polyfluoroaromatics undergo nucleophilic substitution reactions [5], and 2',3',4',5',6'-pentafluoroacetophenone (**7**) has been ring-substituted previously with sodium azide (4'-substitution) [6], sodium sulfide (4'-substitution) [7] and aliphatic or aromatic amines (2'- and 4'-substitutions) [8, 9]. The acetyl group of **7** has been removed by alkoxide [10, 11] and such conditions have also caused **7** to undergo self-condensation instead of nucleophilic displacement of the fluorine [12]. A low yield of such displacement with ethanol has been observed in the presence of excess potassium cyanide [13].

Reaction of **7** with 2',3',4',5',6'-pentafluorobenzyl alcohol (PFBA) in DMF in the presence of potassium carbonate for 3 h at 120 °C was least successful for our purposes, giving little product diversity: 66% of **8**

* Author to whom correspondence should be addressed.



Scheme 1.



Scheme 2.

and 2% of **9**. These results are similar to those observed by Kobrina *et al.* [8], who used nitromethane as the solvent to obtain mainly *para* substitution of **7** with amines. By changing to refluxing toluene, and adding 0.1 g of 18-crown-6 as a phase-transfer catalyst, we obtained much more product diversity: **8** (46%), **9** (16%), **10** (5%) and **11** (2%). All of these products are acetophenone derivatives as desired. It is possible that **10** and **11** are derived from **8** and **9**, rather than via a *para*-ether dimer of PFBA, since we did not find any such dimer in the product mixture.

Nucleophilic substitution of tetrafluoroacetophenone has not been reported. Reaction of 2',3',4',5'-tetrafluoroacetophenone (**12**) with PFBA as above (refluxing toluene, 18-crown-6) was also successful in producing some diversity, as shown here.

The major product was *para*-substituted **13** (71% yield). Whilst no *ortho*-substituted product was observed, the *meta*-substituted product **14** formed in a small amount (2%). Perhaps formation of some **14** is not too surprising, given that hydrogen on a polyfluorophenyl ring tends to direct substitution to the *ortho* and *para* positions [5]. Compound **15** (5%) was also obtained, apparently, via **13**.

Thus, we have added nine new compounds to the category of volatile pentafluorobenzoyloxy acetophenones

to help the development of electrophoric labels for analytical purposes.

Experimental


General

¹H and ¹³C NMR spectra in ppm (δ) were obtained on a Varian XL-300FT instrument; CDCl₃ was used as the solvent and tetramethylsilane (TMS) was used as the internal standard. Infrared (IR) spectra were taken on a Perkin-Elmer model 588B spectrophotometer. Preparative gas chromatographic separations were run on a Varian 3300 instrument and helium was used as the carrier gas; column A: 20% SE-30/Chromosorb W (NAW, 60–80 mesh), 6 ft. \times 1/4 in. aluminum column; column B: 10% SE-30/Chromosorb W (NAW, 60–80 mesh), 3 ft. \times 1/4 in. aluminum column.

3'-Fluoro-4'-pentafluorobenzoyloxyacetophenone (**5**)

Synthesis of 2'-fluorophenyl acetate (**1**)

To a mixture of 2-fluorophenol (5 g, 45 mmol), 50 ml of ether and potassium carbonate (16 g), 5 ml of acetic anhydride was added. After stirring under reflux for 3 h, the reaction mixture was stirred at room temperature overnight. The solid was removed by fil-



Scheme 3.

tration and washed three times with ether. The filtrate was evaporated to remove the solvent and distilled to obtain 1 (4.43 g, 64%). ¹H NMR δ: 2.32 (s, 3H, CH₃); 6.90–7.27 (m, 4H, H-Ar) ppm. ¹³C NMR δ: 20.3 (CH₃CO); 116.6 (⁴J_{C-F}=19 Hz, C5'); 123.7 (C4'/C6'); 124.4 (³J_{C-F}=4 Hz, C6'/C4'); 127.0 (²J_{C-F}=7 Hz, C3'); 138.1 (²J_{C-F}=13 Hz, C1'); 154.1 (J_{C-F}=249 Hz, C2'); 168.3 (C=O) ppm.

Synthesis of 3'-fluoro-4'-hydroxyacetophenone (3)

A mixture consisting of 1 (1.54 g, 10 mmol) and aluminum chloride (1.70 g, 12.5 mmol) in 10 ml of nitrobenzene was stirred at 140–150 °C for 3 h. After cooling to room temperature, 15 ml of ice/water was added followed by 30 ml of 3 N HCl. The water layer was extracted with ether (3 × 30 ml) and the combined organic layers were washed three times with concentrated potassium carbonate solution and then the combined water layer was acidified with 2 N HCl. After extraction three times with ether, the organic layer was dried over anhydrous MgSO₄. Removal of ether by evaporation afforded 1.0 g of 3 (65%). This was used directly for the next step.

Synthesis of 5

A mixture consisting of 3 (1 g, 6.5 mmol) and PFB-Br (6.5 mmol), 30 ml of acetone and 5 g of potassium carbonate was stirred under reflux for 5 h. After addition of 5 g of Celite 521, the reaction mixture was filtered and the residue washed with acetone (3 × 30 ml). The filtrate was then evaporated on a Rotavapor. After recrystallization (hexane/ether), 1.98 g of product (91%) was obtained, m.p. 99–100 °C. ¹H NMR δ: 2.51 (s, 3H, CH₃CO); 5.24 (s, 2H, CH₂O); 7.11 (t, 1H, J₅₋₆=8.8 Hz, ⁴J_{H-F}=8.2 Hz, H-C5'); 7.62 (d, 1H, ³J_{H-F}=11.2 Hz, H-C2'); 7.70 (d, 1H, J₆₋₅=8.8 Hz, H-C6') ppm. ¹³C

NMR δ: 25.9 (CH₃-CO); 58.5 (CH₂O); 109.1 (C1', ²J_{C-F}=17 Hz); 114.4 (C5'); 116.0 (C2', ²J_{C-F}=19 Hz); 125.4 (C6', ⁴J_{C-F}=3 Hz); 131.7 (C1', ³J_{C-F}=5 Hz); 137.4 (C2'' and C6'', J_{C-F}=254 Hz); 141.9 (C4', J_{C-F}=256 Hz); 145.7 (C3'' and C5'', J_{C-F}=251 Hz); 149.8 (C4', ²J_{C-F}=11 Hz); 152.2 (C3', J_{C-F}=249 Hz); 195.5 (C=O) ppm. MS (EI) m/z: 334 (M⁺) MS (ECNI) m/z: 153 ([M-PFBz]⁻, 100%).

3',5'-Difluoro-4'-pentafluorobenzyloxyacetophenone (6)

Synthesis of 2',6'-difluorophenyl acetate (2)

To a stirred mixture consisting of 2,6-difluorophenol (3.6 g, 27.7 mmol), 30 ml of toluene and 30 ml of pyridine was added 3 ml of acetyl chloride at room temperature. After stirring for 5 h, 30 ml of water was added. The organic layer was separated and evaporated. The residue was diluted with 40 ml of ether and then washed with 30 ml of 5% HCl followed by 30 ml of 5% sodium carbonate. After drying over anhydrous MgSO₄, distillation afforded 4.3 g of product (90%). ¹H NMR δ: 2.33 (s, 3H, CH₃CO); 6.94 (t, 2H, J_{H-F}=J_{H-H}=7.9 Hz, H-C3 and H-C5); 7.12 (m, 1H, H-C4) ppm. ¹³C NMR δ: 19.5 (CH₃CO); 111.9 (C3 and C5, ²J_{C-F}=22 Hz, ⁴J_{C-F}=6 Hz); 126.3 (C4, ³J_{C-F}=9 Hz); 155.2 (C2 and C6, J_{C-F}=250 Hz, ³J_{C-F}=4 Hz); 167.2 (C=O) ppm. IR (film) ν_{max} (cm⁻¹): 3010 (w); 1730 (vs) (C=O); 1565 (m); 1440 (s); 1335 (m); 1255 (s); 1210 (m); 1120 (vs) (C-O); 1010 (m); 965 (s); 860 (s); 740 (s); 710 (m); 670 (w).

Synthesis of 3',5'-difluoro-4'-hydroxyacetophenone (4)

A mixture consisting of 2 (4.2 g, 24.4 mmol) and aluminum chloride (6.0 g, 4.5 mmol) in 100 ml of nitrobenzene was stirred at 140–150 °C for 5 h. Work-up was the same as for the synthesis of 3. Product 4 (2.2 g) was obtained (52%). ¹H NMR δ: 2.32 (s, 1H,

OH); 2.57 (s, 3H, CH₃); 7.53 (d, 2H, $J=8.1$ Hz, HC2' and HC6') ppm. ¹³C NMR δ : 25.4 (CH₃); 111.8 ($^2J_{C-F}=22$ Hz, $^4J_{C-F}=8$ Hz, C2' and C6'); 117.8 ($^3J_{C-F}=9$ Hz, C1'); 139.1 ($^2J_{C-F}=16$ Hz, C4'); 151.6 ($J_{C-F}=245$ Hz, $^3J_{C-F}=7$ Hz, C3' and C5'); 196.0 (C=O) ppm.

Synthesis of 6

A mixture consisting of 4 (2.2 g, 12.8 mmol), PFB-Br (12.8 mmol), 30 ml of acetone and 5 g of potassium carbonate was stirred under reflux for 5 h. Work-up was as usual, giving 4 g of product (88%), m.p. 62–63 °C. ¹H NMR δ : 2.58 (s, 3H, CH₃CO); 5.38 (s, 2H, CH₂O); 7.52 (d, 2H, $^3J_{H-F}=8.7$ Hz, H-C2' and H-C6') ppm. ¹³C NMR δ : 25.8 (CH₃CO); 62.3 (CH₂O); 109.5 (C1'', $^2J_{C-F}=17$ Hz); 112.3 (C2' and C6', $^2J_{C-F}=24$ Hz, $^4J_{C-F}=8$ Hz); 132.8 (C1', $^3J_{C-F}=7$ Hz); 137.4 (C2'' and C6'', $J_{C-F}=252$ Hz); 138.2 (C4', $^2J_{C-F}=15$ Hz); 142.0 (C4'', $J_{C-F}=256$ Hz); 145.8 (C3'' and C5'', $J_{C-F}=253$ Hz); 155.5 (C3' and C5', $J_{C-F}=251$ Hz, $^3J_{C-F}=5$ Hz); 194.4 (C=O) ppm. MS (EI) m/z : 352 (M⁺). MS (ECNI) m/z : 171 ([M – PFBz][–], 100%).

4'-Pentafluorobenzoyloxy-2',3',5'-trifluoroacetophenone (13)

A mixture consisting of 12 (0.96 g, 5 mmol), PFBA (1.0 g, 5 mmol), potassium carbonate (7 g), toluene (30 ml) and 18-crown-6 (0.1 g) was stirred under reflux for 8 h. After cooling to room temperature and adding 2 g Celite 521, the reaction mixture was filtered and the residue washed three times with ether. After evaporation of the filtrate on a Rotavapor, the crude products were separated by preparative gas chromatography using column A, giving 1.31 g of product (71%), m.p. 76–77 °C. ¹H NMR δ : 2.65 (d, 3H, $^5J_{H-F}=5.2$ Hz, CH₃–CO); 5.42 (s, 2H, CH₂O); 7.46 (ddd, 1H, $^2J_{H-F}=11.3$ Hz, $^3J_{H-F}=6.1$ Hz, $^4J_{H-F}=2.4$ Hz, H-C6') ppm. ¹³C NMR δ : 31.1 (CH₃CO, $^4J_{C-F}=8$ Hz); 62.6 (CH₂O); 109.1 (C1'', $^2J_{C-F}=19$ Hz); 111.1 (C6', $^2J_{C-F}=21$ Hz, $^3J_{C-F}=3$ Hz); 121.0 (C1'); 137.5 (C2' and C6'', $J_{C-F}=250$ Hz); 142.2 (C4'', $J_{C-F}=249$ Hz); 144.6 (C3', $J_{C-F}=253$ Hz, $^2J_{C-F}=17$ Hz, $^3J_{C-F}=5$ Hz); 145.8 (C3'' and C5'', $J_{C-F}=251$ Hz); 148.2 (C2', $J_{C-F}=254$ Hz, $^2J_{C-F}=13$ Hz, $^4J_{C-F}=3$ Hz); 151.0 (C5', $J_{C-F}=248$ Hz, $^3J_{C-F}=^4J_{C-F}$ 3 Hz); 168.4 (C4'); 192.4 (C=O) ppm. MS (EI) m/z : 370 [M]⁺. MS (ECNI) m/z : 189 ([M – PFBz][–], 100%).

4'-Pentafluorobenzoyloxy-2',3',5',6'-tetrafluoroacetophenone (8)

A mixture consisting of 7 (1.05 g, 5 mmol), PFBA (1.0 g, 5 mmol), potassium carbonate (7 g), toluene (30 ml) and 18-crown-6 (0.1 g) was stirred under reflux for 8 h. Work-up was the same as for the synthesis of 13, giving 0.89 g of product (46%), m.p. 76–77 °C. ¹H NMR δ : 2.62 (t, 3H, $^5J_{H-F}=2.0$ Hz, CH₃CO); 5.41 (s, 2H, CH₂O) ppm. ¹³C NMR δ : 32.3 (CH₃CO); 63.0

(CH₂O); 108.9 (C1'', $^2J_{C-F}=17$ Hz); 114.4 (C1', $^2J_{C-F}=16$ Hz); 137.5 (C2'' and C6'', $J_{C-F}=258$ Hz); 138.6 (C4'); 141.2 (C3' and C5', $J_{C-F}=249$ Hz); 142.4 (C4'', $J_{C-F}=257$ Hz); 144.8 (C2' and C6', $J_{C-F}=263$ Hz); 145.8 (C3'' and C5'', $J_{C-F}=252$ Hz); 195.5 (C=O) ppm. MS (EI) m/z : 388 (M⁺). MS (ECNI) m/z : 207 ([M – PFBz][–], 100%). In addition to 8, 0.31 g (16%) of 2'-pentafluorobenzoyloxy-3',4',5',6'-tetrafluoroacetophenone (9), 142 mg (5%) of 4'-(4''-pentafluorobenzoyloxy-2'',3'',5'',6''-tetrafluorobenzyl)oxy-2',3',5',6'-tetrafluoroacetophenone (10) and 57 mg (2%) of 2'-(4''-pentafluorobenzoyloxy-2'',3'',5'',6''-tetrafluorobenzyl)oxy-3',4',5',6'-tetrafluoroacetophenone (11) were also obtained by preparative gas chromatography separation.

Compound 9: ¹H NMR δ : 2.51 (d, 3H, $^5J_{H-F}=1.6$ Hz, CH₃CO); 5.25 (s, 2H, with fine splitting, CH₂O) ppm. ¹³C NMR δ : 32.3 (CH₃CO); 63.7 (CH₂O); 108.9 (C1'', $^2J_{C-F}=18$ Hz); 120.9 (C1'); 137.5 (C2'' and C6'', $J_{C-F}=253$ Hz); 137.7 (C4' and C5', $J_{C-F}=253$ Hz); 138.2 (C2'); 142.1 (C4'', $J_{C-F}=252$ Hz); 142.2 (C3'/C6', $J_{C-F}=252$ Hz); 143.8 (C6'/C3', $J_{C-F}=254$ Hz); 145.8 (C3'' and C5'', $J_{C-F}=259$ Hz); 194.0 (C=O) ppm.

Compound 10: ¹H NMR δ : 2.62 (t, 3H, $^5J_{H-F}=1.9$ Hz, CH₃CO); 5.38 (s, 2H, CH₂O); 5.39 (s, 2H, CH₂O) ppm. ¹³C NMR δ : 32.5 (CH₃CO); 63.0 (CH₂O); 63.3 (CH₂O); 108.6 (C1'' and C1#, $^2J_{C-F}=18$ Hz); 127.5 (C1'); 137.6 (C2# and C6#, $J_{C-F}=246$ Hz); 137.7 (C4'/C4''); 138.0 (C4''/C4'); 141.2 (C3', C5', C2'' and C6'', $J_{C-F}=256$ Hz); 142.2 (C4#, $J_{C-F}=244$ Hz); 144.8 (C2' and C6', $J_{C-F}=250$ Hz); 145.9 (C3'', C5'', C3# and C5#, $J_{C-F}=251$ Hz); 191.7 (C=O) ppm.

Compound 11: ¹H NMR δ : 2.51 (d, 3H, $^5J_{H-F}=1.5$ Hz, CH₃CO); 5.21 (s, 2H, CH₂O); 5.36 (s, 2H, CH₂O) ppm. ¹³C NMR δ : 32.3 (CH₃CO); 62.9 (CH₂O); 63.5 (CH₂O); 109.1 (C1'' and C1#, $^2J_{C-F}=18$ Hz); 120.9 (C1'); 137.5 (C4', C5, C2# and C6#, $J_{C-F}=250$ Hz); 137.6 (C4''); 138.8 (C2'); 141.5 (C4#, $J_{C-F}=252$ Hz); 142.4 (C2'' and C6'', $J_{C-F}=252$ Hz); 144.0 (C3'/C6', $J_{C-F}=250$ Hz); 145.8 (C3'', C5'', C3# and C5#, $J_{C-F}=251$ Hz); 146.1 (C6'/C3', $J_{C-F}=249$ Hz); 191.7 (C=O) ppm.

Acknowledgment

This work was supported by NIH Grant HG00029, Contribution No. 571 from the Barnett Institute.

References

- 1 S. Abdel-Baky, N. Klempier and R.W. Giese, *Tetrahedron*, 46 (1990) 5859.
- 2 S. Abdel-Baky, K. Allam and R.W. Giese, *Anal. Chem.*, 65 (1993) 498.
- 3 L. Xu and R.W. Giese, *Tetrahedron Lett.*, 34 (1993) 3829.

- 4 (a) S.M. Kelly, *Helv. Chim. Acta*, 72 (1989) 594; (b) R. Martin, *Bull. Soc. Chim. Fr.* (1985) 239; (c) J. Allen and D. Giffard, *J. Labelled Compd. Radiopharm.*, 19 (1982) 301; (d) K.C. Joshi and S.C. Bahel, *J. Indian Chem. Soc.*, 37 (1960) 687.
- 5 (a) R.D. Chambers, *Fluorine in Organic Chemistry*, Wiley, New York, 1973, p. 275; (b) L.S. Kобрina, in P. Tarrant (ed.), *Fluorine Chemistry Reviews*, Marcel Dekker, New York, 1974, Vol. 7, p. 1.
- 6 (a) J.F.W. Keana and S.X. Cai, *J. Org. Chem.*, 55 (1990) 3640; (b) *idem*, *J. Fluorine Chem.*, 43 (1989) 151.
- 7 G.G. Furin, S.A. Krupoder and G.G. Yakobson, *Izv. Sib. Otd. Akad. Nauk SSSR, Ser. Khim. Nauk*, (1976) 147; [*Chem. Abs.*, 86 (1977) 72 110j].
- 8 L.S. Kобрina, G.G. Furin and G.G. Yakobson, *J. Org. Chem. USSR, (Engl. Ed.)*, 6 (1970) 510.
- 9 (a) T.N. Gerasimova, L.L. Gelumbovskaya, I.I. Baturina and E.P. Forkin, *Izv. Sib. Otd. Akad. Nauk SSSR, Ser. Khim. Nauk*, (1973) 88 [*Chem. Abs.*, 79 (1973) 53 161r]; (b) O.P. Sheremet, P.P. Rodionov and S.M. Shein, *Ibid.*, 71 (1975) [*Chem. Abs.*, 84 (1976) 89 725v].
- 10 J.H. James, M.E. Peach and C.R. Williams, *J. Fluorine Chem.*, 27 (1985) 91.
- 11 O.I. Andreevskaya, V.A. Barkhash, I.K. Korobeinicheva, E.D. Krivousova, V.A. Sokolenko, G.G. Yakobson and L.M. Yan'kova, *J. Org. Chem. USSR (Engl. Ed.)*, 6 (1970) 713.
- 12 O.V. Zakharova, V.M. Vlasov and G.G. Yakobson, *J. Org. Chem. USSR (Engl. Ed.)*, 15 (1979) 1964.
- 13 T.N. Vasilevskaya, A.G. Badashkeeva, T.N. Gerasimova, V.A. Barkhash and N.N. Vorozhtsov, Jr., *J. Org. Chem. USSR (Engl. Ed.)*, 6 (1970) 125.