

to the trans isomer, i.e. when the carbonyl and phenyl groups have a trans relationship to each other.

The ^{19}F NMR spectra of **1**, both in CD_2Cl_2 and $\text{THF-}d_8$, gave two multiplets of equal intensity, with a line-shape compatible with an asymmetrically para-disubstituted fluorobenzene, and two signals compatible with symmetrically para-disubstituted fluorobenzenes. This is interpreted as indicating the presence of the cis–trans, trans–trans and cis–cis isomers [6]. It is proposed that the trans–trans resonances lie to the low field of those of the cis–cis [6] isomer and that from the ^{19}F integration the ratio of trans–trans/cis–cis/cis–trans is 74:2:24 in CD_2Cl_2 at ambient temperature. The ^1H spectrum seems more complicated but is in reasonable agreement with the proposed structure and isomers. Compound **2** had a pale pink colour [1] when recrystallized. The colour darkened by leaving the compound dissolved, most probably as a result of oxidation. On standing, a sublimed sample [2] acquired a pale pink colour.

3. Experimental

NMR spectra were obtained with a Varian XL-300 (300 MHz proton frequency) spectrometer. ^1H chemical shifts are reported with reference to SiMe_4 and ^{19}F chemical shifts with respect to C_6F_6 with a reference line position at 163.0 ppm. Downfield shifts were positive. Mass spectroscopy data were obtained with a VG Micromass 7070F instrument with the data system VG 11-250J.

All reactions were performed under N_2 . Reaction mixtures were hydrolysed, after adding ether (30 ml), by addition of 10% hydrogen chloride to give an acidic medium, extracted with ether, dried with anhydrous sodium sulphate, concentrated and chromatographed on silica (230–400 mesh) starting with light petroleum as the elution solvent. When no more could be eluted from the mixture, more and more ether was added to the eluent, ending with pure ether. Recrystallization, when necessary, was performed from a mixture of light petroleum (b.p. 40–60 °C) and ether.

3.1. Sodium formanilide in tetrahydrofuran

Sodium hydride (0.042 mole, 1.00 g in oil) was added slowly to a mixture of formanilide (0.042 mole, 5.09 g) in tetrahydrofuran (20 ml) on an ice/water bath. The mixture was stirred at ambient temperature over night to give a low viscosity, white mixture used in the following reactions.

3.1.1. In tetrahydrofuran/dimethylsulphoxide

I. Dimethylsulphoxide (20 ml) was added to the sodium anilide mixture placed on an ice/water bath and followed by the addition of hexafluorobenzene (0.02 mole, 3.72 g). A green mixture was formed. The ice/water bath was removed. A faint exothermic reaction took place after 20–30 min, and a clear, brown reaction mixture was formed. Work up next morning gave 2.14 g (28%) *N,N'*-diformyl-*N,N'*-diphenyl-

2,3,5,6-tetrafluorophenylene-1,4-diamine (**1**) (nc), m.p. 229–229.5 °C, and 1.3 g (20%) *N,N'*-diphenyl-2,3,5,6-tetrafluorophenylene-1,4-diamine (**2**), m.p. 151–153 °C [1,2].

Found for **1**: m/z : (i.e. 70 eV) (% rel. int.): 390(3), 389(24), 388(100), 361(5), 360(22), 333(17), 332(86), 331(12), 312(7), 311(9), 310(3), 293(4), 292(17), 291(18), 267(3), 266(14), 265(3), 257(17), 255(15), 235(8), 220(10), 208(6), 166(12), 104(6), 77(87). High resolution: $\text{C}_{20}\text{H}_{12}\text{F}_4\text{N}_2\text{O}_2$ requires $M = 388.0835$. Found, 388.0855. NMR (CD_2Cl_2): δ (–CHO) 8.65 (s), 8, 42 (t). ^{19}F NMR (CD_2Cl_2): δ 19.65 (m), 15.93 (m) (amount 24%), 18.59 (small sideband at 18.64. In $\text{THF-}d_8$ only a singlet was present at 18.67) (amount 74%) and 16.96 (s) (amount 2%). IR in KBr: 1703 (s), 1593 (m), 1502 (s), 1492 (w), 1345 (m), 1271 (s), 1217 (s), 1003 (m), 973 (m), 875 (m), 768 (m), 709 (m), 696 (w), 643 (w), 599 (m), 522 (w).

II. Reaction mixture as in I. 1.5 h after addition of hexafluorobenzene the mixture was placed on a 95 °C bath for 2.5 h, cooled on an ice/water bath and hydrolysed with 40 ml 10% HCl. Work up gave 3.91 g (59%) *N,N'*-diphenyl-2,3,5,6-tetrafluorophenylene-1,4-diamine (**2**) [2].

3.1.2. In tetrahydrofuran/dimethylformamide

To the sodium formanilide mixture placed on an ice/water bath dimethyl-formamide (20 ml) and then hexafluorobenzene (0.02 mole, 3.72 g) were added whereupon the bath was removed. After 1 h the mixture was placed on a 95 °C bath for 4.5 h. Gas chromatography (GC) of the crude showed traces of *N,N'*-diformyl-*N,N'*-diphenyl-2,3,5,6-tetrafluorophenylene-1,4-diamine (**1**), together with *N,N'*-diphenyl-2,3,5,6-tetrafluorophenylene-1,4-diamine. Work up gave 1.87 g (28%) *N,N'*-diphenyl-2,3,5,6-tetrafluorophenylene-1,4-diamine (**2**) [2].

3.2. Sodium formanilide in dimethylformamide

Sodium formanilide was formed in dimethylformamide as described above for tetrahydrofuran. Next morning hexafluorobenzene (0.02 mole, 3.72 g) was added all at once to the sodium formanilide mixture placed on a tap-water bath. After 2 h the bath temperature was raised to 80 °C and held for 6 h. Work up gave 3.02 g (45%) *N,N'*-diphenyl-2,3,5,6-tetrafluorophenylene-1,4-diamine (**2**) [2].

3.3. Lithium hydroxide in tetrahydrofuran

Lithium hydroxide (1.5 g) was added during 30 min to a stirred mixture of formanilide (0.042 mole, 5.09 g) in tetrahydrofuran (20 ml) at ambient temperature. Next morning hexafluorobenzene (0.02 mole, 3.72 g) was added all at once. The yellow mixture was hydrolysed after being refluxed slowly for 24 h. GC analysis of the dried ether solution showed only two main compounds present, formanilide and 2,3,4,5,6-pentafluorodiphenylamine. After the solvent was

stripped off, the mixture was chromatographed with light petroleum resulting in 1.0 g (19%) 2,3,4,5,6-pentafluorodiphenylamine (**3**) [1,2].

References

- [1] J. Burdon, J. Castaner and J.C. Tatlow, *J. Chem. Soc.*, (1964) 5017.
- [2] R. Koppang, *J. Organometal. Chem.*, 46 (1972) 193.
- [3] R. Koppang, *J. Fluorine Chem.*, 8 (1976) 389.
- [4] R. Koppang, *J. Polym. Sci.: Polym. Chem. Ed.*, 14 (1976) 2225.
- [5] S. Hammerum and P. Wolkoff, *Adv. Mass Spectrometry*, 7B (1978) 1318.
- [6] R. Koppang and D. Grace, *J. Fluorine Chem.*, 66 (1994) 27.
- [7] A.J.R. Bourn, D.G. Gillies and E.W. Randall, *Tetrahedron*, 20 (1964) 1811.
- [8] A.J.R. Bourn, D.G. Gillies and E.W. Randall, *Tetrahedron*, 22 (1966) 1825.
- [9] R. Koppang and K. Schaumburg, *J. Fluorine Chem.*, 12 (1978) 169.