

FOR THE RECORD

S. Panneerchelvam,¹ M.Sc.; K. Kumara Thevan,¹ Lai KokFai,¹ M. Saravanakumar,¹ V. Sumathy,¹ K. C. Yuvaneswari,¹ and M. N. Norazmi,¹ Ph.D.

Polymorphism of 9 STRs in Ethnic Chinese Population of Malaysia*

POPULATION: Random Chinese population, Malaysia

KEYWORDS: forensic science, DNA typing, database, Malaysia, population genetics

Buccal swabs were collected from random ethnic Chinese individuals living in peninsular Malaysia. DNA was extracted by simple salting out procedure (1). Using STR multiplex primer kits,

¹ School of Health Sciences, Universiti Sains Malaysia, 16150 Kubang Kerian, Kelantan, Malaysia.

* This research was funded by Experimental Applied Research grant scheme, Ministry of Science, Technology and the Environment, Malaysia (09-02-05-3122 EA011).

Promega GeneprintTM (CTT, FFV, and STR III) and following the manufacturer's guidelines 10 ng of DNA was PCR amplified. Allele frequencies were calculated from the numbers of each genotype by gene count method. The randomness of the population was ascertained by subjecting the data to Chi-square test. No deviations from Hardy-Weinberg equilibrium were observed. Heterozygosity and discrimination power were calculated as per the methods already reported (2-4). The allelic distribution in the Chinese population

TABLE 1—Allele frequency for 9 STR loci in ethnic Chinese population of Malaysia (*n* = 102).

Allele	Frequency								
	CSF1PO	TPOX	TH01	F13A01	FESFPS	vWA	D16S539	D7S820	D13S317
3.2	0.2250
4	0.2550
5	0.0650
6	0.0943	0.4350	0.0050	...
7	0.0090	0.0138	0.2453	0.0150	0.0050	0.0100
8	0.0150	0.5321	0.0708	0.0450	0.0930	0.3000
9	0.1080	0.0917	0.4858	...	0.0149	...	0.2000	0.0970	0.1500
9.3	0.0142
10	0.2500	0.0413	0.0802	...	0.0644	...	0.1550	0.1390	0.1500
11	0.2450	0.2982	0.0047	...	0.4059	...	0.2950	0.4030	0.1700
12	0.3140	0.0229	0.0047	...	0.3514	...	0.2050	0.2220	0.1600
13	0.0540	0.0050	0.1634	0.0150	0.0820	0.0320	0.0500
14	0.0050	0.2600	0.0140	0.0050	0.0100
15	0.0650	0.0050
16	0.2150
17	0.2050
18	0.1450
19	0.0800
20	0.0100
21	0.0050
H	87.31	83.66	78.61	82.90	85.08	91.43	90.34	92.95	92.39
PE	68.40	52.29	60.46	59.28	57.85	75.17	73.33	67.80	74.69
PD	0.9269	0.7963	0.8262	0.8983	0.8695	0.9330	0.9295	0.9027	0.8776
Chi	5.46	4.62	6.61	10.87	10.6	7.54	4.96	7.76	11.29
(<i>p</i> < 0.05)	(df 8)	(df 6)	(df 6)	(df 7)	(df 6)	(df 12)	(df 13)	(df 11)	(df 11)
CDP	0.999999998								

H: Heterozygosity; PE: Power of exclusion; PD: Power of discrimination; Chi: Chi square; CDP: Cumulative discrimination power.

of Malaysia is found to be different from that of ethnic Malay and Tamil populations of Malaysia (5,6).

The complete dataset is available to any interested party at <http://www.ppsk.usm.my>.

References

1. Miller SA, Dykes DD, Polesky HF. A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res* 1988;16: 12–5.
2. Nei M, Roychoudhury AK. Sampling variances of heterozygosity and genetic distance. *Genetics* 1974;76:379–90.
3. Jones DA. Blood samples: probability of discrimination. *J Forensic Sci Soc* 1972;12:355–9.
4. Ohno Y, Sebetan IM, Akaishi S. [A simple method for calculating the probability of excluding paternity with any number of co dominant alleles](#). *Forensic Science Int* 1982;19:93–8.

[PubMed]

[PubMed]

[probability of excluding paternity with any number of co dominant alleles](#).

Forensic Science Int 1982;19:93–8.

5. Panneerchelvam S, Nur Haslindawaty, Ravichandran M, Norazmi MN, Zainuddin ZF. Allele frequency distribution for 10 STR loci in the Malay population of Malaysia. *J Forensic Sci* 2003;48(2):451–2.

6. Panneerchelvam S, Kumara Thevan K, Lai KokFai, Saravanakumar M, Sumathy V, Yuvaneswari KC, et al. Allele frequency distribution for 9 STR loci in the Tamil population of Malaysia. *J Forensic Sci*. In Press. 2004.

[PubMed]

[PubMed]

Additional information and reprint requests:

S. Panneerchelvam, M.Sc.
School of Health Sciences
Universiti Sains Malaysia
16150 Kubang Kerian
Kelantan
Malaysia