

Erratum

REFERENCE: Farrell LJ, Kerrigan S, Logan BK. Recommendations for toxicological investigation of drug impaired driving. *J Forensic Sci* 2007;52(5):1214–18.

We have noticed an error in Table 1. The mode for LSD screening survey data should be 0.5 ng/mL and not 0.05 ng/mL as printed.

A corrected Table 1 is provided below.

Laurel J. Farrell, B.A.
Colorado Bureau of Investigation
690 Kipling Steet, Suite 4000
Denver, CO 80215

TABLE 1—Survey data: urine screen and confirmation levels.

Drug/Drug Class	Screen			Confirmation		
	No. of Labs Responding	Cutoff Range (ng/mL)	Mode (ng/mL)	No. of Labs Responding	Cutoff Range (ng/mL)	Mode (ng/mL)
Amphetamines (methamphetamine)	37	20–1100	1000	33	10–1100	50
Barbiturates (secobarbital)	33	5–1000	200	29	5–1000	100
Benzodiazepines (oxazepam)	32	10–300	200	28	5–150	50
Cannabinoids (11-nor-delta-9-carboxy-tetrahydrocannabinol)	37	7–55	50	34	2–50	5
Cocaine metabolite	38	50–500	300	33	15–500	20
Methodone	18	20–2200	300	28	10–300	50
Opiates (morphine)	38	20–2200	300	34	10–500	20
Phencyclidine (PCP)	31	5–200	25	29	5–50	25
Propoxyphene	16	20–2200	300	26	10–300	100
LSD	5	0.5–25	0.5	4	0.1–10	None
Meprobamate	16	5–5000	1000	26	5–5000	500