



ELSEVIER

Journal of Geometry and Physics 32 (2000) 349–363

JOURNAL OF  
GEOMETRY AND  
PHYSICS

## A non-commutative binomial formula

Hjalmar Rosengren<sup>1</sup>

*Centre for Mathematical Sciences, Mathematics (Faculty of Science), University of Lund,  
Box 118, S-221 00 Lund, Sweden*

Received 1 March 1999

---

### Abstract

We prove a binomial formula for variables  $x$  and  $y$  satisfying a quadratic relation  $xy = ax^2 + qyx + by^2$ . Such relations are important in quantum group theory and non-commutative geometry. © 2000 Elsevier Science B.V. All rights reserved.

*Subj. Class.:* Quantum groups

*1991 MSC:* 33D80; 58B30

*Keywords:* Quadratic algebra; Binomial formula; Quantum plane

---

### 1. Introduction

The classical binomial formula is given by

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} y^k x^{n-k}.$$

Since the binomial coefficients are integers, this identity makes sense in an arbitrary ring with 1, provided that  $x$  and  $y$  satisfy the commutation relation  $xy = yx$ .

An important generalization is obtained by considering polynomials in two non-commuting variables  $x$  and  $y$  satisfying

$$xy = qyx, \tag{1}$$

---

<sup>1</sup> *E-mail address:* [hjalmar@maths.lth.se](mailto:hjalmar@maths.lth.se) (H. Rosengren).

where  $q$  is a scalar. In that case one has

$$(x + y)^n = \sum_{k=0}^n \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}} y^k x^{n-k}, \tag{2}$$

where we use the standard notation [5]

$$(a; q)_k = \prod_{i=0}^{k-1} (1 - aq^i).$$

Since the  $q$ -binomial coefficients occurring in (2) are polynomials in  $q$  (the Gaussian polynomials), this identity makes sense for  $x$ ,  $y$ , and  $q$  in an arbitrary ring, assuming also that  $q$  commutes with  $x$  and  $y$ . The  $q$ -binomial formula (2) was first given explicitly by Schützenberger [11].

Recently, Benaoum [2] proved the more general formula

$$(x + y)^n = \sum_{k=0}^n \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}} \prod_{j=0}^{k-1} (1 + [j]_q h) y^k x^{n-k},$$

where

$$[j]_q = 1 + q + \dots + q^{j-1}$$

and where  $x$  and  $y$  satisfy

$$xy = qyx + hy^2 \tag{3}$$

with  $q$  and  $h$  scalars. Assuming that  $q \neq 1$  and  $h \neq q - 1$ , one may rewrite the formula using

$$\prod_{j=0}^{k-1} (1 + [j]_q h) = \left( \frac{1 - q + h}{1 - q} \right)^k (h / (1 - q + h); q)_k.$$

For  $q = 1$  one has the formula [1]

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} \prod_{j=0}^{k-1} (1 + jh) y^k x^{n-k},$$

where

$$xy = yx + hy^2, \tag{4}$$

while the case when  $h = q - 1$  is

$$(x + y)^n = \sum_{k=0}^n \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}} q^{\binom{k}{2}} y^k x^{n-k},$$

where

$$xy = qyx + (q - 1)y^2.$$

Relations such as (1) and (4) play a fundamental role in quantum group theory. Loosely speaking, quantum groups may be constructed as symmetry “groups” of algebras defined by quadratic relations [10]. In the case of the  $q$ -deformed quantum plane (1) this construction gives the standard quantum deformation of  $GL(2)$  [9], while the  $h$ -deformed quantum plane (4) gives the non-standard, or Jordanian, deformation [4,7,12]. From the viewpoint of non-commutative geometry, the  $h$ -deformed plane is in some ways better behaved than the  $q$ -deformed plane [3].

The purpose of this paper is to obtain a binomial formula for variables  $x$  and  $y$  satisfying the more general quadratic relation

$$xy = ax^2 + qyx + by^2. \tag{5}$$

Here  $x$  and  $y$  are elements of some ring with 1, and  $a, q,$  and  $b$  are central elements. For instance one may consider the algebra of polynomials in  $x$  and  $y$  over the complex field, and let  $a, q,$  and  $b$  be scalars. Actually these algebras are all isomorphic to (1) or to (4), though this observation does not simplify our problem.

Note that the elements  $y^i x^j$  need not span the space of polynomials in  $x$  and  $y$ , and thus a binomial formula need not exist in general. As an illustration we try to express  $x^2y$  as a sum of such elements:

$$\begin{aligned} x^2y &= x(ax^2 + qyx + by^2) = ax^3 + qxyx + bxy^2 \\ &= ax^3 + qxyx + b(ax^2 + qyx + by^2)y \\ &= ax^3 + qxyx + abx^2y + qbyxy + b^2y^3. \end{aligned}$$

If, for instance,  $ab = 1$ , the  $x^2y$ -terms will cancel, and there seems to be no such expansion. However, if  $1 - ab$  is invertible we get

$$x^2y = (1 - ab)^{-1}(ax^3 + qxyx + qbyxy + b^2y^3).$$

Using again (5) and simplifying gives

$$x^2y = (1 - ab)^{-1}(a(1 + q)x^3 + q(q + ab)yx^2 + bq(1 + q)y^2x + b^2(1 + q)y^3).$$

In fact, some further computations show that if  $1 - ab$  is invertible, the elements  $(y^i x^j)_{i+j=3}$  span the space of cubic polynomials in  $x$  and  $y$ . Similarly, if both  $1 - ab$  and  $1 - 2ab - abq$  are invertible, the elements  $(y^i x^j)_{i+j=4}$  span the space of quartic polynomials.

To formulate our binomial formula, we define recursively  $\alpha_0 = 0, \beta_0 = 1,$

$$\begin{aligned} \alpha_{n+1} &= q\alpha_n + \beta_n, \\ \beta_{n+1} &= -ab\alpha_n + \beta_n. \end{aligned} \tag{6}$$

The first few cases are

$$\begin{aligned} \alpha_1 &= 1, & \alpha_2 &= 1 + q, & \alpha_3 &= 1 - ab + q + q^2, \\ \beta_1 &= 1, & \beta_2 &= 1 - ab, & \beta_3 &= 1 - 2ab - abq, \end{aligned}$$

where we recognise  $\beta_2$  and  $\beta_3$  from the previous paragraph. We will prove a binomial formula

$$(x + y)^n = \sum_{k=0}^n C_{kn} y^k x^{n-k}$$

under the additional condition that  $\beta_2, \dots, \beta_{n-1}$  are invertible. Assuming that this is the case we define

$$\gamma_k = \frac{\alpha_k}{\beta_k}$$

and introduce the polynomials

$$h_k(x) = \prod_{i=0}^{k-1} (1 + \gamma_i x),$$

the first few cases being

$$h_0(x) = h_1(x) = 1, \quad h_2(x) = 1 + x, \quad h_3(x) = (1 + x) \left( 1 + \frac{1+q}{1-ab} x \right).$$

We are now ready to formulate our main result.

**Theorem 1.1.** *In the notation above, assuming that the elements  $\beta_2, \dots, \beta_{n-1}$  are invertible, the formula*

$$(x + y)^n = \sum_{k=0}^n \frac{h_n(q)}{h_k(q)h_{n-k}(q)} h_k(b)h_{n-k}(a) y^k x^{n-k}$$

holds.

Working in a polynomial algebra, so that  $a$ ,  $q$ , and  $b$  are complex numbers, one may solve the recursion (6). This gives a more explicit version of Theorem 1.1.

**Corollary 1.2.** *For generic values of  $a$ ,  $q$ , and  $b$ ,*

$$\begin{aligned} (x + y)^n &= \sum_{k=0}^n \frac{(Q; Q)_n}{(Q; Q)_k (Q; Q)_{n-k}} \\ &\times \frac{\left( \frac{Q-q+b(1+Q)}{1-qQ+b(1+Q)}; Q \right)_k \left( \frac{Q-q+a(1+Q)}{1-qQ+a(1+Q)}; Q \right)_{n-k}}{\left( \frac{Q-q}{1-qQ}; Q \right)_n} \\ &\times \frac{(1-qQ+b(1+Q))^k (1-qQ+a(1+Q))^{n-k}}{(1-qQ)^n} y^k x^{n-k}, \end{aligned}$$

where  $Q$  satisfies

$$Q + Q^{-1} = \frac{q^2 + 1 - 2ab}{q + ab}. \quad (7)$$

Using the identity

$$(a; Q^{-1})_k = (a^{-1}; Q)_k (-a)^k Q^{-\binom{k}{2}}$$

one may check that the coefficients in Corollary 1.2 are invariant if replacing  $Q$  by  $Q^{-1}$ , and thus are indeed rational functions of  $a, q$ , and  $b$ .

An interesting degenerate case of Corollary 1.2 is when  $Q = 1$ , which happens for  $(q - 1)^2 = 4ab$ . This yields a two-parameter version of the  $h$ -deformed quantum plane (4), with the binomial formula

$$\begin{aligned} (x + y)^n &= \sum_{k=0}^n \binom{n}{k} \frac{\prod_{j=0}^{k-1} (1 + q + (1 - q + 2b)j) \prod_{j=0}^{n-k-1} (1 + q + (1 - q + 2a)j)}{\prod_{j=0}^{n-1} (1 + q + (1 - q)j)} \\ &\quad \times y^k x^{n-k}, \end{aligned} \tag{8}$$

or, if  $q \neq 1, 1 + 2a, 1 + 2b$ ,

$$\begin{aligned} (x + y)^n &= \sum_{k=0}^n \binom{n}{k} \frac{(1 - q + 2b)^k (1 - q + 2a)^{n-k}}{(1 - q)^n} \\ &\quad \times \frac{((1 + q)/(1 - q + 2b))_k ((1 + q)/(1 - q + 2a))_{n-k}}{((1 + q)/(1 - q))_n} y^k x^{n-k}, \end{aligned}$$

where

$$(a)_k = a(a + 1) \cdots (a + k - 1).$$

The plan of the paper is as follows. In Section 2 we prove Theorem 1.1. Our proof is different from the one given for the case  $a = 0$  in [2]. In Section 3 we apply Theorem 1.1 to compute the coefficients in the expansion

$$x^m y^n = \sum_{l=0}^{m+n} C_{lmn} y^l x^{m+n-l},$$

which exists if  $\beta_2, \dots, \beta_{m+n-1}$  are invertible. For generic complex values of  $a, q$ , and  $b$ , the coefficients  $C_{lmn}$  may be expressed as terminating balanced  ${}_4\phi_3$  series [5] of a special type. In Section 4 we derive Corollary 1.2.

## 2. Proof of the binomial formula

In this section we will prove Theorem 1.1. First we observe that if  $x$  and  $y$  satisfy relation (5), and if  $c$  and  $d$  are central invertible elements, then  $cx$  and  $dy$  satisfy a similar relation with  $(a, q, b)$  replaced by  $(ad/c, q, bc/d)$ . Since this does not change  $q$  or  $ab$ , the polynomials  $h_j$  are the same for the two rings. Thus Theorem 1.1 implies

$$(cx + dy)^n = \sum_{k=0}^n \frac{h_n(q)}{h_k(q)h_{n-k}(q)} h_k(bc/d)h_{n-k}(ad/c)d^k c^{n-k} y^k x^{n-k}. \tag{9}$$

With the interpretation

$$d^k h_k(bc/d) = \prod_{i=0}^{k-1} (d + \gamma_i bc) \tag{10}$$

and similarly for  $h_{n-k}$ , this equality makes sense also if  $c$  or  $d$  is not invertible. We will prove Theorem 1.1 by induction on  $n$ , the induction hypothesis being that (9) holds for all central elements  $c$  and  $d$ .

**Lemma 2.1.** *Let  $A_0, B_0, C_0,$  and  $D_0$  be arbitrary central elements, and define recursively*

$$\begin{aligned} A_{n+1} &= A_n + aB_n, \\ B_{n+1} &= -bA_n + qB_n, \\ C_{n+1} &= C_n + aD_n, \\ D_{n+1} &= -bC_n + qD_n. \end{aligned}$$

Then, for all  $n, m \geq 0$ ,

$$(C_n x + D_n y)^m (A_0 x + B_0 y)^n = (A_m x + B_m y)^n (C_0 x + D_0 y)^m.$$

**Proof.** Let  $L_{mn}$  be the statement of the lemma for fixed  $m$  and  $n$ . Then  $L_{m0}$  and  $L_{0n}$  are trivial, and it is also clear that  $L_{mn}$  and  $L_{11}$  together imply  $L_{m+1,n}$  and  $L_{m,n+1}$ . Thus it suffices to prove  $L_{11}$ , that is, the identity

$$(C_1 x + D_1 y)(A_0 x + B_0 y) = (A_1 x + B_1 y)(C_0 x + D_0 y),$$

or equivalently

$$[C_0(x - by) + D_0(ax + qy)](A_0 x + B_0 y) = [A_0(x - by) + B_0(ax + qy)](C_0 x + D_0 y).$$

This follows from the commutation relation (5), written in the form

$$(x - by)y = (ax + qy)x. \quad \square$$

Let us put  $m = 1, A_0 = B_0 = C_0 = 1, D_0 = 0$  in Lemma 2.1. One may check that in this case

$$C_n = \beta_n, \quad D_n = -b\alpha_n,$$

so that the lemma gives

$$(\beta_n x - b\alpha_n y)(x + y)^n = [(a + 1)x + (q - b)y]^n x.$$

Thus, if  $\beta_n$  is invertible,

$$\begin{aligned} (x + y)^{n+1} &= \left( \frac{1}{\beta_n}(\beta_n x - b\alpha_n y) + (1 + b\gamma_n)y \right) (x + y)^n \\ &= \frac{1}{\beta_n} [(a + 1)x + (q - b)y]^n x + (1 + b\gamma_n)y(x + y)^n, \end{aligned}$$

and hence, by the induction hypothesis (9),

$$\begin{aligned}
 (x + y)^{n+1} &= \frac{1}{\beta_n} \sum_{k=0}^n \frac{h_n(q)h_k(b(a + 1)/(q - b))h_{n-k}(a(q - b)/(a + 1))}{h_k(q)h_{n-k}(q)} \\
 &\quad \times (q - b)^k (a + 1)^{n-k} y^k x^{n-k+1} \\
 &\quad + (1 + b\gamma_n) \sum_{k=0}^n \frac{h_n(q)}{h_k(q)h_{n-k}(q)} h_k(b)h_{n-k}(a) y^{k+1} x^{n-k}. \tag{11}
 \end{aligned}$$

The terms in the first sum may be simplified using the following lemma.

**Lemma 2.2.** *In the notation above,*

$$\beta_k h_{k+1}(x) = (1 + x)^k h_k \left( \frac{qx - ab}{1 + x} \right).$$

**Proof.** By induction on  $k$  this reduces to

$$\beta_{k+1} + x\alpha_{k+1} = (1 + x)\beta_k + (qx - ab)\alpha_k,$$

which is equivalent to (6).  $\square$

In particular, choosing  $x = a$  and replacing  $k$  by  $n - k$  gives

$$(a + 1)^{n-k} h_{n-k} \left( \frac{a(q - b)}{a + 1} \right) = \beta_{n-k} h_{n-k+1}(a),$$

while choosing  $x = b(a + 1)/(q - b)$  and replacing  $k$  by  $k - 1$  gives

$$(q - b)^k h_k \left( \frac{b(a + 1)}{q - b} \right) = (q - b) \frac{(q + ab)^{k-1}}{\beta_{k-1}} h_{k-1}(b).$$

Plugging these expressions into (11) we obtain

$$\begin{aligned}
 (x + y)^{n+1} &= h_{n+1}(a)x^{n+1} + h_{n+1}(b)y^{n+1} \\
 &\quad + \sum_{k=1}^n \left( \frac{1}{\beta_n} \frac{h_n(q)}{h_k(q)h_{n-k}(q)} (q - b) \frac{(q + ab)^{k-1}}{\beta_{k-1}} h_{k-1}(b)\beta_{n-k}h_{n-k+1}(a) \right. \\
 &\quad \left. + (1 + b\gamma_n) \frac{h_n(q)}{h_{k-1}(q)h_{n-k+1}(q)} h_{k-1}(b)h_{n-k+1}(a) \right) y^k x^{n-k+1}.
 \end{aligned}$$

Thus, to prove Theorem 1.1, it suffices to prove that the coefficient of  $y^k x^{n-k+1}$  in this sum equals

$$\frac{h_{n+1}(q)}{h_k(q)h_{n-k+1}(q)} h_k(b)h_{n-k+1}(a).$$

Multiplying the corresponding identity with

$$\frac{h_k(q)h_{n-k+1}(q)\beta_n\beta_{k-1}}{h_n(q)h_{n-k+1}(a)h_{k-1}(b)}$$

gives

$$(q - b)(q + ab)^{k-1}(\beta_{n-k} + q\alpha_{n-k}) + (\beta_n + b\alpha_n)(\beta_{k-1} + q\alpha_{k-1}) = (\beta_n + q\alpha_n)(\beta_{k-1} + b\alpha_{k-1}).$$

Writing  $\beta_{n-k} + q\alpha_{n-k} = \alpha_{n-k+1}$  and simplifying gives

$$(q - b)(q + ab)^{k-1}\alpha_{n-k+1} = (q - b)(\alpha_n\beta_{k-1} - \beta_n\alpha_{k-1}).$$

Thus all that remains is to prove the following lemma.

**Lemma 2.3.** For  $i \geq j \geq 0$ ,

$$\alpha_i\beta_j - \alpha_j\beta_i = (q + ab)^j\alpha_{i-j}.$$

**Proof.** Since

$$\begin{pmatrix} \alpha_k \\ \beta_k \end{pmatrix} = A^k \begin{pmatrix} \alpha_0 \\ \beta_0 \end{pmatrix},$$

where

$$A = \begin{pmatrix} q & 1 \\ -ab & 1 \end{pmatrix}, \tag{12}$$

it follows from standard properties of determinants that indeed

$$\det \begin{pmatrix} \alpha_i & \alpha_j \\ \beta_i & \beta_j \end{pmatrix} = \det(A)^j \det \begin{pmatrix} \alpha_{i-j} & \alpha_0 \\ \beta_{i-j} & \beta_0 \end{pmatrix} = (q + ab)^j\alpha_{i-j}. \quad \square$$

### 3. A commutation rule

In this section we will apply our binomial formula to compute the coefficients in the expansion

$$x^m y^n = \sum_{l=0}^{m+n} C_{lmn} y^l x^{m+n-l}.$$

We will see that such an expansion exists if  $\beta_2, \dots, \beta_{m+n-1}$  are invertible. We need Lemma 2.1 when  $A_0 = D_0 = 0, B_0 = C_0 = 1$ . In this case

$$A_n = a\alpha_n, \quad B_n = \delta_n, \quad C_n = \beta_n, \quad D_n = -b\alpha_n,$$

where

$$\delta_n = \alpha_{n+1} - \alpha_n,$$

so that the lemma gives

$$(\beta_n x - b\alpha_n y)^m y^n = (a\alpha_m x + \delta_m y)^n x^m. \tag{13}$$



**Remark 3.1.** One may check that

$$A^n = \begin{pmatrix} \delta_n & \alpha_n \\ -ab\alpha_n & \beta_n \end{pmatrix},$$

where  $A$  is the matrix (12). If  $q + ab$  is invertible this implies

$$A^{-n} = \frac{1}{(q + ab)^n} \begin{pmatrix} \beta_n & -\alpha_n \\ ab\alpha_n & \delta_n \end{pmatrix},$$

so it is natural to define

$$\begin{aligned} \alpha_{-n} &= -(q + ab)^{-n} \alpha_n, \\ \beta_{-n} &= (q + ab)^{-n} \beta_n, \\ \delta_{-n} &= (q + ab)^{-n} \delta_n \end{aligned} \tag{14}$$

for negative integers  $n$ . Moreover,  $A^{m+n} = A^m A^n$  implies

$$\begin{aligned} \alpha_{m+n} &= \delta_m \alpha_n + \alpha_m \beta_n, \\ \beta_{m+n} &= \beta_m \beta_n - ab\alpha_m \alpha_n, \\ \delta_{m+n} &= \delta_m \delta_n - ab\alpha_m \alpha_n. \end{aligned} \tag{15}$$

The next step is to find an expansion of the form

$$x^m = \sum_{k=0}^m D_{kmn} y^k (\beta_n x - b\alpha_n y)^{m-k}.$$

**Lemma 3.2.** If  $c$  and  $d$  are central elements such that  $c\beta_k + ad\alpha_k$  is invertible for  $0 \leq k \leq m - 1$ , then

$$x^m = \sum_{k=0}^m \frac{h_m(q)}{h_k(q)h_{m-k}(q)} \frac{d^k h_k(q - 1 - bc/d)}{c^m h_m(ad/c)} (-1)^k y^k (cx + dy)^{m-k}.$$

This should be interpreted as in (10) if  $c$  or  $d$  is not invertible.

**Proof.** The starting point is the identity

$$-(a + 1)(x + y)y = -a(x + y)^2 - (q - a)y(x + y) + (q - a - b - 1)y^2,$$

which is easily verified using (5). Thus, if  $a + 1$  is invertible, then  $\tilde{x} = x + y$  and  $\tilde{y} = -y$  satisfy relation (5) with  $a, q, b$  replaced by

$$\tilde{a} = -\frac{a}{a + 1}, \quad \tilde{q} = \frac{q - a}{a + 1}, \quad \tilde{b} = \frac{q - a - b - 1}{a + 1}.$$

The corresponding elements  $\tilde{y}_k$  and polynomials  $\tilde{h}_k$  may be computed:

$$\begin{aligned} \tilde{y}_k &= \frac{(1 + a)\alpha_k}{\beta_k + a\alpha_k}, \\ \tilde{h}_k(x) &= \frac{h_k(a + (1 + a)x)}{h_k(a)}. \end{aligned}$$

In particular

$$\tilde{h}_k(\tilde{a}) = \frac{1}{h_k(a)}, \quad \tilde{h}_k(\tilde{q}) = \frac{h_k(q)}{h_k(a)}, \quad \tilde{h}_k(\tilde{b}) = \frac{h_k(q-1-b)}{h_k(a)},$$

so that Theorem 1.1 gives

$$x^m = \sum_{k=0}^m \frac{h_m(q)}{h_k(q)h_{m-k}(q)} \frac{h_k(q-1-b)}{h_m(a)} (-1)^k y^k (x+y)^{m-k}$$

if  $\beta_k + a\alpha_k$  is invertible for  $0 \leq k \leq m-1$ . This proves the lemma for  $c = d = 1$ . The general case follows by replacing  $x, y$  with  $cx, dy$ , as in (9).  $\square$

In particular, when  $c = \beta_n, d = -b\alpha_n$ , (15) gives

$$c\beta_k + ad\alpha_k = \beta_{n+k},$$

and one may also check that

$$q-1-\frac{bc}{d} = q-1+\frac{1}{\gamma_n} = \frac{\delta_n}{\alpha_n}.$$

Thus in this case we obtain

$$x^m = \sum_{k=0}^m \frac{h_m(q)}{h_k(q)h_{m-k}(q)} \frac{(b\alpha_n)^k h_k(\delta_n/\alpha_n)}{\beta_n^m h_m(-ab\gamma_n)} y^k (\beta_n x - b\alpha_n y)^{m-k},$$

if  $\beta_n, \dots, \beta_{m+n-1}$  are invertible. Together with (13) this gives

$$x^m y^n = \sum_{k=0}^m \frac{h_m(q)}{h_k(q)h_{m-k}(q)} \frac{(b\alpha_n)^k h_k(\delta_n/\alpha_n)}{\beta_n^m h_m(-ab\gamma_n)} y^k (a\alpha_{m-k} x + \delta_{m-k} y)^n x^{m-k},$$

and, by (9),

$$\begin{aligned} x^m y^n &= \sum_{k=0}^m \sum_{j=0}^n \frac{h_m(q)}{h_k(q)h_{m-k}(q)} \frac{(b\alpha_n)^k h_k(\delta_n/\alpha_n)}{\beta_n^m h_m(-ab\gamma_n)} \frac{h_n(q)}{h_j(q)h_{n-j}(q)} \\ &\quad \times \delta_{m-k}^j h_j \left( ab \frac{\alpha_{m-k}}{\delta_{m-k}} \right) (a\alpha_{m-k})^{n-j} h_{n-j} \left( \frac{\delta_{m-k}}{\alpha_{m-k}} \right) y^{k+j} x^{m+n-k-j}, \end{aligned}$$

assuming that also  $\beta_2, \dots, \beta_{n-1}$  are invertible.

Thus we have proved the following theorem.

**Theorem 3.3.** *If  $\beta_2, \dots, \beta_{m+n-1}$  are invertible, there is an expansion*

$$x^m y^n = \sum_{l=0}^{m+n} C_{lmn} y^l x^{m+n-l}, \tag{16}$$

where the coefficient  $C_{lmn}$  equals

$$\sum_{\substack{j+k=l \\ 0 \leq j \leq n \\ 0 \leq k \leq m}} \frac{h_m(q)h_n(q)\delta_{m-k}^j h_j \left( ab \frac{\alpha_{m-k}}{\delta_{m-k}} \right) (b\alpha_n)^k h_k \left( \frac{\delta_n}{\alpha_n} \right) (a\alpha_{m-k})^{n-j} h_{n-j} \left( \frac{\delta_{m-k}}{\alpha_{m-k}} \right)}{\beta_n^m h_m(-ab\gamma_n)h_j(q)h_k(q)h_{n-j}(q)h_{m-k}(q)}. \tag{17}$$

**Remark 3.4.** The polynomials  $h_k$  appearing in Theorem 3.3 admit simple expressions in terms of the elements  $\alpha_k, \beta_k, \delta_k$ . From the first identity in (15) it follows that

$$\alpha_n^k h_k \left( \frac{\delta_n}{\alpha_n} \right) = \prod_{j=0}^{k-1} \frac{\alpha_{n+j}}{\beta_j};$$

in particular,

$$h_k(q) = \prod_{j=0}^{k-1} \frac{\alpha_{j+1}}{\beta_j}.$$

Similarly the second identity in (15) gives

$$\beta_n^k h_k(-ab\gamma_n) = \prod_{j=0}^{k-1} \frac{\beta_{n+j}}{\beta_j}.$$

By an extension of (15) to negative indices,

$$\beta_k \delta_n + ab\alpha_k \alpha_n = (q + ab)^n \beta_{k-n} = (q + ab)^k \delta_{n-k},$$

which leads to

$$\delta_n^k h_k \left( ab \frac{\alpha_n}{\delta_n} \right) = (q + ab)^{nk} \prod_{j=0}^{k-1} \frac{\beta_{j-n}}{\beta_j} = (q + ab)^{\binom{k}{2}} \prod_{j=0}^{k-1} \frac{\delta_{n-j}}{\beta_j},$$

where we must use (14) if some indices are negative.

**Remark 3.5.** The coefficients  $C_{lmn}$  are very simple if  $a$  or  $b$  is zero. Let us put  $a = 0, b = h$ , so that we consider relation (3). Then, in (17), only the term with  $j = n$  is non-zero, and such a term exists only if  $l \geq n$  in (16). Moreover, one may check that  $\alpha_n = [n]_q, \beta_n = 1, \delta_n = q^n$  and that

$$\alpha_n^k h_k \left( \frac{\delta_n}{\alpha_n} \right) = \frac{(q^n; q)_k}{(1 - q)^k};$$

in particular,

$$h_k(q) = \frac{(q; q)_k}{(1 - q)^k}.$$

Plugging this into Theorem 3.3 gives

$$x^m y^n = \sum_{l=n}^{m+n} \frac{(q; q)_m (q^n; q)_{l-n}}{(q; q)_{l-n} (q; q)_{m+n-l}} (1-q)^{n-l} q^{n(m+n-l)} h^{l-n} y^l x^{m+n-l},$$

where  $xy = qyx + hy^2$ . The case  $m = 1$  of this identity was used in [2].

**Remark 3.6.** Working in a polynomial algebra over the complex numbers, one may solve the recursion (6) explicitly; cf. Section 4. For generic values of the parameters, the coefficients  $C_{lmn}$  may then be written as terminating balanced  ${}_4\phi_3$  series [5]; for instance, when  $l \leq m$ ,

$$C_{lmn} = b^l a^n \left( \frac{1+Q}{1-qQ} \right)^{l+n} \frac{(Q; Q)_m (Q^n; Q)_l (Q^{m-l}; Q)_n}{(Q; Q)_l (Q; Q)_{m-l} (Q^{m-l} (Q-q) / (1-qQ); Q)_{l+n}} \\ \times {}_4\phi_3 \left[ \begin{matrix} Q^{-l}, Q^{-n}, Q^{m-l} \frac{(Q-q)}{(1-qQ)}, Q^{l+m-l} \frac{(1-qQ)}{(Q-q)} \\ Q^{1-n-l}, Q^{m-l}, Q^{l+m-l} \end{matrix} ; Q, Q \right],$$

where  $Q$  satisfies (7). This kind of sums have very rich symmetry properties [5].

#### 4. Explicit form of the binomial formula

In this section we work in the algebra of polynomials in  $x, y$ , over the complex field, with the relation (5), where  $a, q$ , and  $b$  are scalars. Let us first assume that the matrix

$$A = \begin{pmatrix} q & 1 \\ -ab & 1 \end{pmatrix}$$

is diagonalizable with different eigenvalues  $\lambda_1$  and  $\lambda_2$ . Then  $\alpha_n$  and  $\beta_n$  are linear combinations of  $\lambda_1^n$  and  $\lambda_2^n$ . Since  $\alpha_0 = 0, \alpha_1 = 1$ , we have

$$\alpha_n = \frac{\lambda_1^n - \lambda_2^n}{\lambda_1 - \lambda_2}, \tag{18}$$

where

$$\lambda_1 + \lambda_2 = 1 + q, \quad \lambda_1 \lambda_2 = q + ab.$$

**Remark 4.1.** At this point we may find explicit expressions for  $\alpha_n$  and  $\beta_n$  as polynomials in  $ab$  and  $q$ . Using the symmetric function identity [8]

$$\frac{\lambda_1^{n+1} - \lambda_2^{n+1}}{\lambda_1 - \lambda_2} = \sum_{k=0}^n \lambda_1^k \lambda_2^{n-k} = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n-k}{k} (-1)^k (\lambda_1 \lambda_2)^k (\lambda_1 + \lambda_2)^{n-k} \tag{19}$$

we find that

$$\alpha_n = \sum_{k=0}^{[(n-1)/2]} \binom{n-k-1}{k} (-1)^k (q+ab)^k (1+q)^{n-k-1}.$$

One may then express  $\beta_n$  as a polynomial by

$$\beta_n = \alpha_{n+1} - q\alpha_n.$$

The identity (19) is easily proved using generating functions; write

$$\frac{1}{(1-\lambda_1 t)(1-\lambda_2 t)} = \frac{1}{1-t(\lambda_1+\lambda_2)+t^2\lambda_1\lambda_2}$$

and identify the coefficient of  $t^n$ .

Now let  $Q = \lambda_2/\lambda_1$ . Then  $Q$  satisfies

$$Q + Q^{-1} = \frac{(\lambda_1 + \lambda_2)^2 - 2\lambda_1\lambda_2}{\lambda_1\lambda_2} = \frac{(1+q)^2 - 2(q+ab)}{q+ab} = \frac{1+q^2-2ab}{q+ab},$$

that is, Eq. (7). We may write

$$\lambda_1 = \lambda_1 \frac{1+q}{\lambda_1 + \lambda_2} = \frac{1+q}{1+Q},$$

$$\lambda_2 = Q\lambda_1 = Q \frac{1+q}{1+Q}.$$

Plugging this into (18) gives

$$\alpha_n = \left(\frac{1+q}{1+Q}\right)^{n-1} \frac{1-Q^n}{1-Q},$$

so that

$$\beta_n = \alpha_{n+1} - q\alpha_n = \frac{(1+q)^{n-1}(1-qQ - (Q-q)Q^n)}{(1+Q)^n(1-Q)},$$

$$\gamma_n = \frac{\alpha_n}{\beta_n} = \frac{(1-Q^n)(1+Q)}{1-qQ - (Q-q)Q^n},$$

$$\delta_n = \alpha_{n+1} - \alpha_n = \frac{(1+q)^{n-1}(q-Q + (1-qQ)Q^n)}{(1+Q)^n(1-Q)}.$$

If  $1-qQ, 1-qQ+x(1+Q) \neq 0$ , one may then write

$$\begin{aligned} h_n(x) &= \prod_{i=0}^{n-1} (1+x\gamma_i) = \prod_{i=0}^{n-1} \frac{1-qQ - (Q-q)Q^i + x(1-Q^i)(1+Q)}{1-qQ - (Q-q)Q^i} \\ &= \prod_{i=0}^{n-1} \frac{1-qQ+x(1+Q) - (Q-q+x(1+Q))Q^i}{1-qQ - (Q-q)Q^i} \end{aligned}$$

$$= \left( \frac{1 - qQ + x(1 + Q)}{1 - qQ} \right)^n \frac{(Q - q + x(1 + Q)/1 - qQ + x(1 + Q); Q)_n}{((Q - q)/(1 - qQ); Q)_n}.$$

Inserting this expression into Theorem 1.1 gives Corollary 1.2.

As was remarked in the introduction, the case  $(q - 1)^2 = 4ab$ ,  $Q = 1$  is of special interest. This corresponds to the matrix  $A$  not being diagonalizable. In this case, both eigenvalues equal  $\lambda = (1 + q)/2$ , and one has

$$\begin{aligned} \alpha_n &= \lambda^{n-1} n, \\ \beta_n &= \lambda^n (n + 1) - q \lambda^{n-1} n, \\ \gamma_n &= \frac{2n}{(1 - q)n + q + 1}, \\ h_n(x) &= \prod_{i=0}^{n-1} (1 + x\gamma_i) = \prod_{i=0}^{n-1} \frac{1 + q + (1 - q + 2x)i}{1 + q + (1 - q)i}. \end{aligned}$$

This yields the binomial formula (8).

Finally we make a remark on formal power series identities. The  $q$ -binomial formula (2) is equivalent to the identity

$$e_q(x + y) = e_q(y) e_q(x), \quad xy = qyx,$$

where

$$e_q(x) = \sum_{n=0}^{\infty} \frac{x^n}{(q; q)_n}.$$

A generalization to relation (5) may be obtained from Corollary 1.2 by multiplying with

$$\frac{((Q - q)/(1 - qQ); Q)_n}{(Q; Q)_n}$$

and then summing over  $n$ ; namely,

$$\begin{aligned} & {}_1\phi_0 \left[ \frac{Q - q}{1 - qQ}; Q, x + y \right] \\ &= {}_1\phi_0 \left[ \frac{Q - q + b(1 + Q)}{1 - qQ + b(1 + Q)}; Q, \frac{1 - qQ + b(1 + Q)}{1 - qQ} y \right] \\ &\quad \times {}_1\phi_0 \left[ \frac{Q - q + a(1 + Q)}{1 - qQ + a(1 + Q)}; Q, \frac{1 - qQ + a(1 + Q)}{1 - qQ} x \right], \end{aligned}$$

where

$${}_1\phi_0 [a; Q, x] = \sum_{n=0}^{\infty} \frac{(a; Q)_n}{(Q; Q)_n} x^n.$$

It would be interesting to find generalizations of other formal power series identities such as

$$e_q(x + y - yx) = e_q(x) e_q(y), \quad xy = qyx;$$

cf. [6] for this and other more general identities.

## References

- [1] H.B. Benaoum,  $h$  Analogue of Newton's binomial formula, *J. Phys. A* 31 (1998) L751–L754.
- [2] H.B. Benaoum,  $(q, h)$ -Analogue of Newton's binomial formula, *J. Phys. A* 32 (1999) 2037–2040.
- [3] S. Cho, J. Madore, K.S. Park, Noncommutative geometry of the  $h$ -deformed quantum plane, *J. Phys. A* 31 (1998) 2639–2654.
- [4] E.E. Demidov, Yu.I. Manin, E.E. Mukhin, D.V. Zhdanovich, Nonstandard quantum deformations of  $GL(n)$  and constant solutions of the Yang–Baxter equation, *Progr. Theoret. Phys. Suppl.* 102 (1990) 203–218.
- [5] G. Gasper, M. Rahman, *Basic Hypergeometric Series*, Cambridge University Press, Cambridge, 1990.
- [6] T.H. Koornwinder, Special functions and  $q$ -commuting variables, in: *Special functions,  $q$ -Series and related topics*, *Fields Inst. Commun.* 14, Am. Math. Soc., Providence (1997) 131–166.
- [7] B.A. Kupershmidt, The quantum group  $GL_h(2)$ , *J. Phys. A* 25 (1992) L1239–L1244.
- [8] P.A. MacMahon, *Combinatory Analysis*, vol. 1, Cambridge, 1915.
- [9] Yu.I. Manin, Some remarks on Koszul algebras and quantum groups, *Ann. Inst. Fourier (Grenoble)* 37 (1987) 191–205.
- [10] Yu.I. Manin, *Quantum Groups and Noncommutative Geometry*, Centre de Recherches Mathématiques, Montréal, 1988.
- [11] M.P. Schützenberger, Une interprétation de certaines solutions de l'équation fonctionnelle:  $F(x + y) = F(x)F(y)$ , *C. R. Acad. Sci. Paris* 236 (1953) 352–353.
- [12] S. Zakrzewski, A Hopf star-algebra of polynomials on the quantum  $SL(2, R)$  for a “unitary”  $R$ -matrix, *Lett. Math. Phys.* 22 (1991) 287–289.