



ELSEVIER

Journal of Geometry and Physics 42 (2002) 12–18

JOURNAL OF  
GEOMETRY AND  
PHYSICS

www.elsevier.com/locate/jgp

# Traces for star products on symplectic manifolds<sup>☆</sup>

Simone Gutt<sup>a,b,\*</sup>, John Rawnsley<sup>c</sup>

<sup>a</sup> *Dépt. de Mathématiques, Université Libre de Bruxelles, Campus Plaine, CP 218, BE-1050 Brussels, Belgium*

<sup>b</sup> *Université de Metz, Ile du Saulcy, 57045 Metz, Cedex 01, France*

<sup>c</sup> *Mathematics Institute, University of Warwick, Coventry CV4 7AL, UK*

Received 24 May 2001

---

## Abstract

We give a direct elementary proof of the existence of traces for arbitrary star products on a symplectic manifold. We follow the approach we used in [J. Geom. Phys. 29 (1999) 347], solving first the local problem. A normalisation introduced by Karabegov [Lett. Math. Phys. 45 (1998) 217] makes the local solutions unique and allows them to be pieced together to solve the global problem. © 2002 Elsevier Science B.V. All rights reserved.

MSC: 53D55

Subj. Class.: Differential geometry

Keywords: Star products; Symplectic manifolds

---

## 1. Introduction

In the previous paper [9], we gave lowbrow proofs of some properties of differential star products on symplectic manifolds (in particular, the classification of equivalence classes of such star products) using Čech cohomology methods and the existence of special local derivations which we called  $\nu$ -Euler derivations in [9]. In this note, we present, in a similar spirit, properties of existence and uniqueness of traces for such star products. Our proof of the existence of a trace relies on a canonical way of normalisation of the trace introduced by Karabegov [10], using local  $\nu$ -Euler derivations.

---

<sup>☆</sup> This research was partially supported by an Action de Recherche Concertée de la Communauté française de Belgique.

\* Corresponding author. Present address: Dépt. de Mathématiques, Université Libre de Bruxelles, Campus Plaine, CP 218, BE-1050 Brussels, Belgium.

E-mail addresses: sgutt@ulb.ac.be (S. Gutt), j.rawnsley@warwick.ac.uk (J. Rawnsley).

Let  $*$  be a star product (which we always assume here to be defined by bidifferential operators) on a symplectic manifold  $(M, \omega)$ . In the algebra of smooth functions on  $M$ , consider the ideal  $C_0^\infty(M)$  of compactly supported functions. A *trace* is a  $\mathbb{C}[[\nu]]$ -linear map  $\tau: C_0^\infty(M)[[\nu]] \rightarrow \mathbb{C}[\nu^{-1}, \nu]$  satisfying

$$\tau(u * v) = \tau(v * u).$$

The question of existence and uniqueness of such traces has been solved by the following result.

**Theorem 1.1** (Fedosov [4,5]; Nest and Tsygan [11]). *Any star product on a symplectic manifold  $(M, \omega)$  has a trace which is unique up to multiplication by an element of  $\mathbb{C}[\nu^{-1}, \nu]$ . Every trace is given by a smooth density  $\rho \in C^\infty(M)[\nu^{-1}, \nu]$ :*

$$\tau(u) = \int_M u \rho \frac{\omega^n}{\nu^n n!}.$$

We shall give here an elementary proof of this theorem. The methods use intrinsically that we have a symplectic manifold. For Poisson manifolds, Felder and Shoikhet [6] have shown that the Kontsevich star product also has a trace.

## 2. Traces

Let  $(M, \omega)$  be a connected symplectic manifold,  $N$  the algebra of smooth functions and  $N_c$  the ideal in  $N$  of compactly supported functions. Obviously,  $N_c[[\nu]]$  is an ideal in  $N[[\nu]]$  and any differential star product or equivalence on  $N[[\nu]]$  is determined on  $N_c[[\nu]]$ .

**Definition 2.1.** Let  $*$  be a star product on  $(M, \omega)$  then a *trace* is a  $\mathbb{C}[[\nu]]$ -linear map  $\tau: N_c[[\nu]] \rightarrow \mathbb{C}[\nu^{-1}, \nu]$  satisfying

$$\tau(u * v) = \tau(v * u).$$

**Remark 2.2.** Since any  $\mathbb{C}[\nu^{-1}, \nu]$ -multiple of a trace is a trace, it is not necessary to work with Laurent series, but we do so for two reasons. Firstly, in the formula for the trace of a pseudo-differential operator, a factor of  $\nu^{-n}$  occurs, and secondly, the presence of such a factor simplifies Eq. (3) below.

If  $\tau$  is a non-trivial trace, we take  $u$  in  $N_c$  and can then expand

$$\tau(u) = \nu^r \sum_{s \geq 0} \nu^s \tau_s(u),$$

where each  $\tau_s: N_c \rightarrow \mathbb{R}$  is a linear map and we assume  $\tau_0 \neq 0$ . The condition to be a trace takes the form

$$\tau_k(\{u, v\}) + \tau_{k-1}(C_2^-(u, v)) + \dots + \tau_0(C_{k+1}^-(u, v)) = 0 \tag{1}$$

for  $k = 0, 1, 2, \dots$ , where  $C_r^-$  denotes the antisymmetric part of  $C_r$ .

**Remark 2.3.** In [3], the notion of a *closed star product* was introduced and related to cyclic cohomology of the algebra of functions. The existence of a closed star product was proved in [12]; see also [1,13]. A star product is closed if

$$\int_M C_r^-(u, v)\omega^n = 0 \quad \forall u, v \in N_c \tag{2}$$

for all  $1 \leq r \leq n$ , where  $2n$  is the dimension of  $M$ . If (2) holds for all  $r \geq 1$  then  $u * v$  is said to be *strongly closed*. Thus to be strongly closed is the same as requiring that

$$\tau(u) = \int_M u\omega^n$$

be a trace on  $N_c[[\hbar]]$ .

For  $k = 0$ , Eq. (1) reduces to the condition  $\tau_0(\{u, v\}) = 0$  for all  $u, v$ . In [2], it is shown by elementary means that this implies  $\tau_0$  is a multiple of the integral  $\int_M u\omega^n$  when  $M$  is connected. We give a proof here for completeness.

**Lemma 2.4** (Gelfand and Shilov [7]). *If  $u$  is a compactly supported smooth function on  $\mathbb{R}^N$  with  $\int_{\mathbb{R}^N} u \, d^N x = 0$ , then  $u$  is a sum of derivatives of compactly supported smooth functions.*

**Proof.** For  $N = 1$ , we simply set  $v(x) = \int_{-\infty}^x u(t) \, dt$  and observe that  $v$  obviously vanishes when  $x$  is below the support of  $u$ , and is zero again when  $x$  is large as a consequence of  $\int_{-\infty}^{\infty} u(t) \, dt = 0$ . Thus  $v$  is compactly supported and  $u(x) = dv/dx$ . Moreover, if  $u$  depends smoothly on parameters, so will  $v$  and if  $u$  is compactly supported in the parameters, so is  $v$ .

Now proceed by induction on  $N$ . Clearly  $w(x_1, \dots, x_{N-1}) = \int_{-\infty}^{\infty} u(x_1, \dots, x_{N-1}, t) \, dt$  is compactly supported in  $\mathbb{R}^{N-1}$  and has vanishing integral, so by the inductive assumption  $w(x_1, \dots, x_{N-1}) = \sum_{i=1}^{N-1} (\partial w_i(x_1, \dots, x_{N-1})/\partial x_i)$  for some compactly supported functions  $w_i$  on  $\mathbb{R}^{N-1}$ . Take a compactly supported bump function  $r(t)$  on  $\mathbb{R}$  with  $\int_{\mathbb{R}} r(t) \, dt = 1$  and consider  $u(x_1, \dots, x_N) - w(x_1, \dots, x_{N-1})r(x_N)$ , which is compactly supported in all its variables. Integrating in  $x_N$ , we see that the integral over  $\mathbb{R}$  vanishes, and so  $u(x_1, \dots, x_N) - w(x_1, \dots, x_{N-1})r(x_N) = (\partial v/\partial x_N)(x_1, \dots, x_N)$ . Thus,

$$u = \frac{\partial v}{\partial x_N}(x_1, \dots, x_N) + \sum_{i=1}^{N-1} \frac{\partial w_i(x_1, \dots, x_{N-1})r(x_N)}{\partial x_i},$$

which completes the inductive step. □

**Lemma 2.5** (Bordemann et al. [2]). *Let  $(M, \omega)$  be a connected symplectic manifold. If  $\sigma: N_c \rightarrow \mathbb{R}$  is a linear map with  $\sigma(\{u, v\}) = 0$  for all  $u, v \in N_c$ , then  $\sigma(u) = c \int_M u\omega^n$  for some constant  $c$ .*

**Proof.** Fix  $u \in N_c$  and cover  $M$  by Darboux charts  $U_\alpha$ , such that only finitely many  $U_\alpha$  intersect the support of  $u$ . Take a partition of unity  $\varphi_\alpha$  subordinate to  $U_\alpha$  then only a finite number of  $u\varphi_\alpha$  are non-zero. Thus,  $\sigma(u) = \sum_\alpha \sigma(u\varphi_\alpha)$ .

$u\varphi_\alpha - (\int_M u\varphi_\alpha\omega^n / \int_M \varphi_\alpha\omega^n)\varphi_\alpha$  has vanishing integral on  $U_\alpha$  which can be viewed as an open set in some  $\mathbb{R}^{2n}$ . Thus, by the previous lemma, there are functions  $v_i, w_i$  with compact support, such that

$$u\varphi_\alpha - \frac{\int_M u\varphi_\alpha\omega^n}{\int_M \varphi_\alpha\omega^n}\varphi_\alpha = \sum_i \frac{\partial v_i}{\partial p_i} + \frac{\partial w_i}{\partial q_i}$$

for the Darboux coordinates  $p_i$  and  $q_i$ . But  $\partial v_i / \partial p_i = \{v_i, q_i\}$ , and if we choose a function  $s_i$  of compact support which is identically 1 on the support of  $v_i$  then  $\partial v_i / \partial p_i = \{v_i, s_i q_i\}$  so we see that  $\sigma(\partial v_i / \partial p_i) = 0$  and similarly  $\sigma(\partial w_i / \partial q_i) = 0$ . Thus,  $\sigma(u\varphi_\alpha) = (\int_M u\varphi_\alpha\omega^n / \int_M \varphi_\alpha\omega^n)\sigma(\varphi_\alpha)$ . Hence

$$\sigma(u) = \int_M u \sum_\alpha \frac{\varphi_\alpha}{\int_M \varphi_\alpha\omega^n} \sigma(\varphi_\alpha)\omega^n = \int_M u\rho\omega^n.$$

Since  $\sigma(\{u, v\}) = 0$ ,  $\int_M \{u, v\}\rho\omega^n = 0$  and hence  $\int_M \{\rho, u\}v\omega^n = 0$  for all  $v$ . Thus  $\{\rho, u\} = 0$  for all  $u$  and hence  $\rho = c$ , a constant.  $\square$

Thus, any trace has the form

$$\tau(u) = av^r \left( \int_M u\omega^n + \sum_{k \geq 1} v^k \tau_k(u) \right),$$

where  $a \neq 0$ . We can divide by  $a$  and multiply by  $v^{-n-r}$  to bring  $\tau$  into the form

$$\int_M u \frac{\omega^n}{v^n} + v^{-n} \sum_{k \geq 1} v^k \tau_k(u).$$

Any trace in this form will be said to be *standard* cf. [8].

As observed in [2], this is enough to show that any two traces  $\tau$  and  $\tau'$  for the same star product are proportional. For, if  $\tau$  is standard, the leading term of  $\tau'$  is a multiple  $cv^r$  of the integral, hence is equal to the leading term of  $\tau$  multiplied by  $cv^{r+n}$ . But then  $\tau' - cv^{r+n}\tau$  is a trace which vanishes to at least  $r + 1$  order in  $v$ . This argument can be repeated indefinitely to show that  $\tau' = cv^{r+n}(1 + \sum_k v^k c_k)\tau$ . Remark that, in particular, if  $\tau$  and  $\tau'$  are standard then  $\tau' = (1 + \sum_k v^k c_k)\tau$ . This proves the following theorem.

**Theorem 2.6** (Nest and Tsygan [11]). *On a connected symplectic manifold  $(M, \omega)$ , any two traces are proportional by an element of  $\mathbb{C}[v^{-1}, v]$ .*

### 3. The local case

In the case of  $\mathbb{R}^{2n}$  with its standard constant 2-form  $\Omega$ , then

$$\tau_M(u) = \int_{\mathbb{R}^{2n}} u \frac{\Omega^n}{v^n n!}$$

is a trace on compactly supported functions for the Moyal star product  $*_M$ . The Moyal star product and this trace have an important homogeneity property [10]. If we take a conformal

vector field  $\xi$  on  $\mathbb{R}^{2n}$ , so  $\mathcal{L}_\xi \Omega = \Omega$  then,  $D_M = \xi + \nu(\partial/\partial\nu)$  is a derivation of  $*_M$  and  $\tau_M$  satisfies

$$\tau_M(D_M u) = \nu \frac{\partial}{\partial \nu} \tau_M(u).$$

If  $*$  is any star product defined on an open ball  $U$  in  $\mathbb{R}^{2n}$ , then it is equivalent to the restriction of the Moyal star product by a map  $T = \text{Id} + \sum_{k \geq 1} \nu^k T_k$  with

$$T(u * v) = T(u) *_M T(v),$$

and then we see that

$$\tau(u) = \int_U T(u) \frac{\Omega^n}{\nu^n}$$

is a trace for  $*$ . Each  $T_k$  is a differential operator, so it has a formal adjoint  $T'_k$  so that, if we put  $T' = \text{Id} + \sum_{k \geq 1} \nu^k T'_k$ , then

$$\tau(u) = \int_U u T'(1) \frac{\Omega^n}{\nu^n}$$

for  $u \in C^\infty(U)$ . If we put  $\rho = T'(1) \in C^\infty(U)[\nu^{-1}, \nu]$ , then

$$\tau(u) = \int_U u \rho \frac{\Omega^n}{\nu^n}.$$

If  $\tau$  is standard, then  $\rho = 1 + \sum_{k \geq 1} \nu^k \rho_k$ .

Further  $D = T^{-1} \circ D_M \circ T$  will be a derivation of  $*$  and satisfies the transform of Eq. (3):

$$\tau(Du) = \nu \frac{\partial}{\partial \nu} \tau(u). \tag{3}$$

Here  $D$  has the form  $\xi + \nu(\partial/\partial\nu) + D'$ . Local derivations of this form we give a special name.

**Definition 3.1.** Let  $(M, \omega)$  be a symplectic manifold. Say that a derivation  $D$  on an open set  $U$  of  $N[[\nu]]$ ,  $*$  is  $\nu$ -Euler if it has the form

$$D = \nu \frac{\partial}{\partial \nu} + X + D', \tag{4}$$

where  $X$  is conformally symplectic ( $\mathcal{L}_X \omega = \omega$ ) and  $D' = \sum_{r \geq 1} \nu^r D'_r$  with the  $D'_r$  differential operators on  $U$ .

Note that conformally symplectic vector fields only exist locally in general, so we also cannot ask for global  $\nu$ -Euler derivations. Two local  $\nu$ -Euler derivations defined on the same open set will differ by a  $\nu$ -linear derivation, and so the difference is  $\nu^{-1} \times$  inner. This means that  $\tau \circ D$  will be independent of  $D$  and thus is globally defined as an  $\mathbb{R}$ -linear functional, even if  $D$  is not.

**Definition 3.2.** A standard trace  $\tau$  is *normalised* if it satisfies the analogue of the Moyal homogeneity condition

$$\tau(Du) = v \frac{\partial}{\partial v} \tau(u)$$

on any open set where there are  $v$ -Euler derivations and for any such local  $v$ -Euler derivation  $D$ .

This condition was introduced by Karabegov [10].

It is clear that the pull-back of the Moyal trace by an equivalence with the Moyal star product is normalised, so a normalised trace always exists for any star product on an open ball in  $\mathbb{R}^{2n}$ .

**Proposition 3.3** (Karabegov [10]). *If  $\tau$  and  $\tau'$  are normalised traces for the same star product on an open ball  $U$  in  $\mathbb{R}^{2n}$ , then  $\tau = \tau'$ .*

**Proof.** Neither trace can be zero, and are proportional so  $\tau' = (1 + cv^r + \dots)\tau$ . If  $\tau' \neq \tau$ , then there is a first  $r > 0$  where  $c \neq 0$ . Then we substitute in the normalisation condition to give

$$\begin{aligned} (1 + cv^r + \dots)\tau(Du) &= \tau'(Du) = v \frac{\partial}{\partial v} \tau'(u) = v \frac{\partial}{\partial v} ((1 + cv^r + \dots)\tau(u)) \\ &= (rcv^{r-1} + \dots)\tau(u) + (1 + cv^r + \dots)\tau(Du), \end{aligned}$$

which implies that  $c = 0$ . This contradiction shows that  $\tau' = \tau$ . □

#### 4. The global case

Let  $(M, \omega)$  be a connected symplectic manifold. Then we can cover  $M$  by Darboux charts  $U$  which are diffeomorphic to open balls in  $\mathbb{R}^{2n}$  and such that all non-empty intersections are also diffeomorphic to open balls. Let  $*$  be a star product on  $M$  and then the restriction of  $*$  to  $U$  has a normalised trace with density  $\rho_U \in C^\infty(U)[[v]]$ . If we have two open sets  $U, V$  which overlap, then on the intersection both  $\rho_U$  and  $\rho_V$  will determine normalised traces on an open ball in  $\mathbb{R}^{2n}$ . Since there is only one such trace,  $\rho_U = \rho_V$  on  $U \cap V$ . It follows that there is a globally defined function  $\rho$  on  $M$ , such that  $\rho_U = \rho|_U$ . Set

$$\tau(u) = \int_M u \rho \frac{\omega^n}{v^n}$$

for  $u \in C_c^\infty(M)$ .

Given  $u, v$  in  $C_c^\infty(M)$ , we can find a finite partition of unity  $\varphi_i$  on  $\text{supp } u \cup \text{supp } v$  with supports in the open sets above. Then  $u = \sum_i \varphi_i u$  and  $v = \sum_j \varphi_j v$ , so  $u * v = \sum_{i,j} (\varphi_i u) * (\varphi_j v)$ , so  $\tau(u * v) = \sum_{i,j} \tau((\varphi_i u) * (\varphi_j v)) = \sum_{i,j} \tau((\varphi_j v) * (\varphi_i u)) = \tau(v * u)$ , so  $\tau$  is a trace. A similar partition of unity argument shows that  $\tau$  is normalised.

Combining this with Theorem 2.6 and the fact that the normalised trace we just constructed has a smooth density, we obtain the following theorem.

**Theorem 4.1** (Fedosov, Nest and Tsygan). *On a connected symplectic manifold  $(M, \omega)$ , any differential star product has a unique normalised trace. Any trace is multiple of this and is given by a smooth density.*

**Remark 4.2.** In [11], a proof that any trace has a smooth density is given using cyclic cohomology.

**Corollary 4.3.** *Any trace is invariant under all  $\mathbb{C}[[\nu]]$ -linear automorphisms of the star product.*

**Proof.** A smooth trace is a multiple (in  $\mathbb{C}[\nu^{-1}, \nu]$ ) of a normalised trace. The transform of a  $\nu$ -Euler derivation by a  $\mathbb{C}[[\nu]]$ -linear automorphism is again a  $\nu$ -Euler derivation, thus the transform of a normalised trace by a  $\mathbb{C}[[\nu]]$ -linear automorphism is again a normalised trace, and so is equal to the original normalised trace.  $\square$

## Acknowledgements

We are grateful to Martin Bordemann, Alexander Karabegov and Stefan Waldmann for their helpful comments.

## References

- [1] M. Bordemann, N. Neumaier, S. Waldmann, Homogeneous Fedosov star products on cotangent bundles. II. GNS representations, the WKB expansion, traces, and applications, *J. Geom. Phys.* 29 (1999) 199–234.
- [2] M. Bordemann, H. Romer, S. Waldmann, A remark on formal KMS states in deformation quantization, *Lett. Math. Phys.* 45 (1998) 49–61.
- [3] A. Connes, M. Flato, D. Sternheimer, Closed star products and cyclic cohomology, *Lett. Math. Phys.* 24 (1992) 1–12.
- [4] B.V. Fedosov, Quantization and the index, *Dokl. Akad. Nauk. SSSR* 291 (1986) 82–86.
- [5] B.V. Fedosov, Deformation quantization and index theory, *Mathematical Topics*, Vol. 9, Akademie Verlag, Berlin, 1996.
- [6] G. Felder, B. Shoikhet, Deformation quantization with traces. math. QA/0002057.
- [7] I.M. Gelfand, G.E. Shilov, *Generalized Functions*, Academic Press, New York, 1964.
- [8] V. Guillemin, Star products on compact pre-quantizable symplectic manifolds, *Lett. Math. Phys.* 35 (1995) 85–89.
- [9] S. Gutt, J. Rawnsley, Equivalence of star products on a symplectic manifold, *J. Geom. Phys.* 29 (1999) 347–392.
- [10] A. Karabegov, On the canonical normalization of a trace density of deformation quantization, *Lett. Math. Phys.* 45 (1998) 217–228.
- [11] R. Nest, B. Tsygan, Algebraic index theorem, *Commun. Math. Phys.* 172 (1995) 223–262.
- [12] H. Omori, Y. Maeda, A. Yoshioka, Existence of a closed star product, *Lett. Math. Phys.* 26 (1992) 285–294.
- [13] M.J. Pflaum, A deformation-theoretical approach to Weyl quantization on Riemannian manifolds, *Lett. Math. Phys.* 45 (1998) 277–294.