



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)



Journal of Geometry and Physics 47 (2003) 369–377

JOURNAL OF  
GEOMETRY AND  
PHYSICS

[www.elsevier.com/locate/jgp](http://www.elsevier.com/locate/jgp)

# Geometric proprieties of invariant connections on $SL(n, \mathbb{R})/SO(n)$

Marco A.N. Fernandes<sup>a</sup>, Luiz A.B. San Martin<sup>b,\*</sup>

<sup>a</sup> Instituto de Matemática, Universidade Federal da Bahia, Av. Ademar de Barros s/n,  
40.170-110 Salvador, BA, Brazil

<sup>b</sup> Instituto de Matemática, Universidade Estadual de Campinas, Cx. Postal 6065,  
13.081-970 Campinas, SP, Brazil

Received 4 September 2002

---

## Abstract

This article describes some geometric aspects of a class of affine connections in homogeneous spaces, that emerged in an earlier paper by the authors, related to the geometry of statistical models. We describe the geodesics as well some properties of the curvature of these connections.

© 2002 Elsevier Science B.V. All rights reserved.

*Subj. Class:* Differential geometry; 53C30

*Keywords:* Symmetric spaces;  $\alpha$ -Connections; Invariant connections

---

## 1. Introduction

The  $\alpha$ -connections were introduced in the statistical literature in 1980s (see [1,2]) as a differential geometric tool for studying parametric models. The idea was to refine the classical concept of Fisher information, which is a Riemannian metric attached to a statistical model. Both these geometric objects are defined by means of integrals over some measure space, making them very hard to analyze by the standard differential geometric methods.

In [3] the authors consider a set-up, based on Lie group theory, in which it is possible to take advantage of the symmetries and describe the possible affine connections in homogeneous spaces arising as  $\alpha$ -connections of the so-called transformational statistical models. In particular, we were laid to consider invariant affine connections on symmetric spaces. It was proved in [3] that only those symmetric spaces whose restricted root system are of

---

\* Corresponding author.

*E-mail addresses:* [marcoanf@ufba.br](mailto:marcoanf@ufba.br) (M.A.N. Fernandes), [smartin@ime.unicamp.br](mailto:smartin@ime.unicamp.br) (L.A.B. San Martin).

type  $A_I$  admit such connections which are different from the canonical Riemannian one. In particular, the non-compact symmetric space  $SL(n, \mathbb{R})/SO(n)$  has a one-parameter family of invariant connections, which at the origin  $o = SO(n)$  is given by

$$\nabla_A^\alpha B = \alpha \left( \frac{AB + BA}{2} - \frac{\text{tr}(AB)}{n} I_n \right).$$

Here  $A, B \in \mathfrak{s}$ , the subspace of the symmetric matrices with zero trace, which we identify to the tangent space of  $SL(n, \mathbb{R})/SO(n)$  at  $o$  and  $I_n$  is the identity  $n \times n$  matrix. In [3] some properties of these connections were already discussed. However, many questions related to their geometry remained unsolved. The purpose of this article is to develop these properties further.

When  $\alpha = 1$  we denote the connection simply by  $\nabla_A B$ . This is the only case to be considered, since for general  $\alpha \neq 0$ , the computations are similar. For this connection we describe its geodesics in Section 2. Afterwards, in Section 3, we prove that the curvature tensor  $R(A, B, C)$  and all its covariant derivatives belong to the subspace spanned by  $A$  and  $B$ . We apply this fact to prove the following properties of  $\nabla$ : (i) the Ricci tensor is zero; (ii)  $\nabla$  is not compatible with any Riemannian metric.

Before starting, it is convenient to fix some notations, remind a few of the geometry of the symmetric space  $SL(n, \mathbb{R})/SO(n)$ , more details on the subject can be seen in [4]. Write  $M = G/K = SL(n, \mathbb{R})/SO(n)$ , the symmetric space of the positive definite matrices and  $\mathfrak{sl}(n, \mathbb{R}) = \mathfrak{so}(n) \oplus \mathfrak{s}$ , for the Cartan decomposition of the Lie algebra of  $G$  given by splitting the tangent space at the origin of  $M$ .

The group  $G$  acts transitively in  $M$  by  $g(hK) = (gh)K$ , and for each  $g \in G$ , the map  $g : M \rightarrow M$  defined for  $g(\xi) = g\xi$  is a diffeomorphism that satisfies:

$$(dg)_\xi(\tilde{A}(\xi)) = (\text{Ad}(g)(A))^\sim(g\xi), \tag{1}$$

where

$$\tilde{A}(\xi) = \frac{d}{dt}(\exp(tA)(\xi))|_{t=0},$$

and  $\text{Ad}(g) : \mathfrak{sl}(n, \mathbb{R}) \rightarrow \mathfrak{sl}(n, \mathbb{R})$ ,  $g \in G$  is the adjoint map. Still in relation to the Lie algebra of  $G$ , we know that the roots of  $\mathfrak{sl}(n, \mathbb{R})$  relative at  $\mathfrak{h}$ , Cartan subalgebra formed for the diagonal matrices of trace zero are given by

$$(\lambda_i - \lambda_k)(H) = (H, E_{ii} - E_{kk}) = \text{tr}(H(E_{ii} - E_{kk})),$$

for each  $i \neq k, i, k = 1, \dots, n$ , where  $E_{ik}$  is the basic  $n \times n$  matrix whose  $ik$  entry is 1 and all the others are zero.

## 2. Geodesics

Using a geometric characterization of the  $\alpha$ -connections, made in [3] we shall obtain here a description of the geodesics for the  $\alpha$ -connections in the symmetric space  $M = G/K$  of positive definite matrices.

The  $\alpha$ -connections for a models considered here have ageometric interpretation which were described in [3]. Let  $\mathcal{S}$  be the vector space of all symmetric  $n \times n$  matrices. We have that  $\mathfrak{s} = \{A \in \mathcal{S} : \text{tr } A = 0\}$  is a subspace of codimension one of  $\mathcal{S}$ , complemented by the line spanned by the identity 1. The trace form  $\text{tr}(AB)$  an inner product on  $\mathcal{S}$ . With respect to this inner product the line of scalar matrices is orthogonal to  $\mathfrak{s}$ . We denote by  $\mathcal{S}^+$  the cone of the positive semi-definite matrices in  $\mathcal{S}$ .

There is a natural action of  $\text{Sl}(n, \mathbb{R})$  on  $\mathcal{S}$  given by the law

$$(g, A) \mapsto g \cdot s = gAg^*,$$

where  $g^*$  means transposition of matrix. The induced infinitesimal action of  $\mathfrak{sl}(n, \mathbb{R})$  on  $\mathcal{S}$  is given by the derivative

$$\left. \frac{d}{dt} (e^{tX} s e^{tX^*}) \right|_{t=0} = Xs + sX.$$

Each  $X \in \mathfrak{sl}(n, \mathbb{R})$  induces the linear vector field

$$\tilde{X}(s) = Xs + sX.$$

If  $a \cdot 1, a \neq 0$  is a scalar matrix in  $\mathcal{S}$  then its orbit  $\mathcal{O}(a)$  under  $\text{Sl}(n, \mathbb{R})$  is the subset of matrices with determinant  $a^n$  which are positive definite if  $a > 0$  or negative definite if  $a < 0$ . Since  $g(a \cdot 1)g^* = a \cdot 1$  if and only if  $g$  is an orthogonal matrix, it follows that  $\mathcal{O}(a), a \neq 0$ , identifies with the homogeneous space  $\text{Sl}(n, \mathbb{R})/\text{SO}(n, \mathbb{R})$ . These orbits have codimension one in  $\mathcal{S}$ , and the tangent space  $T_{a1}\mathcal{O}(a)$  is the subspace of matrices with trace zero. Note that the line of scalar matrices complements  $T_{a1}\mathcal{O}(a)$  in  $\mathcal{S}$ . Similarly, one checks easily that

$$\mathcal{S} = T_s\mathcal{O}(a) \oplus [s], \tag{2}$$

where  $[s]$  stands for the line spanned by  $s \in \mathcal{O}(a)$ . From this decomposition we obtain the following connection  $\nabla$  on  $\mathcal{O}(a)$ :

$$(\nabla_X Y)(s) = \text{pr}_s((dY)_s(X(s))). \tag{3}$$

Here  $\text{pr}_s : \mathcal{S} \rightarrow T_s\mathcal{O}(a)$  is the projection coming from the decomposition in (2), and  $X, Y$  are vector fields in  $\mathcal{O}(a)$  with  $Y$  viewed as a mapping  $Y : \mathcal{O}(a) \rightarrow \mathcal{S}$  so that  $(dY)_s$  stands for its differential at  $s$ . The definition of  $\nabla$  is analogous to the Levi-Civita connection of the Riemannian metric induced in an immersed submanifold of an Euclidean space. However here the projection is not orthogonal with respect to  $\text{tr}(AB)$ , since the line  $[s^{-1}]$  is orthogonal to  $T_s\mathcal{O}(a)$  so that  $\text{pr}_s$  is orthogonal if and only if  $s = a \cdot 1$ . Each orbit  $\mathcal{O}(a), a \neq 0$  is diffeomorphic to  $\text{Sl}(n, \mathbb{R})/\text{SO}(n, \mathbb{R})$ . Hence we have a family of connections  $\nabla^a$  in  $\text{Sl}(n, \mathbb{R})/\text{SO}(n, \mathbb{R})$ . It was checked in [3] that  $\nabla^a$  is a  $\alpha$ -connection for each  $a$ .

In discussing geodesics we simplify matters and take  $a = 1$ . The other cases follow analogously. Thus we consider the orbit  $\mathcal{O}(1)$  and put  $\nabla = \nabla^1$ . Due to invariance it is sufficient to find the geodesics going through the origin. Let  $s = s(t), \det s = 1$ , be a geodesic of  $\nabla$ . Then  $\nabla_s \dot{s} = 0$  and therefore by the above description of  $\nabla$  the projection of  $\dot{s}$  is annihilates. This means that  $\dot{s} = es$  for some constant  $e$ . We compute this constant by

taking the second derivative of the equality  $\det s = 1$ . Using a well known formula for the derivative of the determinant we get

$$\operatorname{tr}(s^{-1}\dot{s}) \det s = 0,$$

hence  $\operatorname{tr}(s^{-1}\dot{s}) \det s = 0$ . Taking another derivative and using  $(s^{-1})' = -s^{-1}\dot{s}s^{-1}$ , we get

$$(-\operatorname{tr}(s^{-1}\dot{s}s^{-1}\dot{s}) + \operatorname{tr}(s^{-1}\ddot{s})) \det s + \operatorname{tr}(s^{-1}\dot{s})^2 \det s = 0.$$

But  $\operatorname{tr}(s^{-1}\dot{s}) = 0$  and  $\det(s) = 1$ . Hence

$$\operatorname{tr}(s^{-1}\ddot{s}) = \operatorname{tr}(s^{-1}\dot{s}s^{-1}\dot{s}).$$

Now,  $\ddot{s} = es$ , so that  $\operatorname{tr}(s^{-1}\ddot{s}) = e \operatorname{tr}(1) = ne$ . Therefore, the equation satisfied by the geodesics through the identity is

$$\ddot{s} = \frac{\operatorname{tr}(s^{-1}\dot{s}s^{-1}\dot{s})}{n} s. \tag{4}$$

Since this equation looks hard to integrate explicitly we shall give a geometric description of the trace of the geodesics, and then write down a reparametrization of them.

**Proposition 2.1.** *The traces of the geodesic of  $\nabla$  in  $\mathcal{S}_1$  are the subsets*

$$\mathcal{S}_1 \cap V,$$

where  $V \subset \mathcal{S}$  is a two-dimensional subspace which has non-empty intersection with  $\mathcal{S}_1$ .

**Proof.** We check first that  $\mathcal{S}_1 \cap V$  is a curve, that is, a one-dimensional submanifold in case the intersection is not empty. In fact, denoted by  $P$  the restriction to  $V$  of  $\det$ . Then  $P$  is a polynomial function on  $V$ . Take  $s \in \mathcal{S}_1 \cap V$ . By the well known formula for the differential of  $\det$

$$dP_s(s) = d(\det)_s(s) = \operatorname{tr}(ss^{-1}) \det(s) = n \neq 0.$$

This shows that every  $s \in \mathcal{S}_1 \cap V$  is a regular point of  $P$ . Now,  $\mathcal{S}_1 \cap V$  is a connected component of a level set of  $P$ . Hence the intersection is indeed a one-dimensional submanifold.

Next we verify that  $\mathcal{S}_1 \cap V$  can be entirely parametrized by a curve  $s(t)$  such that its second derivative  $\ddot{s}$  is a multiple of  $s$ , and hence satisfies Eq. (4). For this let  $\tilde{\nabla}$  be the connection  $\mathcal{S}_1 \cap V$  defined analogously to  $\nabla$  by projecting onto the tangent space along the line spanned by  $s \in \mathcal{S}_1 \cap V$ . Note that this is possible because  $dP_s(s) \neq 0$ , so that the line spanned by  $s$  is transversal to the tangent space of  $\mathcal{S}_1 \cap V$  at  $s$ . Now, let  $s : (\alpha, \omega) \in \mathbb{R} \rightarrow \mathcal{S}_1 \cap V$  be a geodesic of  $\tilde{\nabla}$ . Then  $\ddot{s}(t)$  is a multiple of  $s(t)$  for all  $t$ , by definition of  $\tilde{\nabla}$ . Hence,  $s$  satisfies (4), so that it is also a geodesic of  $\nabla$ . The trace of this geodesic  $s$  is the whole  $\mathcal{S}_1 \cap V$ . In fact, suppose that  $\lim_{t \rightarrow \omega} s(t) = s_\infty \in \mathcal{S}_1 \cap V$ . Then by the usual argument we can extend  $s$  with a geodesic going through  $s_\infty$ , concluding the proof.  $\square$

Now, we shall obtain parametrizations of the geodesic curves  $\mathcal{S}_1 \cap V$ . We restrict attention to those subspaces  $V$  containing the identity 1, having in mind that the other subspaces are

obtained by translation. In fact, if the two-dimensional subspace  $V$  meets  $\mathcal{S}_1$ , then for some  $g \in \text{Sl}(n, \mathbb{R})$ ,  $gV$  contains 1, and we can use the equality

$$g(\mathcal{S}_1 \cap V) = \mathcal{S}_1 \cap gV.$$

Thus let  $V$  be such that  $1 \in V$  and  $\dim V = 2$ . As before let  $\mathfrak{s}$  be the subspace of matrices with zero trace. Then there exists  $A \in \mathfrak{s}$  such that  $V$  is spanned by  $\{1, A\}$ . If we take conjugation by an element of  $\text{SO}(n, \mathbb{R})$  we can assume that  $A$  is diagonal, that is

$$A = \text{diag}\{x_1, \dots, x_n\},$$

with  $x_1 + \dots + x_n = 0$ . In this case  $\mathcal{S}_1 \cap V$  becomes the subset of diagonal matrices

$$\text{diag}\{tx_1 + s, \dots, tx_n + s\},$$

satisfying

$$(tx_1 + s) \cdots (tx_n + s) = 1, \quad tx_i + s > 0. \tag{5}$$

To get a parametrization of this curve note that the matrices

$$\text{diag} \left\{ \frac{1}{n} + tx_1, \dots, \frac{1}{n} + tx_n \right\}, \quad \frac{1}{n} + tx_i > 0,$$

belong to the interior of the simplex

$$\Delta = \{(y_1, \dots, y_n) : y_1 + \dots + y_n = 1, \quad y_i \geq 0\}.$$

On the other hand, the map  $(y_1, \dots, y_n) \in \text{int } \Delta$

$$(y_1, \dots, y_n) \in \text{int } \Delta \mapsto \frac{1}{\sqrt[n]{y_1 \cdots y_n}}(y_1, \dots, y_n),$$

is a bijection between  $\Delta$  and the set of diagonal symmetric matrices with  $\det = 1$ . Thus a parametrization of our curve is given by

$$t \mapsto \frac{1}{\sqrt[n]{(1 + tx_1) \cdots (1 + tx_n)}}(1 + tx_1, \dots, 1 + tx_n). \tag{6}$$

Its domain is the largest interval such that  $1 + tx_i > 0$  for all  $i$ . At this point we re-order if necessary the basis so that the matrix  $A = \text{diag}(x_1, \dots, x_n)$  satisfies  $x_1 \geq \dots \geq x_n$ . In this case  $x_1 > 0$  and  $x_n < 0$  and the domain of definition of the above parametrization becomes

$$\left( -\frac{1}{x_1}, -\frac{1}{x_n} \right).$$

Now we use the parametrization (6) to write down some further equations related to the geodesic of the given curve. Another parametrization of (6) is obtained by writing  $t = \phi(u)$ . The reparametrization is a geodesic if and only if the second derivative is a multiple of the curve. Thus we write  $s(u)$  as

$$s = \frac{1}{R}(1 + \phi x_1, \dots, 1 + \phi x_n),$$

where  $R = \sqrt[n]{(1 + \phi x_1) \cdots (1 + \phi x_n)}$ . In order to perform the computations we write

$$l_i = \log s_i,$$

where  $s_i, i = 1, \dots, n$  are the coordinates of  $s$ . We have

$$l'_i = \frac{s'_i}{s_i}, \quad l''_i = \frac{s''_i}{s_i} - \left(\frac{s'_i}{s_i}\right)^2. \tag{7}$$

Now the condition  $\ddot{s} = cs$  for  $s$  to be a geodesic means that  $s''_i/s_i$  is independent of the index  $i$ . By the expressions (7) this happens if and only if  $l''_i + (l'_i)^2$  does not depend on  $i = 1, \dots, n$ . A straightforward computation yields:

- $l_i = \log(1 + \phi x_i) - \frac{1}{n}(\log(1 + \phi x_1) + \cdots + \log(1 + \phi x_n)),$
- $l'_i = \phi' \left( \frac{x_i}{1 + \phi x_i} - \frac{1}{n} \left( \frac{x_1}{1 + \phi x_1} + \cdots + \frac{x_n}{1 + \phi x_n} \right) \right),$
- $l''_i = \phi'' \left( \frac{x_i}{1 + \phi x_i} - \frac{1}{n} \left( \frac{x_1}{1 + \phi x_1} + \cdots + \frac{x_n}{1 + \phi x_n} \right) \right) + (\phi')^2 \left( -\frac{x_i^2}{(1 + \phi x_i)^2} + \frac{1}{n} \left( \frac{x_1^2}{(1 + \phi x_1)^2} + \cdots + \frac{x_n^2}{(1 + \phi x_n)^2} \right) \right),$
- $(l'_i)^2 = (\phi')^2 \left( \frac{x_i^2}{(1 + \phi x_i)^2} - \frac{2}{n} \frac{x_i}{1 + \phi x_i} \left( \frac{x_1}{1 + \phi x_1} + \cdots + \frac{x_n}{1 + \phi x_n} \right) + \frac{1}{n^2} \left( \frac{x_1}{1 + \phi x_1} + \cdots + \frac{x_n}{1 + \phi x_n} \right)^2 \right).$

Looking at these expressions we see that the term of  $l''_i + (l'_i)^2$  which depends explicitly on  $i$  is given by

$$\frac{x_i}{1 + \phi x_i} \left( \phi'' - \frac{2(\phi')^2}{n} \left( \frac{x_1}{1 + \phi x_1} + \cdots + \frac{x_n}{1 + \phi x_n} \right) \right).$$

Therefore the reparametrization  $\phi$  turns the curve into a geodesic if and only if it satisfies the second order differential equation

$$\phi'' = \frac{2(\phi')^2}{n} \left( \frac{x_1}{1 + \phi x_1} + \cdots + \frac{x_n}{1 + \phi x_n} \right). \tag{8}$$

Taking logarithms this equation is written as

$$(\log \phi')' = \left( \frac{2}{n} \log(1 + \phi x_1) \cdots (1 + \phi x_n) \right)'$$

Hence (8) is equivalent to

$$\phi' = ((1 + \phi x_1) \cdots (1 + \phi x_n))^{2/n} + c, \tag{9}$$

where the constant  $c$  accounts for the initial condition in the second derivative in (8). If we choose  $\phi$  so that  $\phi(0) = 0$  and  $\phi'(0) = 1$  we arrive at the equation for the geodesics.

**Proposition 2.2.** *The geodesics of the connection  $\nabla = \nabla^{1/4}$  starting at the identity matrix 1 in the direction of the matrix*

$$A = \text{diag}\{x_1, \dots, x_n\}, \quad x_1 + \dots + x_n = 0,$$

is given by

$$\gamma(u) = \frac{1}{\sqrt{\phi'(u)}}(1 + \phi(u)x_1, \dots, 1 + \phi(u)x_n),$$

where  $\phi$  is the solution of the first order differential equation

$$\phi' = ((1 + \phi x_1) \cdots (1 + \phi x_n))^{2/n},$$

with  $\phi(0) = 0$ .

It is convenient to make the following remark about the domain of definition of the first order equation (9): first if some  $x_i = 0$  then the  $i$ th term does not appear, hence we assume that  $x_i \neq 0$  for every  $i$ . In this case the equation is not Lipschitz in  $\phi$  when  $\phi = -1/x_i$ . Therefore, if we take  $x_1 \geq \dots \geq x_n$  the domain of definition of the equation is  $(-1/x_1, -1/x_n)$ , which is precisely the domain of the original parametrization.

### 3. The curvature tensor

Given a differentiable manifold  $M$  with an affine connection  $\nabla$ , a tensor of the type  $(r, s)$  is a map

$$T : \underbrace{\chi(M) \times \dots \times \chi(M)}_{r \times} \rightarrow \underbrace{\chi(M) \times \dots \times \chi(M)}_{s \times},$$

that is linear in each component  $\chi(M)$  considered as module on  $C^\infty(M)$ . The covariant derivative of  $T$ ,  $\nabla T$  is the tensor of type  $(r + 1, s)$  defined by

$$\begin{aligned} (\nabla T)(A_1, \dots, A_r) &= (\nabla_A T)(A_1, \dots, A_r) = \nabla_A(T(A_1, \dots, A_r)) \\ &= \sum_{i=1}^r T(A_1, \dots, \nabla_A A_i, \dots, A_r), \end{aligned}$$

for  $A_1, \dots, A_r \in \chi(M)$ . The second covariant derivative of  $T$ ,  $\nabla^2 T = \nabla(\nabla T)$  is then a tensor of the type  $(r + 2, s)$  given by

$$(\nabla^2 T)(A_1, \dots, A_r, B) = (\nabla_B(\nabla T))(A_1, \dots, A_r),$$

$A_1, \dots, A_r, B \in \chi(M)$ . In general, the  $m$ th covariant derivative,  $\nabla^m T$  is inductively defined by  $\nabla(\nabla^{m-1} T)$ .

Now, using the usual formula for the curvature

$$R(A, B, C) = \nabla_A \nabla_B C - \nabla_B \nabla_A C - \nabla_{[A, B]} C, \quad A, B, C \in \mathfrak{s},$$

a direct computation yields for  $R(A, B, C)$  the following expression

$$\frac{1}{n}(\text{tr}(AC)B - \text{tr}(BC)A + \text{tr}([A, B]C)I_n) - \frac{1}{[A, B]C} - C[A, B].$$

Note that if we restrict  $R$  to the totally geodesic submanifold of the diagonal matrices of  $M$ , then

$$R(A, B, C) = \frac{1}{n}(\text{tr}(AC)B - \text{tr}(BC)A). \tag{10}$$

Our next objective it is enough to compute the covariant derivatives of  $R$  in this submanifold, that is, we want to compute  $\nabla^m R(A, B, C)$ , for  $A, B, C \in \mathfrak{h}$ . For this we introduce the tensor of the type  $(r, 0)$ ,  $T_r : \mathfrak{s} \times \dots \times \mathfrak{s} \rightarrow C^\infty(M)$ , defined by

$$T_r(A_1, \dots, A_r) = \text{tr}(A_1 \dots A_r).$$

An easy computation shows that

$$\begin{aligned} &\nabla T_r(A_1, \dots, A_{r+1}) \\ &= -rT_{r+1}(A_1, \dots, A_{r+1}) + \frac{1}{n} \sum_{i=1}^r (T_2 \otimes T_{r-1})(A_i, A_{r+1}, A_1, \dots, \hat{A}_i, A_r). \end{aligned}$$

Also, if we put  $S(A_1, \dots, A_r, A_{r+1}) = T_r(A_1, \dots, A_r)A_{r+1}$ , then we get

$$(\nabla T_r)(A_1, \dots, A_r, A_{r+2})A_{r+1} = (\nabla S)(A_1, \dots, A_{r+1}, A_{r+2}).$$

Using these notations we arrive at the following formulas:

- $R(A, B, C) = (1/n)(T_2(A, C)B - T_2(B, C)A)$ .
- $(\nabla R)(A, B, C, D) = -(2/n)(T_3(A, C, D)B - T_3(B, C, D)A)$ .
- $-(n/6)(\nabla^2 R)(A, B, C, D, E)$  is given by

$$\left\{ \left( T_4 - \frac{1}{3n} U \right) (A, C, D, E) \right\} B + \left\{ \left( T_4 - \frac{1}{3n} U \right) (B, C, D, E) \right\} A,$$

where  $U(A, C, D, E)$  is the tensor

$$T_2(A, E)T_2(C, D) + T_2(D, E)T_2(A, C) + T_2(C, E)T_2(A, D).$$

We can proceed successively and compute the covariant derivatives of any order. We shall refrain ourselves to develop a general formula for these derivatives. But it is clear from these formulas that the following statement holds.

**Proposition 3.1.** *If  $A$  and  $B$  are zero trace diagonal matrices then the covariant derivatives  $\nabla^m R$  belong to the subspace spanned by  $A$  and  $B$ , for all  $m \in \mathbb{N}$ , where  $\nabla^0 R = R$ .*



In the sequel we shall obtain some applications of the formulas obtained so far.

First let us consider the *Ricci tensor*. For a general connection this is the  $(2, 0)$ -tensor defined by

$$\text{Ric}(A, B) = \text{tr}(C \mapsto R(A, B, C)), \quad A, B, C \in \chi(M).$$

In our case if  $A, B, C \in \mathfrak{s}$  then the map  $C \mapsto R(A, B, C)$  is an element of  $\mathfrak{gl}(\mathfrak{s})$ , having trace zero. Hence,  $\text{Ric} \equiv 0$ .

As a second application we ask weather there exists a Riemannian metric  $g$  compatible with the affine connection  $\nabla$ . Recall that this holds if

$$A g(B, C) = g(\nabla_A B, C) + g(B, \nabla_A C), \quad A, B, C \in \chi(M).$$

It is known that for a connection compatible with a given metric, the Lie algebra of the holonomy group in a point of the manifold is a subalgebra of  $\mathfrak{so}(n)$  (see [5]). On the other hand, such Lie algebra is spanned by

$$(\nabla^m R)(A, B, C_1, \dots, C_m), \quad A, B, C_1, \dots, C_m \in \chi(M), \quad m = 0, 1, 2, \dots$$

We shall use these facts to prove that  $\nabla$  is not compatible with any metric.

For this choose  $A, B, C, D, E \in \mathfrak{h}$  satisfying  $\text{tr}(BC) = \text{tr}(BE) = 0, DB = A$ . Then by Proposition 3.1, we have

$$\langle (\nabla R)(A, B, C, D), E \rangle = \langle (\nabla R)(A, B, E, D), C \rangle.$$

Equivalently,  $(\nabla R)(A, B, C)$  is a non-zero self-adjoint operator of  $\mathfrak{h}$ . Hence, it has real eigenvalues, showing that this operator cannot belong to  $\mathfrak{so}(n)$ . This is enough to prove that  $\nabla$  is not compatible with a Riemannian metric.

We note that by (10)

$$\langle R(A, B, C), D \rangle = \langle C, R(A, B, D) \rangle,$$

that is,  $R(A, B) \in \mathfrak{so}(n)$ , so that we in fact need the covariant derivative of the curvature.

## Acknowledgements

Research partially supported by CNPq Grant No. 301060/94-0.

## References

- [1] S. Amari, Differential–geometrical methods in statistics, in: Lecture Notes in Statistics, vol. 28, Springer, Berlin, 1985.
- [2] O.E. Barndorff-Nielsen, Differential and integral geometry in statistical inference, in: Lecture Notes—Monograph Series, vol. 10, Institute of Mathematical Statistics, California, 1987.
- [3] M.A.N. Fernandes, L.A.B. San Martin, Fisher information and  $\alpha$ -connections for a class of transformational models, Diff. Geom. Appl. 12 (2000) 165–184.
- [4] S. Helgason, Differential Geometry, Lie Groups and Symmetric Spaces, Academic Press, New York, 1978.
- [5] S. Kobayashi, K. Nomizu, Foundations of Differential Geometry, vol. 2, Wiley, New York, 1962.