



ELSEVIER

Available online at www.sciencedirect.com



Journal of Geometry and Physics 48 (2003) 546–579

JOURNAL OF
GEOMETRY AND
PHYSICS

www.elsevier.com/locate/jgp

Classification of polarized deformation quantizations

Joseph Donin

Department of Mathematics and Computer Science, Bar-Ilan University, 52900 Ramat-Gan, Israel

Received 18 June 2002; received in revised form 6 March 2003

Abstract

We give a classification of polarized deformation quantizations (PDQs) on a symplectic manifold with a (complex) polarization.

Also, we establish a formula which relates the characteristic class of a PDQ to its Fedosov class and to the Chern class of the polarization.

© 2003 Elsevier Science B.V. All rights reserved.

MSC: 53D55

JGP SC: Quantum mechanics

Keywords: Deformation quantization; Star-product; Symplectic manifold; Polarization

1. Introduction

Let (M, ω) be a smooth symplectic manifold, \mathcal{T}_M its *complexified* tangent bundle. It is known that classes of deformation quantizations (DQs) on (M, ω) are in one-to-one correspondence with their Fedosov classes, the elements of $\omega + tH^2(M, \mathbb{C}[[t]])$. The set $\omega + tH^2(M, \mathbb{C}[[t]])$ may be interpreted in the following way. Let \mathcal{X} be the set of formal closed 2-forms on M of the form $\omega + t\omega_1 + t^2\omega_2 + \dots$. Let $\text{Aut}(M)$ be the group of formal automorphisms of M of the form e^{tX} , where $X = X_0 + tX_1 + t^2X_2 + \dots$, $X_i \in \mathcal{T}_M$, is a formal vector field. Since ω is nondegenerate, it is easy to see that the orbit of an element $\omega_t \in \mathcal{X}$ under the action of $\text{Aut}(M)$ is $\omega_t + t \cdot d(\Gamma(M, \mathcal{T}_M^*))$. This implies that $\omega + tH^2(M, \mathbb{C}[[t]])$ can be identified with the set of orbits in \mathcal{X} under the $\text{Aut}(M)$ -action. Therefore, the equivalence classes of DQs on (M, ω) are in one-to-one correspondence with the orbits in \mathcal{X} .

E-mail address: donin@macs.biu.ac.il (J. Donin).

In the paper, we extend that picture to polarized deformation quantizations (PDQs).

Let (M, ω, \mathcal{P}) be a polarized symplectic manifold (PSM), i.e. \mathcal{P} is a Lagrangian integrable subbundle of the complexified tangent bundle to M . A PDQ on (M, ω, \mathcal{P}) is a pair $(\mathbb{A}_t, \mathbb{O}_t)$, where \mathbb{A}_t is a DQ on (M, ω) , i.e. \mathbb{A}_t is a sheaf of noncommutative algebras being a deformation of the sheaf \mathcal{C}_M^∞ of smooth functions on M , and \mathbb{O}_t is a subsheaf of commutative t -adically complete subalgebras of \mathbb{A}_t such that $\mathbb{O}_0 = \mathcal{O}_{\mathcal{P}}$, the sheaf of functions constant along \mathcal{P} .

Let \mathcal{Y} denote the set of pairs $(\omega_t, \mathcal{P}_t)$, where $\omega_t \in \mathcal{X}$ and \mathcal{P}_t is a polarization of ω_t such that $\mathcal{P}_0 = \mathcal{P}$. Our result is that the equivalence classes of PDQs on (M, ω, \mathcal{P}) are in one-to-one correspondence with the orbits in \mathcal{Y} under the $\text{Aut}(M)$ -action. Let us describe this correspondence more precisely.

First, we show that any PDQ is equivalent to a polarized star-product (PSP). By a PSP we mean a triple, $(\mathcal{C}_M^\infty[[t]], \mu_t, \mathcal{O}_t)$, where $(\mathcal{C}_M^\infty[[t]], \mu_t)$ is a star-product (SP), $\mathcal{O}_t = \mathcal{O}_{\mathcal{P}_t}$, the algebra of functions from $\mathcal{C}_M^\infty[[t]]$ constant along a deformed polarization \mathcal{P}_t , and the multiplication μ_t satisfies the condition: $\mu_t(f, g) = fg$ (the usual multiplication) for $f \in \mathcal{O}_t$, $g \in \mathcal{C}_M^\infty[[t]]$.

Further, we assign to any PSP $(\mathcal{C}_M^\infty[[t]], \mu_t, \mathcal{O}_t)$ a pair $(\omega_t, \mathcal{P}_t) \in \mathcal{Y}$ in the following way. We put $\mathcal{P}_t = \mathcal{P}_{\mathcal{O}_t}$, the sheaf of formal vector fields annihilating \mathcal{O}_t . The form ω_t is equal, locally, to $\sum_i dy_i \wedge dx_i$, where $x_i \in \mathcal{O}_t$, $y_i \in \mathcal{C}_M^\infty[[t]]$, $i = 1, \dots, (1/2)\dim M$, are Darboux coordinates with respect to $[,] = [,]_{\mu_t}$, the commutator of μ_t ; namely, $[x_i, x_j] = [y_i, y_j] = 0$, $[y_i, x_j] = \delta_{ij}$. It turns out that ω_t is well defined, i.e. does not depend on the choice of local Darboux coordinates. Denote the constructed map from PSPs to \mathcal{Y} by τ .

The map τ turns out to descend to an isomorphism between the set of classes of PDQs and the set $[\mathcal{Y}]$ of orbits in \mathcal{Y} . So, we obtain the following commutative diagram of maps:

$$\begin{array}{ccc} \{\text{PSPs}\} & \xrightarrow{\tau} & \mathcal{Y} \\ \downarrow & & \downarrow \\ \{\text{classes of PDQs}\} & \rightarrow & [\mathcal{Y}], \end{array} \quad (1.1)$$

where the left downward arrow is an epimorphism and the bottom arrow is an isomorphism of sets.

We prove that the top arrow τ is an epimorphism with the following properties:

- (1) Two PSPs are equivalent if and only if their images with respect to τ lie on the same orbit.
- (2) Two PSPs $(\mathcal{C}_M^\infty[[t]], \mu_t, \mathcal{O}_t)$ and $(\mathcal{C}_M^\infty[[t]], \tilde{\mu}_t, \tilde{\mathcal{O}}_t)$ have the same image with respect to τ if and only if $\mathcal{O}_t = \tilde{\mathcal{O}}_t$ and $[,]_{\mu_t} = [,]_{\tilde{\mu}_t}$.
- (3) Let $(\omega_t, \mathcal{P}_t) = \tau(\mathcal{C}_M^\infty[[t]], \mu_t, \mathcal{O}_t)$. Then, the 2-form

$$\theta_t = \omega_t + \frac{t}{2} \text{tr}(\nabla^2|_{\mathcal{P}_t}) \quad (1.2)$$

represents the Fedosov class of the SP $(\mathcal{C}_M^\infty[[t]], \mu_t)$. Here ∇ is a connection on M preserving ω_t , \mathcal{P}_t , and flat on \mathcal{P}_t along \mathcal{P}_t . We prove that such a connection always exists and for it $\text{tr}(\nabla^2|_{\mathcal{P}_t})$ belongs to $\Gamma(M, d\mathcal{P}_t^\perp)$.

By definition, ω_t belongs to $\omega + t\Gamma(M, d\mathcal{P}_t^\perp)$. Hence, it follows from (1.2) that the Fedosov class of the SP ($\mathcal{C}_M^\infty[[t]]$, μ_t) can be represented by a 2-form belonging to $\omega + t\Gamma(M, d\mathcal{P}_t^\perp)$, as well. In particular, both $(M, \omega_t, \mathcal{P}_t)$ and $(M, \theta_t, \mathcal{P}_t)$ are formal PSMs that are deformations of (M, ω, \mathcal{P}) .

Another consequence of (1.2) is the following one. Let \mathbb{A}_t be a DQ on (M, ω) . Suppose its Fedosov class $\text{cl}_F(\mathbb{A}_t)$ is represented by a 2-form θ_t that has a polarization \mathcal{P}_t . Then \mathbb{A}_t can be extended to a PDQ $(\mathbb{A}_t, \mathbb{O}_t)$, i.e. there exists a commutative subalgebra $\mathbb{O}_t \subset \mathbb{A}_t$ isomorphic to $\mathcal{O}_{\mathcal{P}_t}$.

There is the following interpretation of the image $[\omega_t]$ of element ω_t in the $\mathbb{C}[[t]]$ -module $\Gamma(M, d\mathcal{P}_t^\perp)/d(\Gamma(M, \mathcal{P}_t^\perp))$. Let $F(\mu_t, \mathcal{O}_t) = \{a \in \mathcal{C}_M^\infty[[t]] : [a, \mathcal{O}_t]_{\mu_t} \subset \mathcal{O}_t\}$. Then, there is the following exact sequence of \mathcal{O}_t -module and Lie algebra sheaves:

$$0 \rightarrow \mathcal{O}_t \rightarrow F(\mu_t, \mathcal{O}_t) \rightarrow \text{Der}(\mathcal{O}_t) \rightarrow 0. \quad (1.3)$$

According to Beilinson and co-workers [3,4], $F(\mu_t, \mathcal{O}_t)$ is called an \mathcal{O}_t -extension of $\text{Der}(\mathcal{O}_t)$. Equivalence classes of such extensions are described by their extension classes that are elements of $\Gamma(M, d\mathcal{P}_t^\perp)/d(\Gamma(M, \mathcal{P}_t^\perp))$. We show that $[\omega_t]$ is just the extension class of (1.3).

Analogously, $-\text{tr}(\nabla^2|_{\mathcal{P}_t})$ is the extension class of the \mathcal{O}_t -extension

$$0 \rightarrow \mathcal{O}_t \rightarrow \tilde{T}_{\det(\mathcal{P}_t)} \rightarrow \text{Der}(\mathcal{O}_t) \rightarrow 0, \quad (1.4)$$

where $\tilde{T}_{\det(\mathcal{P}_t)}$ is the sheaf of \mathcal{O}_t -differential operators of order at most one on the \mathcal{O}_t -line bundle $\det(\mathcal{P}_t)$.

Note that $-\text{tr}(\nabla^2|_{\mathcal{P}_t})$ divided by $2\pi\sqrt{-1}$ represents the first Chern class of \mathcal{P} [15]. Therefore, formula (1.2) gives a relation of the extension class of a PDQ to its Fedosov class and to the Chern class of the polarization.

Among results related to ours we mention the following.

In [22], Reshetikhin and Yakimov considered the case of a real polarization on M defined by a Lagrangian fiber bundle $M \rightarrow B$.

In [12], Karabegov constructed SPs with separation of variables on Kähler manifolds. This case corresponds to two polarizations on M defined by holomorphic and anti-holomorphic vector fields. In the case of quantization on a Kähler manifold with separation of variables the class of ω_t in $H^2(M, \mathbb{C}[[t]])$ coincides with the class defined by Karabegov in [12]. A formula relating the Karabegov and Fedosov classes in the case of Kähler manifolds is found in [13], see also [14,19]. In the general case of the quantization with separation of variables the analogous formula was conjectured in [1].

An important case of PSMs is the cotangent bundle to a smooth manifold with the natural symplectic structure. Polarized quantizations on such manifolds, the so-called standard-ordered SPs, was investigated by Bordemann et al. in [6], Bordemann et al. in [8] and Pflaum in [20].

Our proof of the existence of a PSP associated with any orbit in \mathcal{Y} uses the Fedosov method adapted for the case with polarization. The analogous method was applied by Bordemann and Waldmann [7], for constructing a quantization with separation of variables on a Kähler manifold.

Another approach to proving a formula relating the Fedosov and extension classes is found in [5]; that approach uses the Deligne classes of DQs [9]. Unfortunately, there was a

deficiency in proving Lemma 4.3 of [5] relating the extension and Deligne classes, however, the proof can be modified using arguments of the present paper.

The paper is organized as follows.

In Section 2, we study cohomologies of the differential Hochschild complex on M in presence of a distribution. Also, we prove a version of the Kostant–Hochschild–Rosenberg theorem for functions constant along a distribution. We use these results latter in proving that any PDQ is equivalent to a PSP.

In Section 3, we introduce a notion of \mathbb{C} -symplectic manifold, which is convenient for our consideration. This notion is a generalization of the notion of symplectic manifold. Namely, we suppose that symplectic form ω on a \mathbb{C} -symplectic manifold is a *complex* one and, locally, there exist *complex* Darboux coordinates with respect to ω . For a usual symplectic manifold, when ω is real, such coordinates exist by the Darboux theorem. In this section, we establish some facts on \mathbb{C} -symplectic manifolds with polarization. By a polarization of ω we mean a Lagrangian subbundle, \mathcal{P} , of the *complexified* tangent bundle on M such that, locally on M , there exist Darboux coordinates $x_i, y_i, i = 1, \dots, (1/2) \dim M$, where $x_i \in \mathcal{O}_{\mathcal{P}}$ for all i . Thus, (pseudo-) Kähler manifolds as well as purely real polarizations are included in our consideration. Note that from an analog of “Dolbeault Lemma” proved in [21] one can derive sufficient conditions for \mathcal{P} to be a complex polarization of ω .

In Section 4, we study properties of formal (or deformed) PSMs. In Section 5, we prove the existence of a polarized symplectic connection on a formal PSM, $(M, \omega_t, \mathcal{P}_t)$, and with the help of such a connection we introduce a characteristic class of a PSM.

In Section 6, we prove some technical statements related to deformations of Poisson brackets on M . Such deformations appear, in particular, as commutators of SPs.

In Section 7, we study properties of PDQs. In particular, we prove the important fact that any PDQ is equivalent to a PSP.

In Section 8, we define the extension class of a PDQ. Besides, we assign to any PSP an element of \mathcal{Y} , and to any class of PDQs an orbit in \mathcal{Y} . We prove that the later assignment is a monomorphism that, actually, is an isomorphism, as we show in the next section.

In Section 9, we prove that each element of \mathcal{Y} corresponds to a PSP. To this end, we adapt the Fedosov method for constructing a PSP corresponding to a given pair $(\omega_t, \mathcal{P}_t) \in \mathcal{Y}$. By this method, a polarized symplectic connection, ∇ , extends to a Fedosov connection on the bundle of Weyl algebras on M . This connection has two scalar curvatures: the Weyl curvature, θ_t , and the Wick curvature that turns out to be just ω_t . We show that these curvatures differ from each other by $(t/2) \operatorname{tr}(\nabla^2|_{\mathcal{P}_t})$, which immediately proves (1.2). Note that in order to consider Weyl and Wick curvatures simultaneously on the same bundle of Weyl algebras, we realize latter as a quotient of the tensor algebra over the cotangent bundle to M . This realization was used in [10] to give an algebraic version of Fedosov’s construction.

In Section 10, we formulate the main theorem collecting the results of the paper and give some corollaries.

2. Complex distributions

For a smooth manifold M , we will denote by \mathcal{C}_M^∞ the sheaf of *complex valued* smooth functions on M and by $\mathcal{T}_M^\mathbb{C} = \mathcal{T}_M \otimes_{\mathbb{R}} \mathbb{C}$ the complexified tangent bundle on M .

We say that a set of smooth functions x_i , $i = 1, \dots, \dim M$, given on an open subset $U \subset M$ form a system of (complex) coordinates on U , if dx_i are linearly independent at each point of U .

Since any 1-form on U can be uniquely written as $\sum_i a_i dx_i$, one can define vector fields $\partial/\partial x_i \in \mathcal{T}_M^{\mathbb{C}}$ in the following way. If f is a function on U and $df = \sum_i a_i dx_i$, then $(\partial/\partial x_i)f = a_i$.

Let Q be a subbundle of a complex vector bundle E on M . We denote by Q^{\perp} the subbundle of E^* , the complex dual to E , orthogonal to Q . If sections e_i form a local frame in Q , we set $(e_i)^{\perp} = Q^{\perp}$.

A (complex) distribution on a manifold M is a subbundle of $\mathcal{T}_M^{\mathbb{C}}$.

Definition 2.1. A distribution \mathcal{P} is said to be *integrable* if, locally on M , there exist (complex valued) functions f_1, \dots, f_k such that df_1, \dots, df_k give a local frame in \mathcal{P}^{\perp} , i.e. df_i are linearly independent at each point and $\mathcal{P} = (df_i)^{\perp}$.

An integrable distribution \mathcal{P} is obviously involutive, i.e. $[\mathcal{P}, \mathcal{P}] \subset \mathcal{P}$.

Let \mathcal{P} be an integrable distribution on M . We will denote by $\mathcal{O}_{\mathcal{P}}$ the sheaf of functions on M constant along \mathcal{P} , i.e. $f \in \mathcal{O}_{\mathcal{P}}$ if and only if $Xf = 0$ for any vector field $X \in \mathcal{P}$.

2.1. The Kostant–Hochschild–Rosenberg theorem in presence of a distribution

Let M be a smooth manifold. Let \mathcal{D}^n be the sheaf of n -differential operators on M and \mathcal{D}^{\bullet} the corresponding Hochschild complex with differential d . Let $\wedge^{\bullet}\mathcal{T}$ be the complex of sheaves of polyvector fields on M with zero differential, $\mathcal{T} = \mathcal{T}_M^{\mathbb{C}}$.

There is the following “smooth” version of the Kostant–Hochschild–Rosenberg theorem [16, Theorem 4.6.1.1.].

Proposition 2.2. *The natural embedding*

$$\wedge^{\bullet}\mathcal{T} \rightarrow \mathcal{D}^{\bullet} \tag{2.1}$$

is a quasiisomorphism of complexes. Moreover, if $\varphi \in \mathcal{D}^n$ is a Hochschild cocycle, then its alternation $\text{Alt}(\varphi)$ is a polyvector field of $\wedge^{\bullet}\mathcal{T}$ cohomological to φ .

Proof. Arguments of this proof will be used also in proving the next proposition. The proposition is local on M , so it is enough to prove it replacing M by an open set $U \subset M$ having complex coordinates x_i , $i = 1, \dots, \dim M$. Any differential operator on U may be uniquely presented as a polynomial in $\partial/\partial x_i$ with coefficients being smooth functions on U . Hence, \mathcal{D}^{\bullet} coincides over U with the complex $\mathcal{C}^{\bullet}(\mathcal{T})$. Here, for any vector bundle E , we denote by $\mathcal{C}^{\bullet}(E)$ the complex $(\otimes^{\bullet}\text{Sym}(E), d)$ with differential of the form

$$d : \otimes^n \text{Sym}(E) \rightarrow \otimes^{n+1} \text{Sym}(E),$$

$$\begin{aligned} d(a_1 \otimes \cdots \otimes a_n) &= 1 \otimes a_1 \otimes \cdots \otimes a_n + \sum_{i=1}^n (-1)^i a_1 \otimes \cdots \otimes \Delta a_i \otimes \cdots \otimes a_n \\ &\quad + (-1)^{n+1} a_1 \otimes \cdots \otimes a_n, \end{aligned}$$

where Δ is the comultiplication in the symmetric algebra $\text{Sym}(E)$ generated by the rule $\Delta(a) = a \otimes 1 + 1 \otimes a$ for $a \in E$.

One has the following well-known statement (see, for example, the proof of Theorem 4.6.1.1. in [16]).

Lemma 2.3. *Let E be a line bundle over M . Then the conclusion of Proposition 2.2 holds for the map*

$$\wedge^\bullet E \rightarrow \mathcal{C}^\bullet(E). \quad (2.2)$$

Applying this lemma to $E = \mathcal{T}$ we prove the proposition.

Let (M, \mathcal{P}) be a smooth manifold with integrable distribution. We call an n -chain $v \in \mathcal{D}^n$ *polarized* if $v(a_1, \dots, a_n) = 0$ whenever $a_1, \dots, a_n \in \mathcal{O}_{\mathcal{P}}$. We call v *strongly polarized*, if $v(a_1, \dots, a_n) = 0$ whenever $a_1, \dots, a_{n-1} \in \mathcal{O}_{\mathcal{P}}$ and $a_n \in \mathcal{C}_M^\infty$.

Proposition 2.4. *Let $v \in \mathcal{D}^2$ be a polarized Hochschild 2-cochain such that dv is strongly polarized. Then, there exists a polarized differential operator b such that $v + db$ is strongly polarized.*

Proof. Since subsheaves of polarized and strongly polarized cochains are subsheaves of \mathcal{C}_M^∞ -modules, it is enough to prove the proposition locally on M . Thus, let U be an open set with coordinates $x_i, i = 1, \dots, \dim M$ such that $\partial/\partial x_i, i = 1, \dots, k$, form a local frame in \mathcal{P} . Let \mathcal{Q} be the subbundle of \mathcal{T} generated by $\partial/\partial x_i, i = k+1, \dots, n$. Thus, $\mathcal{T} = \mathcal{Q} \oplus \mathcal{P}$ on U . There is the natural isomorphism of complex $\mathcal{C}^\bullet(\mathcal{T})$ with the tensor product of complexes $\mathcal{C}^\bullet(\mathcal{Q}, \mathcal{P}) = \mathcal{C}^\bullet(\mathcal{Q}) \otimes \mathcal{C}^\bullet(\mathcal{P})$, so that we can identify $\mathcal{C}^\bullet(\mathcal{T})$ with $\mathcal{C}^\bullet(\mathcal{Q}, \mathcal{P})$. Similarly, we identify the complex $\wedge^\bullet(\mathcal{T})$ with $\wedge^\bullet(\mathcal{Q}, \mathcal{P}) = \wedge^\bullet(\mathcal{Q}) \otimes \wedge^\bullet(\mathcal{P})$.

Thus, the map (2.2) generates the map of complexes

$$\wedge^\bullet(\mathcal{Q}, \mathcal{P}) \rightarrow \mathcal{C}^\bullet(\mathcal{Q}, \mathcal{P}). \quad (2.3)$$

Complex $\mathcal{C}^\bullet(\mathcal{Q}, \mathcal{P})$ decomposes obviously into a direct sum of subcomplexes $\mathcal{C}_{k,l}^\bullet(\mathcal{Q}, \mathcal{P})$, where $\mathcal{C}_{k,l}^\bullet(\mathcal{Q}, \mathcal{P})$ consists of elements of total degree k with respect to \mathcal{P} and l with respect to \mathcal{Q} . The same is true for $\Lambda^\bullet(\mathcal{Q}, \mathcal{P})$. Due to Proposition 2.2, the map (2.3) is a quasiisomorphism of bigraded complexes.

It is clear that an element of $\mathcal{C}^n(\mathcal{Q}, \mathcal{P})$ is polarized if it is a sum of tensor monomials having degree > 0 with respect to \mathcal{P} . An element of $\mathcal{C}^n(\mathcal{Q}, \mathcal{P})$ is strongly polarized if it is a sum of tensor monomials of the form $a_1 \otimes \dots \otimes a_n$, where $a_1 \otimes \dots \otimes a_{n-1}$ is polarized.

Thus, we may suppose that the given v is a polarized Hochschild cochain on U belonging to $\mathcal{C}^2(\mathcal{Q}, \mathcal{P})$. It can be written as $v = v_0 + v_1$, where $v_0 = \sum(a_i \otimes c_i)$, the sum of all tensor monomials in v such that $a_i \in \mathcal{C}^1(\mathcal{Q}), c_i \in \mathcal{C}^1(\mathcal{P})$. Let us set $b = \sum a_i c_i$. It is clear that $v' = v + db$ does not contain tensor monomials of that type. The proposition will be proved if we show that v' is strongly polarized. Let us prove this.

Suppose $v' = v'_0 + v'_1$, where v'_0 is not strongly polarized and v'_1 is strongly polarized. Then v'_0 has the form

$$v'_0 = \sum(a_i \otimes b_i)(1 \otimes c_i), \quad (2.4)$$

where $(a_i \otimes b_i) \in \otimes^2 \text{Sym}(\mathcal{Q})$, and elements $c_i \in \text{Sym}(\mathcal{P})$ are linearly independent. Besides, b_i are of degree > 0 .

The element $d\nu = d\nu' = dv'_0 + dv'_1$ is strongly polarized by hypothesis of the proposition, dv'_1 being the coboundary of a strongly polarized element ν'_1 is strongly polarized, too. All summands of dv'_0 with first two factors being of degree 0 with respect to \mathcal{P} are $(d(a_i \otimes b_i) + a_i \otimes b_i \otimes 1)(1 \otimes 1 \otimes c_i)$. These summands are not strongly polarized. Hence, $\sum_i (d(a_i \otimes b_i) + a_i \otimes b_i \otimes 1)(1 \otimes 1 \otimes c_i) = 0$. Since elements $1 \otimes 1 \otimes c_i$ are linearly independent, it follows that $d(a_i \otimes b_i) = -a_i \otimes b_i \otimes 1$ for all i , which is only possible if $a_i \otimes b_i = 0$ for all i . Equality (2.4) implies that $v'_0 = 0$. Therefore, ν' is equal to ν'_1 which is strongly polarized. \square

2.2. Differential operators in presence of a distribution

Let (M, \mathcal{P}) be a smooth manifold with integrable distribution. Let z_i, y_j be complex coordinates on an open set $U \subset M$ such that $\mathcal{P} = (dz_i)^\perp$. Vector fields $\partial/\partial y_j$ form a local frame in \mathcal{P} , since, by definition, $\mathcal{O}_{\mathcal{P}}$ consists of functions $a \in \mathcal{C}_M^\infty$ such that da has the form $\sum_i a_i dz_i$. Since da is closed, $\partial a_i / \partial y_j = 0$ for all i, j , which implies that $a_i = \partial a / \partial z_i \in \mathcal{O}_{\mathcal{P}}$ for all i .

Let \mathcal{Q} be a subbundle in $\mathcal{T} = \mathcal{T}_M^\mathbb{C}$ generated by $\partial/\partial z_j$, so that $\mathcal{T} = \mathcal{Q} \oplus \mathcal{P}$ on U .

The vector bundle \mathcal{T}/\mathcal{P} may be considered as the sheaf of derivations from $\mathcal{O}_{\mathcal{P}}$ to \mathcal{C}_M^∞ , $\text{Der}(\mathcal{O}_{\mathcal{P}}, \mathcal{C}_M^\infty)$. Locally, such derivations can be presented in the form $\sum b_i \partial/\partial z_i$, $b_i \in \mathcal{C}_M^\infty$, i.e. as sections of \mathcal{Q} . Denote by $\text{Der}(\mathcal{O}_{\mathcal{P}})$ the $\mathcal{O}_{\mathcal{P}}$ -submodule of $\text{Der}(\mathcal{O}_{\mathcal{P}}, \mathcal{C}_M^\infty)$ consisting of operators which take $\mathcal{O}_{\mathcal{P}}$ to itself. It is clear that $\text{Der}(\mathcal{O}_{\mathcal{P}}, \mathcal{C}_M^\infty) = \mathcal{C}_M^\infty \otimes_{\mathcal{O}_{\mathcal{P}}} \text{Der}(\mathcal{O}_{\mathcal{P}})$. Locally, elements of $\text{Der}(\mathcal{O}_{\mathcal{P}})$ have the form $\sum_i a_i \partial/\partial z_i$, $a_i \in \mathcal{O}_{\mathcal{P}}$.

Let $\wedge^\bullet(\mathcal{T}/\mathcal{P})$ denote the complex of sheaves of polyvector fields on M from $\mathcal{O}_{\mathcal{P}}$ to \mathcal{C}_M^∞ .

Let $\mathcal{D}^\bullet(\mathcal{O}_{\mathcal{P}}, \mathcal{C}_M^\infty)$ denote the restriction of the Hochschild complex \mathcal{D}^\bullet to $\mathcal{O}_{\mathcal{P}}$. Hence, the sheaf $\mathcal{D}^n(\mathcal{O}_{\mathcal{P}}, \mathcal{C}_M^\infty)$ may be considered as the sheaf of n -differential operators from $\mathcal{O}_{\mathcal{P}}$ to \mathcal{C}_M^∞ . Locally, elements of $\mathcal{D}^1(\mathcal{O}_{\mathcal{P}}, \mathcal{C}_M^\infty)$, the sheaf of differential operators from $\mathcal{O}_{\mathcal{P}}$ to \mathcal{C}_M^∞ , can be presented as polynomials in $\partial/\partial z_i$ with smooth coefficients. Therefore, locally on M , complex $\mathcal{D}^\bullet(\mathcal{O}_{\mathcal{P}}, \mathcal{C}_M^\infty)$ is isomorphic to the complex $\mathcal{C}^\bullet(\mathcal{Q})$ (see previous subsection).

We will need the following version of the Kostant-Hochschild-Rosenberg theorem.

Proposition 2.5. *The natural embedding*

$$\wedge^\bullet(\mathcal{T}/\mathcal{P}) \rightarrow \mathcal{D}^\bullet(\mathcal{O}_{\mathcal{P}}, \mathcal{C}_M^\infty) \quad (2.5)$$

is a quasiisomorphism of complexes. Moreover, if $\varphi \in \mathcal{D}^n(\mathcal{O}_{\mathcal{P}}, \mathcal{C}_M^\infty)$ is a cocycle, then $\text{Alt}(\varphi)$ is a polyvector field of $\wedge^n(\mathcal{T}/\mathcal{P})$ cohomological to φ .

Proof. As follows from above, embedding (2.5) is locally isomorphic to the embedding

$$\wedge^\bullet \mathcal{Q} \rightarrow \mathcal{C}^\bullet(\mathcal{Q}).$$

Now the proposition follows from Lemma 2.3 when $E = \mathcal{Q}$. \square

Remark. All conclusions of Propositions 2.2, 2.4 and 2.5 remain true for global sections of the corresponding sheaves, since they are sheaves of \mathcal{C}_M^∞ -modules.

2.3. Differential forms in presence of a distribution

Let (M, \mathcal{P}) be a smooth manifold with integrable distribution.

The sheaf $\mathcal{P}^\perp = (\mathcal{T}/\mathcal{P})^*$ of differential forms on M which being applied to vector fields from \mathcal{P} give zero, may be written as $\mathcal{C}_M^\infty d\mathcal{O}_{\mathcal{P}}$.

Denote $\Omega_{\mathcal{O}_{\mathcal{P}}}^1 = \text{Hom}_{\mathcal{O}_{\mathcal{P}}}(\text{Der}(\mathcal{O}_{\mathcal{P}}), \mathcal{O}_{\mathcal{P}})$, the sheaf of 1-forms on $\mathcal{O}_{\mathcal{P}}$. It is clear that $\Omega_{\mathcal{O}_{\mathcal{P}}}^1 = \mathcal{O}_{\mathcal{P}} d\mathcal{O}_{\mathcal{P}}$.

Denote by $\Omega_{\mathcal{O}_{\mathcal{P}}}^{1\text{cl}}$ the subsheaf of closed forms of $\Omega_{\mathcal{O}_{\mathcal{P}}}^1$.

Lemma 2.6.

(a) *The sequence of sheaves*

$$0 \rightarrow \mathbb{C} \rightarrow \mathcal{O}_{\mathcal{P}} \xrightarrow{d} \Omega_{\mathcal{O}_{\mathcal{P}}}^{1\text{cl}} \rightarrow 0 \quad (2.6)$$

is exact.

(b) *The sequence of sheaves*

$$0 \rightarrow \Omega_{\mathcal{O}_{\mathcal{P}}}^{1\text{cl}} \rightarrow \mathcal{P}^\perp \xrightarrow{d} d\mathcal{P}^\perp \rightarrow 0 \quad (2.7)$$

is exact.

Proof. Let functions z_i, y_j form a local basis on M and $\mathcal{P} = (dz_i)^\perp$. Let us prove (a). It is sufficient to establish the exactness at the third term of (2.6). Let $\alpha = \sum_i a_i dz_i \in \Omega_{\mathcal{O}_{\mathcal{P}}}^{1\text{cl}}$. Since α is closed, there exists, locally, $f \in \mathcal{C}_M^\infty$ such that $df = \alpha$. Since α does not contain terms of the form $g dy_j$, one has $\partial f / \partial y_j = 0$ for all j . Hence, $f \in \mathcal{O}_{\mathcal{P}}$.

To prove (b), it is sufficient to establish the exactness at \mathcal{P}^\perp . To this end, suppose $\beta = \sum_i b_i dz_i \in \mathcal{P}^\perp$ and $d\beta = 0$. Since β is closed, $\partial b_i / \partial y_j = 0$ for all i, j . This means that all $b_i \in \mathcal{O}_{\mathcal{P}}$. Thus, $\beta \in \Omega_{\mathcal{O}_{\mathcal{P}}}^1$ and closed, i.e. $\beta \in \Omega_{\mathcal{O}_{\mathcal{P}}}^{1\text{cl}}$. \square

Proposition 2.7. *Let (M, \mathcal{P}) be a smooth manifold with an integrable distribution. Then, there is a natural isomorphism*

$$H^1(M, \Omega_{\mathcal{O}_{\mathcal{P}}}^{1\text{cl}}) \simeq \frac{\Gamma(M, d\mathcal{P}^\perp)}{d(\Gamma(M, \mathcal{P}^\perp))}. \quad (2.8)$$

Proof. This is an immediate consequence of the cohomological exact sequence for (2.7) and of $H^i(M, \mathcal{P}^\perp) = 0$ for $i > 0$. \square

3. \mathbb{C} -symplectic manifolds and their polarizations

3.1. \mathbb{C} -symplectic manifolds

Definition 3.1. By a \mathbb{C} -symplectic manifold we mean a pair (M, ω) , where M is a smooth manifold and ω a closed nondegenerate complex 2-form on M satisfying the following condition: each point of M has a neighborhood U and complex coordinates x_i, y_i ,

$i = 1, \dots, (1/2) \dim M$, on U such that the form ω on U can be presented as

$$\omega = \sum_i dy_i \wedge dx_i. \quad (3.1)$$

If the form ω is real, such a presentation is possible by the Darboux theorem. One has

$$\{x_i, x_j\} = \{y_i, y_j\} = 0, \quad \{y_i, x_j\} = \delta_{ij} \quad (3.2)$$

for all i, j , where $\{\cdot, \cdot\}$ is the Poisson bracket inverse to ω . We call such functions x_i, y_i *Darboux coordinates* with respect to ω (or $\{\cdot, \cdot\}$).

In what follows we only deal with \mathbb{C} -symplectic manifolds, so we simply call them symplectic ones.

3.2. Polarization

Definition 3.2. Let (M, ω) be a symplectic manifold. We call a (complex) distribution \mathcal{P} on M a polarization of ω , if, locally, there exist (complex) Darboux coordinates, x_i, y_i , with respect to ω such that $\mathcal{P} = (dx_i)^\perp$, i.e. $x_i \in \mathcal{O}_\mathcal{P}$.

We call the triple (M, ω, \mathcal{P}) a PSM.

It follows that a polarization of ω is, in particular, an integrable distribution and a Lagrangian subbundle with respect to ω .

Proposition 3.3. Let (M, ω, \mathcal{P}) be a PSM. Then $\omega \in \Gamma(M, d\mathcal{P}^\perp)$.

Proof. Let x_i, y_i be local Darboux coordinates on M such that $\mathcal{P} = (dx_i)^\perp$ and $\omega = \sum dy_i \wedge dx_i$. Then, locally, $\omega = d(\sum y_i dx_i)$ and $\sum y_i dx_i \in \mathcal{P}^\perp$. \square

Proposition 3.4. Let (M, ω, \mathcal{P}) be a PSM. Then, $\mathcal{O}_\mathcal{P}$ is a maximal commutative Lie subalgebra in \mathcal{C}_M^∞ with respect to the Poisson bracket ω^{-1} .

Proof. It follows from **Definition 3.2** that, locally, the bracket ω^{-1} can be written in the form

$$\{\cdot, \cdot\} = \sum_i \bar{\partial}_i \wedge \partial_i, \quad (3.3)$$

where $\partial_i = \{y_i, \cdot\}$, $\bar{\partial}_i = \{\cdot, x_i\}$. The module $\mathcal{O}_\mathcal{P}$ consists, locally, of elements $g \in \mathcal{C}_M^\infty$ such that $\{g, x_i\} = \bar{\partial}_i g = 0$ for all i . Putting two such elements g_1, g_2 in (3.3), we obtain that $\{g_1, g_2\} = 0$. Therefore, $\mathcal{O}_\mathcal{P}$ is commutative. The maximality of $\mathcal{O}_\mathcal{P}$ is obvious. Indeed, if $a \in \mathcal{C}_M^\infty$ commutes with $\mathcal{O}_\mathcal{P}$, then, in particular, $\{x_i, a\} = 0$ for all i , hence $a \in \mathcal{O}_\mathcal{P}$. \square

4. Deformations of a PSM

4.1. Formal everything

Let t be a formal parameter. We will consider on M formal functions, formal vector fields, formal forms, etc., which are elements of $\mathcal{C}_M^\infty[[t]]$, $\mathcal{T}_M^\mathbb{C}[[t]]$, $\wedge^k(\mathcal{T}_M^\mathbb{C})^*[[t]]$, etc. In

the formal case, all sheaves on M and their morphisms will be sheaves and morphisms of $\mathbb{C}[[t]]$ -modules.

Let \mathcal{B} be a sheaf on M . We call the map $\sigma : \mathcal{B}[[t]] \rightarrow \mathcal{B}$, $b_0 + tb_1 + \dots \mapsto b_0$, the *symbol map*. For a subsheaf $\mathcal{F}_t \subset \mathcal{B}[[t]]$, we denote $\mathcal{F}_0 = \sigma(\mathcal{F}_t) \subset \mathcal{B}$.

Let \mathcal{F}_t be a subsheaf of $\mathcal{B}[[t]]$. We call \mathcal{F}_t a *t-regular* subsheaf if it is complete in t -adic topology, and $tb \in \mathcal{F}_t$, $b \in \mathcal{B}[[t]]$ implies $b \in \mathcal{F}_t$.

For a *t-regular* subsheaf $\mathcal{F}_t \subset \mathcal{B}[[t]]$ the natural map $\mathcal{F}_t/t\mathcal{F}_t \rightarrow \mathcal{F}_0$ is an isomorphism. In this case we call \mathcal{F}_t a *deformation* of \mathcal{F}_0 .

Let \mathcal{B} be a vector bundle on M . Then, a subsheaf $\mathcal{F}_t \subset \mathcal{B}[[t]]$ is called a (formal) subbundle, if it is a *t-regular* subsheaf of $\mathcal{C}_M^\infty[[t]]$ -modules.

All notions and statements above carry over to the formal case.

For example, a deformation of a distribution \mathcal{P} on M is a subbundle, \mathcal{P}_t , of $\mathcal{T}_M^\mathbb{C}[[t]]$ such that $\mathcal{P}_0 = \mathcal{P}$. A system of (formal) coordinates on an open set $U \subset M$ is a set of formal functions $x_i = x_{i0} + tx_{i1} + \dots$, $i = 1, \dots, \dim M$, $\mathbb{C}[[t]]$ -linearly independent at each point of U . A formal PSM is a triple $(M, \omega_t, \mathcal{P}_t)$, where ω_t is a formal symplectic form on M (i.e. a closed 2-form of the form $\omega_t = \omega_0 + t\omega_1 + \dots$ with nondegenerate ω_0) and \mathcal{P}_t is a polarization of ω_t in sense of [Definition 3.2](#), i.e. locally, there exist formal Darboux coordinates x_i, y_i with respect to ω_t such that $\mathcal{P}_t = (\mathrm{d}x_i)^\perp$.

We say that $(M, \omega_t, \mathcal{P}_t)$ is a deformation of a PSM (M, ω, \mathcal{P}) , if $(M, \omega_0, \mathcal{P}_0) = (M, \omega, \mathcal{P})$.

The following proposition, which follows from [Propositions 6.1](#) and [6.2](#), shows that any deformation of (polarized) symplectic manifold is a formal (polarized) symplectic manifold.

Proposition 4.1.

- (a) Let ω_t be a formal closed 2-form. If ω_0 admits, locally, Darboux coordinates, then there exist their lifts being formal Darboux coordinates for ω_t .
- (b) Let \mathcal{P}_t be an integrable distribution and Lagrangian subbundle of $\mathcal{T}_M^\mathbb{C}[[t]]$ with respect to ω_t . Let \mathcal{P}_0 be a polarization of ω_0 . Then \mathcal{P}_t is a polarization of ω_t .

4.2. Local structure of deformed polarizations

It is clear that formal vector fields $tX_1 + t^2X_2 + \dots \in t\mathcal{T}_M^\mathbb{C}[[t]]$ form a sheaf of pro-nilpotent Lie algebras. Therefore, elements e^{tX} , $X \in \mathcal{T}_M^\mathbb{C}[[t]]$, form a sheaf of pro-unipotent Lie groups of formal automorphisms of M .

Let x_i be formal coordinates on U and a_i , $i = 1, \dots, \dim M$, arbitrary formal functions on U . Then there exists a derivation, D , of $\mathcal{C}_M^\infty[[t]]$ that takes x_i to a_i . Such a derivation is $D = \sum_i a_i \partial/\partial x_i$. This implies the following lemma.

Lemma 4.2. *Let x_i, x'_i , $i = 1, \dots, \dim M$, be two systems of formal coordinates on an open set $U \subset M$, and $x_i = x'_i \bmod t$. Then, there exists a formal automorphism on U that takes x_i to x'_i .*

Proposition 4.3.

- (a) Let \mathcal{P}_t be an integrable distribution on M that is a deformation of a distribution \mathcal{P} . Then, locally, there exists a formal vector field, X , such that e^{tX} gives an isomorphism \mathcal{P}_t with $\mathcal{P}[[t]]$.

- (b) Let $(M, \omega_t, \mathcal{P}_t)$ be a deformation of a PSM (M, ω, \mathcal{P}) . Then, each point of M has a neighborhood, U , and a formal vector field X on U such that e^{tX} gives an isomorphism of $(M, \omega_t, \mathcal{P}_t)|_U$ with the trivial deformation $(U, \omega, \mathcal{P}[[t]])$.

Proof.

- (a) Locally, there exist functions $x_{it} = x_{i0} + tx_{i1} + \dots, i = 1, \dots, k$, such that $\mathcal{P}_t = (dx_{it})^\perp$ and hence $\mathcal{P} = (dx_{i0})^\perp$. Let us add functions $x_{(k+1)0}, \dots, x_{n0}$ in such a way that all $x_{i0}, i = 1, \dots, n$, form a coordinate system. According to Lemma 4.2, there exists a formal automorphism which takes coordinates $x_{it}, i = 1, \dots, k, x_{j0}, j = k+1, \dots, n$, to coordinates $x_{i0}, i = 1, \dots, n$. This formal automorphism gives obviously an isomorphism of \mathcal{P}_t onto $\mathcal{P}[[t]]$.
- (b) Let U be an open set in M where Darboux coordinates $x_{it} = x_{i0} + tx_{i1} + \dots, y_{it} = y_{i0} + tx_{i1} + \dots, i = 1, \dots, (1/2)\dim M$, exist, and $\mathcal{P}_t = (dx_{it})^\perp$. Then, x_{i0}, y_{i0} are Darboux coordinates with respect to $\omega_0 = \omega$ such that $\mathcal{P} = (dx_{i0})^\perp$. By Lemma 4.2, there exists a formal vector field X on U such that the formal automorphism e^{tX} takes coordinates x_{it}, y_{it} to x_{i0}, y_{i0} . This e^{tX} satisfies the conclusion of the proposition. \square

Let \mathcal{P} be an integrable distribution. Denote by $\mathcal{O}_{\mathcal{P}}$ the sheaf of functions constant along \mathcal{P} . Let \mathcal{P}_t be a deformation of \mathcal{P} . It follows from the previous proposition that, locally, the pair $(\mathcal{C}_M^\infty[[t]], \mathcal{O}_{\mathcal{P}_t})$ is isomorphic to the pair $(\mathcal{C}_M^\infty[[t]], \mathcal{O}_{\mathcal{P}}[[t]])$, hence $\mathcal{O}_{\mathcal{P}_t}$ is a t -regular subalgebra of $\mathcal{C}_M^\infty[[t]]$. One holds the following inverse statement.

Proposition 4.4. *Let \mathcal{P} be an integrable distribution on M . Let \mathcal{O}_t be a t -regular subalgebra of $\mathcal{C}_M^\infty[[t]]$ such that $\mathcal{O}_0 = \mathcal{O}_{\mathcal{P}}$. Then, there exists a deformation, \mathcal{P}_t , of \mathcal{P} such that $\mathcal{O}_t = \mathcal{O}_{\mathcal{P}_t}$.*

Proof. Let us prove that $d\mathcal{O}_t \subset (\mathcal{T}_M^\mathbb{C})^*[[t]]$ is t -regular. Let $b \in (\mathcal{T}_M^\mathbb{C})^*[[t]]$ and $tb \in d\mathcal{O}_t$. Then, there exists $a = a_0 + ta' \in \mathcal{O}_t$ such that $da = tb$. It follows that a_0 is a constant, so $ta' \in \mathcal{O}_t$. Since \mathcal{O}_t is t -regular, $a' \in \mathcal{O}_t$, too. Therefore, $b = da' \in d\mathcal{O}_t$, so that $d\mathcal{O}_t$ is a t -regular submodule in $(\mathcal{T}_M^\mathbb{C})^*[[t]]$. This implies that $\mathcal{C}_M^\infty[[t]]d\mathcal{O}_t$ is a subbundle in $(\mathcal{T}_M^\mathbb{C})^*[[t]]$, so $\mathcal{P}_t = (d\mathcal{O}_t)^\perp$ is a subbundle of $\mathcal{T}_M^\mathbb{C}[[t]]$. Moreover, $\mathcal{P}_0 = \mathcal{P}$. Therefore, \mathcal{P}_t is a deformation of \mathcal{P} . One has $\mathcal{O}_t \subset \mathcal{O}_{\mathcal{P}_t}$. Since these two subalgebras are t -regular and coincide at $t = 0$, one has $\mathcal{O}_t = \mathcal{O}_{\mathcal{P}_t}$. \square

The last proposition gives a one-to-one correspondence between deformations of an integrable distribution \mathcal{P} and deformations of $\mathcal{O}_{\mathcal{P}}$.

4.3. Action of formal automorphisms on a symplectic form

Let (M, ω_t) be a formal symplectic manifold, $\mathcal{T} = \mathcal{T}_M^\mathbb{C}$.

The well-known formula for the Lie derivative

$$L_X = i(X)d + di(X)$$

implies

$$L_X \omega_t = d\alpha(X), \quad (4.1)$$

where $\alpha : \mathcal{T}[[t]] \rightarrow \mathcal{T}^*[[t]]$ is a map defined by $X \mapsto \omega_t(X, \cdot)$. Since α is an isomorphism, we have the following lemma.

Lemma 4.5. *The orbit of ω_t by the action of the group of formal automorphisms e^{tX} , $X \in \Gamma(M, \mathcal{T}[[t]])$, is $\omega_t + t d(\Gamma(M, \mathcal{T}^*[[t]]))$.*

The lemma shows that the orbit of ω_t corresponds to the cohomology class of ω_t in $\omega_0 + t H^2(M, \mathbb{C}[[t]])$.

Let $(M, \omega_t, \mathcal{P}_t)$ be a formal PSM and $X \in \Gamma(M, \mathcal{P}_t)$. Since $\alpha(\mathcal{P}_t) = \mathcal{P}_t^\perp$, one has $L_X \omega_t = d\alpha(X) \in \Gamma(M, d\mathcal{P}_t^\perp)$. The argument as above implies.

Lemma 4.6. *By the action of the group generated by e^{tX} , $X \in \Gamma(M, \mathcal{P}_t)$, the orbit of ω_t is $\omega_t + t d(\Gamma(M, \mathcal{P}_t^\perp))$.*

The lemma shows that the orbit of ω_t corresponds to the cohomology class of ω_t in $\Gamma(M, d\mathcal{P}_t^\perp)/d(\Gamma(M, \mathcal{P}_t^\perp))$.

5. Polarized symplectic connection and characteristic class of a PSM

5.1. Polarized symplectic connection

Let (M, ω, \mathcal{P}) be a (formal) PSM. Denote, for shortness, $\mathcal{T} = \mathcal{T}_M^\mathbb{C}[[t]]$.

Definition 5.1. We call a connection, ∇ , on M a \mathcal{P} -symplectic connection if

- (a) it preserves ω and is torsion free, i.e. is a symplectic connection;
- (b) it preserves \mathcal{P} , i.e. $\nabla_X(\mathcal{P}) \subset \mathcal{P}$ for any $X \in \mathcal{T}$;
- (c) it is flat on \mathcal{P} along \mathcal{P} , i.e. for any $X, Y \in \mathcal{P}$ one has $(\nabla_X \nabla_Y - \nabla_Y \nabla_X - \nabla_{[X, Y]})(\mathcal{P}) = 0$.

Proposition 5.2. *Let (M, ω, \mathcal{P}) be a (formal) PSM. Then, there exists a \mathcal{P} -symplectic connection on M .*

Proof. Let functions $a_1, \dots, a_{2n}, 2n = \dim M$, form local Darboux coordinates on an open set $U \subset M$ and be such that $a_i \in \mathcal{O}_{\mathcal{P}}$ for $i = 1, \dots, n$. Let $X_i = X_{a_i}$ be the corresponding Hamiltonian vector fields. Then, vector fields $X_i, i = 1, \dots, n$, form a local frame in \mathcal{P} . Also, all X_i commute and form a local frame in \mathcal{T} . Let ∇ be the standard flat connection on U associated with coordinates a_i . This connection is defined on U by the rule $\nabla_{X_i} X_j = 0$.

It is easy to see that ∇ is a \mathcal{P} -symplectic connection on U . Moreover, since $X_f \in \mathcal{P}$ is equivalent to $df \in \mathcal{P}^\perp$, the connection ∇ satisfies the following property for Hamiltonian vector fields:

$$\nabla_{X_f} X_g = 0 \quad \text{for } X_f, Y_g \in \mathcal{P}. \quad (5.1)$$

Now, let us prove the existence of a global connection.

Let $\{U_\alpha\}$ is an open covering of M such that on each U_α there is a \mathcal{P} -symplectic connection ∇_α as above. Then $\nabla_\alpha - \nabla_\beta$ defined on $U_\alpha \cap U_\beta$ form a Čech cocycle $\psi_{\alpha,\beta} \in \text{Hom}_{C_M^\infty[[t]]}(\mathcal{T} \otimes \mathcal{T}, \mathcal{T})$, $\psi_{\alpha,\beta}(X, Y) = \nabla_\alpha X Y - \nabla_\beta X Y$.

Elements $\psi_{\alpha,\beta}$ satisfy the following properties. As follows from (5.1),

$$\psi_{\alpha,\beta}(X, Y) = 0 \quad \text{for } X, Y \in \mathcal{P}. \quad (5.2)$$

Since all ∇_α are torsion free, $\psi_{\alpha,\beta}$ are symmetric. Since all ∇_α preserve \mathcal{P} , one has $\psi_{\alpha,\beta}(X, Y) \in \mathcal{P}$ for $Y \in \mathcal{P}$.

In addition, $\psi_{\alpha,\beta}$ considered as elements from $\text{Hom}(\mathcal{T}, \text{Hom}(\mathcal{T}, \mathcal{T}))$, $X \mapsto \psi_{\alpha,\beta}(X, \cdot)$, belong to $\text{Hom}(\mathcal{T}, \mathfrak{sp}(\mathcal{T}))$, where $\mathfrak{sp}(\mathcal{T})$ consists of endomorphisms of \mathcal{T} preserving ω .

Since all the properties above are $C_M^\infty[[t]]$ -linear, one can find tensors $\psi_\alpha \in \text{Hom}(\mathcal{T} \otimes \mathcal{T}, \mathcal{T})$ satisfying all of them and such that $\psi_\alpha - \psi_\beta = \psi_{\alpha,\beta}$. Then $\nabla = \nabla_\alpha - \psi_\alpha = \nabla_\beta - \psi_\beta$ is a globally defined connection. Flatness of ∇ on \mathcal{P} along \mathcal{P} follows from the fact that for all α tensors ψ_α satisfy property (5.2), i.e. $\psi_\alpha(X, Y) = 0$ for $X, Y \in \mathcal{P}$. Also, ∇ is torsion free because all ψ_α are symmetric. Therefore, ∇ satisfies the proposition. \square

5.2. Characteristic class of a PSM

Let ∇ be a \mathcal{P}_t -symplectic connection on a formal PSM $(M, \omega_t, \mathcal{P}_t)$. Let us denote by $\nabla^2|_{\mathcal{P}_t}$ the curvature of ∇ restricted to \mathcal{P}_t . Then, $\text{tr}(\nabla^2|_{\mathcal{P}_t})$ is a closed 2-form on M representing, up to a constant factor, the first Chern class of \mathcal{P}_0 .

Lemma 5.3. *Let ∇ be a \mathcal{P}_t -symplectic connection. Then $\text{tr}(\nabla^2|_{\mathcal{P}_t})$ belongs to $\Gamma(M, d\mathcal{P}_t^\perp)$. If ∇_1 is another \mathcal{P}_t -symplectic connection, then $\text{tr}(\nabla_1^2|_{\mathcal{P}_t})$ differs from $\text{tr}(\nabla^2|_{\mathcal{P}_t})$ by an element of $d(\Gamma(M, \mathcal{P}_t^\perp))$.*

Proof. Follows from flatness of \mathcal{P}_t along \mathcal{P}_t with respect to the connections. \square

The lemma allows us to consider the element of $\Gamma(M, d\mathcal{P}_t^\perp)/d(\Gamma(M, \mathcal{P}_t^\perp))$ represented by the form $\text{tr}(\nabla^2|_{\mathcal{P}_t})$, where ∇ is a \mathcal{P}_t -symplectic connection, as a characteristic class of the PSM $(M, \omega_t, \mathcal{P}_t)$.

Due to Proposition 2.7, one can also consider this class as an element of $H^1(M, \Omega_{\mathcal{O}_{\mathcal{P}_t}}^{\text{cl}})$.

6. Deformations of Poisson brackets

In this section, we prove three technical statements which we use through the paper.

Let $\pi_0 = \{\cdot, \cdot\}$ be a nondegenerate Poisson bracket on a smooth manifold M of dimension $2n$. We say that a formal sum $\pi_t = \pi_0 + t\pi_1 + \dots$ is a deformation of π_0 if all π_i are *bidifferential operators* on M and π_t defines a Lie algebra structure on the sheaf $C_M^\infty[[t]]$. We will also denote π_t by $[\cdot, \cdot]$. Let us recall the symbol map $\sigma : C_M^\infty[[t]] \rightarrow C_M^\infty$, $a = a_0 + ta_1 + \dots \mapsto a_0$. We call a a lift of a_0 . We say that functions $\hat{x}_i, \hat{\xi}_i, i = 1, \dots, n$, on an open set $U \subset M$ form Darboux coordinates with respect to $[\cdot, \cdot]$, if $[\hat{x}_j, \hat{x}_k] = [\hat{\xi}_j, \hat{\xi}_k] = 0$,

$[\hat{\xi}_j, \hat{x}_k] = \delta_{jk}$ for all j, k . It is clear that then functions $x_i = \sigma(\hat{x}_i)$, $\xi_i = \sigma(\hat{\xi}_i)$ form Darboux coordinates with respect to $\{\cdot, \cdot\}$.

Proposition 6.1. *Let $[\cdot, \cdot]$ be a deformation of a Poisson bracket $\{\cdot, \cdot\}$ on M . Let $\hat{x}_1^{(i)}, \dots, \hat{x}_n^{(i)}, \hat{\xi}_1^{(i)}, \dots, \hat{\xi}_n^{(i)} \in \mathcal{C}_M^\infty[[t]]$, $i = 1, 2$, be two systems of Darboux coordinates with respect to $[\cdot, \cdot]$ on a contractible open set U in M satisfying*

$$\sigma(\hat{x}_j^{(1)}) = \sigma(\hat{x}_j^{(2)}), \quad \sigma(\hat{\xi}_j^{(1)}) = \sigma(\hat{\xi}_j^{(2)}).$$

Then, there exists $B \in \mathcal{C}_M^\infty[[t]]$ on U such that the automorphism $\Phi = \exp(t \cdot \text{ad}(B))$, where $\text{ad}(B) = [B, \cdot]$, satisfies $\Phi(\hat{x}_j^{(1)}) = \hat{x}_j^{(2)}$ and $\Phi(\hat{\xi}_j^{(1)}) = \hat{\xi}_j^{(2)}$.

Proof. Let $B_0 = 0$. Assume that B_m is such that the automorphism $\Phi_m = \exp(t \cdot \text{ad}(B_m))$ satisfies the conclusion of the proposition modulo t^{m+1} . This assumption is valid for $m = 0$.

Then, $\hat{x}_j^{(2)} = \Phi_m(\hat{x}_j^{(1)}) + t^{m+1}y_j \bmod t^{m+2}$, $\hat{\xi}_j^{(2)} = \Phi_m(\hat{\xi}_j^{(1)}) + t^{m+1}\eta_j \bmod t^{m+2}$ for suitable $y_j, \eta_j \in \mathcal{C}_M^\infty$. The Darboux commutation relations for $\hat{x}_1^{(i)}, \dots, \hat{x}_n^{(i)}, \hat{\xi}_1^{(i)}, \dots, \hat{\xi}_n^{(i)}$ imply that the functions $y_1, \dots, y_n, \eta_1, \dots, \eta_n$ satisfy $\{x_j, y_k\} - \{x_k, y_j\} = 0$, $\{\xi_j, \eta_k\} - \{\xi_k, \eta_j\} = 0$, $\{\xi_j, y_k\} - \{x_k, \eta_j\} = 0$, where $x_j = \sigma(\hat{x}_j^{(1)}) = \sigma(\hat{x}_j^{(2)})$, $\xi_j = \sigma(\hat{\xi}_j^{(1)}) = \sigma(\hat{\xi}_j^{(2)})$. Equivalently, the differential form $\alpha = \sum_j y_j dx_j + \eta_j d\xi_j$ is closed. By the Poincaré lemma there exists $f \in \mathcal{C}_M^\infty$ such that $df = \alpha$, equivalently $y_j = \{\xi_j, f\}$, and $\eta_j = \{f, x_k\}$.

There exists $B_{m+1} \in \mathcal{C}_M^\infty$ such that

$$\exp(t \cdot \text{ad}(B_{m+1})) = \exp(\text{ad}(t^{m+1}f)) \circ \exp(t \cdot \text{ad}(B_m))$$

and $B_{m+1} = B_m \bmod t^{m+1}$. The limit $B = \lim_{m \rightarrow \infty} B_m$ exists and satisfies the conclusions of the proposition. \square

Proposition 6.2. *Let $[\cdot, \cdot]$ be a deformation of a Poisson bracket $\{\cdot, \cdot\}$ on M and \mathcal{O}_t a t -adically complete submodule in $\mathcal{C}_M^\infty[[t]]$ being a commutative Lie subalgebra with respect to $[\cdot, \cdot]$. Let functions $x_i \in \mathcal{O}_0$, $\xi_i \in \mathcal{C}_M^\infty$ form Darboux coordinates with respect to $\{\cdot, \cdot\}$ on a contractible open set $U \subset M$. Then, there exist their lifts $\hat{x}_i \in \mathcal{O}_t$, $\hat{\xi}_i \in \mathcal{C}_M^\infty[[t]]$ on U which are Darboux coordinates with respect to $[\cdot, \cdot]$.*

Proof. Since, by definition, $\mathcal{O}_t \rightarrow \sigma(\mathcal{O}_t)$ is surjective, we choose arbitrary lifts $\hat{x}_j \in \mathcal{O}_t$ of x_j . Note that $\hat{x}_j, \hat{\xi}_j$ satisfy the both conclusions modulo t .

Let $\hat{x}_{j,1} := \hat{x}_j$, $\hat{\xi}_{j,1} := \xi_j$. Suppose that $m \geq 2$ and $\hat{x}_j \in \mathcal{O}_t$, $\hat{\xi}_{j,m} \in \mathcal{C}_M^\infty[[t]]$ satisfies the conclusion of the proposition modulo t^{m+1} . The assumption on $x_{j,m}, \xi_{j,m}$ implies that

$$[\hat{x}_j, \hat{x}_k] = 0, \quad [\hat{\xi}_{j,m}, \hat{x}_k] = \delta_{jk} + y_{jk}t^{m+1} \bmod t^{m+2},$$

$$[\hat{\xi}_j, \hat{\xi}_k] = t^{m+1}z_{jk} \bmod t^{m+2}$$

for suitable $y_{jk}, z_{jk} \in \mathcal{C}_M^\infty$. The Jacobi identity and the commutation relations for x_j, ξ_j imply that

$$\begin{aligned}\{y_{jk}, x_l\} - \{y_{jl}, x_k\} &= 0, & \{z_{lj}, x_k\} + \{\xi_j, y_{lk}\} - \{\xi_l, y_{jk}\} &= 0, \\ \{z_{jk}, \xi_l\} + \{z_{kl}, \xi_j\} + \{z_{lj}, \xi_k\} &= 0.\end{aligned}$$

These identities say that the differential form

$$\alpha = y_{jk} dx_j \wedge d\xi_k + z_{jk} dx_j \wedge dx_k$$

is closed.

By the Poincaré lemma there exists a 1-form $\beta = \sum a_i d\xi_i + \sum b_i dx_i$ on U such that $d\beta = \alpha$.

Note that $\{x_k, a_i\} = 0$ for all pairs i, k , since α contains no terms of the form $f d\xi_i \wedge d\xi_k$. Hence, $a_i \in \sigma(\mathcal{O}_t)$. Let \hat{a}_i be a lift of a_i in \mathcal{O}_t .

It is easy to check that $\hat{x}_{i,m+1} = x_{i,m} + t^{m+1} \hat{a}_i$ and $\hat{\xi}_{i,m+1} + t^{m+1} b_i$ satisfy the conclusions of the proposition modulo t^{m+2} . Hence, the limits $\hat{x}_i = \lim_{m \rightarrow \infty} \hat{x}_{i,m}$, $\hat{\xi}_i = \lim_{m \rightarrow \infty} \hat{\xi}_{i,m}$ exist and satisfy the conclusions of the proposition. \square

We will need a more strong assertion.

Proposition 6.3. *Let $[\cdot, \cdot]$ be a deformation of a Poisson bracket $\{\cdot, \cdot\}$ on M and \mathcal{O}_t a t -regular submodule in $\mathcal{C}_M^\infty[[t]]$, which is a commutative Lie subalgebra with respect to $[\cdot, \cdot]$. Let functions $x_i \in \mathcal{O}_t$, $\xi_i \in \mathcal{C}_M^\infty[[t]]$ form Darboux coordinates modulo t^k , $k > 0$, with respect to $[\cdot, \cdot]$ on a contractible open set $U \subset M$. Then, there exist functions $a_i \in \mathcal{O}_t$, $b_i \in \mathcal{C}_M^\infty[[t]]$ on U such that the functions $x_i + t^k a_i$, $\xi_i + t^k b_i$ form Darboux coordinates with respect to $[\cdot, \cdot]$.*

Proof. The same as of Proposition 6.2. \square

7. Deformation quantization on a polarized symplectic manifold

7.1. Deformation quantization

Let us recall some definitions and facts about the DQ on a smooth manifold M , see [2,9,11].

Definition 7.1.

- (a) Let \mathcal{C}_M^∞ be the sheaf of smooth complex valued functions on M . A *formal deformation* of \mathcal{C}_M^∞ is a sheaf of $\mathbb{C}[[t]]$ -algebras, \mathbb{A}_t , with an epimorphism $\sigma : \mathbb{A}_t \rightarrow \mathcal{C}_M^\infty$ of $\mathbb{C}[[t]]$ -algebras (called the *symbol map*) satisfying the condition: there exists an isomorphism of $\mathbb{C}[[t]]$ -modules $\mathbb{A}_t \rightarrow \mathcal{C}_M^\infty[[t]]$ commuting with symbol maps. (Recall that the symbol map $\sigma : \mathcal{C}_M^\infty[[t]] \rightarrow \mathcal{C}_M^\infty$ takes $f_0 + tf_1 + \dots$ to f_0 .)

- (b) Two formal deformations \mathbb{A}_{t1} and \mathbb{A}_{t2} of \mathcal{C}_M^∞ are equivalent if there exists a map of sheaves of $\mathbb{C}[[t]]$ -algebras $\mathbb{A}_{t1} \rightarrow \mathbb{A}_{t2}$, commuting with symbol maps.

Suppose \mathbb{A}_t is a formal deformation of \mathcal{C}_M^∞ . The formula

$$\{a, b\} = \sigma \left(\frac{1}{t} [\tilde{a}, \tilde{b}] \right), \quad (7.1)$$

where a and b are locally defined functions on M and \tilde{a}, \tilde{b} are their lifts with respect to σ , gives a well-defined Poisson bracket on \mathcal{C}_M^∞ . It is clear that equivalent formal deformations have the same Poisson bracket.

Definition 7.2. A DQ on a symplectic manifold (M, ω) is a formal deformation of \mathcal{C}_M^∞ whose Poisson bracket is equal to ω^{-1} .

Definition 7.3.

- (a) An SP on (M, ω) is the structure of an associative algebra on the sheaf $\mathcal{C}_M^\infty[[t]]$ with the multiplication of the form

$$f * g = \sum_{i \geq 0} t^i \mu_i(f, g), \quad f, g \in \mathcal{C}_M^\infty, \quad (7.2)$$

where all μ_i , $i > 0$, are bidifferential operators on M vanishing on constants, i.e. $\mu_i(f, g) = 0$ if f or g is a constant, $\mu_0(f, g) = fg$, and $\mu_1(f, g) - \mu_1(g, f) = \{f, g\}$, the Poisson bracket inverse to ω .

- (b) Two SPs $(\mathcal{C}_M^\infty[[t]], \mu')$ and $(\mathcal{C}_M^\infty[[t]], \mu'')$ on (M, ω) are equivalent if there exists a power series $B = 1 + tB_1 + \dots$, where B_i are differential operators vanishing on constants, such that $\mu''(f, g) = B\mu'(B^{-1}f, B^{-1}g)$.

It is clear that any SP $(\mathcal{C}_M^\infty[[t]], \mu)$ defines a DQ with the natural symbol map $\mathcal{C}_M^\infty[[t]] \rightarrow \mathcal{C}_M^\infty$, $f_0 + tf_1 + \dots \mapsto f_0$.

Proposition 7.4. *The above assignment gives a one-to-one correspondence between the equivalence classes of SPs and DQs.*

Proof. Let \mathbb{A}_t be a DQ. Let us prove that it is equivalent to an SP. By definition of DQ, there exists an isomorphism of $\mathbb{C}[[t]]$ -modules $\mathcal{C}_M^\infty[[t]] \rightarrow \mathbb{A}_t$ commuting with symbol maps. Let $\mu = \mu_0 + t\mu_1 + \dots$ be the multiplication in the sheaf $\mathcal{C}_M^\infty[[t]]$ being the pullback of the multiplication in \mathbb{A}_t . In order to prove that each μ_i is a bidifferential operator it is enough to prove, according to the Peetre theorem, that $\text{supp}(\mu(f, g)) \subset \text{supp}(f) \cap \text{supp}(g)$ for any functions f and g on M . But this is obvious because if, for example, $f = 0$ on an open set U , then $\mu(f, g) = 0$ on U , since μ is a map of sheaves. The same argument proves that two equivalent DQs correspond to equivalent SPs. \square

Example 7.5 (Moyal–Weyl SP). Let U be an open set in a symplectic manifold (M, ω) , in which there exist Darboux coordinates, x_i, y_i , $i = 1, \dots, (1/2)\dim M$, so that $\omega =$

$\sum_i dy_i \wedge dx_i$. Then, $\omega^{-1} = \sum_i \partial/\partial y_i \wedge \partial/\partial x_i$ and for $f, g \in C_U^\infty$ the multiplication formula

$$f \otimes g \mapsto m \exp \left(\frac{t}{2} \sum_i (\partial/\partial y_i \otimes \partial/\partial x_i - \partial/\partial x_i \otimes \partial/\partial y_i) \right) (f \otimes g), \quad (7.3)$$

where m is the usual multiplication of functions, defines an SP on (U, ω) . This SP is called the Moyal–Weyl SP.

The following statement is well-known, see for example [17].

Proposition 7.6. *Locally, any SP on (M, ω) is equivalent to the Moyal–Weyl SP.*

7.2. Polarized DQ

Definition 7.7.

- (a) A PDQ on a PSM (M, ω, \mathcal{P}) is a pair $(\mathbb{A}_t, \mathcal{O}_t)$ where \mathbb{A}_t is a DQ on (M, ω) and \mathcal{O}_t is a t -adically complete commutative subalgebra in \mathbb{A}_t such that $\sigma(\mathcal{O}_t) = \mathcal{O}_{\mathcal{P}}$, the sheaf of functions constant along \mathcal{P} .
- (b) Two PDQs $(\mathbb{A}_{t1}, \mathcal{O}_{t1})$ and $(\mathbb{A}_{t2}, \mathcal{O}_{t2})$ of (M, ω, \mathcal{P}) are equivalent if there exists an equivalence map of DQs $\mathbb{A}_{t1} \rightarrow \mathbb{A}_{t2}$ which takes \mathcal{O}_{t1} to \mathcal{O}_{t2} .

Since any DQ is equivalent to an SP, any PDQ is equivalent to a triple $(C_M^\infty[[t]], \mu_t, \mathcal{O}_t)$, where $(C_M^\infty[[t]], \mu_t)$ is an SP and \mathcal{O}_t is a commutative subalgebra in $(C_M^\infty[[t]], \mu_t)$.

Proposition 7.8. *Let $(\mathbb{A}_t, \mathcal{O}_t)$ be a PDQ. Then,*

- (a) \mathcal{O}_t is a maximal commutative subalgebra in \mathbb{A}_t ;
- (b) \mathcal{O}_t is a t -regular subalgebra in \mathbb{A}_t .

Proof.

- (a) Locally, the bracket $[a, b] = (1/t)(ab - ba)$, $a, b \in \mathbb{A}_t$, is a deformation of the Poisson bracket ω^{-1} . Hence, the statement easily follows from Proposition 3.4.
- (b) Follows from (a). □

The following definition will play an auxiliary role in the paper.

Definition 7.9. A weakly PSP (wPSP) on a PSM (M, ω, \mathcal{P}) is a triple $(C_M^\infty[[t]], \mu_t, \mathcal{O}_t)$, where $(C_M^\infty[[t]], \mu_t)$ is an SP, \mathcal{O}_t is a t -adically complete $\mathbb{C}[[t]]$ -submodule in $C_M^\infty[[t]]$ satisfying the conditions:

- (a) $\sigma(\mathcal{O}_t) = \mathcal{O}_{\mathcal{P}}$;
- (b) \mathcal{O}_t is a commutative subalgebra in $C_M^\infty[[t]]$ with respect to the usual multiplication in $C_M^\infty[[t]]$;
- (c) μ_t being restricted to \mathcal{O}_t coincides with the usual multiplication.

Definition 7.10. We say that a wPSP $(\mathcal{C}_M^\infty[[t]], \mu_t, \mathcal{O}_t)$ is a PSP, if for any $a \in \mathcal{O}_t$, $b \in \mathcal{C}_M^\infty[[t]]$ the product $\mu_t(a, b)$ coincides with the usual product in $\mathcal{C}_M^\infty[[t]]$.

Thus, both wPSP and PSP are particular cases of a PDQ. We are going to prove that, in fact, any PDQ is equivalent to a PSP. But first we prove the following lemma.

Lemma 7.11. Any PDQ is equivalent to a wPSP.

Proof. Since any DQ is equivalent to an SP, it is enough to prove the following. Let \mathcal{O}_t be a commutative subalgebra in an SP $(\mathcal{C}_M^\infty[[t]], \mu_t)$ such that the triple $(\mathcal{C}_M^\infty[[t]], \mu, \mathcal{O}_t)$ forms a PDQ, in particular, $\mu_t(a, b) = \mu_t(b, a)$ for $a, b \in \mathcal{O}_t$. Then, there exists a differential operator $D_t = 1 + tD_1 + \dots$ on M such that the multiplication $\tilde{\mu}_t(f, g) = D_t^{-1}\mu(D_t f, D_t g)$ being restricted to $D_t^{-1}\mathcal{O}_t$ is the usual multiplication.

Suppose μ_t being restricted to \mathcal{O}_t coincides with the usual multiplication modulo t^n . Then, $\mu(a, b) = ab + t^n v(a, b) \bmod t^n$ for $a, b \in \mathcal{O}_t$. It follows from Proposition 7.8 (b) that being considered modulo t , the bilinear form v defines a Hochschild cocycle $\bar{v} \in \Gamma(M, \mathcal{D}^2(\mathcal{O}_P, \mathcal{C}_M^\infty))$ (see Proposition 2.5).

Since \bar{v} is commutative, it follows from Proposition 2.5 that it is a coboundary. Therefore, there exists a differential operator $\tilde{D} \in \Gamma(M, \mathcal{D}^1(\mathcal{O}_P, \mathcal{C}_M^\infty))$ such that $d_{\text{Hoch}} \tilde{D} = \bar{v}$.

Let \tilde{D} be a lift of \tilde{D} to a differential operator on M . Let $D_n = 1 + t^n \tilde{D}$. It is easy to see that $D_n^{-1}\mathcal{O}_t$ is a commutative subalgebra in the SP $(\mathcal{C}_M^\infty[[t]], \tilde{\mu}_t)$ with $\tilde{\mu}_t(a, b) = D_n^{-1}\mu_t(D_n a, D_n b)$, and $\tilde{\mu}_t(a, b) = ab \bmod t^{n+1}$ for $a, b \in D_n^{-1}\mathcal{O}_t$.

By induction, we construct such differential operators D_n for all n . Let $D = \prod_{n=1}^\infty D_n$, $\mathcal{O}'_t = D^{-1}\mathcal{O}_t$, and $\mu'_t(a, b) = D^{-1}\mu_t(Da, Db)$. Then D gives an isomorphism between PDQs $(\mathcal{C}_M^\infty[[t]], \mu_t, \mathcal{O}_t)$ and $(\mathcal{C}_M^\infty[[t]], \mu'_t, \mathcal{O}'_t)$, and the second triple is a wPSP, which proves the proposition. \square

Proposition 7.12. For any wPSP $(\mathcal{C}_M^\infty[[t]], \mu_t, \mathcal{O}_t)$, there exists a differential operator $D = 1 + tD_1 + \dots$ on M such that $Df = f$ for $f \in \mathcal{O}_t$ and the multiplication $\mu'_t(a, b) = D^{-1}\mu_t(Da, Db)$ defines a PSP $(\mathcal{C}_M^\infty[[t]], \mu'_t, \mathcal{O}_t)$.

Proof. It is obvious that $\mu = \mu_t$ defines a PSP modulo t . Proceeding by induction, we assume that there exists a wPSP multiplication μ' equivalent to μ and being a PSP modulo t^n with respect to \mathcal{P}_t . The proposition will be proved if we find a differential operator, D_n , such that $D_n(f) = 0$ for all $f \in \mathcal{O}_0$ and the multiplication

$$\mu''(a, b) = D^{-1}\mu'(Da, Db), \quad (7.4)$$

where $D = 1 + t^n D_n$ defines a PSP modulo t^{n+1} with respect to \mathcal{P}_t .

Let

$$\mu' = \mu_0 + t\mu'_1 + \dots + t^{n-1}\mu'_{n-1} + t^n v \bmod t^{n+1}.$$

By our assumption, elements $\mu'_1, \dots, \mu'_{n-1}$ are strongly polarized and v is polarized with respect to \mathcal{O}_t (see the definition before Proposition 2.4). It follows from associativity of μ' that

$$d_{\text{Hoch}} v(a, b, c) = \sum_{i+j=n} (\mu'_i(a, \mu'_j(b, c)) - \mu'_i(\mu'_j(a, b), c)).$$

It is easy to check that each term in the right hand side is strongly polarized, since all μ'_i , $i = 1, \dots, n - 1$, are such. Therefore, ν satisfies the hypothesis of [Proposition 2.4](#). Hence, there exists a polarized differential operator D_n such that $\nu + d_{\text{Hoch}} D_n$ is a strongly polarized bidifferential operator. It is obvious that the multiplication μ'' defined in [\(7.4\)](#) with D_n just constructed is as required. \square

Corollary 7.13. Any PDQ is equivalent to a PSP.

Proof. Follows from [Lemma 7.11](#) and [Proposition 7.12](#). \square

Proposition 7.14.

- (a) Let $(\mathcal{C}_M^\infty[[t]], \mu_t, \mathcal{O}_t)$ be a PSP on (M, ω, \mathcal{P}) . Let $\mathcal{P}_t = (d\mathcal{O}_t)^\perp$. Then, \mathcal{P}_t is a deformation of \mathcal{P} and $\mathcal{O}_t = \mathcal{O}_{\mathcal{P}_t}$.
- (b) Let $(\mathcal{C}_M^\infty[[t]], \tilde{\mu}_t, \tilde{\mathcal{O}}_t)$ be another PSP on (M, ω, \mathcal{P}) equivalent to $(\mathcal{C}_M^\infty[[t]], \mu_t, \mathcal{O}_t)$ as a PDQ. Then, there exists a formal automorphism of M which takes \mathcal{P}_t to $\tilde{\mathcal{P}}_t = (d\tilde{\mathcal{O}}_t)^\perp$.
- (c) Let $(\mathcal{C}_M^\infty[[t]], \mu_t, \mathcal{O}_t)$ and $(\mathcal{C}_M^\infty[[t]], \tilde{\mu}_t, \mathcal{O}_t)$ are two equivalent PSPs with the same \mathcal{O}_t . Let $D = 1 + tD_1 + \dots$ gives an equivalence. Then, there exists a decomposition, $D = D'e^{tX}$, where D' is a differential operator identical on \mathcal{O}_t and X is a formal vector field on M taking \mathcal{O}_t to itself.

Proof.

- (a) Follows from [Proposition 4.4](#).
- (b) Let us put $X_0 = 0$. Then the automorphism $e^{tX_0} = \text{Id}$ takes \mathcal{P}_t to $\tilde{\mathcal{P}}_t$ modulo t . Suppose we have constructed a formal vector field X_k such that the formal automorphism $e^{tX_{k-1}}$ of M takes \mathcal{P}_t to $\tilde{\mathcal{P}}_t$ modulo t^k . Then, replacing \mathcal{P}_t by $e^{tX_{k-1}}\mathcal{P}_t$ we may assume that \mathcal{P}_t and $\tilde{\mathcal{P}}_t$ coincide modulo t^k . The proposition will be proved, if we show that it is possible to find a vector field Y such that $e^{t^k Y}$ takes \mathcal{P}_t to $\tilde{\mathcal{P}}_t$ modulo t^{k+1} .

Since, by our assumption, the SPs $(\mathcal{C}_M^\infty[[t]], \mu_t)$ and $(\mathcal{C}_M^\infty[[t]], \tilde{\mu}_t)$ are equivalent and coincide modulo t^k , there exists a differential operator $1 + t^k D_k$ realizing that equivalence. Since $1 + t^k D_k$ takes \mathcal{O}_t to $\tilde{\mathcal{O}}_t$ and on the both of these subalgebras the respecting multiplications μ_t and $\tilde{\mu}_t$ are trivial, D_k being restricted to \mathcal{O}_0 is a derivation from \mathcal{O}_0 to \mathcal{C}_M^∞ . Let Y be an extension of that restricted D_k to a derivation on \mathcal{C}_M^∞ . It is clear that $e^{t^k Y}$ takes \mathcal{P}_t to $\tilde{\mathcal{P}}_t$ modulo t^{k+1} .

- (c) The operator D being restricted to \mathcal{O}_t is a formal automorphism of \mathcal{O}_t . Since formal automorphisms form a pro-unipotent group, there exists $X' \in \text{Der}(\mathcal{O}_t)$ such that D being restricted to \mathcal{O}_t is equal to $e^{tX'}$. Let $X \in \text{Der}(\mathcal{C}_M^\infty)$ be a lift of X' . We put $D' = De^{-tX}$ which is obviously identical on \mathcal{O}_t . \square

Example 7.15 (Moyal–Wick PSP). Let (M, ω, \mathcal{P}) be a PSM. Let U be an open set in M where there exist Darboux coordinates, $x_i, y_i, (dx_i)^\perp = \mathcal{P}, i = 1, \dots, (1/2)\dim M$, so that $\omega = \sum_i dy_i \wedge dx_i$. Then, $\omega^{-1} = \sum_i \partial/\partial y_i \wedge \partial/\partial x_i$ and for $f, g \in C_U^\infty$ the multiplication

formula

$$f \otimes g \mapsto m \exp \left(t \sum_i \partial/\partial y_i \otimes \partial/\partial x_i \right) (f \otimes g), \quad (7.5)$$

where m is the usual multiplication of functions, defines a PSP on (U, ω, \mathcal{P}) . This PSP is called the Moyal–Wick PSP.

Note that the functions a_i, f_i satisfying the Darboux relations with respect to the Poisson bracket ω^{-1} also satisfy the Darboux relations with respect to the deformed bracket $(1/t)[\cdot, \cdot]$, where $[\cdot, \cdot]$ is the commutator of the Moyal–Wick PSP.

Let us remark that the Moyal–Weyl SP from [Example 7.5](#) constructed with help of the same Darboux coordinates gives just a wPSP but not a PSP.

Proposition 7.16. *Locally, any PSP on (M, ω, \mathcal{P}) is equivalent to the Moyal–Wick PSP.*

Proof. Let us prove that any two PSPs are locally equivalent. Since any SPs are locally equivalent, we may suppose that there are given two PSPs, $(\mathcal{C}_M^\infty[[t]], \mu_t, \mathcal{O}_t)$ and $(\mathcal{C}_M^\infty[[t]], \mu_t, \tilde{\mathcal{O}}_t)$, on (M, ω, \mathcal{P}) with the same multiplication and different polarizations, and we have to prove that they are, locally, equivalent. Let x_i, y_i are Darboux coordinates with respect to ω such that $(dx_i)^\perp = \mathcal{P}$. By [Proposition 6.2](#), there exist their lifts x_{it}, y_{it} and x'_{it}, y'_{it} which satisfy the Darboux relations with respect to the bracket $[a, b] = (1/t)(\mu_t(a, b) - \mu_t(b, a))$, and $x_{it} \in \mathcal{O}_t, x'_{it} \in \tilde{\mathcal{O}}_t$. By [Proposition 6.1](#), there exists, locally, an inner automorphism of the SP $(\mathcal{C}_M^\infty[[t]], \mu_t)$ that takes x_{it}, y_{it} to x'_{it}, y'_{it} . It follows that this automorphism takes \mathcal{O}_t to $\tilde{\mathcal{O}}_t$. \square

8. Characteristic classes of PDQs and PSPs

8.1. Extension class associated with a PDQ

Let $(\mathbb{A}_t, \mathbb{O}_t)$ be a PDQ on a PSM (M, ω, \mathcal{P}) . Since any PDQ is equivalent to a PSP, the sheaf \mathbb{O}_t is isomorphic to $\mathcal{O}_{\mathcal{P}_t}$ for some deformed distribution \mathcal{P}_t . Thus, the sheaves $\text{Der}(\mathbb{O}_t)$ and $\Omega_{\mathbb{O}_t}^{1\text{cl}}$ are well defined (see [Section 2](#)). Let

$$F(\mathbb{A}_t, \mathbb{O}_t) = \{b \in \mathbb{A}_t; [b, \mathbb{O}_t] \subset \mathbb{O}_t\}, \quad (8.1)$$

where $[\cdot, \cdot]$ denotes the commutator in \mathbb{A}_t . It is clear that $F(\mathbb{A}_t, \mathbb{O}_t)$ is a sheaf of Lie algebras with the bracket $(1/t)[\cdot, \cdot]$ and the center \mathbb{O}_t . Moreover, any element $b \in F(\mathbb{A}_t, \mathbb{O}_t)$ determines the derivation $(1/t)[b, \cdot]$ of \mathbb{O}_t and, due to [Proposition 6.2](#), this correspondence defines an epimorphism $\sigma : F(\mathbb{A}_t, \mathbb{O}_t) \rightarrow \text{Der}(\mathbb{O}_t)$.

We consider $F(\mathbb{A}_t, \mathbb{O}_t)$ as a left \mathbb{O}_t -module with respect to multiplication in \mathbb{A}_t . As a Lie algebra sheaf, $F(\mathbb{A}_t, \mathbb{O}_t)$ is an extension of $\text{Der}(\mathbb{O}_t)$.

Thus, we have the following exact sequence of Lie algebras and \mathcal{O}_t -modules:

$$0 \rightarrow \mathbb{O}_t \xrightarrow{\iota} F(\mathbb{A}_t, \mathbb{O}_t) \xrightarrow{\sigma} \text{Der}(\mathbb{O}_t) \rightarrow 0. \quad (8.2)$$

According to the terminology of Beilinson and co-workers [3,4], $F(\mathbb{A}_t, \mathbb{O}_t)$ is called an \mathbb{O}_t -extension of $\text{Der}(\mathbb{O}_t)$.

We say that a map of Lie algebras and \mathbb{O}_t -modules, $s : \text{Der}(\mathbb{O}_t) \rightarrow F(\mathbb{A}_t)$, given on an open set of M is a splitting of (8.2), if $s\sigma = \text{id}$. Since $(\mathbb{A}_t, \mathbb{O}_t)$ can be realized as a PSP and, locally, there exist Darboux coordinates with respect to $(1/t)[\cdot, \cdot]$, the sequence (8.2) locally splits (see the next subsection, where splittings are presented with help of Darboux coordinates).

Lemma 8.1. *Let s and s' are two splittings of (8.2) over an open set of M . Then, $s - s' \in \Omega_{\mathbb{O}_t}^{\text{lcl}}$.*

Proof. Direct calculation. \square

Let us define an extension class of (8.1) in the following way. Let $\{U_\alpha\}$ be an open covering of M such that over each U_α there is a splitting, s_α , of $F(\mathbb{A}_t, \mathbb{O}_t)$. By Lemma 8.1, $f_{\alpha,\beta} = s_\beta - s_\alpha$ is a section of $\Omega_{\mathbb{O}_t}^{\text{lcl}}$ over $U_\alpha \cap U_\beta$. We define $\text{cl}_{\text{Ext}}(F(\mathbb{A}_t, \mathbb{O}_t))$ as the element of $H^1(M, \Omega_{\mathbb{O}_t}^{\text{lcl}})$ represented by the collection $\{f_{\alpha,\beta}\}$. One can prove that given \mathbb{O}_t , the extension class $\text{cl}_{\text{Ext}}(F(\mathbb{A}_t, \mathbb{O}_t))$ determines a \mathbb{O}_t -extension of $\text{Der}(\mathbb{O}_t)$ up to equivalence.

We will denote the element $\text{cl}_{\text{Ext}}(F(\mathbb{A}_t, \mathbb{O}_t))$ by $\text{cl}_{\text{Ext}}(\mathbb{A}_t, \mathbb{O}_t)$ and call it the *extension class* of the PDQ $(\mathbb{A}_t, \mathbb{O}_t)$.

In the next subsection, the extension class of a PSP, $(\mathcal{C}_M^\infty[[t]], \mu_t, \mathcal{O}_t)$, will be represented as an element of $\Gamma(M, d\mathcal{P}_t^\perp)/d(\Gamma(M, \mathcal{P}_t^\perp))$, $\mathcal{P}_t = \mathcal{P}_{\mathcal{O}_t}$, with the help of a characteristic 2-form associated with that PSP.

8.2. Characteristic 2-form associated with a PSP

Let $(\mathcal{C}_M^\infty[[t]], \mu_t, \mathcal{O}_t)$ be a PSP that we denote for shortness by (μ_t, \mathcal{O}_t) . We denote $\mathcal{P}_t = \mathcal{P}_{\mathcal{O}_t}$. Then the extension $F(\mu_t, \mathcal{O}_t)$ coincides as a left \mathcal{O}_t -module with an \mathcal{O}_t -submodule of $\mathcal{C}_M^\infty[[t]]$ with respect to the usual multiplication in $\mathcal{C}_M^\infty[[t]]$. In this case, the local splittings of $F(\mu_t, \mathcal{O}_t)$ are differential forms of $\text{Hom}_{\mathcal{O}_t}(\text{Der}(\mathcal{O}_t), \mathcal{C}_M^\infty[[t]]) = \mathcal{P}^\perp$. These forms can be described explicitly by Darboux coordinates.

Let $\{U_\alpha\}$ be an open covering of M such that each U_α has Darboux coordinates $x_{\alpha i}, y_{\alpha i}$, $x_{\alpha i} \in \mathcal{O}_t$, $i = 1, \dots, n$, with respect to the bracket $[a, b] = (1/t)(\mu_t(a, b) - \mu_t(b, a))$ (in particular, $[y_{\alpha i}, x_{\alpha j}] = \delta_{ij}$). By Proposition 6.2 such a covering exists. Then, on U_α , the \mathcal{O}_t -submodule $F(\mu_t, \mathcal{O}_t) \subset \mathcal{C}_M^\infty[[t]]$ is equal to $\mathcal{O}_t \oplus \mathcal{O}_t y_{\alpha 1} \oplus \dots \oplus \mathcal{O}_t y_{\alpha n}$.

Splittings s_α may be taken by the condition $x_{\alpha i} \mapsto y_{\alpha i}$, and the corresponding forms are

$$s_\alpha = \sum_i y_{\alpha i} dx_{\alpha i}. \quad (8.3)$$

By Lemma 8.1, $ds_\alpha = ds_\beta$ on $U_\alpha \cap U_\beta$. Therefore, forms s_α define the global 2-form

$$\omega_t \in \omega_0 + t\Gamma(M, d\mathcal{P}_t^\perp), \quad \omega_t = ds_\alpha = \sum_i dy_{\alpha i} \wedge dx_{\alpha i}. \quad (8.4)$$

Lemma 8.2. *This form represents the extension class $\text{cl}_{\text{Ext}}(\mu_t, \mathcal{O}_t)$ by the isomorphism (2.8).*

Proof. Clear. \square

Lemma 8.1 shows that if we take other splittings of $F(\mu_t, \mathcal{O}_t)$, the procedure above gives the same form ω_t . Therefore, in the case of PSP, it is well defined not only the class $\text{cl}_{\text{Ext}}(\mu_t, \mathcal{O}_t)$ but also its representative in $\omega_0 + t\Gamma(M, d\mathcal{P}_t^\perp)$ that we call the *characteristic 2-form* of the PSP and denote it by $\text{cl}_{\text{PSP}}(\mu_t, \mathcal{O}_t)$.

Given an SP ($\mathcal{C}_M^\infty[[t]]$, μ_t), let us set $[a, b]_{\mu_t} = (1/t)(\mu_t(a, b) - \mu_t(b, a))$, $a, b \in \mathcal{C}_M^\infty[[t]]$. The bracket $[\cdot, \cdot]_{\mu_t}$ is a deformation of the initial Poisson bracket on M .

Proposition 8.3. *Let (μ_t, \mathcal{O}_t) , $(\tilde{\mu}_t, \mathcal{O}_t)$ be PSPs with the same \mathcal{O}_t . Then, $\text{cl}_{\text{PSP}}(\mu_t, \mathcal{O}_t) = \text{cl}_{\text{PSP}}(\tilde{\mu}_t, \mathcal{O}_t)$ if and only if $[\cdot, \cdot]_{\mu_t} = [\cdot, \cdot]_{\tilde{\mu}_t}$.*

Proof. If $\text{cl}_{\text{PSP}}(\mu_t, \mathcal{O}_t) = \text{cl}_{\text{PSP}}(\tilde{\mu}_t, \mathcal{O}_t)$, then forms (8.3) can be taken to be the same. This implies that there exist common Darboux coordinates with respect to $[\cdot, \cdot]_{\mu_t}$ and $[\cdot, \cdot]_{\tilde{\mu}_t}$. \square

Proposition 8.4. *Let (μ_t, \mathcal{O}_t) , $(\tilde{\mu}_t, \mathcal{O}_t)$ be two PSPs with the same \mathcal{O}_t . Let $\mathcal{P}_t = \mathcal{P}_{\mathcal{O}_t}$ and $\Gamma\mathcal{P}_t$ denote the module of global sections of \mathcal{P}_t . Then, the 2-forms $\text{cl}_{\text{PSP}}(\mu_t, \mathcal{O}_t)$, $\text{cl}_{\text{PSP}}(\tilde{\mu}_t, \mathcal{O}_t)$ are lying on the same orbit of $e^{t\Gamma\mathcal{P}_t}$ if and only if there exists a formal differential operator $D = 1 + tD_1 + \dots$ identical on \mathcal{O}_t such that $\tilde{\mu}_t(a, b) = D^{-1}\mu_t(Da, Db)$.*

Proof. Let us denote $\omega_t = \text{cl}_{\text{PSP}}(\mu_t, \mathcal{O}_t)$, $\tilde{\omega}_t = \text{cl}_{\text{PSP}}(\tilde{\mu}_t, \mathcal{O}_t)$. Assume such D exists. Let s_α be forms from (8.3) corresponding to (μ_t, \mathcal{O}_t) . Let \tilde{s}_α be forms obtained from s_α by applying D . **Lemmas 8.1 and 2.6** imply that there exist $f_{\alpha, \beta} \in \mathcal{O}_t$ on $U_\alpha \cap U_\beta$ such that $s_\alpha - s_\beta = df_{\alpha, \beta}$. Since D acts on \mathcal{O}_t trivially, one has $\tilde{s}_\alpha - \tilde{s}_\beta = df_{\alpha, \beta}$, too. This means that $\tilde{s}_\alpha - s_\alpha$ does not depend on α and give a global form, b , of \mathcal{P}_t^\perp . Since $\omega_t = ds_\alpha$, $\tilde{\omega}_t = d\tilde{s}_\alpha$, one has $\tilde{\omega}_t = \omega_t + db$. By **Lemma 4.6**, there exists a formal automorphism, e^{tY} , $Y \in \Gamma\mathcal{P}_t$, taking ω_t to $\tilde{\omega}_t$.

Conversely, let us suppose that ω_t and $\tilde{\omega}_t$ are lying on the same orbit of $e^{t\Gamma\mathcal{P}_t}$. Assume that we have found a differential operator identical on \mathcal{O}_t which transforms $\tilde{\mu}_t$ to a multiplication $\tilde{\mu}'_t$ that is equal to μ_t modulo t^k , i.e.

$$\tilde{\mu}'_t - \mu_t = t^k v + \dots \quad (8.5)$$

By the previous part of the proof, the corresponding form $\tilde{\omega}'_t$ is also lying on the same orbit as ω_t . The proposition will be proved if we find a differential operator $1 + t^k D_k$, $D_k(\mathcal{O}_t) = 0$, which transforms $\tilde{\mu}'_t$ to μ_t modulo t^{k+1} . Let us prove that.

Since $\tilde{\mu}'_t = \mu_t \bmod t^k$, we can choose, by **Proposition 6.3**, systems of Darboux coordinates with respect to $[\cdot, \cdot]_{\tilde{\mu}'_t}$ and $[\cdot, \cdot]_{\mu_t}$ that coincides modulo t^k . It follows that $\tilde{\omega}'_t = \omega_t \bmod t^k$. Hence, there is a formal automorphism $e^{t^k X}$, $X \in \mathcal{P}_t$, which takes $\tilde{\omega}'_t$ to ω_t . Applying $e^{t^k X}$ to $\tilde{\mu}'_t$, we obtain a multiplication that is still equal to μ_t modulo t^k but whose characteristic form is equal to ω_t . Thus, we come to the situation when the multiplications $\tilde{\mu}'_t$ and μ_t have the same characteristic form ω_t .

By **Proposition 8.3**, $\tilde{\mu}'_t$ and μ_t have the same commutator. This implies that bidifferential operator v in (8.5) is commutative. Moreover, it is a Hochschild strongly polarized cocycle (see **Section 2**). Therefore, there exists a polarized differential operator D_k such that

$d_{\text{Hoch}} D = v$. It follows that transformation $1 + t^k D_k$ applying to $\tilde{\mu}'_t$ gives a multiplication equal to μ_t modulo t^{k+1} and identical on \mathcal{O}_t . \square

8.3. Characteristic pairs for PSPs and PDQs

Let (M, ω, \mathcal{P}) be a PSM. Let us denote by $\text{Aut}(M)$ the group of formal automorphisms of M and by $\mathcal{Y} = \mathcal{Y}(M, \omega, \mathcal{P})$ the set of pairs $(\omega_t, \mathcal{P}_t)$, where $\omega_t = \omega_0 + t\omega_1 + \dots$ is a formal symplectic form being a deformation of $\omega = \omega_0$ and \mathcal{P}_t is a polarization of ω_t being a deformation of \mathcal{P} . The group $\text{Aut}(M)$ naturally acts on \mathcal{Y} .

It is natural to assign to a PSP (μ_t, \mathcal{O}_t) on (M, ω, \mathcal{P}) a pair $(\omega_t, \mathcal{P}_t) \in \mathcal{Y}$, where $\omega_t = \text{cl}_{\text{PSP}}(\mu_t, \mathcal{O}_t)$ and $\mathcal{P}_t = \mathcal{P}_{\mathcal{O}_t}$. Hence, we obtain the map

$$\tau : \{\text{PSPs}\} \rightarrow \mathcal{Y}.$$

Proposition 8.5. *Two PSPs (μ_t, \mathcal{O}_t) and $(\tilde{\mu}_t, \tilde{\mathcal{O}}_t)$ are equivalent if and only if $\tau(\mu_t, \mathcal{O}_t)$ and $\tau(\tilde{\mu}_t, \tilde{\mathcal{O}}_t)$ are lying on the same orbit in \mathcal{Y} under the $\text{Aut}(M)$ -action.*

Proof. Let (μ_t, \mathcal{O}_t) and $(\tilde{\mu}_t, \tilde{\mathcal{O}}_t)$ be equivalent. Let us prove that the pairs $\tau(\mu_t, \mathcal{O}_t)$ and $\tau(\tilde{\mu}_t, \tilde{\mathcal{O}}_t)$ are lying on the same orbit. By Proposition 7.14 (b) and (c), one can find a formal automorphism of M such that after its applying we come to the situation when $(\tilde{\mu}_t, \tilde{\mathcal{O}}_t)$ turns into a PSP, $(\tilde{\mu}_t, \mathcal{O}_t)$, with the same \mathcal{O}_t as in (μ_t, \mathcal{O}_t) and the equivalence morphism from (μ_t, \mathcal{O}_t) to $(\tilde{\mu}_t, \mathcal{O}_t)$ is given by a differential operator identical on \mathcal{O}_t . Now the statement follows from Proposition 8.4.

Conversely, suppose that for (μ_t, \mathcal{O}_t) and $(\tilde{\mu}_t, \tilde{\mathcal{O}}_t)$ the corresponding pairs $(\omega_t, \mathcal{P}_t)$, $(\tilde{\omega}_t, \tilde{\mathcal{P}}_t)$ lie on the same orbit. Let us prove that those PSPs are equivalent. Let B be a formal automorphism of M taking $\tilde{\mathcal{P}}_t$ to \mathcal{P}_t . Applying B to $(\tilde{\mu}_t, \tilde{\mathcal{O}}_t)$, we come to the case when $\tilde{\mathcal{P}}_t = \mathcal{P}_t$. Therefore, we may suppose that $(\tilde{\omega}_t, \tilde{\mathcal{P}}_t) = (\tilde{\omega}_t, \mathcal{P}_t)$. Since the pairs $(\omega_t, \mathcal{P}_t)$, $(\tilde{\omega}_t, \mathcal{P}_t)$ lie on the same orbit, there exists $X \in \Gamma \mathcal{P}_t$ such that $\tilde{\omega}_t = e^{tX} \omega_t$. Now the statement follows from Proposition 8.4. \square

Let us denote by $[\mathcal{Y}]$ the set of orbits in $\mathcal{Y} = \mathcal{Y}(M, \omega, \mathcal{P})$.

Corollary 8.6. *The map τ induces the embedding*

$$\bar{\tau} : \{\text{classes of PDQs}\} \rightarrow [\mathcal{Y}].$$

Proof. Let $(\mathbb{A}_t, \mathbb{O}_t)$ be a PDQ on (M, ω, \mathcal{P}) . Then, by Corollary 7.13, there exists a PSP, (μ_t, \mathcal{O}_t) , equivalent to $(\mathbb{A}_t, \mathbb{O}_t)$. We put $\bar{\tau}(\mathbb{A}_t, \mathbb{O}_t) = [\tau(\mu_t, \mathcal{O}_t)]$, the orbit passing through the pair $\tau(\mu_t, \mathcal{O}_t)$. By Propositions 7.14 and 8.5, this map is correctly defined, i.e. does not depend on the choice of a PSP equivalent to $(\mathbb{A}_t, \mathbb{O}_t)$. Proposition 8.5 also shows that being descended to equivalence classes of PDQs $\text{cl}_{\text{PSP}} Q$ is embedding. \square

In the next section, we will prove that any element of \mathcal{Y} is equal to $\tau(\mu_t, \mathcal{O}_t)$ for a PSP (μ_t, \mathcal{O}_t) , which implies that the map $\bar{\tau}$ is, in fact, an isomorphism.

9. Existence of PDQs and relation between the extension and Fedosov classes of a PDQ

Let (M, ω_0) be a symplectic manifold. It is known that all equivalence classes of DQs on M with the Poisson bracket ω_0^{-1} can be obtained by the Fedosov method. According to this method, starting with a symplectic connection, one constructs a flat connection, D (called the Fedosov connection) on the Weyl algebra that we define as a quotient of the tensor algebra over the cotangent bundle to M . A quantized algebra, \mathbb{A}_t , is realized as the subalgebra of flat sections in the Weyl algebra. The Weyl curvature of D (see below), being a closed scalar 2-form of the form $\theta_t = \omega_0 + t\omega_1 + \dots$, defines the Fedosov class

$$\text{cl}_F(\mathbb{A}_t) = [\theta_t] \in [\omega_0] + tH^2(M, \mathbb{C}[[t]]). \quad (9.1)$$

It is also known that the correspondence $\mathbb{A}_t \mapsto \text{cl}_F(\mathbb{A}_t)$ is a bijection between the set of equivalence classes of DQs on (M, ω_0) and the set $[\omega_0] + tH^2(M, \mathbb{C}[[t]])$ [11,18,23].

Let (M, ω, \mathcal{P}) be a PSM and $(\omega_t, \mathcal{P}_t) \in \mathcal{Y}(M, \omega, \mathcal{P})$ a deformation of the pair (ω, \mathcal{P}) , see Section 8.3. We adapt the Fedosov method to construct a PSP, (μ_t, \mathcal{O}_t) , such that $\tau(\mu_t, \mathcal{O}_t) = (\omega_t, \mathcal{P}_t)$. We start with a \mathcal{P}_t -symplectic connection, ∇ , and construct the Fedosov connection on the same Weyl algebra. We will see that by realizing the Fedosov scheme in presence of a polarization, the form ω_t appears as a so-called Wick curvature of the Fedosov connection. Moreover, ω_t differs from the Weyl curvature of that Fedosov connection by the form $(t/2) \text{tr}(\nabla^2|_{\mathcal{P}_t})$.

9.1. Some notations

Let \mathcal{E} be a formal vector bundle over M , i.e. a free $\mathcal{C}_M^\infty[[t]]$ -module of finite rank over M . Denote by $T^k(\mathcal{E})$ the k th tensor power of \mathcal{E} over $\mathcal{C}_M^\infty[[t]]$ and by $T(\mathcal{E})$ the corresponding tensor algebra completed in the $\{\mathcal{E}, t\}$ -adic topology. Similarly, we define the completed symmetric algebra $S(\mathcal{E})$. For a subbundle \mathcal{P} of \mathcal{E} , we denote by $\text{sym}_{\mathcal{P}} : S(\mathcal{P}) \rightarrow T(\mathcal{E})$ the natural embedding of $\mathcal{C}_M^\infty[[t]]$ -modules defined by symmetrization.

Let $\Lambda(\mathcal{E})$ be the exterior algebra of \mathcal{E} over $\mathcal{C}_M^\infty[[t]]$. We will consider $T(\mathcal{E}) \otimes \Lambda(\mathcal{E})$ as a graded super-algebra regarding a section $x \in T(\mathcal{E}) \otimes \Lambda^k(\mathcal{E})$ of degree k even (odd) if k is even (odd).

Denote by $\delta_{T(\mathcal{E})}$ the continuous $\mathcal{C}_M^\infty[[t]]$ -linear derivation of $T(\mathcal{E}) \otimes \Lambda(\mathcal{E})$ defined by the map $T^1(\mathcal{E}) \otimes 1 \rightarrow 1 \otimes \Lambda^1(\mathcal{E})$, $v \otimes 1 \mapsto 1 \otimes v$, $v \in \mathcal{E}$. It is clear that $\delta_{T(\mathcal{E})}$ is a derivation of degree 1 and $\delta_{T(\mathcal{E})}^2 = 0$. It is easy to see that for any subbundle $\mathcal{P} \subset \mathcal{E}$, the map $\delta_{T(\mathcal{E})}$ being restricted to $S(\mathcal{P}) \otimes \Lambda(\mathcal{P})$ via the embedding $\text{sym}_{\mathcal{P}} \otimes \text{id}_{\mathcal{P}}$ gives a derivation of the algebra $S(\mathcal{P}) \otimes \Lambda(\mathcal{P})$; we denote it by $\delta_{\mathcal{P}}$.

On the algebra $S(\mathcal{P}) \otimes \Lambda(\mathcal{P})$, there is another derivation, $\delta_{\mathcal{P}}^*$, of degree -1 generated by the map $1 \otimes \Lambda^1(\mathcal{P}) \rightarrow S^1(\mathcal{P}) \otimes 1$, $1 \otimes v \rightarrow v \otimes 1$, $v \in \mathcal{P}$. It is easy to check that $(\delta_{\mathcal{P}}^*)^2 = 0$ and $[\delta_{\mathcal{P}}, \delta_{\mathcal{P}}^*] = \delta_{\mathcal{P}}\delta_{\mathcal{P}}^* + \delta_{\mathcal{P}}^*\delta_{\mathcal{P}} = \deg$, where \deg is the derivation assigning to an element $x \in S^p(\mathcal{P}) \otimes \Lambda^q(\mathcal{P})$ the element $(p+q)x$.

Let \mathcal{E} be presented as a direct sum of $\mathcal{C}_M^\infty[[t]]$ -submodules, $\mathcal{E} = \mathcal{P} \oplus \mathcal{Q}$. It is obvious that the derivations $\delta_{\mathcal{P}}$, $\delta_{\mathcal{P}}^*$, $\delta_{\mathcal{Q}}$, $\delta_{\mathcal{Q}}^*$ induce derivations on the algebra $\mathbf{S}(\mathcal{P}, \mathcal{Q}) = (S(\mathcal{P}) \otimes S(\mathcal{Q})) \otimes (\Lambda(\mathcal{P}) \otimes \Lambda(\mathcal{Q}))$ that we will identify in the natural way with the algebra $(S(\mathcal{P}) \otimes S(\mathcal{Q})) \otimes \Lambda(\mathcal{E})$. We put $\delta_{\mathcal{P}, \mathcal{Q}} = \delta_{\mathcal{P}} + \delta_{\mathcal{Q}}$ and $\delta_{\mathcal{P}, \mathcal{Q}}^* = \delta_{\mathcal{P}}^* + \delta_{\mathcal{Q}}^*$.

Let us define the operator $\delta_{\mathcal{P}, \mathcal{Q}}^{-1}$ on $\mathbf{S}(\mathcal{P}, \mathcal{Q})$ in the following way. We put $\delta_{\mathcal{P}, \mathcal{Q}}^{-1}(x) = 0$ for $x \in \mathcal{C}_M^\infty[[t]]$ and $\delta_{\mathcal{P}, \mathcal{Q}}^{-1}(x) = (1/(p+r+q))\delta_{\mathcal{P}, \mathcal{Q}}^*(x)$ for $x \in (S^p(\mathcal{P}) \otimes S^r(\mathcal{Q})) \otimes \Lambda^q(\mathcal{E})$, $p+r+q > 0$. There is the obvious relation

$$\delta_{\mathcal{P}, \mathcal{Q}}\delta_{\mathcal{P}, \mathcal{Q}}^{-1} + \delta_{\mathcal{P}, \mathcal{Q}}^{-1}\delta_{\mathcal{P}, \mathcal{Q}} = \text{projection on } \mathbf{S}^+(\mathcal{P}, \mathcal{Q}) \text{ along } \mathcal{C}_M^\infty[[t]], \quad (9.2)$$

where $\mathbf{S}^+(\mathcal{P}, \mathcal{Q})$ is the closure of $\oplus_{p+r+q>0}(S^p(\mathcal{P}) \otimes S^r(\mathcal{Q})) \otimes \Lambda^q(\mathcal{E})$.

One has the embedding

$$\text{sym}_{\mathcal{P}} \otimes \text{sym}_{\mathcal{Q}} \otimes \text{id} : (S(\mathcal{P}) \otimes S(\mathcal{Q})) \otimes \Lambda(\mathcal{E}) \rightarrow T(\mathcal{E}) \otimes \Lambda(\mathcal{E}). \quad (9.3)$$

It is clear that $\delta_{\mathcal{P}, \mathcal{Q}}$ coincides with the restriction of $\delta_{T(\mathcal{E})}$ to $\mathbf{S}(\mathcal{P}, \mathcal{Q})$ via this embedding.

9.2. The Fedosov algebra

Let $\varphi : \mathcal{E} \otimes \mathcal{E} \rightarrow \mathcal{C}_M^\infty[[t]]$ be a $\mathcal{C}_M^\infty[[t]]$ -linear skew-symmetric nondegenerate form and I the closed ideal in $T(\mathcal{E})$ generated by relations

$$x \otimes y - y \otimes x = t\varphi(x, y). \quad (9.4)$$

We call $\mathcal{W}(\mathcal{E}) = T(\mathcal{E})/I$ the Weyl algebra and $\mathbf{W} = \mathbf{W}(\mathcal{E}) = \mathcal{W} \otimes \Lambda(\mathcal{E})$ the Fedosov algebra over \mathcal{E} . The derivation $\delta_{T(\mathcal{E})}$ on $T(\mathcal{E}) \otimes \Lambda(\mathcal{E})$ induces a derivation on \mathbf{W} . Indeed, $\delta_{T(\mathcal{E})}$ applied to the both sides of (9.4) gives zero. We denote this derivation by δ .

Let $\mathcal{E} = \mathcal{P} \oplus \mathcal{Q}$ be a decomposition into $\mathcal{C}_M^\infty[[t]]$ -modules.

Define the Wick map, $\mathbf{w}_{\mathcal{P}, \mathcal{Q}}$, as the composition $\mathbf{S}(\mathcal{P}, \mathcal{Q}) \rightarrow T(\mathcal{E}) \otimes \Lambda(\mathcal{E}) \rightarrow \mathbf{W}$, where the first map is (9.3) and the second one is the projection. By the PBW theorem $\mathbf{w}_{\mathcal{P}, \mathcal{Q}}$ is an isomorphism of $\mathcal{C}_M^\infty[[t]]$ -modules.

Due to the isomorphism $\mathbf{w}_{\mathcal{P}, \mathcal{Q}}$, the operators $\delta_{\mathcal{P}, \mathcal{Q}}$ and $\delta_{\mathcal{P}, \mathcal{Q}}^{-1}$ are carried over from $\mathbf{S}(\mathcal{P}, \mathcal{Q})$ to \mathbf{W} . We retain for them the same notation. Note that while $\delta_{\mathcal{P}, \mathcal{Q}}$ does not depend on the decomposition of \mathcal{E} and coincides with the derivation δ induced from $T(\mathcal{E}) \otimes \Lambda(\mathcal{E})$, the operator $\delta_{\mathcal{P}, \mathcal{Q}}^{-1}$ is not a derivation and does depend on the decomposition. In particular, one can suppose that the decomposition is trivial, $\mathcal{E} = \mathcal{E} \oplus 0$. In this case we denote $\delta_{\mathcal{E}}^{-1} = \delta_{\mathcal{E}, 0}^{-1}$.

Proposition 9.1. *One has*

$$H(\mathbf{W}, \delta) = \mathcal{C}_M^\infty[[t]].$$

Moreover, if $x \in \mathcal{W}(\mathcal{E}) \otimes \Lambda^{k>0}(\mathcal{E})$, then $y = \delta_{\mathcal{P}, \mathcal{Q}}^{-1}x$ is such that $\delta y = x$ for any decomposition $\mathcal{E} = \mathcal{P} \oplus \mathcal{Q}$.

Proof. Follows from (9.2). □

9.3. Lie subalgebras in \mathcal{W}

Let $\mathcal{E} = \mathcal{P} \oplus \mathcal{Q}$ be a decomposition. We say that $x \in \mathbf{W}$ has $\mathbf{w}_{\mathcal{P}, \mathcal{Q}}$ -degree (p, q) if $\mathbf{w}_{\mathcal{P}, \mathcal{Q}}^{-1}(x) \in (S^p(\mathcal{P}) \otimes S^q(\mathcal{Q})) \otimes \Lambda(\mathcal{E})$. We say that $x \in \mathbf{W}$ has $\mathbf{w}_{\mathcal{P}, \mathcal{Q}}$ -degree k if $\mathbf{w}_{\mathcal{P}, \mathcal{Q}}^{-1}(x) \in$

$\oplus_{p+q=k} (S^p(\mathcal{P}) \otimes S^q(\mathcal{Q})) \otimes \Lambda(\mathcal{E})$. We define the $\mathbf{w}_{\mathcal{E}}$ -degree as the $\mathbf{w}_{\mathcal{E},0}$ -degree for the trivial decomposition $\mathcal{E} = \mathcal{E} \oplus 0$.

Let \mathfrak{g} be a sheaf of Lie algebras acting on \mathcal{E} . We call a $C_M^\infty[[t]]$ -linear map $\lambda : \mathfrak{g} \rightarrow \mathcal{W}$ a realization of \mathfrak{g} , if it is a Lie algebra morphism (\mathcal{W} is considered as a Lie algebra with respect to the commutator $(1/t)[\cdot, \cdot]$) and for any $x \in \mathfrak{g}$ and $v \in \mathcal{E}$ one has $x(v) = (1/t)[\lambda(x), v]$. It is easy to check that any two realizations differ by a Lie algebra morphism of \mathfrak{g} to the center of \mathcal{W} , therefore, if \mathfrak{g} is a sheaf of semisimple Lie algebras, there is not more than one realization of \mathfrak{g} .

Denote by $\mathfrak{sp}(\mathcal{E})$ the sheaf of symplectic Lie algebras with respect to φ . Since $\mathfrak{sp}(\mathcal{E})$ is semisimple, there is a unique realization $\rho_{\mathcal{E}} : \mathfrak{sp}(\mathcal{E}) \rightarrow \mathcal{W}$. The image of this realization consists of elements having $\mathbf{w}_{\mathcal{E}}$ -degree 2.

Let $\mathcal{E} = \mathcal{P} \oplus \mathcal{Q}$ be a decomposition into Lagrangian subsheaves. Denote by $\mathfrak{sp}(\mathcal{P}, \mathcal{E})$ the subsheaf of $\mathfrak{sp}(\mathcal{E})$ preserving \mathcal{P} . It is easy to check that $\mathfrak{sp}(\mathcal{P}, \mathcal{E})$ can be realized as the subset of elements of \mathcal{W} having $\mathbf{w}_{\mathcal{P}, \mathcal{Q}}$ -degree $(1, 1)$ or $(2, 0)$. Denote this realization by $\rho_{\mathcal{P}, \mathcal{E}} : \mathfrak{sp}(\mathcal{P}, \mathcal{E}) \rightarrow \mathcal{W}$. On the other hand, $\mathfrak{sp}(\mathcal{P}, \mathcal{E})$ is realized in \mathcal{W} by $\rho_{\mathcal{E}}$.

Lemma 9.2. *Let $b \in \mathfrak{sp}(\mathcal{P}, \mathcal{E})$. Then*

$$\rho_{\mathcal{E}}(b) = \rho_{\mathcal{P}, \mathcal{E}}(b) + \frac{t}{2} \text{trace}(\bar{b}),$$

where \bar{b} is b restricted to \mathcal{P} .

Proof. Direct computation using the fact that $\rho_{\mathcal{P}, \mathcal{E}}(\bar{b})$ is the component of $\rho_{\mathcal{P}, \mathcal{E}}(b)$ of $\mathbf{w}_{\mathcal{P}, \mathcal{Q}}$ -degree $(1, 1)$ in any decomposition $\mathcal{E} = \mathcal{P} \oplus \mathcal{Q}$. \square

9.4. Filtrations on \mathcal{W}

We define two decreasing filtrations on \mathcal{W} numbered by nonnegative integers.

The T -filtration $F_\bullet^T \mathcal{W}$ is defined as follows. We ascribe to the elements of \mathcal{E} -degree 1 and to t -degree 2. Then $F_n^T \mathcal{W}$ consists of elements of \mathcal{W} having the leading term of total degree $\geq n$.

The \mathcal{P} -filtration, $F_\bullet^{\mathcal{P}} \mathcal{W}$, is firstly defined on $S(\mathcal{P}) \otimes S(\mathcal{Q})$ by the subsets $F_n^{\mathcal{P}} = \bigoplus_{i \geq n} S^i(\mathcal{P}) \otimes S(\mathcal{Q})$, $n = 0, 1, \dots$, and carried over to \mathcal{W} via the Wick isomorphism.

We extend those filtrations to \mathbf{W} in the natural way standing, for example, $F_n^T \mathbf{W} = F_n^T \mathcal{W} \otimes \Lambda(\mathcal{E})$. We will use the following mnemonic notation. To point out, for example, that a section $x \in \mathbf{W}$ belongs to $F_n^T \mathbf{W}$ we write $F^T(x) \geq n$.

In the following we denote $\mathbf{S}(\mathcal{P}) = S(\mathcal{P}) \otimes \Lambda(\mathcal{E})$ embedded in $\mathbf{S}(\mathcal{P}, \mathcal{Q})$ as $(S(\mathcal{P}) \otimes 1) \otimes \Lambda(\mathcal{E})$.

Proposition 9.3. *Let $\mathcal{E} = \mathcal{P} \oplus \mathcal{Q}$ be a decomposition of \mathcal{E} into Lagrangian subsheaves. Then*

- (a) *The Wick map $\mathbf{w} = \mathbf{w}_{\mathcal{P}, \mathcal{Q}} : \mathbf{S}(\mathcal{P}, \mathcal{Q}) \rightarrow \mathbf{W}$ has the following property: for $a \in \mathbf{S}(\mathcal{P})$ and arbitrary $c \in \mathbf{S}(\mathcal{P}, \mathcal{Q})$ one has $\mathbf{w}(ac) = \mathbf{w}(a)\mathbf{w}(c)$.*

The filtrations on \mathbf{W} have the properties:

- (b) for $x, y \in \mathbf{W}$, if $F^{\mathcal{P}}(x) \geq k$, then $F^{\mathcal{P}}(xy) \geq k$;
- (c) $F^{\mathcal{P}}(\delta_{\mathcal{P}, \mathcal{Q}}^{-1}x) \geq F^{\mathcal{P}}(x)$;
- (d) $F^T(\delta_{\mathcal{P}, \mathcal{Q}}^{-1}x) > F^T(x)$.

Proof. Clear. □

9.5. Fedosov's construction in the Wick case

Let $(M, \omega_t, \mathcal{P}_t)$ be a formal PSM. We will write for shortness $\omega = \omega_t$, $\mathcal{P} = \mathcal{P}_t$. Let us denote $\mathcal{T} = \mathcal{T}_M^{\mathbb{C}}[[t]]$. It is easy to prove that there exists on M a Lagrangian subbundle $\mathcal{Q} \subset \mathcal{T}$ such that $\mathcal{T} = \mathcal{P} \oplus \mathcal{Q}$.

In the following, we set $\mathcal{E} = \mathcal{T}^*$ and consider the Fedosov algebra $\mathbf{W} = \mathbf{W}(\mathcal{E})$ with respect to φ being the Poisson bracket inverse to ω . The decomposition $\mathcal{E} = \mathcal{P}^\perp \oplus \mathcal{Q}^\perp$ is Lagrangian with respect to this φ .

Let ∇ be a \mathcal{P} -symplectic connection on M (see Section 5). Then the induced connection $\nabla : \mathcal{E} \rightarrow \mathcal{E} \otimes \Lambda^1 \mathcal{E}$ on \mathcal{E} preserves \mathcal{P}^\perp , i.e. $\nabla(\mathcal{P}^\perp) \subset \mathcal{P}^\perp \otimes \Lambda^1(\mathcal{E})$, and is flat on \mathcal{P}^\perp along \mathcal{P} , i.e. for any $X, Y \in \mathcal{P}$ one has $(\nabla_X \nabla_Y - \nabla_Y \nabla_X - \nabla_{[X, Y]})(\mathcal{P}^\perp) = 0$.

We will identify \mathcal{P} with \mathcal{P}^\perp and \mathcal{Q} with \mathcal{Q}^\perp by the isomorphism $x \mapsto \omega(x, \cdot)$ between \mathcal{T} and \mathcal{E} . Thus, we allow us to write $\mathcal{E} = \mathcal{P} \oplus \mathcal{Q}$.

The connection ∇ gives a derivation of the Fedosov algebra \mathbf{W} , which is an extension of the de Rham differential on functions. Analogously, ∇ gives derivations of the algebras $T(\mathcal{E}) \otimes \Lambda(\mathcal{E})$, $S(\mathcal{E}) \otimes \Lambda(\mathcal{E})$, and $(S(\mathcal{P}) \otimes S(\mathcal{Q})) \otimes \Lambda(\mathcal{E})$. These derivations commute with the maps (9.3) and $w_{\mathcal{P}, \mathcal{Q}}$.

For convenience, we will mark the elements of the Fedosov algebra lying in $\mathcal{E} \otimes 1$ by letters with hat over them (\hat{x}), while by dx we will denote the copy of \hat{x} lying in $1 \otimes \Lambda^1 \mathcal{E}$.

Let $\omega = \omega_{ij} dx_i \wedge dx_j$ in some local coordinates. It is easy to check that for $\tilde{\delta} = \omega_{ij} \hat{x}_i \otimes dx_j$ one has

$$\delta = \frac{1}{t} \text{ad}(\tilde{\delta}), \quad \tilde{\delta}^2 = t\omega. \quad (9.5)$$

Since the torsion of ∇ is equal to zero,

$$\nabla(\tilde{\delta}) = 0. \quad (9.6)$$

Since ∇^2 is a $\mathcal{C}_M^{\infty}[[t]]$ -linear derivation of degree 0 preserving \mathcal{P} , there is an element $R \in \rho_{\mathcal{P}, \mathcal{E}}(\mathfrak{sp}(\mathcal{P}, \mathcal{E})) \otimes \Lambda^2(\mathcal{E})$ such that $\nabla^2 = (1/t) \text{ad}(R)$. In particular, one has to be

$$F^{\mathcal{P}}(R) \geq 1. \quad (9.7)$$

According to Fedosov [11], we also define $R^F \in \rho_{\mathcal{E}}(\mathfrak{sp}(\mathcal{P}, \mathcal{E})) \otimes \Lambda^2(\mathcal{E})$ satisfying $\nabla^2 = (1/t) \text{ad}(R^F)$. Equality (9.6) implies that

$$\delta(R) = \delta(R^F) = 0. \quad (9.8)$$

Following to Fedosov, we will consider connections on \mathbf{W} of the form

$$D = \nabla + \frac{1}{t} \text{ad}(\gamma), \quad \gamma \in \mathcal{W} \otimes \Lambda^1(\mathcal{E}). \quad (9.9)$$

We define the Wick curvature of D as

$$\Omega_D = R + \nabla(\gamma) + \frac{1}{t}\gamma^2.$$

According to Fedosov, we also define the Weyl (or Fedosov) curvature of D as

$$\Omega_D^F = R^F + \nabla(\gamma) + \frac{1}{t}\gamma^2. \quad (9.10)$$

Since by Lemma 9.2 $R^F = R + (t/2)\text{tr}(\nabla^2|_{\mathcal{P}})$, we have

$$\Omega_D^F = \Omega_D + \frac{t}{2}\text{tr}(\nabla^2|_{\mathcal{P}}). \quad (9.11)$$

One checks

$$D^2 = \frac{1}{t}\text{ad}(\Omega_D) = \frac{1}{t}\text{ad}(\Omega_D^F). \quad (9.12)$$

Let us take γ in the form

$$\gamma = \tilde{\delta} + r, \quad r \in \mathcal{W} \otimes \Lambda^1(\mathcal{E}), \quad F^T(r) \geq 3. \quad (9.13)$$

Then the connection D has the form

$$D = \nabla + \delta + \frac{1}{t}\text{ad}(r). \quad (9.14)$$

Using (9.5) and (9.6), we obtain that its Wick curvature is

$$\Omega_D = R + \nabla(\tilde{\delta} + r) + \frac{1}{t}(\tilde{\delta} + r)^2 = \omega + \delta r + R + \nabla r + \frac{1}{t}r^2. \quad (9.15)$$

Proposition 9.4. *There exists an element $r \in \mathcal{W}(\mathcal{E}) \otimes \Lambda^1(\mathcal{E})$ such that*

- (a) $F^T(r) \geq 3$;
- (b) $F^{\mathcal{P}}(r) \geq 1$;
- (c) *the connection $D = \nabla + \delta + (1/t)\text{ad}(r)$ is flat, i.e. $D^2 = 0$;*
- (d) *for its Wick curvature one has*

$$\Omega_D = \omega;$$

- (e) *its Weyl curvature Ω_D^F belongs to $\Gamma(M, d\mathcal{P}^\perp)$ and there is the formula*

$$\Omega_D^F = \omega + \frac{t}{2}\text{tr}(\nabla^2|_{\mathcal{P}}). \quad (9.16)$$

Proof. First of all, we apply the Fedosov method [11, Theorem 5.2.2], to find r satisfying d). According to (9.15), r must obey the equation

$$\delta r = - \left(R + \nabla r + \frac{1}{t}r^2 \right). \quad (9.17)$$

Let us look for r being the limit of the sequence, $r = \lim r_k$, where $r_k \in \mathcal{W}(\mathcal{E}) \otimes \Lambda^1(\mathcal{E})$, $k = 3, 4, \dots$, and $F^T(r_k - r_{k-1}) \geq k$. As in Lemma 5.2.3 of [11], using Proposition 9.3 (d) and the fact that $F^T(R) \geq 2$, such r_k can be calculated recursively:

$$r_3 = -\delta_{\mathcal{P}, \mathcal{Q}}^{-1}(R), \quad r_{k+3} = -\left(r_3 + \delta_{\mathcal{P}, \mathcal{Q}}^{-1}\left(\nabla r_{k+2} + \frac{1}{t}r_{k+2}^2\right)\right). \quad (9.18)$$

Hence (a) and (d) are proved.

Let us prove that $F^{\tilde{\mathcal{P}}}(r_k) \geq 1$ for all k . The inequality $F^{\mathcal{P}}(r_3) \geq 1$ follows from the fact that $F^{\mathcal{P}}(R) \geq 1$ and from Proposition (9.3) (c). Suppose that $F^{\tilde{\mathcal{P}}}(r_i) \geq 1$ for $i < k+3$, $k > 0$. Then $F^{\mathcal{P}}(\nabla r_{k+2}) \geq 1$ because ∇ preserves \mathcal{P} . On the other hand, $F^{\mathcal{P}}(r_{k+2}^2) \geq 1$ because of Proposition (9.3) (b), therefore, as follows from (9.18), $F^{\mathcal{P}}(r_{k+3}) \geq 1$ as well. Hence, r being the limit of the convergent sequence r_k satisfies the conditions (a), (b), and (d) of the proposition.

(c) obviously follows from (d) and (9.12).

(e) follows immediately from (d), (9.11), and Lemma 5.3. \square

Proposition 9.5. *Let $\tilde{\nabla}$ be another \mathcal{P} -symplectic connection on M . Let $\tilde{r} \in \mathcal{W}(\mathcal{E}) \otimes \Lambda^1(\mathcal{E})$, satisfy the conclusions of Proposition 9.4, in particular, the connection $\tilde{D} = \tilde{\nabla} + \delta + (1/t) \text{ad}(\tilde{r})$ is flat. Then, there exists an element $B \in \mathcal{W}(\mathcal{E})$ such that*

- (a) $F^T(B) \geq 3$;
- (b) $F^{\mathcal{P}}(B) \geq 1$;
- (c) $e^{(1/t) \text{ad} B} D = \tilde{D}$.

Proof. Note that $\nabla - \tilde{\nabla}$ can be presented as $(1/t) \text{ad } R_0$, where $R_0 \in \rho_{\mathcal{P}, \mathcal{E}}(\mathfrak{sp}(\mathcal{P}, \mathcal{E}))$. Therefore, $F^T(R_0) \geq 2$ and $F^{\mathcal{P}}(R_0) \geq 1$. Let us put $R_1 = r - \tilde{r}$. Then, $F^T(R_1) \geq 3$ and $F^{\mathcal{P}}(R_1) \geq 1$. We have

$$\tilde{D} = D - \frac{1}{t} \text{ad}(R_0 + R_1).$$

Since $\Omega_D = \Omega_{\tilde{D}} = \omega$, using (9.6) we obtain

$$\delta(R_0) = 0.$$

It follows that the element $B_0 = \delta_{\mathcal{P}, \mathcal{Q}}^{-1}(R_0)$ is such that $\delta(B_0) = R_0$ and $F^T(B_0) \geq 3$, $F^{\mathcal{P}}(B_0) \geq 1$.

Replacing D by $e^{(1/t) \text{ad} B_0} D$, we obtain

$$\tilde{D} = D - \frac{1}{t} \text{ad}(R'_0 + R'_1),$$

where $F^T(R'_0) \geq 3$, $F^{\mathcal{P}}(R'_0) \geq 1$ and $F^T(R'_1) \geq 4$, $F^{\mathcal{P}}(R'_1) \geq 1$.

Proceeding by induction on F^T -filtration, we obtain a sequence $B_i \in \mathcal{W}(\mathcal{E})$ with increasing F^T -filtration such that $\prod_{i=0}^{\infty} e^{(1/t) \text{ad} B_i}(D) = \tilde{D}$.

Since elements $e^{(1/t) \text{ad} B'}$, $B' \in \mathcal{W}(\mathcal{E})$, $F^T(B') \geq 3$, form a pro-unipotent Lie group, there exists an element $B \in \mathcal{W}(\mathcal{E})$, $F^T(B) \geq 3$, $F^{\mathcal{P}}(B) \geq 1$, such that $\prod_{i=0}^{\infty} e^{(1/t) \text{ad} B_i} = e^{(1/t) \text{ad} B}$. \square

Let D be a connection satisfying [Proposition 9.4](#). Denote by \mathcal{W}_D the subsheaf of \mathcal{W} consisting of flat sections a , i.e. such that $Da = 0$. Since D is a derivation of \mathbf{W} , it is clear that \mathcal{W}_D is a sheaf of subalgebras. Let $\sigma = \text{id} - (\delta\delta_{\mathcal{P}, \mathcal{Q}}^{-1} + \delta_{\mathcal{P}, \mathcal{Q}}^{-1}\delta)$. Then, as follows from [\(9.2\)](#), $\sigma : \mathcal{W} \rightarrow \mathcal{C}_M^\infty[[t]]$ is a projection, where $\mathcal{C}_M^\infty[[t]]$ is considered as the center of the algebra \mathcal{W} .

Proposition 9.6.

- (a) *The map $\sigma : \mathcal{W}_D \rightarrow \mathcal{C}_M^\infty[[t]]$ is a bijection.*
- (b) *The inverse map $\eta : \mathcal{C}_M^\infty[[t]] \rightarrow \mathcal{W}_D$ has the form $\eta(f) = f + \hat{f}$, there $F^T(\hat{f}) > F^T(f)$.*
- (c) *If $\text{d}f \in \mathcal{P}$, then $F^{\mathcal{P}}(\hat{f}) \geq 1$.*
- (d) *If $\text{d}f \in \mathcal{P}$, then $\sigma(\eta(f)\eta(g)) = fg$ for any $g \in \mathcal{C}_M^\infty[[t]]$.*

Proof. Again, we apply the Fedosov iteration procedure. According to [\[11\]](#), Theorem 5.2.4, we look for $\eta(f)$ as a limit, $\eta(f) = \lim a_k$, there $a_k \in \mathcal{W}$ can be calculated recursively:

$$a_0 = f, \quad a_{k+1} = a_0 + \delta_{\mathcal{P}, \mathcal{Q}}^{-1} \left(\nabla a_k + \frac{1}{t} \text{ad } r(a_k) \right). \quad (9.19)$$

Put $\hat{f} = \eta(f) - a_0$. As in [\[11\]](#), Theorem 5.2.4, one proves that such $\eta(f)$ and \hat{f} satisfy (a) and (b). Now observe that $a_1 - a_0 = \delta_{\mathcal{P}, \mathcal{Q}}^{-1}(1 \otimes \text{d}f)$ and if $\text{d}f \in \mathcal{P}$, then $F^{\mathcal{P}}(a_1 - a_0) \geq 1$. By induction, we conclude that $F^{\mathcal{P}}(a_k - a_0) \geq 1$ for all $k \geq 1$. Therefore, $F^{\mathcal{P}}(a - a_0) \geq 1$ as well, which proves (c).

Let us prove (d). We have $\eta(f)\eta(g) = f\eta(g) + \hat{f}\eta(g)$. Since by (c) $F^{\mathcal{P}}(\hat{f}) \geq 1$, $F^{\mathcal{P}}(\hat{f}\eta(g)) \geq 1$ as well. It follows that $\sigma(\hat{f}\eta(g)) = 0$ and $\sigma(\eta(f)\eta(g)) = \sigma(f\eta(g)) = fg$, because σ is a $\mathcal{C}_M^\infty[[t]]$ -linear map and $\sigma(\eta(g)) = g$. \square

9.6. Existence of PSPs

Let (M, ω, \mathcal{P}) be a PSM. Recall that in [Section 8.3](#) we assigned to any PSP (μ_t, \mathcal{O}_t) on (M, ω, \mathcal{P}) an element $\tau(\mu_t, \mathcal{O}_t) \in \mathcal{Y}(M, \omega, \mathcal{P})$, which is a pair $(\omega_t, \mathcal{P}_t)$ being a deformation of the pair (ω, \mathcal{P}) . The form ω_t represents the extension class $\text{cl}_{\text{Ext}}(\mu_t, \mathcal{O}_t)$.

We are going to show that any element of $\mathcal{Y}(M, \omega, \mathcal{P})$ corresponds to a PSP.

Proposition 9.7.

- (a) *For any pair $(\omega_t, \mathcal{P}_t) \in \mathcal{Y}(M, \omega, \mathcal{P})$, there exists a PSP, (μ_t, \mathcal{O}_t) , such that*
- $\tau(\mu_t, \mathcal{O}_t) = (\omega_t, \mathcal{P}_t).$ (9.20)
- (b) *The Fedosov class of the corresponding SP ($\mathcal{C}_M^\infty[[t]]$, μ_t) is represented by the form of $\omega + t\Gamma(M, \text{d}(\mathcal{P}_t^\perp))$ equal to*

$$\theta_t = \omega_t + \frac{t}{2} \text{tr}(\nabla^2|_{\mathcal{P}_t}), \quad (9.21)$$

where ∇ is a \mathcal{P}_t -symplectic connection on the formal symplectic manifold $(M, \omega_t, \mathcal{P}_t)$.

Proof. Let \mathbf{W} be the Fedosov algebra on M corresponding to the symplectic form ω_t . Let ∇ be a \mathcal{P}_t -symplectic connection on M corresponding to ω_t and D be the flat connection on \mathbf{W} constructed in [Proposition 9.4](#) (c). Let \mathcal{W}_D be the sheaf of flat sections of \mathbf{W} . Define an SP $(\mathcal{C}_M^\infty[[t]], \mu_t)$ on M carrying over the multiplication from \mathcal{W}_D to $\mathcal{C}_M^\infty[[t]]$ via the map σ from [Proposition 9.6](#). Point (d) of that proposition shows that, in fact, this SP presents the PSP $(\mathcal{C}_M^\infty[[t]], \mu_t, \mathcal{O}_{\mathcal{P}_t})$. We are going to prove that this SP is as required. In the following we identify \mathcal{W}_D with the corresponding PSP via σ .

Let us prove (9.20). Let (U_α) be an open covering of M such that on each U_α there exist formal Darboux coordinates $x_{\alpha i}, y_{\alpha i}, z_{\alpha i} \in \mathcal{O}_{\mathcal{P}_t}$ with respect to ω_t .

Denote by ∇_α the standard flat \mathcal{P}_t -symplectic connections over U_α such that the forms $dx_{\alpha i}, dy_{\alpha i}$ are flat sections in \mathcal{P}_t^\perp . Then, the connections $D_\alpha = \nabla_\alpha + \delta$ satisfy on U_α the statements (c)–(e) of [Proposition 9.6](#). Let \mathcal{W}_{D_α} be the SP on U_α constructed in [Proposition 9.6](#) via flat sections of D_α . It is easy to see that \mathcal{W}_{D_α} coincides with the Moyal–Wick PSP with respect to the Darboux coordinates $x_{\alpha i}, y_{\alpha i}$ (see [Example 7.15](#)), so that $x_{\alpha i}, y_{\alpha i}$ are also Darboux coordinates for the commutator $(1/t)[\cdot, \cdot]$ in \mathcal{W}_{D_α} .

Since D and D_α have the same Wick curvature ω_t , there exist, by [Proposition 9.5](#), elements $B_\alpha \in \mathcal{W}$ such that $e^{(1/t)\text{ad } B_\alpha} D_\alpha = D$. It is clear that $e^{(1/t)\text{ad } B_\alpha}$ acting on \mathcal{W} takes \mathcal{W}_{D_α} to \mathcal{W}_D , and point (b) of that Proposition implies that it is identical on $\mathcal{O}_{\mathcal{P}_t}$.

Let $\Psi_{\alpha, \beta} = e^{-(1/t)\text{ad } B_\alpha} e^{(1/t)\text{ad } B_\beta}$. These may be considered as isomorphisms gluing SPs \mathcal{W}_{D_α} on $U_\alpha \cap U_\beta$ to a global SP on M . This global SP is obviously isomorphic to \mathcal{W}_D . We see, that functions $x_{\alpha i}, y_{\alpha i}$ form local Darboux coordinates corresponding to that SP. Therefore, the characteristic 2-form $\text{cl}_{\text{PSP}}(\mathcal{W}_D)$ (see [Section 8.3](#)) is locally represented as $dy_{\alpha i} \wedge dx_{\alpha i}$. On the other hand, this form is equal to ω_t , since from very beginning the functions $x_{\alpha i}, y_{\alpha i}$ was chosen as Darboux coordinates for ω_t . Hence, $\tau(\mathcal{W}_D) = (\omega_t, \mathcal{P}_t)$.

(b) follows from [Proposition 9.4](#) (e) and [Lemma 5.3](#). \square

10. The main theorem and corollaries

Let (M, ω, \mathcal{P}) be a PSM. Denote by \mathcal{Y} the set of pairs $(\omega_t, \mathcal{P}_t)$, where $\omega_t = \omega + t\omega_1 + \dots$ is a deformed symplectic form and $\mathcal{P}_t, \mathcal{P}_0 = \mathcal{P}$, its polarization. Let $\text{Aut}(M)$ be the group of formal automorphisms of M .

Theorem 10.1.

- (a) *The equivalence classes of PDQs on (M, ω, \mathcal{P}) are in one-to-one correspondence with the orbits in \mathcal{Y} under the $\text{Aut}(M)$ -action.*
- (b) *Let the pair $(\omega_t, \mathcal{P}_t)$ be a point on the orbit corresponding to a PDQ $(\mathbb{A}_t, \mathbb{O}_t)$. Then, $(\mathbb{A}_t, \mathbb{O}_t)$ is isomorphic to a PSP, $(\mathcal{C}_M^\infty[[t]], \mu_t, \mathcal{O}_t)$, where \mathcal{O}_t consists of functions constant along \mathcal{P}_t and the multiplication μ_t satisfies the condition*

$$\mu_t(f, g) = fg \quad \text{for } f \in \mathcal{O}_t, \quad g \in \mathcal{C}_M^\infty[[t]].$$

- (c) *The form ω_t represents the extension class $\text{cl}_{\text{Ext}}(\mathbb{A}_t, \mathbb{O}_t) \in H^1(M, \Omega_{\mathcal{O}_t}^{1\text{cl}})$, associated with $(\mathbb{A}_t, \mathbb{O}_t)$.*

- (d) Under the hypothesis of (b), the Fedosov class of the DQ \mathbb{A}_t can be represented by a form θ_t that is a deformation of ω and polarized by \mathcal{P}_t . It is defined by the formula

$$\theta_t = \omega_t + \frac{t}{2} \text{tr}(\nabla^2|_{\mathcal{P}_t}), \quad (10.1)$$

where ∇ is a \mathcal{P}_t -symplectic connection on the formal symplectic manifold $(M, \omega_t, \mathcal{P}_t)$.

Proof. Parts (a) and (b) follow from Proposition 8.5, Corollary 8.6, and Proposition 9.7 (a). Part (c) follows from Lemma 8.2. Part (d) is the same as Proposition 9.7 (b). \square

Remark 10.2. We have interpreted the form ω_t from (10.1) as a representative of the extension class associated to a PDQ (see Section 8). The form $\text{tr}(\nabla^2|_{\mathcal{P}_t})$ is the curvature form of the connection induced by $\nabla|_{\mathcal{P}_t}$ on the complex line bundle $\det(\mathcal{P}_t)$. Actually, $\det(\mathcal{P}_t)$ can be presented as a line \mathcal{O}_t -bundle, i.e. as a locally free sheaf of \mathcal{O}_t -modules of rank one. Indeed, $\det(\mathcal{P}_t) = \mathcal{C}_M^\infty[[t]] \otimes_{\mathcal{O}_t} \mathcal{L}$, where $\mathcal{L} = \Omega_{\mathcal{O}_t}^n$, $n = (1/2) \dim M$. The form $-\text{tr}(\nabla^2|_{\mathcal{P}_t})$, as well as minus curvature form of any other connection on \mathcal{L} , can be interpreted as a representative of the extension class of an \mathcal{O}_t -extension of $\text{Der}(\mathcal{O}_t)$ associated with \mathcal{L} . Indeed, let $\tilde{T}_{\mathcal{L}}$ denote the sheaf of \mathcal{O}_t -differential operators on \mathcal{L} of order at most one. Then $\tilde{T}_{\mathcal{L}}$ is equipped with the left \mathcal{O}_t -module structure and with the Lie bracket being the commutator of differential operators. Moreover, $\tilde{T}_{\mathcal{L}}$ represents in the natural way an \mathcal{O}_t -extension of $\text{Der}(\mathcal{O}_t)$. Splittings of this extension are flat connection on \mathcal{L} . Let d_α be local flat connections on \mathcal{L} on some open covering $\{U_\alpha\}$ of M . Then, $d_\alpha - d_\beta$ are closed 1-forms of $\Omega_{\mathcal{O}_t}^1$ that form a Čech cocycle. Hence, there exist smooth 1-forms $f_\alpha \in \mathcal{P}_t^\perp$ such that $d_\alpha - d_\beta = f_\alpha - f_\beta$. Differential operators $d_\alpha - f_\alpha$ form a global connection on \mathcal{L} , $\nabla_{\mathcal{L}}$, with the curvature locally equal to $-df_\alpha$. On the other hand, by definition (see Section 8), the extension class of $\tilde{T}_{\mathcal{L}}$ is represented by the form $df_\alpha \in d\mathcal{P}_t^\perp$, i.e. $-\nabla_{\mathcal{L}}^2$. Therefore, projecting the equality (10.1) to $\Gamma(M, d\mathcal{P}_t^\perp)/d(\Gamma(M, \mathcal{P}_t^\perp))$, we obtain

$$[\theta_t] = \text{cl}_{\text{Ext}}(\mu_t, \mathcal{O}_t) - \frac{t}{2} \text{cl}_{\text{Ext}}(\tilde{T}_{\det(\mathcal{P}_t)}). \quad (10.2)$$

Details are left to the reader.

Remark 10.3. Note that the form $-(1/2\pi\sqrt{-1}) \text{tr}(\nabla^2|_{\mathcal{P}_t})$ represents the first Chern class of \mathcal{P} [15]. Therefore, projecting (10.1) to $H^2(M, \mathbb{C}[[t]])$ we obtain the formula

$$[\theta_t] = [\omega_t] - \pi\sqrt{-1}tc_1(\mathcal{P}), \quad (10.3)$$

where $[\theta_t]$ and $[\omega_t]$ denote the corresponding classes in $H^2(M, \mathbb{C}[[t]])$. It is easy to see that after normalization of these classes and of the parameter t as in [15,19], formula (10.3) coincides with the analogous formula of those papers for the quantization with separation of variables on Kähler manifolds.

Corollary 10.4. Let \mathbb{A}_t be a DQ on (M, ω) . Suppose its Fedosov class $\text{cl}_F(\mathbb{A}_t)$ is represented by the form θ_t that has a polarization \mathcal{P}_t . Then \mathbb{A}_t can be extended to a PDQ $(\mathbb{A}_t, \mathbb{O}_t)$, where \mathbb{O}_t is isomorphic to $\mathcal{O}_{\mathcal{P}_t}$.

Proof. Let ∇ be a \mathcal{P}_t -symplectic connection on the formal symplectic manifold $(M, \theta_t, \mathcal{P}_t)$. Let $(\mathcal{C}_M^\infty[[t]], \mu_t, \mathcal{O}_t)$ be a PSP such that

$$\tau(\mathcal{C}_M^\infty[[t]], \mu_t, \mathcal{O}_t) = \left(\theta_t - \frac{t}{2} \text{tr}(\nabla^2|_{\mathcal{P}_t}), \mathcal{P}_t \right).$$

By (10.1), $\text{cl}_F(\mathcal{C}_M^\infty[[t]], \mu_t) = [\theta_t] = \text{cl}_F(\mathbb{A}_t)$, therefore SPs \mathbb{A}_t and $(\mathcal{C}_M^\infty[[t]], \mu_t)$ are equivalent. \square

Remark 10.5. All constructions of the paper can be extended to the case when M is a formal manifold, M_λ , which is M endowed with the function sheaf $\mathcal{C}_M^\infty[[\lambda]]$, λ a formal parameter. A formal PSM is a triple, $(M_\lambda, \omega_\lambda, \mathcal{P}_\lambda)$, where ω_λ is a formal symplectic form on M_λ and \mathcal{P}_λ its polarization. The above construction of a PSP applied to a formal PSM $(M_\lambda, \omega_\lambda, \mathcal{P}_\lambda)$ gives the following proposition.

Proposition 10.6. *Let $(\mathbb{A}_t, \mathbb{O}_t)$ be a PDQ and $(\omega_t, \mathcal{P}_t)$ an element on the orbit corresponding to $(\mathbb{A}_t, \mathbb{O}_t)$. Suppose that the pair $(\omega_\lambda, \mathcal{P}_\lambda)$ obtains from the pair $(\omega_t, \mathcal{P}_t)$ by replacing t with λ . Then, there exists on $(M_\lambda, \omega_\lambda, \mathcal{P}_\lambda)$ a PSP*

$$(\mathcal{C}_M^\infty[[\lambda]][[t]], \mu_{\lambda,t}, \mathcal{O}_{\lambda,t})$$

such that $(\mathbb{A}_t, \mathbb{O}_t)$ is equivalent to the diagonal PSP $(\mathcal{C}_M^\infty[[t]], \mu_{t,t}, \mathcal{O}_{t,t})$ obtained by the substitution $\lambda = t$.

Acknowledgements

I thank J. Bernstein, P. Bressler, B. Fedosov, A. Karabegov, and A. Mudrov for helpful discussions. I am also grateful to the Max Planck Institute for Mathematics for hospitality and very stimulating working atmosphere. Joseph Donin was partially supported by Israel Academy of Sciences Grant no. 8007/99-01.

References

- [1] A. Astashkevich, On Karabegov's Quantization of Semisimple Coadjoint Orbits: Advances in Geometry, Progr. Math., vol. 172, Birkhäuser, Boston, MA, 1999, 1–18.
- [2] F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowicz, D. Sternheimer, Deformation theory and quantization, Ann. Phys. 111 (1978) 61–110.
- [3] A. Beilinson, J. Bernstein, A proof of Jantzen conjectures. I. M. Gelfand Seminar, 1–50, Adv. Soviet Math., 16, Part 1, American Mathematical Society, Providence, RI, 1993.
- [4] A. Beilinson, D. Kazhdan, Geometric quantization, preprint.
- [5] P. Bressler, J. Donin, Polarized deformation quantization, math.QA/0007186.
- [6] M. Bordemann, N. Neumaier, M.J. Pflaum, S. Waldmann, On representations of star product algebras over cotangent spaces on Hermitian line bundles, math.QA/9811055.
- [7] M. Bordemann, S. Waldmann, A Fedosov star product of the Wick type for Kähler manifolds, Lett. Math. Phys. 41 (3) (1997) 243–253.
- [8] M. Bordemann, N. Neumaier, S. Waldmann, Homogeneous Fedosov star products on cotangent bundles. I. Weyl and standard ordering with differential operator representation, Commun. Math. Phys. 198 (2) (1998) 363–396.

- [9] P. Deligne, Déformations de l'Algébre des Fonctions d'une Variété Symplectique: Comparaison entre Fedosov et De Wilde, Lecomte, Selecta Mathematica 1 (4) (1995) 667–697 (New Series).
- [10] J. Donin, On the quantization of Poisson brackets, Adv. Math. 127 (1) (1997) 73–93.
- [11] B. Fedosov, Deformation Quantization and Index Theory, Academie Verlag, vol. 9.
- [12] A. Karabegov, Deformation quantization with separation of variables on Kähler manifold, Commun. Math. Phys. 180 (1996) 745–755.
- [13] A. Karabegov, Cohomological classification of deformation quantizations with separation of variables, Lett. Math. Phys. 43 (1998) 347–357.
- [14] A. Karabegov, M. Schlichenmaier, Almost Kähler Deformation Quantization, math.QA/0102169.
- [15] S. Kobayashi, K. Nomizu, Foundations of Differential Geometry, vol. II, Interscience, New York, 1969.
- [16] M. Kontsevich, Deformation quantization of Poisson manifolds, I, q-alg/9709040.
- [17] A. Lichnerowicz, Existence and equivalence of twisted products on a symplectic manifold, Lett. Math. Phys. 3 6 (1979) 495–502.
- [18] R. Nest, B. Tsygan, Algebraic index theorem for families, Adv. Math. 113 (1995) 151–205.
- [19] N. Neumaier, Universality of Fedosov's Construction for Star Products of Wick Type on Semi-Käler Manifolds, math.QA/0204031.
- [20] M.J. Pflaum, Deformation quantization on cotangent bundles, Coherent states, differential and quantum geometry (Białowieża, 1997), Rep. Math. Phys. 43 (1–2) (1999) 291–297.
- [21] J.H. Rawnsley, Polarization and diagonal quantization, Trans. AMS 230 (1977) 235–255.
- [22] N. Reshetikhin, M. Yakimov, Deformation quantization of Lagrangean fiber bundles, preprint, math/9907164.
- [23] P. Xu, Fedosov's \star -products and quantum momentum maps, Commun. Math. Phys. 197 (1998) 167–197.