



ELSEVIER

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Journal of Geometry and Physics 56 (2006) 1235–1256

JOURNAL OF
GEOMETRY AND
PHYSICS

www.elsevier.com/locate/jgp

Surfaces in the four-space and the Davey–Stewartson equations

Iskander A. Taimanov*

Institute of Mathematics, pr. Koptyuga, 4, 630090 Novosibirsk, Russia

Received 6 February 2004; received in revised form 25 June 2005; accepted 28 June 2005

Available online 3 August 2005

Abstract

We show that any equation from the Davey–Stewartson hierarchy induces an infinite family of geometrically different deformations of tori in \mathbb{R}^4 preserving the Willmore functional. We expose a derivation of the Weierstrass representation for surfaces in the four-space, which is not unique in difference from the case of surfaces in the three-space. This non-uniqueness implies that the spectral curve of a torus in \mathbb{R}^4 is not uniquely defined as a complex curve formed by the Floquet multipliers. © 2005 Elsevier B.V. All rights reserved.

MSC: 53C42; 35Q99; 53A05; 58F07

JGP SC: Differential geometry; Classical integrable systems; Integrable PDEs; Global analysis; Analysis on manifolds; Real differential geometry

Keywords: Surfaces in the four-space; The Davey–Stewartson equations; The spectral curve

1. Introduction

The Weierstrass representation for surfaces in \mathbb{R}^3 [8,13] was generalized for surfaces in \mathbb{R}^4 in [12] (see also [4]). This paper uses the quaternion language and the explicit formulas for such a representation were written by Konopelchenko in [9] for constructing surfaces which admit soliton deformation governed by the Davey–Stewartson equations.

* Tel.: +7 3833333496; fax: +7 3833332895.

E-mail address: taimanov@math.nsc.ru.

This generalizes his results from [8] where he introduced the formulas for inducing surfaces in the three-space which involve a Dirac type equations and defined for such surfaces a deformation governed by the modified Novikov–Veselov (mNV) equations.

It was shown in [13] that the formulas for inducing surfaces in \mathbb{R}^3 [8] describe all surfaces and that the modified Novikov–Veselov equation deforms tori into tori preserving the Willmore functional which naturally arises and plays an important role in this representation. The spectral curve of the corresponding Dirac operator is invariant under this deformation. The global Weierstrass representation at least for real analytic surfaces could be obtained by an analytic continuation from a local representation. Thus, a moduli space of immersed tori is embedded into the phase space of an integrable system with the Willmore functional and, moreover, the spectral curve as conservation quantities.

Looking forward to understand the spectral curves for tori in \mathbb{R}^4 we consider in this paper the analogous problems for surfaces in \mathbb{R}^4 and show that this case is very different from the three-dimensional case, in particular, by the following features which were overlooked until recently:

- for tori in \mathbb{R}^4 every equation from the Davey–Stewartson (DS) hierarchy describes not one but infinitely many geometrically different soliton deformations;
- the multipliers on the spectral curve for a torus in \mathbb{R}^4 are not uniquely defined and different complex curves in \mathbb{C}^2 (the spectral curves immersed via the multipliers) are invariants of different DS deformations.

The reason for that is quite clear and consists basically in the non-uniqueness of a Weierstrass representation.

A surface in \mathbb{R}^3 is constructed in terms of one vector function ψ (spinor) which is a lift of the Gauss mapping into non-vanishing spinors. Such a lift is defined up to a sign by fixing a conformal parameter on the surface. This function ψ satisfies a Dirac equation.

A surface in \mathbb{R}^4 is constructed in terms of two vector functions ψ and φ which form again a lift of the Gauss mapping. However, in this case by fixing a conformal parameter one defines a lift only up to a gauge transformation given by e^f where f is any smooth function. Moreover not every lift satisfies the Dirac equations and the lifts meeting these equations are defined up to gauge transformations e^h where h is a any holomorphic function.

In particular, given a Weierstrass representation of a surface $\Sigma \subset \mathbb{R}^4$ and a domain $W \subset \Sigma$ we can replace a representation of the domain by gauge-equivalent using a transformation e^h where h is a holomorphic function on W which is not analytically continued onto the surface. Thus, we obtain a representation of a domain, which is not continued (i.e. expanded to a representation of a surface). This also makes a difference with the case of surfaces in \mathbb{R}^3 .

Another important point is that the DS equations contain the additional potentials, which are defined by resolving the constraint equations. Such resolutions are not unique and we have to choose the potentials carefully to make the DS deformations geometric: for some special choices of the additional potentials the corresponding DS deformations map tori into tori preserving the Willmore functional. However, in general this is not the case and we show how to achieve that in Section 4.

2. Explicit formulas for a representation and soliton deformations

Let us recall the explicit formulas for inducing a surface and its soliton deformation via the Davey–Stewartson equation.

The following proposition is derived by straightforward computations.

Proposition 1 ([9]). *Let vector functions ψ and φ be defined in a simply connected domain $W \subset \mathbb{C}$ (with a complex parameter z) and meet the Dirac equations*

$$D\psi = 0, \quad D^\vee\varphi = 0,$$

where

$$D = \begin{pmatrix} 0 & \partial \\ -\bar{\partial} & 0 \end{pmatrix} + \begin{pmatrix} U & 0 \\ 0 & \bar{U} \end{pmatrix}, \quad D^\vee = \begin{pmatrix} 0 & \partial \\ -\bar{\partial} & 0 \end{pmatrix} + \begin{pmatrix} \bar{U} & 0 \\ 0 & U \end{pmatrix}.$$

Then the 1-forms

$$\eta_k = f_k dz + \bar{f}_k d\bar{z}, \quad k = 1, 2, 3, 4,$$

with

$$\begin{aligned} f_1 &= \frac{i}{2}(\bar{\varphi}_2\bar{\psi}_2 + \varphi_1\psi_1), & f_2 &= \frac{1}{2}(\bar{\varphi}_2\bar{\psi}_2 - \varphi_1\psi_1), & f_3 &= \frac{1}{2}(\bar{\varphi}_2\psi_1 + \varphi_1\bar{\psi}_2), \\ f_4 &= \frac{i}{2}(\bar{\varphi}_2\psi_1 - \varphi_1\bar{\psi}_2) \end{aligned} \tag{1}$$

are closed and the formulas

$$x^k = x^k(0) + \int \eta_k, \quad k = 1, 2, 3, 4, \tag{2}$$

define a surface in \mathbb{R}^4 (here the integral is taken over any path in W and by the Stokes theorem does not depend on a choice of path).

The induced metric equals

$$e^{2\alpha} dz d\bar{z} = (|\psi_1|^2 + |\psi_2|^2)(|\varphi_1|^2 + |\varphi_2|^2) dz d\bar{z} \tag{3}$$

and the norm of the mean curvature vector $\mathbf{H} = \frac{2x_{z\bar{z}}}{e^{2\alpha}}$ meets the equality

$$|U| = \frac{|\mathbf{H}|e^\alpha}{2}. \tag{4}$$

For $U = \bar{U}$ and $\psi = \pm\varphi$ these formulas reduce to the Weierstrass representation for surfaces in \mathbb{R}^3 .

The existence of a local representation of any surface in \mathbb{R}^4 by these formulas is not proved in [9] although it was indicated in [12] that the Weierstrass representation for surfaces in

\mathbb{R}^3 is generalized for surfaces in \mathbb{R}^4 and involves in this case two vector functions ψ and φ and a complex valued potential U .

We expose such a derivation in the next section revealing some features not taking place in the three-dimensional case. Remark that for Lagrangean surfaces in \mathbb{R}^4 this representation was discovered in other terms by Hélein and Romon [7].

Let

$$L = \begin{pmatrix} 0 & \partial \\ -\bar{\partial} & 0 \end{pmatrix} + \begin{pmatrix} -p & 0 \\ 0 & q \end{pmatrix}.$$

Let us consider deformations of this operator which take the form of Manakov's L , A , B -triple:

$$L_t + [L, A_n] - B_n L = 0 \quad (5)$$

or

$$[L, \partial_t - A_n] + B_n L = 0.$$

Notice that if L meets (5), then the solution of the equation

$$L\psi = 0$$

is evolved as follows:

$$\psi_t = A_n \psi.$$

The following two propositions are proved by straightforward computations.

Proposition 2. For

$$A_2 = \begin{pmatrix} -\partial^2 - v_1 & q\bar{\partial} - q_{\bar{z}} \\ -p\partial + p_z & \bar{\partial}^2 + v_2 \end{pmatrix},$$

$$B_2 = \begin{pmatrix} \partial^2 + \bar{\partial}^2 + (v_1 + v_2) & -(p + q)\bar{\partial} + q_{\bar{z}} - 2p_{\bar{z}} \\ (p + q)\partial - p_z + 2q_z & -(\partial^2 + \bar{\partial}^2) - (v_1 + v_2) \end{pmatrix},$$

where

$$v_{1\bar{z}} = -2(pq)_z, \quad v_{2z} = -2(pq)_{\bar{z}},$$

Eq. (5) takes the form

$$p_t = p_{zz} + p_{\bar{z}\bar{z}} + (v_1 + v_2)p, \quad q_t = -q_{zz} - q_{\bar{z}\bar{z}} - (v_1 + v_2)q. \quad (6)$$

Proposition 3. For

$$A_3 = \begin{pmatrix} \partial^3 + \frac{3}{2}v_1\partial - 3w_1 & q\bar{\partial}^2 - q_z\bar{\partial} + q_{z\bar{z}} + \frac{3}{2}v_2q \\ p\partial^2 - p_z\partial + p_{zz} + \frac{3}{2}v_1p & \bar{\partial}^3 + \frac{3}{2}v_2\bar{\partial} - 3w_2 \end{pmatrix},$$

$$B_3 = \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix},$$

where

$$b_{11} = -b_{22} = \bar{\partial}^3 - \partial^3 - \frac{3}{2}(v_1\partial - v_2\bar{\partial}) + 3(w_1 - w_2),$$

$$b_{12} = -(p + q)\bar{\partial}^2 - \frac{3}{2}(p + q)v_2 - (3p_z - q_z)\bar{\partial} - (3p_{z\bar{z}} + q_{z\bar{z}}),$$

$$b_{21} = -(p + q)\partial^2 - \frac{3}{2}(p + q)v_1 - (3q_z - p_z)\partial - (3q_{zz} + p_{zz})$$

and

$$v_{1\bar{z}} = -2(pq)_z, \quad v_{2z} = -2(pq)_{\bar{z}}, \quad w_{1\bar{z}} = (pq_z)_z, \quad w_{2z} = (qp_{\bar{z}})_{\bar{z}},$$

Eq. (5) takes the form ¹

$$p_t = p_{zzz} + p_{z\bar{z}\bar{z}} + \frac{3}{2}(v_1p_z + v_2p_{\bar{z}}) - 3(\partial^{-1}[(qp_{\bar{z}})_{\bar{z}}] + \bar{\partial}^{-1}[(qp_z)_z])p,$$

$$q_t = q_{zzz} + q_{z\bar{z}\bar{z}} + \frac{3}{2}(v_1q_z + v_2q_{\bar{z}}) - 3(\partial^{-1}[(pq_{\bar{z}})_{\bar{z}}] + \bar{\partial}^{-1}[(pq_z)_z])q. \tag{7}$$

Eqs. (6) and (7) are called the Davey–Stewartson equations. In fact these are the equations DSII₂ and DSII₃ from the DSII hierarchy. The equation DSII_n takes the form (5) where A_n equals

$$A_n = \begin{pmatrix} (-1)^{n+1}\partial^n & 0 \\ 0 & \bar{\partial}^n \end{pmatrix} + \dots$$

here by ... we denote terms of lower order.

¹ Eq. (5) after a formal substitution of A and B reduces to the system

$$p_t = p_{zzz} + p_{z\bar{z}\bar{z}} + \frac{3}{2}(v_1p_z + v_2p_{\bar{z}}) + 3\left(w_1 - w_2 + \frac{1}{2}v_{1z}\right)p,$$

$$q_t = q_{zzz} + q_{z\bar{z}\bar{z}} + \frac{3}{2}(v_1q_z + v_2q_{\bar{z}}) - 3\left(w_1 - w_2 - \frac{1}{2}v_{2\bar{z}}\right)q,$$

For $n = 1$ we have

$$A_1 = \begin{pmatrix} \partial & q \\ p & \bar{\partial} \end{pmatrix}, \quad B_1 = \begin{pmatrix} \bar{\partial} - \partial & -(p + q) \\ -(p + q) & \partial - \bar{\partial} \end{pmatrix},$$

and the DSII₁ equations are

$$p_t = p_z + p_{\bar{z}}, \quad q_t = q_z + q_{\bar{z}}.$$

Remark that the DSI hierarchy is a hierarchy of nonlinear equations obtained from the DSII hierarchy by replacing the variables z, \bar{z} by real-valued variables x, y .

Let us consider the reduction of the DSII hierarchy for the case

$$p = -u, \quad q = \bar{u}. \tag{8}$$

Eq. (6) is not compatible however the substitution

$$A_2 \rightarrow iA_2, \quad B_2 \rightarrow iB_2$$

into (5) gives a reduction of (6) compatible with (8):

$$u_t = i(u_{zz} + u_{\bar{z}\bar{z}} + 2(v + \bar{v})u), \quad v_{\bar{z}} = (|u|^2)_z. \tag{9}$$

The substitution of (8) into (7) gives

$$\begin{aligned} u_t &= u_{zzz} + u_{\bar{z}\bar{z}\bar{z}} + 3(vu_z + \bar{v}u_{\bar{z}}) + 3(w + w')u, & v_{\bar{z}} &= (|u|^2)_z, \\ w_{\bar{z}} &= (\bar{u}u_z)_z, & w'_z &= (\bar{u}u_{\bar{z}})_{\bar{z}}. \end{aligned} \tag{10}$$

For brevity we shall call Eqs. (9) and (10) by the DS₂ and DS₃ equations, respectively.

In difference with (9) the DS₃ equation is compatible with the constraint $u = \bar{u}$ and for real-valued potentials it reduces to the modified Novikov–Veselov equation:

$$u_t = u_{zzz} + u_{\bar{z}\bar{z}\bar{z}} + 3(vu_z + \bar{v}u_{\bar{z}}) + \frac{3}{2}(v_z + \bar{v}_{\bar{z}})u, \quad v_{\bar{z}} = (u^2)_z. \tag{11}$$

Notice that A_n depends on two functional parameters which are p and q and put

$$A_n^+ = A, \quad \text{for } p = -u, \ q = \bar{u}, \quad A_n^- = A, \quad \text{for } p = -\bar{u}, \ q = u.$$

Now let us recall the definition of the DS deformations of a surface introduced in [9].

Proposition 4 ([9]). *Let a surface Σ be defined by the formulas (1) and (2) for some ψ^0, φ^0 and let $U(z, \bar{z}, t)$ be a deformation of the potential described by Eq. (9) or (10). Then the formulas (1) and (2) and the equations*

$$\psi_t = iA_2^+ \psi, \quad \varphi_t = -iA_2^- \varphi, \quad \psi_t = A_3^+ \psi, \quad \varphi_t = A_3^- \varphi \tag{12}$$

with $\psi_{t=0} = \psi^0, \varphi_{t=0} = \varphi^0$, define deformations of the surface governed by Eqs. (9) and (10), respectively.

The proof of this proposition is as follows. Since the deformation of $U = u$ is described by Eq. (5), the vector functions ψ and φ meeting $\mathcal{D}\psi = 0$ and $\mathcal{D}^\vee\varphi = 0$ are deformed via Eq. (12) and for any t they meet again the Dirac equations. Therefore, by Proposition 1 they define a surface Σ_t via the Weierstrass formulas (1) and (2). Thus, we have a deformation Σ_t such that $\Sigma_0 = \Sigma$.

Any equation of the DSII hierarchy defines such a deformation (for n even we have to substitute $A_n \rightarrow iA_n$ to preserve the reduction $p = -\bar{q}$). We write down only two equations, DS_2 and DS_3 , because they resemble the main properties of others and do not involve very large expressions.

For $u = \bar{u}$ such a deformation reduces to the mNV deformation defined in [8] and studied in [13,14,5,10,2].

3. The Weierstrass representation

An oriented two-plane in \mathbb{R}^4 is defined by a positively oriented orthonormal basis

$$e_1 = (e_{1,1}, \dots, e_{1,4}), \quad e_2 = (e_{2,1}, \dots, e_{2,4})$$

which is defined up to rotations. There is a one-to-one correspondence

$$\{(e_1, e_2)\} \leftrightarrow (y_1 : y_2 : y_3 : y_4), \quad y_k = e_{1,k} + ie_{2,k}, \quad k = 1, 2, 3, 4,$$

between the moduli space of such planes (which is the Grassmannian $\tilde{G}_{4,2}$) and points of the quadric $Q \subset \mathbb{C}P^3$ defined by the equations

$$y_1^2 + y_2^2 + y_3^2 + y_4^2 = 0.$$

In terms of another homogeneous coordinates y'_1, \dots, y'_4 such that

$$y_1 = \frac{i}{2}(y'_1 + y'_2), \quad y_2 = \frac{1}{2}(y'_1 - y'_2), \quad y_3 = \frac{1}{2}(y'_3 + y'_4),$$

$$y_4 = \frac{i}{2}(y'_3 - y'_4)$$

this quadric is written as

$$y'_1 y'_2 = y'_3 y'_4.$$

Therefore, the correspondence

$$y'_1 = a_2 b_2, \quad y'_2 = a_1 b_1, \quad y'_3 = a_2 b_1, \quad y'_4 = a_1 b_2$$

establishes a biholomorphic equivalence

$$\tilde{G}_{4,2} = \mathbb{C}P^1 \times \mathbb{C}P^1,$$

where $(a_1 : a_2)$ and $(b_1 : b_2)$ are homogeneous coordinates on the copies of $\mathbb{C}P^1$. This mapping $\mathbb{C}P^1 \times \mathbb{C}P^1 \rightarrow Q \subset \mathbb{C}P^3$ is called the Segré mapping.

Let $r : W \rightarrow \mathbb{R}^4$ be an immersion of a surface with a conformal parameter $z \in W \subset \mathbb{C}$. The conformality condition reads

$$\langle r_z, r_z \rangle = \sum_{k=1}^4 (x_z^k)^2 = 0,$$

where $x_z^k = \frac{\partial x^k}{\partial z}$, $k = 1, 2, 3, 4$. The Gauss map takes the form

$$G : W \rightarrow \tilde{G}_{4,2}, \quad Q \in W \rightarrow (x_z^1(Q) : \dots : x_z^4(Q)).$$

By using the equivalence $\tilde{G}_{4,2} = \mathbb{C}P^1 \times \mathbb{C}P^1$, decompose G into two maps

$$G = (G_\psi, G_\varphi),$$

where

$$G_\psi = (\psi_1 : \bar{\psi}_2) \in \mathbb{C}P^1, \quad G_\varphi = (\varphi_1 : \bar{\varphi}_2) \in \mathbb{C}P^1$$

and rewrite the formulas for x_z^k in terms of these maps as follows:

$$\begin{aligned} x_z^1 &= \frac{i}{2}(\bar{\varphi}_2 \bar{\psi}_2 + \varphi_1 \psi_1), & x_z^2 &= \frac{1}{2}(\bar{\varphi}_2 \bar{\psi}_2 - \varphi_1 \psi_1), & x_z^3 &= \frac{1}{2}(\bar{\varphi}_2 \psi_1 + \varphi_1 \bar{\psi}_2), \\ x_z^4 &= \frac{i}{2}(\bar{\varphi}_2 \psi_1 - \varphi_1 \bar{\psi}_2). \end{aligned} \tag{13}$$

We have

$$dx^k = \eta_k, \quad k = 1, 2, 3, 4,$$

where the forms η_k take the same shapes as in Proposition 1.

This decomposition is not unique and functions ψ and φ are defined up to gauge transformations

$$\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \rightarrow \begin{pmatrix} e^f \psi_1 \\ e^{\bar{f}} \psi_2 \end{pmatrix}, \quad \begin{pmatrix} \varphi_1 \\ \varphi_2 \end{pmatrix} \rightarrow \begin{pmatrix} e^{-f} \varphi_1 \\ e^{-\bar{f}} \varphi_2 \end{pmatrix}. \tag{14}$$

By choosing a representative ψ for G_ψ we fix a function φ .

The formula (1) gives exactly a gauge transformation between different lifts of the Gauss mapping $G = (G_\psi, G_\varphi)$ to non-vanishing spinors, i.e. to $(\mathbb{C}^2 \setminus \{0\}) \times (\mathbb{C}^2 \setminus \{0\})$, which we mention in the introduction.

The closedness conditions for the forms η_k are

$$(\bar{\varphi}_2 \psi_1)_{\bar{z}} = (\bar{\varphi}_1 \psi_2)_z, \quad (\bar{\varphi}_2 \bar{\psi}_2)_{\bar{z}} = -(\bar{\varphi}_1 \bar{\psi}_1)_z \tag{15}$$

and they do not have the form of Dirac equations $\mathcal{D}\psi = 0$ and $\mathcal{D}^\vee\varphi = 0$ for arbitrary representatives (ψ_1, ψ_2) and (φ_1, φ_2) of the mappings G_ψ and G_φ . Such representatives have to be found by solving some differential equations.

Let us start with the following lift for $G_\psi = (\psi_1 : \bar{\psi}_2)$ which is correctly defined up to a ± 1 multiple:

$$s_1 = e^{i\theta} \cos \eta, \quad s_2 = \sin \eta.$$

We look for a pair of functions ψ_1, ψ_2 meeting two conditions:

- (1) $G_\psi = (\psi_1 : \bar{\psi}_2) = (s_1 : \bar{s}_2)$;
- (2) $\mathcal{D}\psi = 0$ for some potential U .

By the first condition, ψ has the form

$$\psi_1 = e^g s_1, \quad \psi_2 = e^{\bar{g}} s_2.$$

The second condition is written as

$$\partial(e^{\bar{g}} \sin \eta) + U e^{g+i\theta} \cos \eta = 0, \quad \bar{\partial}(e^{g+i\theta} \cos \eta) = \bar{U} e^{\bar{g}} \sin \eta.$$

These equations are rewritten as follows:

$$U = -\frac{e^{\bar{g}}}{e^{g+i\theta} \cos \eta} (\bar{g}_z \sin \eta + \eta_z \cos \eta),$$

$$U = \frac{e^{\bar{g}}}{e^{g+i\theta} \sin \eta} (\bar{g}_z \cos \eta - i\theta_z \cos \eta - \eta_z \sin \eta),$$

which imply

$$g_{\bar{z}} + i\theta_{\bar{z}} \cos^2 \eta = 0, \quad U = -e^{\bar{g}-g-i\theta} (i\theta_z \sin \eta \cos \eta + \eta_z).$$

The $\bar{\partial}$ -problem for g is solved by the well-known means and its solution is defined up to holomorphic functions. Therefore, the potential U is defined up to a multiplication by $e^{\bar{h}-h}$ where h is an arbitrary holomorphic function.

It is derived by straightforward computations that $\mathcal{D}\psi = 0$ implies that the condition (15) takes the form of the Dirac equation $\mathcal{D}^\vee\varphi = 0$.

Thus, the following theorem is derived.

Theorem 1. *Let $r : W \rightarrow \mathbb{R}^4$ be an immersed surface with a conformal parameter z and let $G_\psi = (e^{i\theta} \cos \eta : \sin \eta)$ be one of the components of its Gauss map.*

There exists another representative ψ of this mapping $G_\psi = (\psi_1 : \bar{\psi}_2)$ such that it meets the Dirac equation

$$\mathcal{D}\psi = 0$$

with some potential U .

A vector function $\psi = (e^{g+i\theta} \cos \eta, e^{\bar{g}} \sin \eta)$ is defined from the equation

$$g_{\bar{z}} = -i\theta_{\bar{z}} \cos^2 \eta, \quad (16)$$

up to holomorphic functions and the corresponding potential U is defined up by the formula

$$U = -e^{\bar{g}-g-i\theta}(i\theta_z \sin \eta \cos \eta + \eta_z)$$

up to multiplications by $e^{\bar{h}-h}$ where h is an arbitrary holomorphic function.

Given the function ψ , a function φ which represents another component G_φ of the Gauss map meets the equation

$$\mathcal{D}^\vee \varphi = 0.$$

Different representations (lifts) of the Gauss mapping G of the surface W are related by gauge transformations of the form

$$\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \rightarrow \psi' = \begin{pmatrix} e^h \psi_1 \\ e^{\bar{h}} \psi_2 \end{pmatrix}, \quad \begin{pmatrix} \varphi_1 \\ \varphi_2 \end{pmatrix} \rightarrow \varphi' = \begin{pmatrix} e^{-h} \varphi_1 \\ e^{-\bar{h}} \varphi_2 \end{pmatrix},$$

$$U \rightarrow U' = e^{\bar{h}-h} U, \quad (17)$$

where h is an arbitrary holomorphic function on W .

Since we decompose the Gauss map into two components G_ψ and G_φ with functions ψ and φ meeting the Dirac equations, we obtain the surface by integrating the differentials dx^k given by (13). The formulas (3) and (4) for the metric and the potential are obtained by simple computations. We have

Corollary 1. Every oriented surface in \mathbb{R}^4 admits a Weierstrass representation given by Proposition 1.

However, the non-uniqueness of such a representation leads to the following conclusion.

Corollary 2. For any surface $r : W \rightarrow \mathbb{R}^4$ where W is an open subset of \mathbb{C} and any subdomain $V \subset W$ such that $V \neq W$ there is a Weierstrass representation of $r|_V : V \rightarrow \mathbb{R}^4$ which is not analytically continued (i.e. expanded) onto W .

Proof. For that take a Weierstrass representation of W , restrict it onto V and take a gauge equivalent representation of V corresponding to a transformation (17) where $h : V \rightarrow \mathbb{C}$ is a holomorphic function which is not analytically continued onto W . Then this representation of V is not expanded onto W . This proves the corollary. \square

The following theorem is clear.

Theorem 2. Given a Weierstrass representation of an immersed closed oriented surface Σ into \mathbb{R}^4 , the corresponding functions ψ and φ are sections of the \mathbb{C}^2 -bundles E and E^\vee

over Σ which are as follows:

(1) E and E^\vee split into sums of pair-wise conjugate line bundles

$$E = E_0 \oplus \bar{E}_0, \quad E^\vee = E_0^\vee \oplus \bar{E}_0^\vee$$

such that ψ_1 and $\bar{\psi}_2$ are sections of E_0 and φ_1 and $\bar{\varphi}_2$ are sections of E_0^\vee ;

(2) the pairing of sections of E_0 and E_0^\vee is a $(1, 0)$ form on Σ : if

$$\alpha \in \Gamma(E_0), \quad \beta \in \Gamma(E_0^\vee),$$

then

$$\alpha\beta \, dz$$

is a correctly defined 1-form on Σ ;

(3) the Dirac equation $\mathcal{D}\psi = 0$ implies that U is a section of the same line bundle E_U as

$$\frac{\partial\gamma}{\alpha} \in \Gamma(E_U), \quad \text{for } \alpha \in \Gamma(E_0), \gamma \in \Gamma(\bar{E}_0)$$

and the quantity $U\bar{U} \, dz \wedge d\bar{z}$ is a correctly defined $(1, 1)$ -form on Σ whose integral equals

$$\int_{\Sigma} U\bar{U} \, dz \wedge d\bar{z} = -\frac{i}{2}\mathcal{W}(\Sigma)$$

where $\mathcal{W}(\Sigma) = \int_{\Sigma} |\mathbf{H}|^2 \, d\mu$ is the Willmore functional of Σ .

Notice that for surfaces in \mathbb{R}^3 functions ψ are sections of spinor bundles [13] and the gauge transformation (17) shows that for surfaces \mathbb{R}^4 this is not necessarily a case.

It is derived from Proposition 1 that, given a Riemann surface Σ , such bundles E, E^\vee , and E_U and solutions ψ and φ to the equations $\mathcal{D}\psi = 0$ and $\mathcal{D}^\vee\varphi = 0$, one may construct an immersion of the universal covering of Σ into \mathbb{R}^4 . The Gauss mapping of this immersion descends through Σ . Let us give a criterion for converting such an immersion into an immersion of the surface Σ_0 .

Proposition 5. *Let Σ be an oriented closed surface, let $\tilde{\Sigma}$ be its universal covering and let (ψ, φ) define an immersion of $\tilde{\Sigma}$ into \mathbb{R}^4 via (1) and (2). Then such an immersion converts into an immersion of Σ if and only if*

$$\int_{\Sigma} \bar{\psi}_1\bar{\varphi}_1 \, d\bar{z} \wedge \omega = \int_{\Sigma} \bar{\psi}_1\varphi_2 \, d\bar{z} \wedge \omega = \int_{\Sigma} \psi_2\bar{\varphi}_1 \, dz \wedge \omega = \int_{\Sigma} \psi_2\varphi_2 \, dz \wedge \omega = 0 \tag{18}$$

for any holomorphic differential ω on Σ .

Proof. Let $g \geq 1$ be the genus of Σ . Then there is a basis $\alpha_1, \dots, \alpha_g, \beta_1, \dots, \beta_g$ of 1-cycles on Σ such that $\Sigma = \tilde{\Sigma}/\Gamma$ and the fundamental domain for the action of $\Gamma = \pi_1(\Sigma)$ is a domain Ω on $\tilde{\Sigma}$ whose boundary has the form

$$\partial\Omega = \alpha_1\beta_1\alpha_1^{-1}\beta_1^{-1}, \dots, \alpha_g\beta_g\alpha_g^{-1}\beta_g^{-1}.$$

Denote by dx^1, \dots, dx^4 the closed 1-forms on $\tilde{\Sigma}$ induced by the immersion into \mathbb{R}^4 and denote by V^1, \dots, V^4 the period vectors

$$V^k = \left(\int_{\alpha_1} dx^k, \dots, \int_{\alpha_g} dx^k, \int_{\beta_k} dx^k, \dots, \int_{\beta_g} dx^k \right), \quad k = 1, 2, 3, 4.$$

An immersion of $\tilde{\Sigma}$ converts into an immersion of Σ if and only if

$$V^1 = V^2 = V^3 = V^4 = 0.$$

Given a holomorphic form ω on Σ , pull it back onto the universal covering $\tilde{\Sigma}$ and compute the integral

$$\int_{\partial\Omega} x^k \omega = \sum_{j=1}^g \left(\int_{\alpha_j} \omega \int_{\beta_j} dx^k - \int_{\alpha_j} dx^k \int_{\beta_j} \omega \right) = \sum_{j=1}^g \left(V_{j+g}^k \int_{\alpha_j} \omega - V_j^k \int_{\beta_j} \omega \right)$$

which is by the Stokes theorem equals

$$\int_{\partial\Omega} x^k \omega = \int_{\Omega} x_{\bar{z}}^k d\bar{z} \wedge \omega.$$

A Riemann surface Σ has a basis $\omega_1, \dots, \omega_g$ for holomorphic differentials normalized by the condition

$$\int_{\alpha_j} \omega_k = \delta_{ij}.$$

In this event the β -periods matrix

$$B_{jk} = \int_{\beta_j} \omega_k$$

is symmetric with positive imaginary part: $\text{Im } B > 0$. This implies that the conditions

$$\sum_{j=1}^g \left(V_{j+g}^k \int_{\alpha_j} \omega_l - V_j^k \int_{\beta_j} \omega_l \right) = 0, \quad l = 1, \dots, g,$$

for a vector V^k with real entries are satisfied if and only if $V^k = 0$. Therefore, an immersion of $\tilde{\Sigma}$ converts into an immersion of a closed surface Σ if and only if

$$\int_{\Omega} x_z^k d\bar{z} \wedge \omega = 0$$

for any $k = 1, \dots, 4$ and any holomorphic differential ω on Σ . It follows from (1) that that is equivalent to the equalities (18). This proves the proposition. \square

4. Deformations of tori via the Davey–Stewartson equations

Let us look what Theorem 1 gives us for tori.

Theorem 3. *Let Σ be a torus in \mathbb{R}^4 which is conformally equivalent to \mathbb{C}/Λ and z is a conformal parameter.*

Then there are vector functions ψ, φ and a function U such that

- (1) ψ and φ give a Weierstrass representation of Σ ;
- (2) the potential U of this representation is Λ -periodic;
- (3) such functions ψ, φ , and U are defined up to gauge transformations

$$\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \rightarrow \begin{pmatrix} e^h \psi_1 \\ e^{\bar{h}} \psi_2 \end{pmatrix}, \quad \begin{pmatrix} \varphi_1 \\ \varphi_2 \end{pmatrix} \rightarrow \begin{pmatrix} e^{-h} \varphi_1 \\ e^{-\bar{h}} \varphi_2 \end{pmatrix}, \quad U \rightarrow U e^{\bar{h}-h} U, \tag{19}$$

where

$$h(z) = a + bz, \quad \text{Im}(b\gamma) \in \pi\mathbb{Z}, \quad \text{for all } \gamma \in \Lambda.$$

Therefore, such representations are parameterized by a \mathbb{Z}^2 lattice formed by admissible values of b .

Proof. By Theorem 1, given a Weierstrass representation of a torus (with a fixed conformal parameter) the Dirac equations are satisfied if and only if

$$U = -e^{\bar{g}-g-i\theta}(i\theta_z \sin \eta \cos \eta + \eta_z), \quad \psi_1 = e^{i\theta+g} \cos \eta, \quad \psi_2 = e^{\bar{g}} \sin \eta$$

where g meets Eq. (16). The component of the Gauss mapping $G_\psi = (e^{i\theta} \cos \eta : \sin \eta)$ is Λ -periodic and U takes the form $U = e^{\bar{g}-g} U_0$ where

$$U_0 = i\theta_z \sin \eta \cos \eta + \eta_z$$

is periodic with respect to Λ . By (16), $g_{\bar{z}}$ is periodic and a general solution to (16) takes the form

$$g = h(z) + c\bar{z} + f(z, \bar{z}),$$

where $h(z)$ is an arbitrary holomorphic function, f is some periodic function, and

$$c = - \int_M i\theta_{\bar{z}} \cos^2 \eta \, dx \, dy.$$

Hence, U is a Λ -periodic function if and only if $g = a + bz + c\bar{z}$ (modulo periodic functions) and $(\bar{g}(z) - g(z)) \in 2\pi\mathbb{Z}$ for $z \in \Lambda$. The latter conditions are easily resolved and b is defined up to b' such that $(b'\bar{\gamma} - b'\gamma) \in 2\pi\mathbb{Z}$ for all $\gamma \in \Lambda$. This proves the theorem. \square

Now let us look for the DS deformations of tori. Exactly in this case in difference with high genera compact surfaces the constraints for defining additional potentials v, a, b could be globally resolved. As in [13] we are interested in two problems:

- when a deformation from the DS hierarchy deforms a torus into tori?
- when the “Willmore functional” $\int_{\Sigma} u\bar{u} \, dz \wedge d\bar{z}$ is preserved by such a deformation.

For that we have to resolve the constraint equations for v, w , and w' carefully choosing specific solutions.

We consider only the DS_2 and DS_3 equations since we do not know until recently a general recursion procedure for writing down explicit formulas for higher equations. However, it is a general point in the soliton theory that the first nontrivial equations in the hierarchy usually resemble the main properties of higher equations.

Since we derive for tori that there is a representation with a double-periodic potential we can look for solutions of the DS equations with such an initial data. In addition we have to define such resolutions of constraints (i.e. the additional potentials v, a, b) to obtain the equations with double-periodic coefficients and, hence, double-periodic solutions.

We do not discuss the existence of a solution assuming that it exists which, in particular, for short times follows from the Cauchy–Kovalevskaya theorem.

4.1. The DS_2 deformation

We have

$$\psi_t = iA_2^+ \psi, \quad \varphi_t = -iA_2^- \varphi, \quad v_{\bar{z}} = (|u|^2)_z.$$

When we work with compact surfaces we have to resolve the constraint equation for v globally. Moreover, for tori we would like to save the periodicity of the integrands of the Weierstrass representation, i.e. terms which are of the form $\psi_2\varphi_2$ or similar to it, we have to have a periodic potential v .

This is quite easy: the constraint for v is uniquely resolved by an inversion of the $\bar{\partial}$ -operator on a torus assuming that $\int v dz \wedge d\bar{z} = 0$. The reason is the same as for the mNV deformation (see [13]): since the right hand-side $(|u|^2)_z$ of the constraint equation is a derivative of a periodic function its Fourier decomposition does not contain a non-zero term and the $\bar{\partial}$ -operator is inverted term by term of the Fourier decomposition. Thus, we have:

$$v = \bar{\partial}^{-1} \partial(|u|^2), \quad \int_{\Sigma} v dz \wedge d\bar{z} = 0. \tag{20}$$

Here and in the sequel we identify Σ with a fundamental domain for the lattice Λ such that the torus is conformally equivalent to \mathbb{C}/Λ .

Theorem 4. *The DS_2 equation*

$$u_t = i(u_{zz} + u_{\bar{z}\bar{z}} + 2(v + \bar{v})u)$$

where v is defined by (20) induces a deformation of tori (with a fixed periodic potential of their Weierstrass representations) into tori.

Proof. We have to prove that the closedness conditions (18) are preserved. We show that only for one of them because for others the proofs are basically the same. There is one dimensional family of holomorphic differential on a torus generated by dz . We have to prove that

$$J = \int_{\Sigma} \psi_2 \varphi_2 dz \wedge d\bar{z} = 0$$

implies that $J_t = 0$.

We have

$$\psi_{2t} = i[(u\partial - u_z)\psi_1 + (\bar{\partial}^2 + \bar{v})\psi_2], \quad \varphi_{2t} = -i[(\bar{u}\partial - \bar{u}_z)\varphi_1 + (\bar{\partial}^2 + \bar{v})\varphi_2].$$

Substituting that into

$$J_t = \int_{\Sigma} (\psi_{2t}\varphi_2 + \psi_2\varphi_{2t})dz \wedge d\bar{z}$$

(notice that the integrand is correctly defined as a function on Σ , i.e. it is double-periodic, although ψ_2 and φ_2 are not periodic) we obtain

$$J_t = i \int_{\Sigma} [(u\psi_{1z} - u_z\psi_1)\varphi_2 + (\psi_{2\bar{z}\bar{z}} + \bar{v}\psi_2)\varphi_2 - (\bar{u}\varphi_{1z} - \bar{u}_z\varphi_1)\psi_2 - (\varphi_{2\bar{z}\bar{z}} + \bar{v}\varphi_2)\psi_2]dz \wedge d\bar{z}.$$

Integrating by parts some of terms and canceling terms repeated with different signs we derive

$$J_t = i \int_{\Sigma} [(u\psi_{1z} - u_z\psi_1)\varphi_2 - (\bar{u}\varphi_{1z} - \bar{u}_z\varphi_1)\psi_2] dz \wedge d\bar{z}.$$

Now an integration by parts implies that

$$\begin{aligned} J_t &= i \int_{\Sigma} [u\psi_{1z}\varphi_2 + u(\psi_1\varphi_2)_z - \bar{u}\varphi_{1z}\psi_2 - \bar{u}(\psi_2\varphi_1)_z] dz \wedge d\bar{z} \\ &= \int_{\Sigma} [2(u\varphi_2)\psi_{1z} + (u\psi_1)\varphi_{2z} - 2(\bar{u}\psi_2)\varphi_{1z} - (\bar{u}\varphi_1)\psi_{2z}] dz \wedge d\bar{z}. \end{aligned}$$

By using the Dirac equations replace the terms in brackets to exclude the potentials u and \bar{u} from the integrand:

$$J_t = 2i \int_{\Sigma} (\varphi_{1\bar{z}}\psi_{1z} - \varphi_{1z}\psi_{1\bar{z}}) dz \wedge d\bar{z}.$$

An integration by parts implies $J_t = 0$ which proves the theorem. \square

Remark that we never use in the proof that $J = 0$ for the initial torus. Therefore, we proved more: the DS_2 deformation preserves the translational periods of surfaces with double-periodic Gauss mapping.

Since we proved that tori are preserved it is reasonable to speak about the conservation laws for the DS deformations. We have

Theorem 5. *The DS_2 deformation of tori preserves the Willmore functional.*

Proof. We have

$$\begin{aligned} \frac{d}{dt} \int_{\Sigma} |u|^2 dz \wedge d\bar{z} &= \int_{\Sigma} (u_t\bar{u} + u\bar{u}_t) dz \wedge d\bar{z} = i \int_{\Sigma} [\bar{u}(u_{zz} + u_{\bar{z}\bar{z}} + 2(v + \bar{v})u) \\ &\quad - u(\bar{u}_{zz} + \bar{u}_{\bar{z}\bar{z}} + 2(v + \bar{v})\bar{u})] dz \wedge d\bar{z} = i \int_{\Sigma} (\bar{u}u_{zz} - u\bar{u}_{zz} + \bar{u}u_{\bar{z}\bar{z}} - u\bar{u}_{\bar{z}\bar{z}}) dz \wedge d\bar{z}. \end{aligned}$$

By integrating by parts the last integral, we easily derive that

$$\frac{d}{dt} \int_{\Sigma} |u|^2 dz \wedge d\bar{z} = 0$$

which proves the theorem. \square

Remark 1. By Proposition 2, there are two potentials v_1 and v_2 which are coming into the equation. We choose an additional constraint $v = v_1 = \bar{v}_2$. Without this constraint Theorem 4 does not hold as one can see from its proof.

In fact we assumed more, that $\int v_{\Sigma} dz \wedge d\bar{z}$ vanishes. This was not important: if we put $v \rightarrow v + f(t)$ then the deformation reduces to a composition of the initial one and some evolution tangent to a torus, i.e. this results in adding a diffeomorphism of a torus which does not affect the geometric picture.

4.2. The DS_3 deformation

For this deformation we have

$$\psi_t = A_3^+ \psi, \quad \varphi_t = A_3^- \varphi$$

and many constraints which we have to resolve. As in the case of the DS_2 deformation we put

$$v_{\bar{z}} = (|u|^2)_z, \quad \int_{\Sigma} v dz \wedge d\bar{z} = 0 \tag{21}$$

and in addition we choose w and w' as follows:

$$w = \bar{\partial}^{-1}(\bar{u}u_z), \quad w' = \bar{\partial}^{-1}(\bar{u}u_{\bar{z}}). \tag{22}$$

These functions satisfy the constraint equations (10) however they are very specific particular solutions to them.

Moreover, we have to resolve the constraint equations for w_1 and w_2 which are different for A^+ and A^- . We put

$$\begin{aligned} w_1^+ &= w - v_z, & w_2^+ &= -w', & w_1^- &= -w, & w_2^- &= w' - \bar{v}_{\bar{z}}, \\ v_1^{\pm} &= 2v, & v_2^{\pm} &= 2\bar{v}. \end{aligned} \tag{23}$$

The reasonings for these choices we shall explain later.

Theorem 6. The DS_3 equation

$$u_t = u_{zzz} + u_{\bar{z}\bar{z}\bar{z}} + 3(vu_z + \bar{v}u_{\bar{z}}) + 3(w + w')u$$

induces a deformation of tori (with fixed periodic potential of their Weierstrass representations) into tori.

Proof. We again demonstrate that only for one of the closedness conditions. Let us take the same as in the proof of Theorem 4. We have

$$\begin{aligned} \psi_{2t} &= (-u\partial^2 + u_z\partial - u_{zz} - 3vu)\psi_1 + (\bar{\partial}^3 + 3\bar{v}\bar{\partial} + 3w')\psi_2, \\ \varphi_{2t} &= (-\bar{u}\partial^2 + \bar{u}_z\partial - \bar{u}_{zz} - 3v\bar{u})\varphi_1 + (\bar{\partial}^3 + 3\bar{v}\bar{\partial} - 3(w' - \bar{v}_z))\varphi_2. \end{aligned}$$

Substitute that into

$$J_t = \int_{\Sigma} (\psi_{2t}\varphi_2 + \psi_2\varphi_{2t}) dz \wedge d\bar{z}.$$

An integration by parts shows that

$$\int_{\Sigma} [(\psi_{2z\bar{z}\bar{z}\bar{z}} + 3\bar{v}\psi_{2\bar{z}})\varphi_2 + (\varphi_{2z\bar{z}\bar{z}\bar{z}} + 3\bar{v}\varphi_{2\bar{z}} + 3\bar{v}_z\varphi_2)\psi_2] dz \wedge d\bar{z} = 0$$

and we are left to prove that

$$\begin{aligned} \int_{\Sigma} [(-u\psi_{1zz} + u_z\psi_{1z} - (u_{zz} + 3vu)\psi_1)\varphi_2 \\ + (-\bar{u}\varphi_{1z\bar{z}} + \bar{u}_z\varphi_{1z} - (\bar{u}_{z\bar{z}} + 3v\bar{u})\varphi_1)\psi_2] dz \wedge d\bar{z} = 0. \end{aligned}$$

Let us rewrite the left-hand side of this formula as

$$\begin{aligned} J_t = \int_{\Sigma} [(-u\psi_1)_{z\bar{z}} + 3u_z\psi_{1z} - 3vu\psi_1)\varphi_2 + (-\bar{u}\varphi_1)_{z\bar{z}} + 3\bar{u}_z\varphi_{1z} \\ - 3v\bar{u}\varphi_1\psi_2] dz \wedge d\bar{z} \end{aligned}$$

which by the Dirac equations $\mathcal{D}\psi = \mathcal{D}^\vee\varphi = 0$, equals to

$$\begin{aligned} J_t &= \int_{\Sigma} [(\psi_{2z\bar{z}\bar{z}\bar{z}} + 3u_z\psi_{1z})\varphi_2 - 3v\psi_1\varphi_{1\bar{z}} + (\varphi_{2z\bar{z}\bar{z}\bar{z}} + 3\bar{u}_z\varphi_{1z})\psi_2 - 3v\psi_{1\bar{z}}\varphi_1] dz \wedge d\bar{z} \\ &= \int_{\Sigma} (\psi_{2z\bar{z}\bar{z}\bar{z}}\varphi_2 + \psi_2\varphi_{2z\bar{z}\bar{z}\bar{z}}) dz \wedge d\bar{z} + 3 \int_{\Sigma} (u_z\psi_{1z}\varphi_2 + \bar{u}_z\psi_2\varphi_{1z} - v(\psi_1\varphi_1)_{\bar{z}}) dz \wedge d\bar{z}. \end{aligned}$$

An integration by parts shows that the first summand vanishes and, by (21), we have

$$\begin{aligned} J_t &= 3 \int_{\Sigma} (u_z\psi_{1z}\varphi_2 + \bar{u}_z\psi_2\varphi_{1z} + (|u|^2)_z\psi_1\varphi_1) dz \wedge d\bar{z} \\ &= 3 \int_{\Sigma} (u_z\psi_{1z}\varphi_2 + \bar{u}_z\psi_2\varphi_{1z} + u_z\bar{u}\psi_1\varphi_1 + \bar{u}_z u\psi_1\varphi_1) dz \wedge d\bar{z} \\ &= 3 \int_{\Sigma} [u_z(\psi_{1z}\varphi_2 - \psi_1\varphi_{2z}) + \bar{u}_z(\varphi_{1z}\psi_2 - \varphi_1\psi_{2z})] dz \wedge d\bar{z} \\ &= 3 \int_{\Sigma} [u(\psi_1\varphi_{2z\bar{z}} - \psi_{1z\bar{z}}\varphi_2) + \bar{u}(\varphi_1\psi_{2z\bar{z}} - \varphi_{1z\bar{z}}\psi_2)] dz \wedge d\bar{z} \\ &= 3 \int_{\Sigma} (-\psi_{2z}\varphi_{2z\bar{z}} - \psi_{1z\bar{z}}\varphi_{1\bar{z}} - \varphi_{2z}\psi_{2z\bar{z}} - \psi_{1\bar{z}}\varphi_{1z\bar{z}}) dz \wedge d\bar{z}. \end{aligned}$$

Now by integrating by parts we easily derive from the last formula that

$$J_t = 0.$$

Analogous reasonings show that other closedness conditions (15) are also preserved by the flow. This proves theorem. \square

As in the case of the DS_2 flow we actually show that the DS_3 deformation preserves the translational periods of surfaces with double-periodic Gauss mapping.

As in the case of the DS_2 deformation the Willmore functional is preserved, i.e. the following theorem holds.

Theorem 7. *The DS_3 deformation with the additional potentials of the form (21–23) preserves the Willmore functional.*

Proof. We have

$$\begin{aligned} \frac{d}{dt} \int_{\Sigma} |u|^2 dz \wedge d\bar{z} &= \int_{\Sigma} (u_t \bar{u} + u \bar{u}_t) dz \wedge d\bar{z} = \int_{\Sigma} [(u_{zzz} + u_{\bar{z}\bar{z}\bar{z}} + 3(vu_z + \bar{v}u_{\bar{z}}) \\ &\quad + 3(w + w')u) \bar{u} + u(\bar{u}_{zzz} + \bar{u}_{\bar{z}\bar{z}\bar{z}} + 3(\bar{v}\bar{u}_{\bar{z}} + \bar{v}\bar{u}_z) \\ &\quad + 3(\bar{w} + \bar{w}')\bar{u})] dz \wedge d\bar{z} = \int_{\Sigma} [(u_{zzz}\bar{u} + u\bar{u}_{zzz}) + (u_{\bar{z}\bar{z}\bar{z}}\bar{u} + u\bar{u}_{\bar{z}\bar{z}\bar{z}})] dz \wedge d\bar{z} \\ &\quad + 3 \int_{\Sigma} (v(|u|^2)_z + \bar{v}(|u|^2)_{\bar{z}}) dz \wedge d\bar{z} + 3 \int_{\Sigma} (w + \bar{w} + w' + \bar{w}')|u|^2 dz \wedge d\bar{z}. \end{aligned}$$

An integration by parts shows that the first integral vanishes, and, by (21), the second integral equals to

$$3 \int_{\Sigma} (vv_{\bar{z}} + \bar{v}\bar{v}_z) dz \wedge d\bar{z} = 0$$

(here we use that the function v is double-periodic). We are left to prove that

$$\int_{\Sigma} (w + \bar{w} + w' + \bar{w}')|u|^2 dz \wedge d\bar{z} = 0.$$

We have

$$w + \bar{w}' = v_z, \quad w' + \bar{w} = \bar{v}_{\bar{z}}.$$

Therefore, the investigated integral is rewritten as

$$\begin{aligned} \int_{\Sigma} (v_z + \bar{v}_{\bar{z}})|u|^2 dz \wedge d\bar{z} &= - \int_{\Sigma} (v(|u|^2)_z + \bar{v}(|u|^2)_{\bar{z}}) dz \wedge d\bar{z} \\ &= - \int_{\Sigma} (vv_{\bar{z}} + \bar{v}\bar{v}_z) dz \wedge d\bar{z} = 0. \end{aligned}$$

This proves the theorem. \square

Remark 2. As for the DS_2 deformation we choose the additional potentials carefully to make the DS_3 deformation geometric. For general potentials meeting the constraint equations it is not the case. For instance, we may add any constants c and c' to w and w' but the proof of [Theorem 7](#) shows that if $k = c + c' + \bar{c} + \bar{c}' \neq 0$ then the Willmore functional is evolved as

$$\mathcal{W}_t = 3k\mathcal{W}.$$

and is not preserved.

The most interesting feature of the DS deformations is that they are defined only for surfaces with fixed potentials of their Weierstrass representations. Indeed, for a torus we may take another gauge-equivalent potential

$$u \rightarrow u' = e^{az - \bar{a}\bar{z}}u \tag{24}$$

and apply the DS deformation for a torus with the potential. In this case the deformation would be completely different geometrically. It is noticeable from the deformation of $|u|^2$ which is, by [Proposition 1](#), is a geometric quantity.

Let us demonstrate that for the DS_3 deformation. The additional potentials defined by (21)–(23) are the same as for u but at $t = 0$ the deformation of $|u'|^2$ is different from the deformation of $|u|^2$ and it is as follows:

$$\frac{d|u'|^2}{dt} = \frac{d|u|^2}{dt} + 6 \operatorname{Re} [a^2(u_z\bar{u} + u\bar{u}_z) + a(u_{zz}\bar{u} - u\bar{u}_{zz})].$$

Although the first additional term is simple and equals to

$$3 \left(a^2 \frac{\partial |u|^2}{\partial z} + \bar{a}^2 \frac{\partial |u|^2}{\partial \bar{z}} \right),$$

i.e. could come from one-parametric diffeomorphism group of the surface, the second term involves the second derivatives and does not have such a form.

We conclude that

- the DS deformations are correctly defined only for surfaces with fixed potentials of their Weierstrass representations and for different choices of the potentials such deformations are geometrically different.

By [Theorem 3](#), for tori such deformations are parameterized by a \mathbb{Z}^2 lattice.

If we shall speak on local deformations then the gauge group is much larger (a gauge transformation is determined by a holomorphic function) and local deformations would be very different for different choices of gauge-equivalent potentials.

5. The spectral curve

It is reasonable to define the spectral curve for a torus in \mathbb{R}^4 as the spectral curve of a double-periodic operator \mathcal{D} coming in its Weierstrass representation.

Let us recall the definition of the spectral curve of a double-periodic Dirac operator \mathcal{D} ([15]).

For that consider all formal solutions to the equation

$$\mathcal{D}\psi = 0$$

meeting in addition the following periodicity conditions:

$$\psi(z + \gamma_j) = e^{2\pi i(k_1 \operatorname{Re} \gamma_j + k_2 \operatorname{Im} \gamma_j)} \psi(z) = \mu(\gamma_j) \psi(z), \quad j = 1, 2.$$

where $z \in \mathbb{C}$ and γ_1, γ_2 generate the periods lattice $A \subset \mathbb{C}$. Such solutions ψ are called Floquet eigenfunctions (on the zero level of energy), the quantities k_1, k_2 are called the quasimomenta of ψ , and $(\mu_1, \mu_2) = \mu(\gamma_1), \mu(\gamma_2)$ are the multipliers of ψ .

The quasimomenta satisfy some analytic relation (called in solid physics the dispersion relation):

$$P(k_1, k_2) = 0$$

which defines a complex curve Q_0 in \mathbb{C}^2 invariant with respect to translations

$$k \rightarrow k + \gamma^*, \quad \gamma^* = (\gamma_1^*, \gamma_2^*) \in A^*,$$

where $A^* \subset \mathbb{C} = \mathbb{R}^2 \subset \mathbb{C}^2$ is the dual lattice to A .

We say that the complex curve $\Gamma = Q_0/A^*$ is the spectral curve of \mathcal{D} (on the zero energy level). This definition originates in the definition of such a curve for a two-dimensional Schrödinger operator [3]. The mapping $\mathcal{M} : \Gamma \rightarrow \mathbb{C}^2$ formed by the multipliers $\mathcal{M} = (\mu_1, \mu_2)$ is called the multiplier mapping.

The spectral genus of a torus is defined as the geometric genus of the normalization of Γ .

For tori in \mathbb{R}^3 it appears that such a curve together with \mathcal{M} contains an important information about the conformal geometry of a torus. Our conjecture confirmed in [6] reads that the pair (Γ, \mathcal{M}) is preserved by conformal transformations of the ambient space \mathbb{R}^3 which map the torus into \mathbb{R}^3 . The discussion of other properties of the spectral curve can be found in [16].

For tori in \mathbb{R}^4 the situation is slightly different: the curve Γ is defined up to biholomorphic equivalences however the multiplier mappings depend on the choice of a potential: the gauge transformation (24) acts on ψ and \mathcal{M} as follows:

$$\psi \rightarrow e^{-az} \psi, \quad (\mu_1, \mu_2) \rightarrow (e^{-a\gamma_1} \mu_1, e^{-a\gamma_2} \mu_2).$$

This is rather reasonable. In the fundamental paper [11] by Novikov the spectral curve of an operator with a potential deformed via some soliton equation was considered as a conservation law itself for this equation. Since we show in Section 4 that there are infinitely many geometrically different soliton deformations of a torus in \mathbb{R}^4 described by the same DS equation, these different curves are just the values of the same conservation law for different solutions.

Acknowledgements

The work was supported by RFBR (grant 03-01-00403) and Max-Planck-Institute on Mathematics in Bonn. We thank U. Abresch for comments and discussions.

References

- [2] F. Burstall, F. Pedit, U. Pinkall, Schwarzian derivatives and flows of surfaces. Differential geometry and integrable systems (Tokyo, 2000), Contemp. Math., vol. 308, American Mathematical Society, Providence, RI, 2002, pp. 39–61.
- [3] B.A. Dubrovin, I.M. Krichever, S.P. Novikov, The Schrödinger equation in a periodic field and Riemann surfaces, Soviet Math. Dokl. 17 (1976) 947–952.
- [4] D. Ferus, K. Leschke, F. Pedit, U. Pinkall, Quaternionic holomorphic geometry: Plücker formula. Dirac eigenvalue estimates and energy estimates of harmonic 2-tori, Invent. Math. 146 (2001) 505–593.
- [5] O. Garay, J. Langer, Taimanov's surface evolution and Bäcklund transformations for curves, Conform. Geom. Dyn. 3 (1999) 37–49.
- [6] P.G. Grinevich, M.U. Schmidt, Conformal invariant functionals of immersions of tori into \mathbb{R}^3 , J. Geom. Phys. 26 (1997) 51–78.
- [7] F. Hélein, P. Romon, Weierstrass representation of Lagrangian surfaces in four dimensional space using spinors and quaternions, Comment. Math. Helv. 75 (2000) 668–680.
- [8] B.G. Konopelchenko, Induced surfaces and their integrable dynamics, Stud. Appl. Math. 96 (1996) 9–52.
- [9] B.G. Konopelchenko, Weierstrass representations for surfaces in 4D spaces and their integrable deformations via DS hierarchy, Ann. Global Anal. Geom. 16 (2000) 61–74.
- [10] J. Langer, Recursion in curve geometry, New York J. Math. 5 (1999) 25–51.
- [11] S.P. Novikov, A periodic problem for the Korteweg–de Vries equation, Funct. Anal. Appl. 8 (1974) 236–246.
- [12] F. Pedit, U. Pinkall, Quaternionic analysis on Riemann surfaces and differential geometry, Doc. Math., J. DMV Extra ICM II (1998) 189–200.
- [13] I.A. Taimanov, Modified Novikov–Veselov equation and differential geometry of surfaces, Am. Math. Soc. Transl. Ser. 2 179 (1997) 133–151.
- [14] I.A. Taimanov, Surfaces of revolution in terms of solitons, Ann. Global Anal. Geom. 12 (1997) 419–435.
- [15] I.A. Taimanov, The Weierstrass representation of closed surfaces in \mathbb{R}^3 , Funct. Anal. Appl. 32 (1998) 258–267.
- [16] I.A. Taimanov, Dirac operators and conformal invariants of tori in three-space, Proc. Steklov Inst. Math. 244 (2004).