

Alvie L. Davis, David L. Tabb, Janet K. Swan and Tommy J. McCord


Department of Chemistry, Abilene Christian University, Abilene, Texas 79699

Received April 18, 1980

The 3-methyl and 4-methyl derivatives of 3-amino-3,4-dihydro-1-hydroxycarbostyryl were synthesized by the reductive cyclization of α -methyl- β -(*o*-nitrophenyl)alanine and α -amino- β -(*o*-nitrophenyl)butyric acid hydrohalides, respectively, under conditions of catalytic hydrogenation in acidic solution. The free bases of the latter two *o*-nitroaromatic amino acids were also catalytically hydrogenated under neutral conditions to yield the respective α -methyl- β -(*o*-aminophenyl)alanine and α -amino- β -(*o*-aminophenyl)butyric acid which were converted to the corresponding lactams, 3-methyl- and 4-methyl-3-amino-3,4-dihydrocarbostyryls. α -Methyl- β -(*o*-nitrophenyl)alanine was obtained by acid hydrolysis of 5-methyl-5-(*o*-nitrobenzyl)hydantoin which was prepared by treatment of *o*-nitrophenylacetone with potassium cyanide and ammonium carbonate. α -Amino- β -(*o*-nitrophenyl)butyric acid was synthesized by condensation of α -bromo-*o*-nitroethylbenzene with diethyl acetamidomalonate, followed by acid hydrolysis of the condensation product. The 4-methylated compounds were obtained as synthetic mixtures of two diastereomeric racemates in nearly the same amounts as shown by nmr spectral analysis. Unlike the demethylated parent compound, 3-amino-3,4-dihydro-1-hydroxycarbostyryl, neither the 3-methyl nor 4-methyl analog was found to possess any antibacterial activity.

J. Heterocyclic Chem., 17, 1405 (1980).


In this laboratory we have been concerned for some time with the modification of the 3-amino-3,4-dihydro-1-hydroxycarbostyryl molecule (I) and the effects that these modifications have on biological activity (1-6) and on chemical rearrangement (7-9). Among the analogs of I




reported previously, the 5-, 6-, 7-, and 8-chlorosubstituted derivatives (II) have been studied most extensively for their antibacterial activities (5,6) and for their rearrangement in hydrohalic acids (9). Such studies have shown that the position of the chloro group on the benzene ring has a marked effect on the level of inhibitory activity against microbial growth and on the orientation of the halide nucleophile in the heteroaromatic rearrangement.

In order to obtain further information on the relationship between structural modification and biological activity, some addition analogs of I which contain a methyl group at positions 3 and 4 in the heterocyclic portion of the carbostyryl ring have been synthesized for microbiological assay. Therefore, the purpose of this paper is to describe the synthesis of the 3-methyl (III) and 4-methyl (IV) analogs of I and to disclose the finding that both methylated analogs are devoid of antibacterial activity. For purposes of comparison, the present study also includes the synthesis of the related lactams, the 3-methyl (V) and 4-methyl (VI) substituted 3-amino-3,4-dihydrocarbostyryls.

The syntheses of the 3-methyl cyclic hydroxamate III



and 3-methyl lactam V are depicted in Scheme I. *o*-Nitrophenylacetone was treated with potassium cyanide and ammonium carbonate in aqueous ethanol to give 5-methyl-5-(*o*-nitrobenzyl)hydantoin (VII), which was hydrolyzed in concentrated hydrobromic acid to afford α -methyl- β -(*o*-nitrophenyl)alanine hydrobromide (VIII). Reductive cyclization of the latter compound VIII to yield the desired 3-amino-3,4-dihydro-1-hydroxy-3-methylcarbostyryl hydrobromide (III) was effected under acidic conditions of catalytic hydrogenation in the presence of platinum on carbon, sulfided catalyst. The hydrobromide salt VIII was converted to the corresponding free base (IX) by neutralization with aqueous sodium hydroxide solution. Catalytic hydrogenation of the free base (IX) in the presence of platinum on carbon catalyst gave *o*-amino- α -methylphenylalanine (X). The *o*-aminoaromatic amino acid (X)



meric mixture is shown in Scheme II. *o*-Nitroethylbenzene was brominated with *N*-bromosuccinimide in the presence of benzoyl peroxide to yield α -bromo-*o*-nitroethylbenzene (XI). This compound was obtained as a low-melting solid which gave an acceptable elemental analysis, whereas no analytical data was reported in a previous paper which described its isolation as an oil (10). Condensation of XI with ethyl acetamidomalonnate (Et AAM) in ethanolic sodium ethoxide gave ethyl 2-acetamido-2-[α -(*o*-nitrophenyl)]malonnate (XII). The latter compound XII was hydrolyzed in a refluxing mixture of concentrated hydrochloric acid and glacial acetic acid to afford α -amino- β -(*o*-nitrophenyl)butyric acid hydrochloride (XIII). Compound XIII was reductively cyclized by catalytic hydrogenation under acidic conditions in the presence of platinum on carbon catalyst to afford 3-amino-3,4-dihydro-1-hydroxy-4-methylcarbostryl hydrochloride (IV).

The 4-methyl analog (VI) of the lactam was also prepared as a mixture of diastereomers by two different methods as presented in Scheme II. The hydrochloride salt of α -amino- β -(*o*-nitrophenyl)butyric acid was converted to the free base XIV prior to catalytic hydrogenation under neutral conditions using platinum on carbon as catalyst whereby α -amino- β -(*o*-aminophenyl)butyric acid (XV) was produced. The latter compound XV underwent lactamization in acidic solution to form 3-amino-3,4-dihydro-4-methylcarbostryl hydrochloride (VI). The other method leading to VI resulted from acid hydrolysis of

rapidly cyclized in acidic solution to give the lactam, 3-amino-3,4-dihydro-3-methylcarbostryl (V).

The 4-methyl analog (IV) of the carbostryl hydroxamic acid was obtained as a mixture of two diastereomeric racemates. The synthetic method leading to the diastereo-




Figure 1. The nmr spectrum of the diastereomeric 3-amino-3,4-dihydro-1-hydroxy-4-methylcarbostryl (IV).

3-acetamido-3-carboethoxy-3,4-dihydro-4-methylcarbo-
styryl (XVI), which was produced by reductive cyclization
of XII under conditions of catalytic hydrogenation.

The nmr spectra were used to determine the presence
and the relative amounts of the two diastereomers in the
synthetic mixtures of the carbostyryl compounds IV and
VI. As shown in Figure 1, the nmr spectrum of IV shows
two distinct doublets in the δ 1 to 2 region resulting from
the magnetic nonequivalence of the methyl groups as
based on the chemical shifts of the protons. The absorp-
tion of each methyl group appears as a doublet as a result
of coupling with the neighboring proton at position 4.
Since the relative intensities of the nmr absorptions
resulting from each diastereomer were observed to be
nearly the same, it was concluded that the synthetic
mixture contained almost equal amounts of the two
diastereomers.

Neither the 3-methyl (III) nor the 4-methyl (IV) analogs
of 3-amino-3,4-dihydro-1-hydroxycarbostyryl were active
against the growth of *Escherichia coli* 9723 and *Lacto-
bacillus plantarium* 8014 even at concentrations of 200
 $\mu\text{g./ml.}$ Therefore, substitution of a methyl group at posi-
tion 3 or 4 of 3-amino-3,4-dihydro-1-hydroxycarbostyryl
nullifies growth inhibitory activity in the microorganisms
studied.

In previous work, the various chloro-substituted carbo-
styryl hydroxamates were uniformly more effective anti-
microbial agents than the corresponding lactams (5,6).
This difference in antimicrobial activity was ascribed to
the presence of the hydroxamate function as part of the
heterocyclic ring. However, such is not the case with the
methylated carbostyryls which are inactive as growth in-
hibitors in spite of the hydroxamate group of III and IV. It
appears that methylation of the 3 position of I sterically in-
terferes with the 3-amino group and that methylation of
the 4 position substantially alters some key conformational
feature in the molecule to markedly decrease its affinity
for an enzyme or receptor site which is necessary for its in-
hibitory effect on microbial growth. A study describing the
separation and preparation of the two diastereomeric
4-methylated 3-amino-3,4-dihydro-1-hydroxycarbostyryls
and the 3-amino-3,4-dihydrocarbostyryls by preparative
liquid chromatography is forthcoming.

EXPERIMENTAL

General.

Melting points were determined on a Thomas-Hoover capillary
melting point apparatus and are uncorrected. Infrared spectra were
recorded on a Beckman Model IR-10 spectrophotometer (potassium
bromide) and were calibrated with polystyrene film. Nmr spectra were
recorded on a Perkin-Elmer Model R12-B spectrometer at 60 MHz. The
nmr spectrum of the diastereomeric mixture of 3-amino-3,4-dihydro-1-
hydroxy-4-methylcarbo-
styryl (30 mg.) was recorded in about 0.5 ml. of
20% deuterium chloride in deuterium oxide solution with 1% sodium
2,2-dimethyl-2-silapentane-5-sulfinate (DSS) as the reference standard.

Microanalyses were performed by M-H-W Laboratories, Phoenix,
Arizona.

5-Methyl-5-(*o*-nitrobenzyl)hydantoin (VII).

A suspension of 11.2 g. (0.0621 mole) of (2-nitrophenyl)acetone, 29.13
g. (0.303 mole) ammonium carbonate, and 8.45 g. (0.130 mole) potassium
cyanide in 200 ml. of 50% aqueous ethanol was heated at 55-60° in a
water bath for 4 hours. The solution was acidified with concentrated
hydrochloric acid, reduced to dryness *in vacuo*, and the resulting residue
was washed with water and benzene to give 12.57 g. (81.2%) of product.
An analytical sample, m.p. 188-190°, was obtained by recrystallization
from ethanol.

Anal. Calcd. for $\text{C}_{11}\text{H}_{11}\text{N}_3\text{O}_4$: C, 53.01; H, 4.45; N, 16.86. Found: C,
53.13; H, 4.55; N, 17.00.

o-Nitro- α -methylphenylalanine Hydrobromide (VIII).

A 6.0 g. (0.024 mole) sample of 5-methyl-5-(*o*-nitrobenzyl)hydantoin
(VII) was refluxed in 150 ml. of redistilled (48%) hydrogen bromide for
48 hours. The hot solution was treated with darco, filtered through celite,
and cooled to give 5.1 g. (69.7%) product. Recrystallization from 50%
aqueous alcohol gave an analytical sample, m.p. 243-246°.

Anal. Calcd. for $\text{C}_{10}\text{H}_{12}\text{N}_2\text{O}_4 \cdot \text{HBr}$: C, 39.36; H, 4.29; N, 9.18.
Found: C, 39.13; H, 4.44; N, 9.03.

o-Amino- α -methylphenylalanine (X).

A 400 mg. (0.0018 mole) sample of *o*-nitro- α -methylphenylalanine (IX),
m.p. 229° dec., prepared from VIII by treatment of an aqueous solution
with sodium hydroxide, was dissolved in 40 ml. of 75% aqueous
methanol and hydrogenated at 3.67 kg./cm² of hydrogen pressure in the
presence of 70 mg. of 5% platinum on carbon catalyst for 4 hours. The
catalyst was removed by filtration, and the filtrate was concentrated *in
vacuo* to about 3 ml. and cooled at -17° overnight to give 240 mg.
(66.3%) of product, m.p. 153-154°.

Anal. Calcd. for $\text{C}_{10}\text{H}_{14}\text{N}_2\text{O}_2 \cdot \frac{1}{2}\text{H}_2\text{O}$: C, 59.10; H, 7.44; N, 13.78.
Found: C, 58.94; H, 7.33; N, 13.87.

3-Amino-3,4-dihydro-1-hydroxy-3-methylcarbo- styryl Hydrobromide (III).

A 500 mg. (0.0016 mole) sample of *o*-nitro- α -methylphenylalanine
hydrobromide (VIII) was dissolved in 3 ml. of 50% aqueous methanol,
0.5 ml. of 48% hydrogen bromide was added, and the solution was
hydrogenated at 3.67 kg/cm² of hydrogen pressure in the presence of 50
mg. of 5% platinum on carbon sulfided catalyst for 3 hours. The
resulting precipitate was dissolved in water and the catalyst was removed
by filtration. The filtrate was reduced in volume *in vacuo* until cloudiness
appeared and 20 ml. of hydrogen bromide was added to help facilitate
precipitation. After chilling, the solution was filtered, and the resulting
precipitate was washed with acetone to give 250 mg. (58.0%) of product,
m.p. 315-316° dec. The ir spectrum showed major absorption bands at
2950 (broad), 1660, 1595, 1515, 1500, 1470, 1425, 1345, 755, and 695
cm⁻¹.

Anal. Calcd. for $\text{C}_{10}\text{H}_{12}\text{N}_2\text{O}_2 \cdot \text{HBr}$: C, 43.97; H, 4.80; N, 10.26.
Found: C, 43.94; H, 4.94; N, 10.14.

3-Amino-3,4-dihydro-3-methylcarbo- styryl Hydrochloride (V).

To a 200 mg. (0.00098 mole) sample of *o*-amino- α -methylphenylalanine
(X), dissolved in water was added 1 ml. of concentrated hydrochloric
acid. The volume of the solution was reduced *in vacuo* until precipitation
started to occur, and the solution was chilled to give 205 mg. (98.4%)
product. Recrystallization from methanol-ether gave an analytical sam-
ple, m.p. 311-313° dec. The ir spectrum showed major absorption bands at
3420, 3050 (broad), 1715, 1605, 1510, 1405, 1370, 1335, 1300, 1275,
780 and 765 cm⁻¹.

Anal. Calcd. for $\text{C}_{10}\text{H}_{12}\text{N}_2\text{O} \cdot \text{HCl}$: C, 56.47; H, 6.16; N, 13.17.
Found: C, 56.60; H, 6.23; N, 13.04.

α -Bromo-*o*-nitroethylbenzene (XI).

To a refluxing solution of 45 g. (0.30 mole) of freshly distilled *o*-nitro-
ethylbenzene and 150 ml. of anhydrous carbon tetrachloride were added
54 g. (0.30 mole) of *N*-bromosuccinimide and 1 g. of benzoyl peroxide.

The reaction mixture was treated with 15 g. of F-20 alumina, filtered, and the filtrate was passed through a 1 × 10 cm column charged with 40 g. of F-20 alumina. The column was eluted with 100 ml. of anhydrous carbon tetrachloride and the combined elements were distilled *in vacuo* to remove the unreacted starting material and residual carbon tetrachloride and then chilled to yield 62.4 g. (90.4%) of product, m.p. 10-12°.

Anal. Calcd. for $C_8H_8BrNO_2$: C, 41.76; H, 3.50; N, 6.09. Found: C, 41.57; H, 3.37; N, 6.06.

Ethyl 2-Acetamido-2-[α -(*o*-nitrophenethyl)]malonate (XII).

To a 57 g. (0.26 mole) sample of ethyl acetamidomalonnate dissolved in 500 ml. of anhydrous ethanol containing 6.0 g. (0.26 mole) of sodium was added 60 g. (0.26 mole) of α -bromo-*o*-nitroethylbenzene (XI), and the reaction mixture was stirred at 50° for 5 hours. Water was added to reaction mixture, and the resulting precipitate was filtered, air-dried, and washed with ether. Additional product was recovered from work-up of the filtrates and washings to give a total of 81.1 g. (85.1%) of product, m.p. 190-191°.

Anal. Calcd. for $C_{17}H_{22}N_2O_7$: C, 55.74; H, 6.05; N, 7.65. Found: C, 56.00; H, 5.99; N, 7.50.

α -Amino- β -(*o*-nitrophenyl)butyric acid Hydrochloride (XIII).

A 20 g. (0.055 mole) sample of ethyl acetamido-2-[α -(*o*-nitrophenethyl)]malonate (XII) was suspended in 180 ml. of concentrated hydrochloric acid and 120 ml. of glacial acetic acid, and the solution was refluxed for 16 hours. The reaction mixture was taken to dryness *in vacuo* and upon repeated addition and evaporation of anhydrous ethanol there was obtained 12.14 g. (84.7%) of product, m.p. 203-209°. A small portion was recrystallized from ethanol-acetone (1:1) to give an analytical sample.

Anal. Calcd. for $C_{10}H_{12}N_2O_4 \cdot HCl$: C, 46.07; H, 5.03; N, 10.75. Found: C, 45.82; H, 4.89; N, 10.63.

α -Amino- β -(*o*-aminophenyl)butyric Acid (XV).

A 1.0 g. (0.0045 mole) sample of the free base XIV of α -amino- β -(*o*-nitrophenyl)butyric acid, m.p. 201-202°, prepared by treatment of XIII with NaOH, was dissolved in 30 ml. of 95% aqueous methanol and hydrogenated at 3.67 kg./cm² of hydrogen pressure in the presence of 100 mg. of 5% platinum on carbon catalyst for 3 hours. The catalyst was removed by filtration, and the filtrate was reduced in volume *in vacuo* to give 0.71 g. (81.2%) of product, m.p. 180-181°.

Anal. Calcd. for $C_{10}H_{14}N_2O_2$: C, 61.82; H, 7.26; N, 14.42. Found: C, 61.69; H, 7.28; N, 14.28.

3-Amino-3,4-dihydro-1-hydroxy-4-methylcarbostyryl Hydrochloride (IV).

A 1.0 g. (0.0038 mole) sample of α -amino- β -(*o*-nitrophenyl)butyric acid hydrochloride (XIII) was suspended in 6 ml. of 50% aqueous methanol and 1 ml. of concentrated hydrochloric acid and hydrogenated at 3.67 kg./cm² of hydrogen pressure in the presence of 100 mg. of 5% platinum on carbon catalyst for 3 hours. The catalyst was removed by filtration, and the volume of the filtrate was reduced *in vacuo* almost to dryness. Addition of 25 ml. of acetone and chilling caused precipitation of 0.69 g. (79.4%) of product, m.p. 249-255°. Recrystallization from methanol gave an analytical sample. The ir spectrum showed major absorption bands at 2960 (broad), 1670, 1605, 1580, 1490, 1460, 1400, 970, and 750 cm⁻¹.

Anal. Calcd. for $C_{10}H_{12}N_2O_2 \cdot HCl$: C, 52.52; H, 5.73; N, 12.25. Found: C, 52.59; H, 5.74; N, 12.05.

3-Acetamido-3-carboethoxy-3,4-dihydro-4-methylcarbostyryl (XVI).

A 1.7 g. (0.0046 mole) sample of ethyl 2-acetamido-2-[α -(*o*-nitrophenethyl)]malonate (XII) was suspended in 10 ml. of 95% aqueous ethanol and hydrogenated at 3.67 kg./cm² of hydrogen pressure in the presence of 170 mg. of 5% platinum on carbon catalyst for 3 hours. The catalyst was removed by filtration, and the solution was reduced in volume *in vacuo* to give 1.25 g. (92.6%) of product, m.p. 118-120°.

Anal. Calcd. for $C_{15}H_{18}N_2O_4$: C, 62.06; H, 6.25; N, 9.65. Found: C, 62.10; H, 6.10; N, 9.62.

3-Amino-3,4-dihydro-4-methylcarbostyryl Hydrochloride (VI). Method A.

A suspension of 100 mg. (0.00051 mole) of α -amino- β -(*o*-aminophenyl)butyric acid (XV) dissolved in 5 ml. of 95% aqueous ethanol was treated with 5 drops of concentrated hydrochloric acid to effect solution. The volume of the solution was reduced *in vacuo* almost to dryness, and the addition of 5 ml. of acetone resulted in precipitation of 90 mg. (83%) of product, m.p. 256-258°. The ir spectrum showed major absorption bands at 3410, 2950 (broad), 1695, 1590, 1485, 1410, 1310, and 750 cm⁻¹.

Anal. Calcd. for $C_{10}H_{12}N_2O \cdot HCl$: C, 56.47; H, 6.16; N, 13.17. Found: C, 56.23; H, 6.17; N, 12.96.

Method B.

A 1.0 g. (0.0034 mole) sample of 3-acetamido-3-carboethoxy-3,4-dihydro-4-methylcarbostyryl (XVI) was heated at reflux in 10 ml. of concentrated hydrochloric acid for 3 hours. The volume of the reaction mixture was reduced *in vacuo* almost to dryness and the solution was chilled to give 630 mg. (87.1%) of product. The products from Methods A & B were shown to be identical by a comparison of their infrared spectra.

Acknowledgment.

The support of this work by research grants (R-285 and R-286 from the Robert A. Welch Foundation, Houston, Texas, is gratefully acknowledged.

REFERENCES AND NOTES

- (1) A. L. Davis, O. H. P. Choun, D. E. Cook and T. J. McCord, *J. Med. Chem.*, **7**, 632 (1964).
- (2) A. L. Davis, J. W. Hughes, R. L. Hance, V. L. Gault and T. J. McCord, *ibid.*, **13**, 549 (1970).
- (3) A. L. Davis, D. R. Smith, D. C. Foyt, J. L. Black and T. J. McCord, *ibid.*, **15**, 325 (1972).
- (4) A. L. Davis, D. R. Smith and T. J. McCord, *ibid.*, **16**, 1043 (1973).
- (5) A. L. Davis, W. H. Chambers, D. H. Kelley, D. A. Fell, J. R. Haynes, K. L. Hulme, L. D. Gage and T. J. McCord, *ibid.*, **18**, 752 (1975).
- (6) A. L. Davis, K. L. Hulme, G. T. Wilson and T. J. McCord, *Antimicrob. Agents Chemother.*, **13**, 542 (1978).
- (7) T. J. McCord, J. L. Kreps, J. N. Hubbard and A. L. Davis, *J. Heterocyclic Chem.*, **6**, 937 (1969).
- (8) T. J. McCord, D. H. Kelley, J. A. Rabon, D. C. Foyt and A. L. Davis, *ibid.*, **9**, 119 (1972).
- (9) T. J. McCord, S. F. Kelley, J. A. Rabon, L. D. Gage, L. L. Maples and A. L. Davis, *ibid.*, **13**, 1091 (1976).
- (10) P. M. Kochergin and K. S. Bushueva, *Zh. Obschch. Khim.*, **32**, 3033 (1962).