

B. Sain, J. N. Baruah and J. S. Sandhu\*


Regional Research Laboratory, Jorhat-785006, India

Received March 17, 1983


Az lactone **3a** reacted with furan-2-aldehyde and thiophene-2-aldehyde imines to give corresponding 3-amido- $\beta$ -lactams in good yields.

*J. Heterocyclic Chem.*, **21**, 257 (1984).


Because of several  $\beta$ -lactam antibiotics in clinical practice, extensive efforts have been made to further modify these molecules and have better understanding of structure and activity relationship [1-4]. Over the past several years various groups have been engaged in the synthesis of simple  $\beta$ -lactams [1,3]. These syntheses had been of preparative interest only, until the discovery of the non-classical  $\beta$ -lactam antibiotic nocardicin (**1**) [5]. The isolation of broad spectrum antibiotic thienamycin (**2**) demonstrated that the presence of the thiazole ring and the 6-amido side chain is not absolutely essential for optimal biological activity [6].



Münchone (3-methyl-2,4-diphenyl-oxazolium-5-olat) a mesoionic oxazolone which participates in 1,3-dipolar cycloaddition reactions with a variety of multiplet bonds gave  $\beta$ -lactams when reacted with imines, the formation of these products was explained *via* a ketene intermediate [7]. This mesoionic oxazolone also gave  $\beta$ -lactams when reacted with conjugated imines [8]. In contrast, azlactones ( $\Delta^2$ -oxazolin-5-ones) **3** reacted with conjugated imines **4** to yield  $\alpha$ -pyridones **5** and there was no evidence for the formation of  $\beta$ -lactams or 1,3 dipolar cycloadducts [8-11].



Now we report the reaction of azlactone **3a** with imines derived from furan-2-aldehyde and thiophene-2-aldehyde where corresponding  $\beta$ -lactams **7a-e** were obtained in good yields. The choice of furan as well as thiophene unit annulation with the  $\beta$ -lactam unit was made because of their already established biological importance.



Equimolar quantities of azlactone **3a** [10] and imine **6a** dissolved in dry toluene when stirred at 0° for 1 hour, then allowed to stand overnight at room temperature on working up gave white crystalline solid mp 165° in 80% yield. The ir (potassium bromide) spectrum showed strong band at 1750  $\text{cm}^{-1}$  indicating the presence of  $\beta$ -lactam, the other bands at 1655  $\text{cm}^{-1}$  and 3225  $\text{cm}^{-1}$  correspond to amido CO and NH respectively. The nmr spectrum (270 MHz, deuteriochloroform):  $\delta$  1.05-2.17 (13H, m), 3.50 (1H, m), 5.17 (1H, s), 6.42 (2H, m), 6.52 (1H, broad s), 7.27-7.50 (4H, m), 7.80 (2H, m) was consistent with the structure **7a**. The mass spectrum showed molecular ion at  $m/e$  352 (0.2%) and major fragments at 306 (5%), 277 (7%), 227 (80%), 226 (85%), 175 (20%), 149 (25%), 105 (100%).

Similar  $\beta$ -lactams **7b-e** were prepared and their characteristics are recorded in Tables 1 and 2. In the case of imines where the *N*-aryl group was present the reaction was found to be very slow and no significant amount of  $\beta$ -lactam could be observed. This suggests that the possible mechanistic pathway for this reaction could be the addition of oxazolone in its carbanion form on to C=N bond followed by attack of nucleophilic nitrogen. The reaction appears to be quite general and provides a convenient method for the synthesis of differently substituted 3-amido- $\beta$ -lactams.

#### EXPERIMENTAL

Melting points were taken in open capillary on a Büchi apparatus and are uncorrected. The nmr spectra were recorded on Bruker 270 MHz or Varian A-60T instrument and chemical shift values are recorded in  $\delta$  parts per million relative to internal standard (tetramethylsilane). Mass spectra were recorded on AEI MS 30 instrument. The ir spectra were recorded on Perkin Elmer 237B ir spectrometer in potassium bromide discs or chloroform solutions.

Table 1  
Physical and Analytical Data for  $\beta$ -Lactams

Compound No.	R <sub>1</sub>	R <sub>2</sub>	X	R <sub>3</sub>	Yield %	Mp (°C)	Molecular Formula	Analyses (%) Calcd. (Found)		
								C	H	N
7a	CH <sub>3</sub>	C <sub>6</sub> H <sub>5</sub>	O	C <sub>6</sub> H <sub>11</sub>	80	165	C <sub>21</sub> H <sub>24</sub> N <sub>2</sub> O <sub>3</sub>	71.59 (71.42)	6.82 (6.79)	7.95 (7.82)
7b	CH <sub>3</sub>	C <sub>6</sub> H <sub>5</sub>	O	CH <sub>3</sub>	75	35	C <sub>16</sub> H <sub>16</sub> N <sub>2</sub> O <sub>3</sub>	67.60 (67.85)	5.63 (5.71)	9.86 (9.89)
7c	CH <sub>3</sub>	C <sub>6</sub> H <sub>5</sub>	O	C <sub>2</sub> H <sub>5</sub>	65	Oil	C <sub>17</sub> H <sub>18</sub> N <sub>2</sub> O <sub>3</sub>	68.46 (68.62)	6.04 (6.15)	9.39 (9.21)
7d	CH <sub>3</sub>	C <sub>6</sub> H <sub>5</sub>	S	CH <sub>3</sub>	62	Oil	C <sub>16</sub> H <sub>16</sub> N <sub>2</sub> O <sub>2</sub> S	64.00 (64.18)	5.33 (5.12)	9.33 (9.45)
7e	CH <sub>3</sub>	C <sub>6</sub> H <sub>5</sub>	S	C <sub>2</sub> H <sub>5</sub>	60	Oil	C <sub>17</sub> H <sub>18</sub> N <sub>2</sub> O <sub>2</sub> S	64.97 (64.81)	5.73 (5.81)	8.92 (8.73)

Table 2  
Spectral Data of  $\beta$ -Lactams

Compound No.	IR $\nu$ max cm <sup>-1</sup>	PMR (deuteriochloroform) $\delta$ ppm	Mass Spectra m/e (%)
7a	3225, 1750, 1655	1.05-2.17 (13H, m), 3.50 (1H, m), 5.17 (1H, s), 6.42 (2H, m), 6.52 (1H, broad s), 7.27-7.50 (4H, m), 7.80 (2H, m)	352 (0.2), 306 (5), 277 (7), 277 (80), 226 (85), 175 (20), 149 (25), 105 (100)
7b	3250, 1750, 1660	1.45 (3H, s), 2.60 (3H, s), 5.16 (1H, s), 6.45 (2H, m), 6.53 (1H, broad s), 7.28-7.51 (4H, m), 7.80 (2H, m)	284 (0.5), 238 (80), 209 (50), 175 (20), 171 (40), 149 (20), 148 (70), 105 (100)
7c	3255, 1750, 1660	1.10 (3H, m), 1.42 (3H, s), 2.83 (2H, m), 5.15 (1H, s), 6.45 (2H, m), 6.52 (1H, broad s), 7.25-7.52 (4H, m), 7.82 (2H, m)	298 (0.6), 252 (100), 227 (10), 223 (70), 175 (15), 171 (8), 149 (70), 148 (75), 105 (85)
7d	3300, 1750, 1660	1.45 (3H, s), 2.60 (3H, s), 5.18 (1H, s), 6.80-7.50 (7H, m), 7.80 (2H, m)	300 (0.2), 254 (100), 225 (50), 191 (7), 175 (25), 151 (30), 105 (80)
7e	3300, 1750, 1662	1.10 (3H, m), 1.45 (3H, s), 2.85 (2H, m), 5.18 (1H, s), 6.80-7.50 (7H, m), 7.80 (2H, m)	314 (0.4), 268 (100), 243 (7), 239 (60), 175 (20), 165 (40), 105 (90)

Imines **6a-e** were prepared by known methods and their bp values agreed with those recorded in the literature.

Preparation of 3-Amido- $\beta$ -lactams **7a-e**. General Procedure.

To a stirred solution of imine **6** (0.01 mole) in dry toluene (10 ml) at 0° was added dropwise a cooled solution azlactone **3a** (0.01 mole) in dry toluene (5 ml). The mixture was stirred at 0° for 1 hour and then allowed to stand overnight at room temperature, which resulted in the separation of gummy material. On removal of toluene under vacuum and washing the residue left with benzene-petroleum ether mixture compounds **7a,b** were obtained as white crystalline solids. Compounds **7c-e** were obtained by purifying the residue left after removal of toluene, on thin layer chromatography using silica gel as adsorbent and chloroform methanol 9:1 as developing solvent system.

Acknowledgement.

We thank analytical chemistry division of this laboratory for the spectral analyses.

#### REFERENCES AND NOTES

[1] T. W. Doyle, B. Bealeu, B-Yu. Luh, C. F. Farrari and M. P. Cinnigham, *Can. J. Chem.*, **55**, 468 (1977) and references cited therein.

[2] "Recent Advances in the Chemistry of  $\beta$ -Lactam Antibiotics", G. I. Gregory, ed, Royal Society of Chemistry, London, 1981.

[3] A. K. Mukerjee and A. K. Singh, *Tetrahedron*, **34**, 1731 (1978).

[4] T. W. Doyle, D. B. Belleau, T. T. Conway, C. F. Farrari, D. E. Horning, G. Lim, B-Yu. Lua, A. Mortel, M. Menard and L. R. Morris, *Can. J. Chem.*, **58**, 2508 (1980).

[5] H. Aoki, H. Sakai, M. Kohsaka, T. Konomi, Y. Kubochi, J. Hosada, E. Iguchi and H. Imanaka, *J. Antibiot.*, **29**, 492 (1976).

[6a] G. Albers-Chonberg, B. H. Arison, O. D. Heusens, J. Hirshfield, K. Hoogsteen, E. A. Kazcka, R. E. Rhodes, J. S. Kahan, F. M. Kahan, R. W. Ratcliff, E. A. Walton, L. J. Ruswinble, R. B. Morin and B. G. Christensen, *J. Am. Chem. Soc.*, **100**, 6491 (1978); [b] J. Kahan, F. Kahan, E. Stapely, R. Gogelman and S. Hernandez, U. S. Patent 3,950,357 (1976); *Chem. Abstr.*, **85**, 92190t (1976).

[7] E. Funke and R. Huisgen, *Chem. Ber.*, **104**, 3222 (1971).

[8] B. Sain and J. S. Sandhu, unpublished results.

[9] S. Mohan, B. Kumar and J. S. Sandhu, *Chem. Ind. (London)*, 671 (1971).

[10] B. Sain, G. Thyagarajan and J. S. Sandhu, *Can. J. Chem.*, **58**, 2034 (1980).

[11] B. Sain, J. N. Baruah and J. S. Sandhu, *J. Heterocyclic Chem.*, in press.