

Sodium Periodate Oxidation of Tetrahydro- β -carboline Derivatives [1]

Franco Gatta*

Laboratorio di Chimica del Farmaco, Istituto Superiore di Sanità, Viale Regina Elena 299,
00161 Roma, Italy

Domenico Misiti

Dipartimento Studi Chimica e Tecnologia Sostanze Biologicamente Attive -
Università "La Sapienza" - P. le A. Moro 5,
00185 Roma, Italy

Received September 14, 1988

The oxidation of some 3-(methoxy- and ethoxycarbonyl)tetrahydro- β -carboline derivatives with sodium periodate led to the formation of 1,4-benzodiazonine derivatives or fully aromatic β -carbolines depending on both nature and number of substituents at 1-position.

J. Heterocyclic Chem., **26**, 537 (1989).

Compounds containing the β -carboline structure have recently aroused remarkable interest in neuropharmacology. Indeed several β -carboline-3-carboxylates have been shown to possess high affinity for benzodiazepine-binding brain proteins [2].

As a result of the continuing interest in the chemistry of the β -carbolines and as a further development of our previous work on the selenium-oxidation of tetrahydro- β -carboline nucleus [3], it seemed interesting to investigate the action of sodium periodate on some methyl and ethyl tetrahydro- β -carboline-3-carboxylates.

Sodium periodate has been used to oxidize with decarboxylation tetrahydro- β -carboline-3-carboxylic acid to β -carboline [4]. Since the esterification would avoid the

decarboxylation resulting from sodium periodate treatment, we tried the latter oxidizing agent in the case of some 3-(methoxy- and ethoxycarbonyl)tetrahydro- β -carbolines **1**, in order to obtain the related β -carboline esters.

However we observed that the course of sodium periodate oxidation in aqueous methanol of compounds **1** depended on the nature of the substituents at the 1-position (Scheme 1). So, when $R_1 = H$ and $R_2 = H$, alkyl or cycloalkyl group, by cleavage of the indole double bond, the 5-(methoxy- and ethoxycarbonyl)-1,3-dihydro-2*H*-1,4-benzodiazonin-2-ones **2** were formed. Compounds **1f,g** afforded **2f,g** also together with the 3-(methoxy- and ethoxycarbonyl)- β -carbolines **4f,g**, in an approximate ratio of 1:1.

Scheme 1




Table 1

Compound No.	Yields %	Mp (°C)	Recrystallized from	Molecular formula	Analysis %					
					Calcd.		Found		Found	
					C	H	N	C	H	N
1i	89	133-135	benzene	C ₁₅ H ₂₀ N ₂ O ₂	70.56	7.40	10.29	70.69	7.57	10.29
2a	42	178-180	methanol	C ₁₅ H ₁₈ N ₂ O ₂	63.92	4.95	11.47	63.72	5.07	11.49
2b	48	163-165	methanol	C ₁₄ H ₁₄ N ₂ O ₂	65.10	5.46	10.82	65.19	5.46	10.86
2c	50	197-200	methanol	C ₁₄ H ₁₄ N ₂ O ₂	65.10	5.46	10.82	65.28	5.69	11.09
2d	48	178-180	methanol	C ₁₅ H ₁₆ N ₂ O ₂	66.16	5.92	10.29	66.22	5.82	10.25
2e	55	185-187	methanol	C ₁₅ H ₁₆ N ₂ O ₂	66.16	5.92	10.29	66.13	5.79	10.07
2f	28	177-180	benzene	C ₁₅ H ₂₂ N ₂ O ₂	69.92	6.79	8.58	70.11	7.02	8.34
2g	22	87-89	<i>n</i> -hexane	C ₁₅ H ₂₀ N ₂ O ₂	70.35	6.22	8.64	70.52	6.06	8.50
3h	62	164-166	benzene	C ₁₅ H ₁₈ N ₂ O ₄	62.05	6.25	9.65	62.28	6.20	9.77
3i	60	139-141	benzene	C ₁₆ H ₂₀ N ₂ O ₄	63.14	6.59	9.21	63.38	6.73	9.19
3j	68	158-160	benzene	C ₁₇ H ₂₂ N ₂ O ₄	64.13	6.97	8.80	63.92	7.00	8.56
3k	74	168-170	benzene	C ₁₈ H ₂₂ N ₂ O ₄	65.44	6.71	8.48	65.47	6.66	8.20
4f	34	280-282	ethyl acetate	C ₁₉ H ₂₀ N ₂ O ₂	74.00	6.54	9.09	73.88	6.82	9.02
4g	30	262-264	ethyl acetate	C ₁₉ H ₁₈ N ₂ O ₂	74.49	5.92	9.15	74.38	5.77	9.05
4l	68	258-260 [a]	methanol	C ₁₉ H ₁₄ N ₂ O ₂	75.48	4.67	9.27	75.36	4.55	9.22
4m	75	242-244	methanol	C ₁₈ H ₁₂ N ₂ O ₄	63.38	4.26	10.02	63.21	4.24	9.86

[a] Lit [8] mp 254-255°.

The oxidation of compounds **1h-k** containing two alkyl substituents at the 1-position followed a different course and 5-(methoxy- and ethoxycarbonyl)-2,3,4,5,6,7-hexahydro-1*H*-1,4-benzodiazonine-2,7-diones **3h-k** were obtained.

Finally compounds **1l** (R₁ = phenyl and R₂ = H and **1m** (R₁ = methoxycarbonyl and R₂ = H), under the same oxidation conditions, gave the related β -carbolines **4l,m**.

It is noteworthy that the sodium periodate oxidation accompanied by cleavage of the indole double bond did seem to require in ring C of the tetrahydro- β -carbolines the presence of the unsubstituted NH group at the 2-position and the presence of the ester group at the 3-position: indeed attempts to oxidize 2-acetyltetrahydro- β -carboline [5] and 2-benzyl-3-(methoxycarbonyl)tetrahydro- β -carboline [6] were unsuccessful.

The structure of compounds **2** and **3** was supported by elemental analyses and by nmr and mass spectra.

Compounds **2** showed the following nmr spectral data (deuteriochloroform): a broad signal between δ 9.85 and 9.30 for the amide NH proton and two doublets in the range of 7.60-7.53 and 6.45-6.37 (J = 16 Hz), both integrating for 1 H, representing *trans* H-6 and H-7. Particularly compound **2a** exhibited a two proton sharp singlet at δ 4.23 (in **2b** it appeared at 4.33) for the methylene at the 3-position, while in **2c,d** the methine at the same position gave a quartet at 3.85 and in **2e** and a triplet at 3.44, affording indirect evidence that the double bond is positioned between C-6 and C-7.

Compound **3h** showed a broad amide NH signal at δ 9.23, a multiplet at 3.98 (partially overlapped by the

singlet of the methyl ester signal; in **3i** it clearly appeared at 3.95) assigned to the C-5 methine group, an apparent triplet at 3.00 due to the C-6 methylene group and a broad amine NH signal at 2.75.

The mass spectra were in agreement with the proposed structures. So, for example, the mass fragmentation pattern of the most significant compounds **2a** and **3h** were as follows: **2a**, *m/z* (%) 244 (M⁺, 50), 216 (23), 199 (19), 185 (100), 184 (20), 167 (33), 156 (25); **3h**, *m/z* (%) 262 (M⁺ - 28, 10), 205 (21), 146 (75), 142 (100), 129 (47), 120 (60), 92 (20).

EXPERIMENTAL

Melting points are uncorrected, the ¹H nmr spectra were determined on a T-60 Varian spectrometer with TMS as the internal standard. Electron ionization mass spectra were obtained on a LKB 2091 apparatus. Column chromatographic separations were accomplished on Merck silica gel (70-230 mesh). The drying agent was sodium sulfate. Yields, crystallization solvents, melting points and microanalytical data for all the new compounds described herein are reported in Table 1.

3-(Methoxy- and ethoxycarbonyl)tetrahydro- β -carbolines **1**.

Compounds **1a,b** [7], **1c** [8], **1d** [9], **1e** [10], **1f,l** [11] and **1h,j,k** [3] were synthesized according to the methods described in the literature. Compound **1g** was prepared by a Pictet-Splenger condensation between L-tryptophan methyl ester and 1,2,5,6-tetrahydrobenzaldehyde in refluxing toluene for 24 hours with a Dean-Stark trap to remove water. Compounds **1i,m** were obtained from the corresponding acids [12] by esterification in methanolic or ethanolic hydrogen chloride and converted into free bases. Compounds **1c**, **1d**, **1e**, **1f**, **1g**, **1l** and **1m** were employed as a mixture of *cis/trans* isomers and the new compounds **1g** and **1m** are not reported in Table 1.

Sodium Periodate Oxidation of Compounds **1**: 5-(Methoxy- and ethoxycarbonyl)-1,3-dihydro-2*H*-1,4-benzodiazonin-2-ones **2a-g**, 5-(Methoxy- and ethoxycarbonyl)-2,3,4,5,6,7-hexahydro-1*H*-1,4-benzodiazonine-2,7-diones **3h-k** and 3-(Methoxycarbonyl)- β -carbolines **4f,g,l,m**.

A solution of each compound **1** (0.02 mole) in methanol (250 ml) was added to a stirred solution of sodium periodate (17 g, 0.08 mole) in water (130 ml). The mixture was allowed to stir for 20 hours at room temperature for **1a-e** or refluxed for 4 hours for **1f-m**. The reaction was monitored by tlc, and stopped when starting material was disappeared. Then water (600 ml) was added and the mixture extracted with dichloromethane (3 x 200 ml). The solvent was evaporated and the residue was treated in a different way.

In the case of the oxidation of **1a-k** the residue was purified by chromatography on a silica gel column by eluting with ethyl acetate for (**1a-d**) or an ethyl acetate/*n*-hexane (1:1) mixture for **1e-k**. Compounds **2a-e** and **3h-k** were isolated and then crystallized. Compounds **4f,g** were eluted first followed by **2f,g**: both were recrystallized to give pure samples.

In the case of the oxidation of **1l,m** the residue was directly recrystallized to give pure **4l,m**.

Acknowledgement.

We gratefully acknowledge financial support from the Italian National Research Council (CNR) - Rome. We wish to thank Mr.

R. Piergallini for microanalyses and Mr. P. Montani for technical assistance.

REFERENCES AND NOTES

- [1] A preliminary account of this work was presented as a poster communication at the French-Italian Joint Meeting on Medicinal Chemistry - Piza, September 22-26, 1987.
- [2] C. Braestrup, M. Nielsen, and C. E. Olsen, *Proc. Natl. Acad. Sci.*, **77**, 2286 (1980).
- [3] F. Gatta and D. Misiti, *J. Heterocyclic Chem.*, **24**, 1183, (1987).
- [4] H. Akimoto, K. Okamura, M. Yui, T. Shioiri, K. Kuramoto, Y. Kigugawa, and S. Yamada, *Chem. Pharm. Bull.*, **22**, 2614 (1974).
- [5] Z. J. Veidelek, V. Troka, and M. Protiva, *J. Med. Pharm., Chem.*, **3**, 427 (1961).
- [6] This compound was obtained in 72% yield by reaction of 3-(methoxycarbonyl)tetrahydro- β -carboline with benzyl bromide and sodium hydrogen carbonate in refluxing methanol for 20 hours, mp 138-140°, from benzene-hexane.
- [7] K. P. Lippke, W. G. Schunach, W. Wenning, and W. E. Muller, *J. Med. Chem.*, **26**, 499 (1983).
- [8] H. R. Snyder, C. H. Hansch, L. Katz, S. M. Parmerter, and E. C. Spaeth, *J. Am. Chem. Soc.*, **70**, 219 (1948).
- [9] J. Le Men and C. Fan, *Bull. Soc. Chim. France*, 1866 (1959).
- [10] F. Ungemach, M. DiPierro, R. Weber, and J. Cook, *Tetrahedron Letters*, 3225 (1979).
- [11] D. Soerens, J. Sandri, F. Ungemach, P. Mokry, G. S. Wu, E. Yamanaka, L. Hutchins, M. DiPierro, and J. M. Cook, *J. Org. Chem.*, **44**, 535 (1979).