


## Some Nucleophilic Substitutions In 2-Cyano-3-Nitroimidazo[1,2-*a*]Pyridine

Lucina Arias,<sup>1</sup> Héctor Salgado-Zamora,<sup>1\*</sup> Humberto Cervantes,<sup>2</sup> Elena Campos,<sup>1</sup>  
Alicia Reyes<sup>1</sup> and Edward C. Taylor<sup>3</sup>

<sup>1</sup> Departamento de Química Orgánica, Escuela Nacional de Ciencias Biológicas, IPN. Prol. Carpio y Plan de Ayala s/n México 11340 D.F. <sup>2</sup> Área de Química, Universidad Autónoma Metropolitana (Unidad Azcapotzalco). México 02200 D.F. <sup>3</sup> Department of Chemistry, Princeton University, Princeton N.J. 08544 U.S.A  
Received July 5, 2005



In some nucleophilic substitution reactions of 2-cyano-3-nitroimidazo[1,2-*a*]pyridine, nitrogen (alkylamines, guanidine) and oxygen nucleophiles (alkoxides) underwent substitution of the 2-cyano group, while sulfur nucleophiles (alkylthiols) underwent substitution of the 3-nitro group.


*J. Heterocyclic Chem.*, **43**, 565 (2006).

### Introduction.

The imidazo[1,2-*a*]pyridine ring is a versatile system which has shown an outstanding array of biological activities [1]. The incorporation of the imidazo[1,2-*a*]pyridine moiety in structures investigated in relation to potential pharmacological activities has continued [2]. Electrophiles predominantly attack on the position 3 of the heterocycle [3]. Furthermore *ipso* electrophilic attack on 3-substituted (Br, NO<sub>2</sub>, CHO)-5-methylimidazo[1,2-*a*]pyridines by chlorine (NCS) rendered 3-chloro substituted derivatives albeit in low yields [4]. Nucleophiles attack both positions, 2 and 3, thus the reaction of 2-chloro-3-nitroimidazo[1,2-*a*]pyridine with diethylamine undergoes substitution of the 2-chloro group [5]. While the nitro functional group has been displaced from the 3-position by thiol nucleophiles if an electron-attracting substituent is present at the 2-position [6].

It is well known that the nitro group is a far better leaving group than the cyano group, particularly in aromatic nucleophilic substitutions [7,8]. Thus, the reaction of methyl 3-nitropyridine-4-carboxylate or 3-nitropyridine-4-carbonitrile undergoes substitution of the 3-nitro group [9]. It is also known that the nitrile group undergoes two types of reaction toward the nucleophile, the nucleophilic substitution as well as the nucleophilic addition. The former example is shown in the conversion of some aldehydes to corresponding esters, where the  $\alpha$ -oxonitriles, prepared by hydrocyanation of the aldehydes with hydrogen cyanide followed by oxidation with manganese dioxide, readily gave the corresponding esters by treatment with alcohols [10]. The latter mode, the nucleophilic addition, is shown in the condensation of 2-amino-5-nitrobenzotrile with 4-nitrobenzotrile giving 2-(4-nitrophenyl)-4-amino-6-nitroquinazoline [11]. The *cine*-

Scheme I




substitution in the 2-cyano-5-nitrofurán with carbanions and thiols also shows that the nitro group is a better leaving group than the cyano [12].

### Results and Discussion.

As shown in Scheme 1, 3-nitroimidazo[1,2-*a*]pyridine-2-carbonitrile (**3**) was prepared from ethyl 3-nitroimidazo[1,2-*a*]pyridine-2-carboxylate (**1**) [13] by ammonolysis with aqueous ammonia to give carboxamide **2** [14], followed by dehydration of the latter with phosphoryl oxychloride. The 3-nitro-2-carbonitrile **3** was then subjected to nucleophilic substitution reactions with guanidine in *n*-butanol. Unexpectedly this process yielded 2-guanidinyl-3-nitroimidazo[1,2-*a*]pyridine (**4a**), not 3-guanidinylimidazo[1,2-*a*]pyridine-2-carbonitrile (**5**), *i.e.* cyanide had been displaced in preference to the better leaving nitro group.


Treatment of **3** with excess amounts of some alkyl amines, as shown in Scheme II, in the absence of solvent gave the corresponding 2-alkylamino-3-nitroimidazo[1,2-*a*]pyridines (**4b-g**) in moderate to high yields. Use of microwave in treatment of **3** with cyclohexylamine reduced the reaction time from 2 h to 14 min and the molar ratio from 8.6 equivalents to 3, and improved the reaction yield from 62% to 97%. Treatment of **3** with aromatic amines caused no substitution. Treatment of **3** with excess amounts of two alkoxides, ethoxide and isopropoxide, in ethyl and isopropyl alcohol respectively, gave corresponding 2-alkoxy-3-nitroimidazo[1,2-*a*]pyridines (**6a,b**) in moderate yields.

Treatment of **3** with equal amount of two thiols, propane-1-thiol and ethyl thioglycolate in DMF, gave corresponding 3-alkylthioimidazo[1,2-*a*]pyridine-2-carbonitriles (**7a,b**) in moderate yields. Results for these substitution reactions are summarized in Table 1. It is very interesting that the nitrogen and oxygen nucleophiles showed the substitutions taking place on the 2-cyano (the less favored leaving group), and that sulfur nucleophiles showed the substitution on the 3-nitro (the more favored leaving group).



In the ir spectra of products **4a-g** and **6a-b**, the nitrile band disappeared and the nitro absorption bands appeared at  $\nu$  1300-1360 and 1500-1528  $\text{cm}^{-1}$ . In the  $^1\text{H}$  nmr spectra of **4a-g** and **6a,b**, the H-5 signal showed at low field in the interval  $\delta$  9.0 - 9.2 as has been established for 3-nitro substituted imidazopyridines [5,6] and as a clear indication of the permanence of the 3-nitro group. In the ir spectra of **7a,b**, the cyano bands appeared at 2238-2250  $\text{cm}^{-1}$  and the nitro bands disappeared. In the pmr spectra of **7a,b**, the H-5 signals shifted upfield, thus showing the displacement of the 3-nitro by thiol.

As shown in Scheme 3, treatment of **3** with other nitrogen nucleophiles, such as ammonia, hydroxylamine and phenylhydrazine showed no substitution on the 2-cyano or on the 3-nitro. Thus, treatment of **3** with aqueous ammonia at room temperature simply caused the hydrolysis of the nitrile to give 3-nitroimidazo[1,2-*a*]pyridine-2-carboxamide (**2**). Treatment of **3** with hydroxylamine caused the nucleophilic addition on the nitrile to give 3-nitroimidazo[1,2-*a*]pyridine-2-carboxamide-oxime (**8**). Treatment of **3** with an excess of phenylhydrazine at 70-75  $^\circ\text{C}$  caused the reduction [15] of the 3-nitro to give 3-aminoimidazo[1,2-*a*]pyridine-2-carbonitrile (**9**) in 73% yield.



Although displacement of either group may be mechanistically understandable in terms of an initial Michael type addition to either carbon supporting the leaving group, followed by an elimination process, it is also possible that the site in which the group itself is attached to the imidazopyridine system might be playing a decisive role on the preference for substitution. Therefore further efforts are on the way to determine whether this is the case. It is noteworthy to mention that the nitrile carbon in **3** shows a  $^{13}\text{C}$  nmr shift at  $\delta$  90, unusually high.

Some Nucleophilic Substitutions In 2-Cyano-3-Nitroimidazo[1,2-*a*]Pyridine

Table 1

Nucleophilic cyanide displacement in 2-cyano-3-nitroimidazoimidazo[1,2-*a*]pyridine **3**.

Compound	R	Reaction conditions	Melting point °C	Yield %
<b>4a</b>	NH <sub>2</sub> C(NH)	BuONa/BuOH, 80 °C	222-223	65
<b>4b</b>	PhCH <sub>2</sub>	60-70°C	157-158	77
<b>4c</b>	(CH <sub>3</sub> ) <sub>2</sub> CH	60-70°C	153-155	93
<b>4d</b>	Cyclohexyl	60-70°C	157-159	62
<b>4d</b>	Cyclohexyl	Microwave	157-159	97
<b>4e</b>	<i>n</i> -Butyl	60-70°C	73.5-74.5	72
<b>4f</b>	Cyclopropyl	60-70°C	186-187	88
<b>4g</b>	( <i>S</i> )-(-)-Ph(CH <sub>3</sub> )CH	60-70°C	102-103.5	48
<b>6a</b>	CH <sub>3</sub> CH <sub>2</sub>	EtONa, 1.5 mol/EtOH, 10 mL, 60 °C	147-149	75
<b>6b</b>	(CH <sub>3</sub> ) <sub>2</sub> CH	<i>i</i> PrONa, 1.5 mol/ <i>i</i> PrOH, 10 mL, 60 °C	142-143	70

## Conclusion.

In conclusion, several nitrogen and oxygen nucleophiles have shown a preference for cyanide displacement over the better leaving nitro group in 2-cyano-3-nitroimidazo[1,2-*a*]pyridine. However alkylthiol nucleophiles did displace the nitro group.

## EXPERIMENTAL

Melting points were measured on an Electrothermal melting point apparatus and are uncorrected. The IR spectra were taken on a Perkin Elmer 1600 series FTIR spectrophotometer. <sup>1</sup>H and <sup>13</sup>C NMR spectral data were recorded at 300 and 75 MHz respectively using a Bruker DPX 300 MHz NMR spectrometer. Chemical shifts (δ) are given in parts per million downfield from TMS (δ = 0). Mass spectra were obtained with a Jeol JMSAX505HA instrument.

2-Cyano-3-nitroimidazoimidazo[1,2-*a*]pyridine (**3**).

A suspension of 3-nitroimidazo[1,2-*a*]pyridine-2-carboxamide **2** (8.5 g, 41.6 mmol) in POCl<sub>3</sub> (85 mL) was stirred at 115 °C for 1 hour. After cooling, the mixture was concentrated under vacuum, and treated with ice (100 g). The yellow solid formed was collected by filtration and washed with cold water (50 mL x 2), and recrystallized from DMSO to afford **3**, 96% yield, mp 166-167 °C (lit. 162 °C [16,17]); IR (KBr, cm<sup>-1</sup>) ν 2250 (C≡N); 1629 (C=C); 1480, 1360 (NO<sub>2</sub>); <sup>1</sup>H NMR (deuteriochloroform): δ 9.42 (td, *J* = 7.0, 1.1 Hz, 1H, H-5); 7.95 (td, *J* = 9.1, 1.1 Hz, 1H, H-8); 7.81 (ddd, *J* = 9.1, 7.0, 1.1 Hz, 1H, H-7); 7.5 (td, *J* = 7.0, 1.1 Hz, 1H, H-6); cmr: 145.3, 131.9, 127.2, 125.8, 120.7, 119.6, 118.9, 111.7; ms: *m/z* (%) 188 (M<sup>+</sup>, 38), 78 (100).

2-Guanidinyl-3-nitroimidazo[1,2-*a*]pyridine (**4a**).

To a sodium *n*-butoxide/butanol solution, prepared from sodium metal (147 mg, 6.38 mmol) and *n*-butanol (20 mL) was added guanidine hydrochloride (508 mg, 5.32 mmol), and 2-cyano-3-nitroimidazo[1,2-*a*]pyridine **3** (1.0 g, 5.32 mmol) was added to the guanidine *n*-butanol solution. The mixture was stirred at 80 °C for 1 hour. After cooling, the solid formed was collected by filtration, washed with cold ethanol, and recrystallized from ethanol to afford **4a**, 65% yield, mp 222-223 °C; IR (KBr, cm<sup>-1</sup>) ν 3425, 3335, 3232 (NH); 1627 (C=C); 1482

(NO<sub>2</sub>); <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>): δ 9.37 (td, *J* = 7.0, 1.0 Hz, 1H, H-5); 7.63 (td, *J* = 7.9, 1.0 Hz, 1H, H-7); 7.43 (td, *J* = 8.8, 1.2 Hz, 1H, H-8); 7.14 (bs, 4H, NH<sub>2</sub>); 7.12 (td, *J* = 7.0, 1.2 Hz, 1H, H-6); cmr: δ 160.84, 157.2, 144.5, 133.2, 129.1, 122.03, 115.1, 114.7; ms: *m/z* (%) 220 M<sup>+</sup> (6.3), 120 (100).

Anal. Calcd. for C<sub>8</sub>H<sub>8</sub>N<sub>6</sub>O<sub>2</sub>: C, 43.63; H, 3.66. Found: C, 43.70; H, 3.45.

General Procedure for Preparation of 2-Alkylamino-3-nitroimidazo[1,2-*a*]pyridine **4b-4g**.

A mixture of 2-cyano-3-nitroimidazo[1,2-*a*]pyridine **3** (1.0 g, 5.32 mmol) and alkylamine (16.3 mmol) was stirred at 60-70 °C for 2 hours. After cooling, the solid formed was collected by filtration and washed with cold ethanol, and recrystallized from ethanol to afford the corresponding 2-alkylamino-3-nitroimidazo[1,2-*a*]pyridine **4b-g**.

2-(*N*-Benzyl)amino-3-nitroimidazo[1,2-*a*]pyridine **4b**.

Obtained in 77% yield, mp 157-158 °C; IR (KBr, cm<sup>-1</sup>) ν 3339 (NH); 1605 (C=C); 1469, 1317 (NO<sub>2</sub>); <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>): δ 9.25 (d, *J* = 6.8 Hz, 1H, H-5); 8.52 (t, *J* = 4.5 Hz, 1H, NH); 7.69 (ddd, *J* = 8.5, 7.2, 1.4 Hz, 1H, H-7); 7.47 (d, *J* = 8.8 Hz, 1H, H-8); 7.39-7.18 (m, 5H, -C<sub>6</sub>H<sub>5</sub>); 7.16 (ddd, *J* = 7.2, 7.0, 1.0 Hz, 1H, H-6); 4.73 (d, *J* = 6.5 Hz, 2H, CH<sub>2</sub>); cmr: δ 153.8 (C-2); 146.6 (C-8a); 139.3 (Ar, *ipso*); 134.3 (C-7); 128.7 (C-5); 128.3 (Ar, *m*); 127.3 (Ar, *o*); 126.9 (Ar, *p*); 117.6 (C-3?); 114.7 (C-8); 114.1 (C-6); 45.1 (CH<sub>2</sub>); ms: *m/z* (%) 268 M<sup>+</sup> (15.8), 91 (100).

Anal. Calcd. for C<sub>14</sub>H<sub>12</sub>N<sub>4</sub>O<sub>2</sub>: C, 62.67; H, 4.50. Found: C, 62.43; H, 4.67.

2-(*N*-Isopropyl)amino-3-nitroimidazo[1,2-*a*]pyridine (**4c**).

Obtained in 93% yield, mp 153-155 °C; IR (KBr, cm<sup>-1</sup>) ν 3369 (NH); 1606 (C=C); 1454, 1332 (NO<sub>2</sub>); <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>): δ 9.24 (td, *J* = 6.7, 1.1 Hz, 1H, H-5); 7.73 (td, *J* = 8.0, 1.4 Hz, 1H, H-7); 7.58 (d, *J* = 8.4 Hz, 1H, NH); 7.50 (td, *J* = 8.8, 1.1 Hz, 1H, H-8); 7.16 (td, *J* = 7.0, 1.2 Hz, 1H, H-6); 4.27 (m, 1H, CH); 1.28 (d, *J* = 6.6 Hz, 6H, CH<sub>3</sub>); cmr: δ 153.2, 146.8, 134.4, 128.7, 117.5, 114.7, 114.1, 44.2, 22.3; ms: *m/z* (%) 220 M<sup>+</sup> (76), 78 (100).

Anal. Calcd. For C<sub>10</sub>H<sub>12</sub>N<sub>4</sub>O<sub>2</sub>: C, 54.53; H, 5.49; N, 25.44. Found: C, 54.65; H, 5.74

2-(*N*-Cyclohexyl)amino-3-nitroimidazo[1,2-*a*]pyridine (**4d**).

Obtained in 62% yield, mp 157-159 °C. IR (KBr, cm<sup>-1</sup>) ν 3333 (NH); 1606 (C=C); 1461, 1332 (NO<sub>2</sub>); <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>): δ

9.23 (d,  $J = 7.0$  Hz, 1H, H-5); 7.73 (td,  $J = 8.0, 1.2$  Hz, 1H, H-7); 7.57 (d,  $J = 8.5$  Hz, 1H, NH); 7.51 (d,  $J = 8.8$  Hz, 1H, H-8); 7.16 (t,  $J = 7.0$  Hz, 1H, H-6); 3.93 (m, 1H, CH<sub>cyclohexyl</sub>); 1.18-1.94 (m, 10H, CH<sub>2 (cyclohexyl)</sub>); cmr:  $\delta$  153.1, 146.9, 134.5, 128.7, 117.5, 114.7, 114.1, 51.1, 32.3, 25.1, 24.7; ms;  $m/z$  (%) 260 M<sup>+</sup> (97.5), 243 (100).

Anal. Calcd. For C<sub>13</sub>H<sub>16</sub>N<sub>4</sub>O<sub>2</sub>: C, 59.98; H, 6.19; N, 21.52. Found: C, 59.68; H, 6.34

#### 2-(*N*-1-Butyl)amino-3-nitroimidazo[1,2-*a*]pyridine (4e).

Obtained in 72% yield, mp 73.5-74.5°C. Ir (KBr cm<sup>-1</sup>)  $\nu$  3375 (NH); 1606 (C=C); 1461, 1332 (NO<sub>2</sub>); <sup>1</sup>H nmr (deuteriochloroform):  $\delta$  9.37 (ddd,  $J = 6.8, 1.4, 1.0$  Hz, 1H, H-5); 7.60 (td,  $J = 8.0, 1.5$  Hz, 1H, H-7); 7.46 (bs, 1H, NH); 7.42 (td,  $J = 8.8, 1.1$  Hz, 1H, H-8); 7.03 (td,  $J = 7.0, 1.2$  Hz, 1H, H-6); 3.65 (m, 2H, CH<sub>2 (n-butyl)</sub>); 1.72 (m, 2H, CH<sub>2 (n-butyl)</sub>); 1.47 (m, 2H, CH<sub>2 (n-butyl)</sub>); 0.98 (t,  $J = 7.3$  Hz, 3H, CH<sub>3 (n-butyl)</sub>); cmr:  $\delta$  154.4, 147.2, 133.5, 128.6, 118.0, 114.9, 113.5, 42.1, 31.6, 19.9, 13.6; ms;  $m/z$  (%) 234 M<sup>+</sup> (95), 146.2 (100).

Anal. Calcd. For C<sub>11</sub>H<sub>14</sub>N<sub>4</sub>O<sub>2</sub>: C, 56.39; H, 6.02; N, 23.91. Found: 56.23; H, 5.81

#### 2-(*N*-Cyclopropyl)amino-3-nitroimidazo[1,2-*a*]pyridine (4f).

Obtained in 88% yield, mp 186-187°C. Ir (KBr cm<sup>-1</sup>)  $\nu$  3329 (NH); 1611 (C=C); 1468, 1334 (NO<sub>2</sub>); <sup>1</sup>H nmr (deuteriochloroform):  $\delta$  9.38 (td,  $J = 6.8, 1.1$  Hz, 1H, H-5); 7.62 (td,  $J = 7.9, 1.5$  Hz, 1H, H-7); 7.53 (td,  $J = 8.8, 1.1$  Hz, 1H, H-8); 7.51 (bs, 1H, NH); 7.07 (td,  $J = 6.9, 1.3$  Hz, 1H, H-6); 3.03 (m, 1H, CH<sub>cyclopropyl</sub>); 0.98 (m, 2H, CH<sub>2 (cyclopropyl)</sub>); 0.76 (m, 2H, CH<sub>2 (cyclopropyl)</sub>); cmr:  $\delta$  155.1, 146.9, 133.4, 128.5, 118.1, 115.3, 113.8, 24.5, 7.3; ms;  $m/z$  (%) 218 M<sup>+</sup> (8), 201 (100).

Anal. Calcd. For C<sub>10</sub>H<sub>10</sub>N<sub>4</sub>O<sub>2</sub>: C, 55.04; H, 4.61; N, 25.67. Found: C, 55.41; H, 4.85

#### 2-(*N*-(*S*)-(-)- $\alpha$ -Methylbenzyl)amino-3-nitroimidazo[1,2-*a*]pyridine (4g).

Obtained in 48% yield, mp 102-103°C. Ir (KBr cm<sup>-1</sup>)  $\nu$  3346 (NH); 1601 (C=C); 1457, 1334 (NO<sub>2</sub>); <sup>1</sup>H nmr (deuteriochloroform):  $\delta$  9.35 (td,  $J = 6.8, 1.2$  Hz, 1H, H-5); 7.74 (d,  $J = 8.0$  Hz, 1H, NH) 7.56 (td,  $J = 8.0, 1.4$  Hz, 1H, H-7); 7.45-7.24 (m, 6H, H-8 and Ph) 7.01 (td,  $J = 6.8, 1.2$  Hz, 1H, H-6); 5.40 (m, 1H, CH); 1.69 (d,  $J = 9.2$  Hz, 3H, CH<sub>3</sub>); cmr:  $\delta$  153.4, 147.0, 142.8, 133.3, 128.66, 128.60, 127.43, 125.96, 115.1, 113.6, 51.8, 22.8; ms;  $m/z$  (%) 282 M<sup>+</sup> (27), 265.1 (100).

Anal. Calcd. For C<sub>15</sub>H<sub>14</sub>N<sub>4</sub>O<sub>2</sub>: C, 63.81; H, 4.99; N, 19.84. Found: C, 63.64; H, 4.67

#### General procedure for the synthesis of 2-alkoxy-3 nitroimidazo[1,2-*a*]pyridine (6a,b).

A mixture of 2-cyano-3-nitroimidazo[1,2-*a*]pyridine 3 (1.0 g, 5.32 mmol), sodium alkoxide (8 mmol) and the corresponding alcohol (10 mL) was stirred at ca. 60 °C for 2 hours. After cooling, the solid formed was collected by filtration and washed with cold ethanol, and recrystallized from ethanol to afford the corresponding 2-alkoxy-3-nitroimidazo[1,2-*a*]pyridine 6a,b.

#### 2-Ethoxy-3-nitroimidazo[1,2-*a*]pyridine (6a).

Obtained in 75 % yield, mp 147-149°C. Ir (KBr cm<sup>-1</sup>)  $\nu$  1540 (C=C); 1486, 1327 (NO<sub>2</sub>); 1210, 1022 (C-O-C); <sup>1</sup>H nmr (CDCl<sub>3</sub>):  $\delta$  9.47 (td,  $J = 6.8, 1.0$  Hz, 1H, H-5); 7.69-7.57 (m, 2H, H-8 and H-7); 7.23 (td,  $J = 6.8, 1.6$  Hz, 1H, H-6); 4.71 (c,  $J = 7.2$

Hz, 2H, OCH<sub>2</sub>); 1.56 (t,  $J = 7.2$  Hz, 3H, CH<sub>3</sub>); cmr:  $\delta$  158.7, 142.1, 131.7, 128.1, 116.3, 115.5, 66.5, 14.6; ms;  $m/z$  (%) 207 M<sup>+</sup> (41), 78.1 (100).

Anal. Calcd. For C<sub>9</sub>H<sub>9</sub>N<sub>3</sub>O<sub>3</sub>: C, 52.17; H, 4.37; N, 20.28. Found: 52.39, H, 5.74

#### 2-Isopropoxy-3-nitroimidazo[1,2-*a*]pyridine (6b).

Obtained in 70 % yield, mp 142-143°C. Ir (KBr cm<sup>-1</sup>)  $\nu$  (C=C); (NO<sub>2</sub>); (C-O-C); <sup>1</sup>H nmr (DMSO-d<sub>6</sub>):  $\delta$  9.39 (dd,  $J = 6.6, 1.2$  Hz, 1H, H-5); 7.78 (td,  $J = 8.0, 1.4$  Hz, 1H, H-7); 7.70 (dd,  $J = 7.7, 1.2$  Hz, 1H, H-8); 7.36 (td,  $J = 6.9, 1.6$  Hz, 1H, H-6); 5.44 (m, 1H, CH); 1.48 (d,  $J = 6.3$  Hz, 6H, CH<sub>3</sub>); cmr:  $\delta$  156.1, 140.2, 131.6, 130.6, 126.3, 114.6, 114.3, 71.8, 20.1; ms;  $m/z$  (%) 221 M<sup>+</sup> (51), 222 MH<sup>+</sup> (100).

Anal. Calcd. For C<sub>10</sub>H<sub>11</sub>N<sub>3</sub>O<sub>3</sub>: C, 54.29; H, 5.01; N, 18.99. Found: C, 53.91, H, 4.65.

#### General procedure for the synthesis of compounds 7a,b.

To a well-stirred, cold solution (ice bath) containing 0.2 g (1.06 mmol) of 2-cyano-3-nitroimidazo[1,2-*a*]pyridine (3) and 0.1 mL (1.06 mmol) of methyl thioglycolate in DMF (2.1 mL), 0.09 g of sodium bicarbonate (1.06 mmol) was slowly added. The mixture was stirred for 1 h at low temperature and then at room temperature during 2 h. The solution was poured into ice-water and the precipitated cyanothioester was collected by filtration, washed with water (2x5 mL) and cold ethanol (5 mL) The solid was further purified by re-crystallization from ethanol.

#### Methyl 2-Cyanoimidazo[1,2-*a*]pyridine-3-thioglycolate (7a).

This compound was obtained in 88% yield, mp 96-98°C. Ir (KBr cm<sup>-1</sup>)  $\nu$  2250 (CN); 1742 (C=O); 1436 (C=C); <sup>1</sup>H nmr (DMSO-d<sub>6</sub>):  $\delta$  8.67 (td,  $J = 8.1, 1.1$  Hz, 1H, H-5); 7.76 (td,  $J = 9.2, 1.1$  Hz, 1H, H-8); 7.60 (s, 2H, SCH<sub>2</sub>); 3.57 (m, 2H, OCH<sub>3</sub>); cmr:  $\delta$  168.89, 146.55, 129.04, 125.82, 124.28, 120.23, 117.87, 115.20, 114.22, 52.28, 36.10; ms;  $m/z$  (%) 247 M<sup>+</sup> (31), 174 (100).

Anal. Calcd. For C<sub>11</sub>H<sub>9</sub>N<sub>3</sub>O<sub>2</sub>S: C, 53.43; H, 3.66; N, 16.99. Found: C, 53.38; H, 3.45.

#### 2-Cyano-3-propanethiolimidazo[1,2-*a*]pyridine (7b).

This compound was obtained in 74% yield, mp 77-78 °C. Ir (KBr cm<sup>-1</sup>)  $\nu$  (CN); (C=C); <sup>1</sup>H nmr (deuteriochloroform):  $\delta$  8.42 (td,  $J = 7.0, 1.1$  Hz, 1H, H-5); 7.68 (td,  $J = 9.2, 1.1$  Hz, 1H, H-8); 7.43 (ddd,  $J = 9.2, 6.9, 1.2$  Hz, 1H, H-7); 7.09 (td,  $J = 6.9, 1.1$  Hz, 1H, H-6); 2.78 (t,  $J = 7.4$  Hz, 3H, SCH<sub>2</sub>); 1.6 (m, 2H, CH<sub>2</sub>); 1.03 (t,  $J = 7.4$  Hz, 3H, CH<sub>3</sub>); cmr:  $\delta$  146.76, 133.85, 127.84, 125.08, 124.55, 122.74, 118.76, 114.78, 114.31, 38.03, 23.25, 12.96; ms;  $m/z$  (%) 217 M<sup>+</sup> (21), 175 (100).

Anal. Calcd. For C<sub>11</sub>H<sub>11</sub>N<sub>3</sub>S: C, 60.80; H, 5.10; N, 19.33. Found: C, 60.52; H, 4.86.

#### 2-Amidoxime-3-nitroimidazo[1,2-*a*]pyridine (8).

2-Cyano-3-nitroimidazo[1,2-*a*]pyridine (0.2g, 1.06 mmol), hydroxylamine hydrochloride (0.36 g, 5.32 mmol) and sodium acetate (0.43 g, 5.32 mmol) were suspended in 2-propanol and heated to a reflux temperature for 2 h. The reaction mixture was allowed to cool to room temperature and the formed solid was collected by filtration and washed with cold water. The title compound was isolated as a yellow solid (0.2 g, 85 % yield), mp 178-180 °C. IR (KBr cm<sup>-1</sup>)  $\nu$  3453, 3353 (NH<sub>2</sub>); 1643 (C=N); 1485, 1338 (NO<sub>2</sub>); <sup>1</sup>H nmr (DMSO-d<sub>6</sub>):  $\delta$  9.86 (s, 1H, OH); 9.35

Some Nucleophilic Substitutions In 2-Cyano-3-Nitroimidazo[1,2-*a*]Pyridine

(d,  $J = 6.4$  Hz, 1H, H-5); 7.94 (d,  $J = 8.4$  Hz, 1H, H-8); 7.82 (td,  $J = 7.8, 1.3$  Hz, 1H, H-7); 7.49 (td,  $J = 7.0, 1.2$  Hz, 1H, H-6); 5.92 (s, 2H, NH<sub>2</sub>): cmr  $\delta$  145.9, 143.9, 141.9, 131.3, 127.6, 118.0, 117.3; ms; m/z (%) 221 M<sup>+</sup> (49), 78.1 (100).

Anal. Calcd. For C<sub>8</sub>H<sub>7</sub>N<sub>5</sub>O<sub>3</sub>: C, 43.44; H, 3.18; N, 31.66. Found: C, 43.08; H, 2.87

3-Aminoimidazo[1,2-*a*]pyridine-2-carbonitrile (**9**).

A mixture of 2-cyano-3-nitroimidazo[1,2-*a*]pyridine **3** (0.5 g, 2.65 mmol) and phenylhydrazine (25.4 mmol) was stirred at 60-70 °C for 12 hours. After cooling, the solid formed was collected by filtration, washed with ethyl ether (10 mL), and recrystallized from ethanol to afford the corresponding 3-aminoimidazo[1,2-*a*]pyridine-2-carbonitrile (**9**) (0.3 g, 73 % yield), mp 238-239 °C (dec.), lit. mp 209-211 °C [18].

## Acknowledgment.

Financial support from the Instituto Politécnico Nacional (México) through grants 20030766 and 20040295 is gratefully acknowledged.

## REFERENCES AND NOTES

- [1] A. Gueiffier, M. Lhassani, A. Elhakmaoui, R. Snoeck, G. Andrei, O. Chavignon, J.-C. Teulade, A. Kerbal, E. M. Essassi, J.-C. Debouzy, M. Witvrouw, Y. Balzarini, E. D. Clercq and J.-P. Chapat, *J. Med. Chem.*, **39**, 2856 (1996).
- [2] M. I. Ismail, R. Brun, T. Wenzler, F. A. Tanious, W. D. Wilson and D. W. Boykin, *J. Med. Chem.*, **47**, 3658 (2004).
- [3] W. W. Paudler and H. L. Blewitt, *Tetrahedron*, **21**, 353 (1965).
- [4] T. Ikemoto and M. Wakimasu, *Heterocycles*, **55**, 99 (2000).
- [5] J. P. Paolini and R. K. Robins, *J. Org. Chem.*, **30**, 4085 (1965).
- [6] J. C. Teulade, G. Grassy, R. Escale and J. P. Chapat, *J. Org. Chem.*, **46**, 1026 (1981).
- [7] F. Effenberger, M. Kochand W. Streicher, *Chem. Ber.*, **124**, 163 (1991).
- [8] J. R. Beck, *Tetrahedron*, **34**, 2057 (1978).
- [9] J. Holt, F. Tjosas, J. M. Bakke and A. Fiksdahl, *J. Heterocyclic Chem.*, **41**, 987 (2004).
- [10] E. J. Corey, N. W. Gilman, B. E. Ganem, *J. Am. Chem. Soc.*, **90**, 5616 (1968).
- [11] E. C. Taylor and A. L. Borrer, *J. Org. Chem.*, **26**, 4967 (1961).
- [12] M. Makosza, *Polish J. Chem.*, **66**, 3 (1992).
- [13] J.-C. Teulade, G. Grassy, J.-P. Girard, and J.-P. Chapat. *Eur. J. Med. Chem.*, **13**, 271 (1978).
- [14] H. Salgado-Zamora, B. Rizo, E. Campos, R. Jiménez and A. Reyes. *J. Heterocyclic Chem.*, **41**, 91 (2004).
- [15] R. Walther, *J. Prakt. Chem.* **52**, 141 (1895); **53**, 433 (1896).
- [16] V.E Winkelmann, W. Raether, H. Hartung, and W.H. Wagner. *Arzneim.-Forsch./Drug Res.*, **27**, 82 (1977)
- [17] J.-C. Teulade, G. Grassy, J.-P. Girard and J.-P. Chapat. *Eur. J. Med. Chem.*, **13**, 271 (1978).
- [18] A. Gueiffier, O. Chavignon, S. Mavel, J.-M. Chezal, J.-C. Teulade, I. Blache, and J.-P. Chapat, *Heterocyclic Commun.*, **2**, 241 (1996).