

Neutron-electron EDM correlations in supersymmetry and prospects for EDM searches

Steven A. Abel

*IPPP, University of Durham
DH1 3LE Durham, U.K.
E-mail: s.a.abel@durham.ac.uk*

Oleg Lebedev

*Deutsches Elektronen-Synchrotron DESY
22603 Hamburg, Germany
E-mail: lebedev@mail.desy.de*

ABSTRACT: Motivated by recent progress in experimental techniques of electric dipole moment (EDM) measurements, we study correlations between the neutron and electron EDMs in common supersymmetric models. These include minimal supergravity (mSUGRA) with small CP phases, mSUGRA with a heavy SUSY spectrum, the decoupling scenario and split SUSY. In most cases, the electron and neutron EDMs are found to be observable in the next round of EDM experiments. They exhibit certain correlation patterns. For example, if $d_n \sim 10^{-27} e \text{ cm}$ is found, d_e is predicted to lie in the range $10^{-28} - 10^{-29} e \text{ cm}$.

KEYWORDS: Beyond Standard Model, Supersymmetric Standard Model, CP violation.

Contents

1.	Introduction	1
2.	EDMs in the standard model	3
3.	EDMs in supersymmetry	3
3.1	Small CP phases	4
3.2	Heavy SUSY spectrum	5
3.3	Decoupling	6
3.4	Split SUSY	8
4.	Conclusions	9

1. Introduction

The current EDM limits [1] –[3]¹

$$\begin{aligned}
 |d_n| &< 6 \times 10^{-26} \text{ e cm} , \\
 |d_e| &< 2 \times 10^{-27} \text{ e cm} , \\
 |d_{\text{Hg}}| &< 2 \times 10^{-28} \text{ e cm}
 \end{aligned}
 \tag{1.1}$$

impose severe constraints on CP violating physics beyond the Standard Model and, in particular, supersymmetry. Supersymmetric models accommodating the above bounds usually predict EDMs not far from the current limits which makes the next round of EDM experiments with the sensitivity $d_n \sim 10^{-28} \text{ e cm}$, $d_e \sim 10^{-30} \text{ e cm}$ [4] –[8] particularly interesting. Other EDM experiments such as measurements of the deuteron and muon EDMs [9] would provide important complementary information on CP violation in supersymmetry.

In this work, we address the questions “What are the EDM expectations in SUSY models ?” and “How can one distinguish different sources of EDMs ?”. It is well known that generic SUSY models predict too large EDMs which constitutes the SUSY CP problem [10]. The problem is resolved in certain classes of supersymmetric models, of which we choose four representative types. These include SUSY models with

¹The EDM collaboration has recently announced an improved (preliminary) neutron EDM bound $|d_n| \lesssim 3 \times 10^{-26} \text{ e cm}$ [4].

- small CP phases
- heavy spectrum
- decoupling
- split SUSY

To determine the source of EDMs, we study correlations between the neutron and electron electric dipole moments. In particular, if the neutron EDM is due to the QCD θ -term [11]–[13], one expects very small leptonic EDMs. On the other hand, in supersymmetry both the hadronic and leptonic EDMs are enhanced and there exist certain correlation patterns. Determination of such patterns is the main subject of this work.

The relevant low energy lagrangian describing interactions of electrons, quarks, gluons and photons is given by

$$\mathcal{L} = \theta \frac{\alpha_s}{8\pi} G\tilde{G} + \frac{1}{3}w f^{abc}G^a\tilde{G}^bG^c - \frac{i}{2}d_f \bar{f}(F\sigma)\gamma_5 f - \frac{i}{2}g_s\tilde{d}_q \bar{q}(G\sigma)\gamma_5 q ,$$

where w , d_f , \tilde{d}_q are the Weinberg operator [14] coefficient, the fermion EDM, and the quark chromo-EDM (CEDM), respectively. Here F and G represent the photon and gluon field strengths, respectively. The electron EDM is given by d_e , whereas the neutron EDM is a model-dependent function of θ , w , d_q and \tilde{d}_q . We will use the naive dimensional analysis (NDA) approach [15],²

$$d_n^{\text{NDA}} \sim 2d_d - 0.5d_u + e(0.4\tilde{d}_d - 0.1\tilde{d}_u) + 0.3 \text{ GeV} \times ew , \tag{1.2}$$

where we have set $\theta = 0$. Here the Wilson coefficients d_q, \tilde{d}_q, w are evaluated at the electroweak scale. The chromo-EDM and the Weinberg operator contributions involve considerable uncertainties. Furthermore, the result is sensitive to the quark masses which we choose as $m_u(M_Z) = 2 \text{ MeV}$ and $m_d(M_Z) = 4 \text{ MeV}$. The approach based on QCD sum rules gives a somewhat similar result [17],

$$d_n^{\text{SR}} \simeq 2d_d - 0.5d_u + e(\tilde{d}_d + 0.5\tilde{d}_u) + 0.1 \text{ GeV} \times ew . \tag{1.3}$$

In the Weinberg operator contribution, we have used the (model-dependent) renormalization factor of ref. [15]. In practice, the two approaches usually agree within a factor of 2 (unless w dominates) which suffices for our purposes. We note that there are also neutron models which include the strange quark contribution [18]. This effect is difficult to estimate and involves large uncertainties. We defer a study of such models until a subsequent publication.

²The QCD correction factor for the $d_{u,d}$ contributions has recently been recalculated [16] and found to be a factor of 2 or so smaller than that in [15]

2. EDMs in the standard model

We start by considering EDMs induced by the QCD θ -term. This is the primary source of hadronic EDMs in the Standard Model [11]–[13, 19]. In particular [13],

$$d_n \simeq 3 \times 10^{-16} \theta e \text{ cm} \tag{2.1}$$

with about 50% uncertainty. On the other hand, the electroweak contributions are very small, $d_n \sim 10^{-32} e \text{ cm}$ [20]. To satisfy the experimental bound (1.1), the θ parameter has to be tiny, $\leq \mathcal{O}(10^{-10})$. Such a small value can hardly be explained by the Standard Model, which constitutes the “strong CP problem”. The most popular solution to this problem in extensions of the SM invokes an anomalous Peccei-Quinn symmetry [21], which sets θ to zero. However, this symmetry is expected to be broken by higher dimensional operators generated at the Planck scale [22], such that the resulting θ is finite but small. There are also alternative solutions to the strong CP problem which employ other symmetries [23]. In these models, a finite θ can be induced by radiative corrections. In either case, a small neutron EDM is expected but it is difficult to make any quantitative prediction.

The θ -term also induces nuclear EDMs. In particular, it generates EDMs of the deuteron and the mercury atom. The former receives contributions from both the constituent nucleons and nucleon interactions, and can be evaluated via QCD sum rules with about 50% uncertainty [24]. An EDM of the mercury atom is induced by the Schiff moment which appears due to CP violating isoscalar and isovector pion-nucleon couplings. Despite recent progress in evaluating these contributions [25], there is still an order of magnitude uncertainty in $d_{\text{Hg}}(\theta)$, whereas d_{Hg} induced by the quark (colour-) EDM contributions is understood much better. The result is

$$\begin{aligned} d_D &\simeq -1 \times 10^{-16} \theta e \text{ cm} , \\ |d_{\text{Hg}}| &\sim \mathcal{O}(10^{-18} - 10^{-19}) \theta e \text{ cm}. \end{aligned} \tag{2.2}$$

The above formulae together with suppressed leptonic EDMs provide a correlation pattern for the θ -induced electric dipole moments.

It is important to remember that d_e is not measured directly, but instead is derived from atomic EDMs. In particular, the current EEDM bound is due to the thallium atom EDM measurement [2] and the relation $d_{\text{Tl}} \simeq -585 d_e$. If the θ -term is non-zero, this relation is altered. Although no reliable calculation of $d_{\text{Tl}}(\theta)$ is available, rough estimates [26] of the valence proton EDM contribution give $d_{\text{Tl}} \sim 10^{-20} \theta e \text{ cm}$. On the other hand, the SM electroweak interactions (including Majorana neutrinos) usually induce d_e of order $10^{-38} e \text{ cm}$ [27]. Thus, in the Standard Model with $\theta > 10^{-15}$, d_{Tl} is dominated by the θ -background and is less sensitive to d_e . This illustrates that the effects of the θ -term can be important for atomic systems and should be taken into account.

3. EDMs in supersymmetry

In supersymmetric models, there are additional sources of CP violation associated with complex phases in the SUSY breaking F-terms and, in addition, flavour misalignment

between the Yukawa matrices and the soft breaking terms [28]. EDMs are generated already at the one loop level and typically exceed the experimental bounds by orders of magnitude [10]. This constitutes the SUSY CP problem. The problem is alleviated in certain classes of supersymmetric models. These include models with small SUSY CP phases, those with a heavy SUSY spectrum, the decoupling scenario and split SUSY. In what follows, we study neutron-electron EDM correlations in these types of models.

We note that there are also other possibilities for EDM suppression. For instance, CP violation may have flavour off-diagonal nature due to some symmetry [29, 30]. We defer a study of this option until a subsequent publication. Finally, EDMs may be suppressed due to accidental cancellations among independent terms [31]. This option is however disfavoured by the mercury EDM constraint [32]–[35].

In our numerical analysis, we choose two representative values of $\tan\beta$, 5 and 35, and analyze the NEDM-EEDM correlations separately for SUSY CP violation induced by the phase of the μ -term, ϕ_μ , and CP violation due to the phase of the trilinear A-terms, ϕ_A . We scan over the parameter space of a given model and present our results as scatter plots d_e vs d_n .

3.1 Small CP phases

First, we study the minimal SUGRA model (mSUGRA) with small (10^{-2}) CP phases. The smallness of the physical phases may be due to approximate alignment between the phases of the soft terms as occurs in the dilaton-domination scenario [36] with the Giudice-Masiero mechanism for the μ -term, or due to some approximate symmetry [37].

The model is defined in terms of the following GUT scale parameters,

$$m_0, m_{1/2}, |A|, \tan\beta \tag{3.1}$$

and CP-phases ϕ_A, ϕ_μ . Here m_0 is the universal scalar mass, $m_{1/2}$ is the universal gaugino mass and A is the trilinear parameter. The Higgs potential parameters $|\mu|$ and $|B\mu|$ are found by imposing radiative electroweak symmetry breaking. We assume that the CP phases associated with the gaugino mass and the $B\mu$ -term have been rotated away by appropriate $U(1)_R$ and $U(1)_{PQ}$ transformations. In this case, the physical CP phases are parametrized by ϕ_A and ϕ_μ .

The (C)EDMs are dominated by one-loop diagrams involving gluinos, charginos and neutralinos, figure 1,

$$\begin{aligned} d_q &= d_q^{\tilde{g}} + d_q^{\tilde{\chi}^+} + d_q^{\tilde{\chi}^0}, \\ \tilde{d}_q &= \tilde{d}_q^{\tilde{g}} + \tilde{d}_q^{\tilde{\chi}^+} + \tilde{d}_q^{\tilde{\chi}^0}, \\ d_e &= d_e^{\tilde{\chi}^+} + d_e^{\tilde{\chi}^0}. \end{aligned} \tag{3.2}$$




Figure 1: One loop SUSY contributions to fermion (C)EDMs.

The two loop contributions are considerably smaller. The relevant formulae can be found, for example, in Ibrahim and Nath, ref. [31].

To get a feeling for the size of the EDMs, let us consider a simple approximation $\phi_\mu \sim \phi_A \equiv \phi \ll 1$, $\tan \beta \sim 3$ and assume a single mass scale M for the SUSY parameters (at the electroweak scale). Then,

$$\begin{aligned} d_n &\sim \left(\frac{300 \text{ GeV}}{M}\right)^2 \sin \phi \times 10^{-24} e \text{ cm} , \\ d_e &\sim \left(\frac{300 \text{ GeV}}{M}\right)^2 \sin \phi \times 10^{-25} e \text{ cm} . \end{aligned} \tag{3.3}$$

Both d_n and d_e grow linearly with $\tan \beta$. Clearly, for the SUSY spectrum with electroweak masses, the CP phase has to be of order 10^{-2} . We see that the neutron and electron EDMs differ by about an order of magnitude, however no prediction of their magnitudes can be made since the result depends on how small the phase is. If no d_e at the level $10^{-30} e \text{ cm}$ is found, the CP phase will have to be smaller than 10^{-5} which appears highly unnatural. It would be rather difficult to engineer a robust mechanism which would force the phases in the lagrangian to align with such an extraordinary precision. Thus, one may argue that d_e and d_n in this scenario should not be far below the current experimental limits.

In the same simple approximation, the mercury atom EDM is given by $d_{\text{Hg}} \sim (300 \text{ GeV}/M)^2 \sin \phi \times 10^{-26} e \text{ cm}$. Here d_{Hg} is dominated by the quark CEDMs [32] and we have included an extra factor $\sim 1/4$ due to a recent reevaluation of the nuclear/atomic matrix elements. The deuteron EDM is expected to be similar to the neutron EDM, $d_D \sim d_n$.

Let now turn to our numerical results presented in figures 6, 7. In these plots, we vary $m_0, m_{1/2}, |A|$ randomly in the range $200 \text{ GeV} - 1 \text{ TeV}$ and the phases ϕ_A, ϕ_μ in the range $[-\pi/500, \pi/500]$. In the left plot, ϕ_A is set to zero and, in the right plot, $\phi_\mu = 0$. For non-zero ϕ_μ , there is a clear linear d_e - d_n correlation and d_e is about an order of magnitude below d_n , as expected. d_e is dominated by the chargino diagram, whereas d_n receives comparable contributions from the charginos and gluinos. At higher $\tan \beta$, the EDMs increase linearly. For $\phi_A \neq 0$, the correlation is less pronounced. For a given d_n , the spread of d_e values is about two orders of magnitude. This is because d_n is dominated by the gluino diagrams, whereas d_e is dominated by the neutralino contributions. The former are relatively insensitive to m_0 since the squark masses are dominated by the gluino RG contributions, whereas the latter are sensitive to m_0 through both the slepton masses and the μ -parameter. Thus, fixing d_e does not determine d_n accurately. Note that EDMs induced by ϕ_A do not receive $\tan \beta$ -enhancement.

3.2 Heavy SUSY spectrum

The SUSY contributions to EDMs are suppressed if the entire SUSY spectrum is in the TeV range (eq. (3.3)). Such a possibility is motivated by the strong bound on the Higgs mass which requires the stop mass to be of order 1 TeV. If all SUSY masses are controlled by the same scale, the spectrum is heavy. This scenario can be motivated in various ways, see e.g. ref. [38].

In this class of models, the EDMs are usually dominated by the one loop diagrams of figure 1 and the analysis is very similar to that presented in the previous subsection.




Figure 2: 2 loop EDM contributions. The diagram on the left induces the Weinberg operator. On the right, is an example of the Barr-Zee type diagram (a is a pseudoscalar Higgs).

We note that if the d_e experiments with the $10^{-30} e$ cm sensitivity yield a null result, it would imply that the scale of SUSY masses is 100 TeV (eq. (3.3)). Models with such a high SUSY breaking scale are disfavoured by the gauge coupling unification and naturalness considerations. Thus, again one expects a non-zero result in the next round of EDM experiments.

In our numerical analysis, we study the mSUGRA model (eq. (3.1)) with $m_0, m_{1/2}, |A|$ in the range 2 TeV – 10 TeV and ϕ_A, ϕ_μ in the range $[-\pi, \pi]$. Our results are shown in figures 8, 9. The d_n-d_e correlation patterns are very similar to those in the small phase scenario, namely there is a well defined correlation in the $\phi_\mu \neq 0$ case, whereas for $\phi_A \neq 0$ it is far less pronounced.

3.3 Decoupling

The largest contributions to EDMs come from sfermions of the first two generations. If these are very heavy, > 10 TeV, the most dangerous contributions are suppressed [39]. The third generation is required to be light by naturalness and contributes to EDMs at the 2 loop level. This decoupling scenario can be realized, for instance, in certain types of GUT models where the hierarchy between the first two and third generation masses appears due to RG running [40].

The neutron EDM is dominated by the stop and sbottom contributions to the Weinberg operator [41], whereas the electron EDM is due to the Barr-Zee [42] type 2 loop diagrams [43], figure 2. This class of diagrams also includes graphs with internal charginos and h, H, A Higgs bosons [44]. Thus,

$$\begin{aligned} d_n &\simeq d_n(w) , \\ d_e &= d_e^{\text{Barr-Zee}(\tilde{f}, \tilde{\chi}^+, \tilde{\chi}^0)} . \end{aligned} \quad (3.4)$$

We note that the Barr-Zee type diagrams also contribute to the neutron EDM, but these are suppressed compared to the Weinberg operator [34]. The correlation between d_e and d_n is rather subtle since d_n depends on the gluino and the third generation squark masses, whereas d_e depends on the latter as well as the chargino and Higgs masses. An order of magnitude estimate of the resulting EDMs can be obtained by setting all SUSY masses (apart from the first two generation sfermions) to be $M \gg M_Z$ and the CP phases to be




Figure 3: d_n^{NDA} (left) and d_e (right) in the decoupling scenario, $\phi_\mu \in [\pi/10, \pi/2]$, $\tan\beta = 3$. The GUT scale parameters $A = m_{1/2} = m_{\tilde{q}_3}$ are in GeV and the EDMs are in e cm. The QCD sum rules model gives d_n^{SR} a factor of a few smaller.

given by a single quantity ϕ . Then, for moderate $\tan\beta$ [41]–[44],

$$\begin{aligned} d_n &\sim \left(\frac{300 \text{ GeV}}{M}\right)^2 \tan\beta \sin\phi \times 10^{-25} e \text{ cm} , \\ d_e &\sim \left(\frac{300 \text{ GeV}}{M}\right)^2 \tan\beta \sin\phi \times 10^{-27} e \text{ cm} . \end{aligned} \quad (3.5)$$

Here d_n and d_e differ by almost two orders of magnitude, whereas in other scenarios this hierarchy is usually a factor of ten.

In this approximation, the mercury EDM is less sensitive to the CP phase ϕ . This is because the Weinberg operator contribution to d_{Hg} is negligible and the EDM is generated by the subleading quark CEDMs. When M or ϕ in eq. (3.5) is adjusted to satisfy the current neutron EDM bound, the resulting d_{Hg} is $\leq \mathcal{O}(10^{-29}) e \text{ cm}$, which is likely to be below the reach of the mercury EDM experiments. The deuteron EDM, however, is again found to be similar to d_n .

In our numerical analysis, we use the following GUT input parameters

$$m_3, m_{1/2}, |A|, \tan\beta \quad (3.6)$$

and CP phases ϕ_A, ϕ_μ . Here m_3 is the mass parameter for all the scalars apart from those for the first two sfermion generations, which are assumed to be decoupled. As in the mSUGRA case, $|\mu|$ and $|B\mu|$ are determined by radiative electroweak symmetry breaking. We vary $m_3, m_{1/2}, |A|$ in the range 200 GeV – 1 TeV and ϕ_A, ϕ_μ in the range $[-\pi, \pi]$. Figure 3 shows typical ranges of the d_n and d_e values for order one CP phases³. Clearly, $d_n \sim 10^{-26} e \text{ cm}$ and $d_e \sim 10^{-27} - 10^{-28} e \text{ cm}$ are expected in this scenario. Smaller EDMs would imply that either the CP phases are small or that the SUSY spectrum is heavy, including the third generation sfermions.

The corresponding correlations are presented in figures 10, 11. For $\phi_\mu \neq 0$, d_e and d_n can differ by two or one order of magnitude, depending on the balance between the Weinberg operator and the chargino Barr-Zee contributions. At larger $\tan\beta$, the situation remains the same except the EDMs increase proportionally. Note that since we scan over

³ d_n and d_e are dominated by contributions sensitive to ϕ_μ .

CP phases in the region $[-\pi, \pi]$, we include the possibility that the CP phases are small such that smaller values of the EDMs compared to those in figure 3 are allowed. For $\phi_A \neq 0$, the hierarchy between d_n and d_e increases to three or four orders of magnitude. This is because d_n is dominated by the stop contribution to the Weinberg operator, whereas the leading chargino Barr-Zee contribution to d_e is now absent and d_e is due to the stop Barr-Zee diagram. At larger $\tan\beta$, this hierarchy reduces since unlike the Weinberg operator, Barr-Zee contributions receive $\tan\beta$ -enhancement.

3.4 Split SUSY

This is an extreme version of the decoupling scenario in which the third generation sfermions are decoupled as well and naturalness is abandoned [45, 46]. From the EDM perspective, it provides an interesting framework in which the neutron and electron EDMs are generated by the same type of diagrams and thus are highly correlated.

The EDMs are induced by a version of the Barr-Zee type diagram, figure 2, with the sfermion loop replaced by the chargino loop [44]. Since all the Higgses except for the SM-like h are assumed to be heavy, the chargino loop is attached to the fermion line through h and a photon [46], or W^+ and W^- [47].⁴ The relevant formulae can be found in refs. [44, 47]. Clearly, the contributing diagrams are the same for the electron and the neutron such that the EEDM and NEDM are strongly correlated. We have therefore,

$$\begin{aligned} d_q &= d_q^{\text{Barr-Zee}}(\tilde{\chi}^+, \tilde{\chi}^0), \\ d_e &= d_e^{\text{Barr-Zee}}(\tilde{\chi}^+, \tilde{\chi}^0). \end{aligned} \tag{3.7}$$

Simple estimates are obtained by setting the chargino mass scale M to be much larger than the Higgs mass ~ 100 GeV [46],

$$\begin{aligned} d_n &\sim \left(\frac{300 \text{ GeV}}{M}\right)^2 \frac{\sin\phi}{\tan\beta} \times 10^{-26} e \text{ cm}, \\ d_e &\sim \left(\frac{300 \text{ GeV}}{M}\right)^2 \frac{\sin\phi}{\tan\beta} \times 10^{-27} e \text{ cm}. \end{aligned} \tag{3.8}$$

Here d_n is obtained by rescaling d_e with the factor $m_q/m_e \sim 10$. The $\tan\beta$ suppression can be traced down to the fact that the EDMs are due to the chargino mass matrix rephasing invariant $M_{11}M_{22}M_{12}^*M_{21}^* \propto \sin\beta \cos\beta$ and the SM-like Higgs couplings have no $\tan\beta$ -enhancement.

We note that since no CEDMs are generated at this level, the mercury EDM is suppressed. The deuteron EDM is, as usual, of order d_n .

In our numerical analysis, we vary the GUT scale gaugino masses $M_{1,2,3}$ in the range 200 GeV – 1 TeV and ϕ_μ in the range $[-\pi, \pi]$. We set the EW scale μ -parameter by hand in the range 200 GeV – 1 TeV and also vary m_h in the range 100 GeV – 300 GeV. Typical d_n values are shown in figure 4, whereas the corresponding d_e is found by a simple rescaling.

⁴It has recently been shown [48] that similar diagrams mediated by h and Z are also significant for the neutron EDM. This does not affect our numerical estimates.




Figure 4: d_n in split SUSY. Here ϕ_μ, ϕ_{M_1} are varied in the range $[-\pi, \pi]$; $M_1, |\mu| \in [200 \text{ GeV}, 1 \text{ TeV}]$ and $m_h \in [100 \text{ GeV}, 300 \text{ GeV}]$.

For order one CP phases and low $\tan \beta$, d_n is between 10^{-26} and 10^{-27} e cm. Smaller values are obtained at large $\tan \beta$ or for small CP phases/heavy gauginos.

The d_n-d_e correlations are presented in figure 12. As expected, d_n is almost in one-to-one correspondence with d_e . The small spread in the EDM values comes from an interplay between the chargino and neutralino Barr-Zee diagrams, which contribute in different proportions to d_e and d_n . In the left figure, the broad tail of d_n below 10^{-28} e cm is an artifact of our numerical procedure and appears due to 1-loop contributions of heavy but not completely decoupled sfermions (we took $m_H, m_A = 10^5 \text{ GeV}$ and $m_{\text{sferm}} = 10^7 \text{ GeV}$).

4. Conclusions

We have analyzed correlations between the electron and the neutron EDMs in typical supersymmetric models. Unlike in the case of the θ -term induced electric dipole moments, in SUSY leptonic EDMs are enhanced and are likely to be observed in the next round of experiments.

Assuming that all SUSY CP phases are order one at the GUT scale, lower bounds on the EDMs can be obtained in the decoupling and the split SUSY scenarios:

$$\begin{aligned} \text{decoupling : } & d_e \sim (10^{-1} - 10^{-2}) d_n \gtrsim 10^{-28} \text{ e cm} , \\ \text{split SUSY : } & d_e \sim 10^{-1} d_n \gtrsim 10^{-29} \text{ e cm} , \end{aligned}$$

where the lower bound is saturated for a TeV range SUSY spectrum (apart from the first two generation sfermions) in the decoupling case, and at large $\tan \beta$ in the case of split SUSY.

For mSUGRA with small CP phases or a heavy spectrum, no solid lower bound can be derived. However, non-observation of d_e at the level 10^{-30} e cm would imply that either the CP phases are $\lesssim 10^{-5}$ or the scale of SUSY masses is $\gtrsim 100 \text{ TeV}$. Both of these options

make supersymmetric models very unappealing, so if low energy supersymmetry is indeed realized in nature, one expects $d_e \gtrsim 10^{-30} e \text{ cm}$.

Concerning the d_n-d_e correlations, the main feature of supersymmetry is that it enhances the leptonic EDMs and we observe the following correlations

$$\begin{aligned}
 \text{small phases :} & \quad d_e \sim 10^{-1} d_n , \\
 \text{heavy spectrum :} & \quad d_e \sim 10^{-1} d_n , \\
 \text{decoupling :} & \quad d_e \sim (10^{-1} - 10^{-2}) d_n , \\
 \text{split SUSY :} & \quad d_e \sim 10^{-1} d_n ,
 \end{aligned}$$

where we have assumed that *all* SUSY CP phases are of the same order of magnitude. These relations as well as correlations with d_D and d_{H_g} can help distinguish supersymmetry from other new physics models. We note that these correlations are quite stable to variations in the GUT scale soft breaking parameters. As long as d_e and d_n are generated by similar sets of diagrams, the most important factor for d_e/d_n is m_e/m_q and other effects being subleading.⁵

In this paper, we have focused on the low and moderate $\tan \beta$ regimes where the effects of four-fermion operators (figure 5) are less important [49]–[51] and have neglected certain two loop RG effects [52]. Also, we have not imposed other experimental constraints such as the bound on the Higgs mass, etc. which are expected to restrict the parameter space further. Clearly, such effects will not change the qualitative picture.

Finally, we note that in the decoupling and the small phases scenarios, at least part of the SUSY spectrum is light and can be observed at the LHC. On the other hand, if the CP phases are order one, the entire SUSY spectrum may lie in the multi-TeV range. Then direct discovery of the superpartners at the LHC may prove to be quite challenging, whereas the EDMs can still probe such a possibility.




Figure 5: An example of a four-fermion operator inducing EDMs of composite objects [50]. Here, a pseudoscalar (a) and scalar (b) Higgs exchange induces a CP odd contact interaction $\bar{d}d \bar{e}i\gamma_5 e$ contributing to d_{T1} .

Acknowledgments

We are grateful to P. Harris and M. Pospelov for helpful communications. We would also like to thank D. Doyle for collaboration at the early stage of this project.

⁵In particular, breaking the universality between the GUT scale squark and slepton masses does not affect our results.




Figure 6: d_e vs d_n in mSUGRA with small phases, $\tan\beta = 5$. Left: $\phi_\mu \neq 0$, right: $\phi_A \neq 0$.




Figure 7: As in figure 6, but for $\tan\beta = 35$.




Figure 8: d_e vs d_n in mSUGRA with a heavy spectrum, $\tan\beta = 5$. Left: $\phi_\mu \neq 0$, right: $\phi_A \neq 0$.




Figure 9: As in figure 8, but for $\tan\beta = 35$.




Figure 10: d_e vs d_n in the decoupling scenario, $\tan\beta = 5$. Left: $\phi_\mu \neq 0$, right: $\phi_A \neq 0$.




Figure 11: As in figure 10, but for $\tan\beta = 35$.




Figure 12: d_e vs d_n in split SUSY. Left: $\tan\beta = 5$, right: $\tan\beta = 35$.

References

- [1] P.G. Harris et al., *New experimental limit on the electric dipole moment of the neutron*, *Phys. Rev. Lett.* **82** (1999) 904.
- [2] B.C. Regan, E.D. Commins, C.J. Schmidt and D. DeMille, *New limit on the electron electric dipole moment*, *Phys. Rev. Lett.* **88** (2002) 071805.
- [3] M.V. Romalis, W.C. Griffith and E.N. Fortson, *A new limit on the permanent electric dipole moment of HG- 199*, *Phys. Rev. Lett.* **86** (2001) 2505 [[hep-ex/0012001](#)].
- [4] P. Harris, talk at SUSY'05, Durham, 18-23 July 2005, <http://susy-2005.dur.ac.uk/susy05programme.html>.
- [5] D. Kawall, F. Bay, S. Bickman, Y. Jiang and D. DeMille, *Proof of principle for a high sensitivity search for the electric dipole moment of the electron using the metastable $a(1)[^3\Sigma^+]$ state of pbo*, *Phys. Rev. Lett.* **92** (2004) 133007 [[hep-ex/0309079](#)].

- [6] J.J. Hudson, B.E. Sauer, M.R. Tarbutt and E.A. Hinds, *Measurement of the electron electric dipole moment using YBF molecules*, *Phys. Rev. Lett.* **89** (2002) 023003 [[hep-ex/0202014](#)].
- [7] C.Y. Liu and S.K. Lamoreaux, *A new search for a permanent dipole moment of the electron in a solid state system*, *Mod. Phys. Lett.* **A 19** (2004) 1235;
S.K. Lamoreaux, *Solid state systems for electron electric dipole moment and other fundamental measurements*, [nucl-ex/0109014](#).
- [8] M. van der Grinten, talk at Lepton-Moments, Cape Cod, 9-12 June 2003,
<http://g2pc1.bu.edu/~leptonmom/program.html>;
S. Lamoreaux, *ibid*.
- [9] EDM collaboration, Y.K. Semertzidis et al., *A new method for a sensitive deuteron EDM experiment*, *AIP Conf. Proc.* **698** (2004) 200–204 [[hep-ex/0308063](#)];
Y.K. Semertzidis et al., *Sensitive search for a permanent muon electric dipole moment*, [hep-ph/0012087](#).
- [10] J.R. Ellis, S. Ferrara and D.V. Nanopoulos, *CP-violation and supersymmetry*, *Phys. Lett.* **B 114** (1982) 231;
W. Buchmuller and D. Wyler, *CP-violation and r invariance in supersymmetric models of strong and electroweak interactions*, *Phys. Lett.* **B 121** (1983) 321;
J. Polchinski and M.B. Wise, *The electric dipole moment of the neutron in low-energy supergravity*, *Phys. Lett.* **B 125** (1983) 393.
- [11] V. Baluni, *CP-violating effects in QCD*, *Phys. Rev.* **D 19** (1979) 2227.
- [12] R.J. Crewther, P. Di Vecchia, G. Veneziano and E. Witten, *Chiral estimate of the electric dipole moment of the neutron in quantum chromodynamics*, *Phys. Lett.* **B 88** (1979) 123.
- [13] M. Pospelov and A. Ritz, *Theta induced electric dipole moment of the neutron via QCD sum rules*, *Phys. Rev. Lett.* **83** (1999) 2526 [[hep-ph/9904483](#)];
M. Pospelov and A. Ritz, *Electric dipole moments as probes of new physics*, *Ann. Phys. (NY)* **318** (2005) 119 [[hep-ph/0504231](#)].
- [14] S. Weinberg, *Larger Higgs exchange terms in the neutron electric dipole moment*, *Phys. Rev. Lett.* **63** (1989) 2333.
- [15] R. Arnowitt, J.L. Lopez and D.V. Nanopoulos, *Keeping the demon of SUSY at bay*, *Phys. Rev.* **D 42** (1990) 2423;
R. Arnowitt, M.J. Duff and K.S. Stelle, *Supersymmetry and the neutron electric dipole moment*, *Phys. Rev.* **D 43** (1991) 3085.
- [16] G. Degrassi, E. Franco, S. Marchetti and L. Silvestrini, *QCD corrections to the electric dipole moment of the neutron in the MSSM*, *JHEP* **0511** (2005) 044.
- [17] M. Pospelov and A. Ritz, *Neutron edm from electric and chromoelectric dipole moments of quarks*, *Phys. Rev.* **D 63** (2001) 073015 [[hep-ph/0010037](#)];
D.A. Demir, M. Pospelov and A. Ritz, *Hadronic edms, the Weinberg operator and light gluinos*, *Phys. Rev.* **D 67** (2003) 015007 [[hep-ph/0208257](#)].
- [18] J.R. Ellis and R.A. Flores, *Implications of the strange spin of the nucleon for the neutron electric dipole moment in supersymmetric theories*, *Phys. Lett.* **B 377** (1996) 83 [[hep-ph/9602211](#)];
J. Hisano and Y. Shimizu, *Hadronic edms induced by the strangeness and constraints on supersymmetric CP phases*, *Phys. Rev.* **D 70** (2004) 093001 [[hep-ph/0406091](#)].

- [19] F. Berruto, T. Blum, K. Orginos and A. Soni, *Neutron electric dipole moment with domain wall quarks*, *Nucl. Phys.* **140** (Proc. Suppl.) (2005) 411 [[hep-lat/0411003](#)];
E. Shintani et al., *Neutron electric dipole moment from lattice QCD*, *Phys. Rev. D* **72** (2005) 014504 [[hep-lat/0505022](#)].
- [20] E.P. Shabalin, *Electric dipole moment of quark in a gauge theory with left-handed currents*, *Sov. J. Nucl. Phys.* **28** (1978) 75;
M.B. Gavela et al., *CP-violation induced by penguin diagrams and the neutron electric dipole moment*, *Phys. Lett. B* **109** (1982) 215;
I.B. Khriplovich and A.R. Zhitnitsky, *What is the value of the neutron electric dipole moment in the Kobayashi-Maskawa model?*, *Phys. Lett. B* **109** (1982) 490.
- [21] R.D. Peccei and H.R. Quinn, *CP conservation in the presence of instantons*, *Phys. Rev. Lett.* **38** (1977) 1440.
- [22] S.M. Barr and D. Seckel, *Planck scale corrections to axion models*, *Phys. Rev. D* **46** (1992) 539;
R. Holman et al., *Solutions to the strong CP problem in a world with gravity*, *Phys. Lett. B* **282** (1992) 132 [[hep-ph/9203206](#)].
- [23] R.N. Mohapatra and G. Senjanovic, *Natural suppression of strong p and t noninvariance*, *Phys. Lett. B* **79** (1978) 283;
S.M. Barr, *Solving the strong CP problem without the Peccei-Quinn symmetry*, *Phys. Rev. Lett.* **53** (1984) 329;
A.E. Nelson, *Naturally weak CP-violation*, *Phys. Lett. B* **136** (1984) 387.
- [24] O. Lebedev, K.A. Olive, M. Pospelov and A. Ritz, *Probing CP-violation with the deuteron electric dipole moment*, *Phys. Rev. D* **70** (2004) 016003 [[hep-ph/0402023](#)].
- [25] V.F. Dmitriev and R.A. Sen'kov, *Phys. Atom. Nucl.* **66** (2003) 1940;
J.S.M. Ginges and V.V. Flambaum, *Violations of fundamental symmetries in atoms and tests of unification theories of elementary particles*, *Phys. Rept.* **397** (2004) 63 [[physics/0309054](#)].
- [26] I.B. Khriplovich and S.K. Lamoreaux, *CP violation without strangeness*, Springer, 1997.
- [27] J.P. Archambault, A. Czarnecki and M. Pospelov, *Electric dipole moments of leptons in the presence of Majorana neutrinos*, *Phys. Rev. D* **70** (2004) 073006 [[hep-ph/0406089](#)];
M.E. Pospelov and I.B. Khriplovich, *Electric dipole moment of the W boson and the electron in the Kobayashi-Maskawa model*, *Sov. J. Nucl. Phys.* **53** (1991) 638.
- [28] S. Abel, S. Khalil and O. Lebedev, *The string CP problem*, *Phys. Rev. Lett.* **89** (2002) 121601 [[hep-ph/0112260](#)].
- [29] K.S. Babu, B. Dutta and R.N. Mohapatra, *Solving the strong CP and the SUSY phase problems with parity symmetry*, *Phys. Rev. D* **65** (2002) 016005 [[hep-ph/0107100](#)].
- [30] S. Abel, D. Bailin, S. Khalil and O. Lebedev, *Flavour-dependent CP-violation and natural suppression of the electric dipole moments*, *Phys. Lett. B* **504** (2001) 241 [[hep-ph/0012145](#)];
J.L. Diaz-Cruz and J. Ferrandis, *Solving the SUSY CP problem with flavor breaking F-terms*, *Phys. Rev. D* **72** (2005) 035003 [[hep-ph/0504094](#)].
- [31] T. Falk and K.A. Olive, *Electric dipole moment constraints on phases in the constrained MSSM*, *Phys. Lett. B* **375** (1996) 196 [[hep-ph/9602299](#)];
T. Ibrahim and P. Nath, *The neutron and the electron electric dipole moment in $N = 1$ supergravity unification*, *Phys. Rev. D* **57** (1998) 478 [[hep-ph/9708456](#)];

- T. Ibrahim and P. Nath, *The neutron and the lepton edms in MSSM, large CP-violating phases and the cancellation mechanism*, *Phys. Rev. D* **58** (1998) 111301 [[hep-ph/9807501](#)];
M. Brhlik, G.J. Good and G.L. Kane, *Electric dipole moments do not require the CP-violating phases of supersymmetry to be small*, *Phys. Rev. D* **59** (1999) 115004 [[hep-ph/9810457](#)].
- [32] T. Falk, K.A. Olive, M. Pospelov and R. Roiban, *MSSM predictions for the electric dipole moment of the HG- 199 atom*, *Nucl. Phys. B* **560** (1999) 3 [[hep-ph/9904393](#)].
- [33] V.D. Barger et al., *CP-violating phases in SUSY, electric dipole moments and linear colliders*, *Phys. Rev. D* **64** (2001) 056007 [[hep-ph/0101106](#)].
- [34] S. Abel, S. Khalil and O. Lebedev, *Edm constraints in supersymmetric theories*, *Nucl. Phys. B* **606** (2001) 151 [[hep-ph/0103320](#)].
- [35] S. Abel, S. Khalil and O. Lebedev, *On the EDM cancellations in D-brane models*, *Phys. Rev. Lett.* **86** (2001) 5850 [[hep-ph/0103031](#)].
- [36] V.S. Kaplunovsky and J. Louis, *Model independent analysis of soft terms in effective supergravity and in string theory*, *Phys. Lett. B* **306** (1993) 269 [[hep-th/9303040](#)];
A. Brignole, L.E. Ibáñez and C. Muñoz, *Soft supersymmetry-breaking terms from supergravity and superstring models*, [hep-ph/9707209](#).
- [37] O. Lebedev and S. Morris, *Towards a realistic picture of CP-violation in heterotic string models*, *JHEP* **08** (2002) 007 [[hep-th/0203246](#)].
- [38] A. Falkowski, O. Lebedev and Y. Mambrini, *SUSY phenomenology of KKLT flux compactifications*, *JHEP* **11** (2005) 034 [[hep-ph/0507110](#)].
- [39] P. Nath, *CP-violation via electroweak gauginos and the electric dipole moment of the electron*, *Phys. Rev. Lett.* **66** (1991) 2565;
Y. Kizukuri and N. Oshimo, *Implications of the neutron electric dipole moment for supersymmetric models*, *Phys. Rev. D* **45** (1992) 1806.
- [40] J. Bagger, J.L. Feng and N. Polonsky, *Naturally heavy scalars in supersymmetric grand unified theories*, *Nucl. Phys. B* **563** (1999) 3 [[hep-ph/9905292](#)].
- [41] J. Dai, H. Dykstra, R.G. Leigh, S. Paban and D. Dicus, *CP-violation from three gluon operators in the supersymmetric standard model*, *Phys. Lett. B* **237** (1990) 216.
- [42] S.M. Barr and A. Zee, *Electric dipole moment of the electron and of the neutron*, *Phys. Rev. Lett.* **65** (1990) 21.
- [43] D. Chang, W.-Y. Keung and A. Pilaftsis, *New two-loop contribution to electric dipole moment in supersymmetric theories*, *Phys. Rev. Lett.* **82** (1999) 900 [[hep-ph/9811202](#)].
- [44] D. Chang, W.-F. Chang and W.-Y. Keung, *New constraint from electric dipole moments on chargino baryogenesis in MSSM*, *Phys. Rev. D* **66** (2002) 116008 [[hep-ph/0205084](#)];
A. Pilaftsis, *Higgs-mediated electric dipole moments in the MSSM: an application to baryogenesis and Higgs searches*, *Nucl. Phys. B* **644** (2002) 263 [[hep-ph/0207277](#)].
- [45] N. Arkani-Hamed and S. Dimopoulos, *Supersymmetric unification without low energy supersymmetry and signatures for fine-tuning at the LHC*, *JHEP* **06** (2005) 073 [[hep-th/0405159](#)].
- [46] N. Arkani-Hamed, S. Dimopoulos, G.F. Giudice and A. Romanino, *Aspects of split supersymmetry*, *Nucl. Phys. B* **709** (2005) 3 [[hep-ph/0409232](#)].

- [47] D. Chang, W.-F. Chang and W.-Y. Keung, *Electric dipole moment in the split supersymmetry models*, *Phys. Rev. D* **71** (2005) 076006 [[hep-ph/0503055](#)].
- [48] G. F. Giudice and A. Romanino, *Electric dipole moments in split supersymmetry*, [[hep-ph/0510197](#)].
- [49] S.M. Barr, *Measurable t and p odd electron - nucleon interactions from Higgs boson exchange*, *Phys. Rev. Lett.* **68** (1992) 1822.
- [50] O. Lebedev and M. Pospelov, *Electric dipole moments in the limit of heavy superpartners*, *Phys. Rev. Lett.* **89** (2002) 101801 [[hep-ph/0204359](#)].
- [51] D.A. Demir, O. Lebedev, K.A. Olive, M. Pospelov and A. Ritz, *Electric dipole moments in the MSSM at large $\tan\beta$* , *Nucl. Phys. B* **680** (2004) 339 [[hep-ph/0311314](#)].
- [52] K.A. Olive, M. Pospelov, A. Ritz and Y. Santoso, *CP-odd phase correlations and electric dipole moments*, *Phys. Rev. D* **72** (2005) 075001 [[hep-ph/0506106](#)].