

Jet quenching at finite 't Hooft coupling and chemical potential from AdS/CFT

Néstor Armesto^a, José Daniel Edelstein^{ab} and Javier Mas^a

^a*Departamento de Física de Partículas and IGFAE,
Universidade de Santiago de Compostela,
E-15782 Santiago de Compostela, Spain*

^b*Centro de Estudios Científicos (CECS), Casilla 1469,
Valdivia, Chile*

E-mail: nestor@fpaxp1.usc.es, edels@fpaxp1.usc.es, jamas@fpaxp1.usc.es

ABSTRACT: Following the nonperturbative prescription for the jet quenching parameter recently proposed by Liu, Rajagopal and Wiedemann, we compute the first correction in the inverse 't Hooft coupling corresponding to string α' corrections in the dual background. We also consider the introduction of a chemical potential for the $U(1)^3$ gauged R-symmetry. While the former mildly diminishes the jet quenching parameter –this suggesting a smooth interpolation between the strong coupling and perturbative results–, the latter generically increases its value. We comment on the extension of this setup to quarks of finite mass.

KEYWORDS: AdS-CFT Correspondence, Black Holes in String Theory, Jets, Thermal Field Theory.

Contents

1. Introduction and summary	1
2. The jet quenching parameter	3
3. \hat{q} at finite coupling	7
4. \hat{q} at finite chemical potential	8

1. Introduction and summary

The experimental program at the Relativistic Heavy Ion Collider (RHIC) in Brookhaven National Laboratory [1], has provided considerable insight into the properties of strongly interacting matter at high energy densities. Phenomenological analysis has established several striking features of such substance. First, the results on elliptic flow are well described by hydrodynamical models only if the shear viscosity is taken very low. The medium behaves like a strongly coupled plasma which resembles a liquid more than the gas of quasi-free partons long expected to be the state of matter at such energy densities, known as the Quark Gluon Plasma (QGP). As a second indication of this behavior, high energy partons traversing the medium are strongly quenched. This phenomenon is usually characterized, in models of medium-induced radiation, by the so-called quenching parameter (or transport coefficient) \hat{q} [2]. This parameter has the meaning of the average squared transverse momentum transferred from the medium to the traversing parton, per unit mean free path (see the reviews [3, 4]).

Phenomenological models differ in the detailed framework for calculating the radiative energy loss [2, 5], in the treatment of the geometry and dilution of the medium [6], as well as in the consideration of flow-induced radiation [7] and of additional elastic scattering [8]. The extracted values of the transport coefficient are $\hat{q} \sim 1 \div 15 \text{ GeV}^2/\text{fm}$, substantially larger than those found in studies of hadron production in DIS on nuclear targets, see e.g. [9]. While the lower bound is compatible with expectations from perturbative QCD [10], higher values demand additional non-perturbative mechanisms. Therefore, and while waiting for upcoming both experimental and phenomenological efforts, it is of uttermost importance to get further information on the possible values of \hat{q} in the strong coupling limit.

The traditional tool for such studies, namely lattice QCD, cannot be presently applied to determining the jet quenching parameter. In contrast, AdS/CFT duality [11] provides a powerful calculational framework where quantum properties of supersymmetric Yang-Mills theories at strong 't Hooft coupling $\lambda = g_{\text{YM}}^2 N_c$ and large number of colors N_c , are translated into classical computations in a gravitational background. The applications of

AdS/CFT techniques to thermal deformations of gauge theories started in [12]. There, the thermodynamics of the black brane geometry was conjectured to describe the behavior of the dual quantum field theory at the Hawking temperature of the black hole. In the limit of a flat horizon, the dual field theory lives in an unconfined phase at strong coupling. Placing probe charges in such a background corresponds to the insertion of sources for very massive quarks in the dual theory. Following this lore, a number of important results were derived concerning the $q\bar{q}$ potential, including features like confinement and screening both at zero and finite temperature [13]. They typically involve a Wilson line stretching either in a timelike or a spacelike direction; rotating Wilson lines were also examined as putative duals to high spin mesons. More recently, the study of thermodynamical properties was extended to encompass near equilibrium magnitudes [14]. A not minor surprise came out with the finding of a universal ratio between the shear viscosity and the entropy density, $\eta/s = 1/(4\pi)$ [15] for quantum field theories admitting a holographic dual description. This ratio was conjectured to set a universal bound on physical thermal field theories. The data at RHIC suggest that the values for the QGP are compatible with the lower bound, this strongly supporting the use of AdS/CFT to describe such a system.

Motivated by these successful applications of AdS/CFT to the study of strongly coupled phenomena in thermal gauge theories, Liu, Rajagopal and Wiedemann (LRW) recently proposed a scheme to determine the jet quenching parameter [16].¹ In their construction, it is central to use the identification [4] of this parameter with the coefficient in the exponent of an adjoint Wilson loop computed along a rectangular contour \mathcal{C} with a large distance L^- along the light-cone, and a spacelike separation L , $L \ll L^-$:

$$\langle W^A(\mathcal{C}) \rangle \equiv \exp \left[-\frac{1}{4} \hat{q} L^- L^2 \right]. \quad (1.1)$$

At large N_c this Wilson loop can be expressed in terms of the Wilson loop for the fundamental representation, $\langle W^A(\mathcal{C}) \rangle \simeq \langle W^F(\mathcal{C}) \rangle^2$. In turn, the AdS/CFT correspondence tells us that this fundamental Wilson loop can be computed [13] evaluating the classical Nambu-Goto action S for a string ending on the boundary along the previous contour,

$$\langle W^F(\mathcal{C}) \rangle = \exp[-S(\mathcal{C})]. \quad (1.2)$$

The result in [16], obtained in a near extremal D3 background corresponding to $\mathcal{N} = 4$ SUSY QCD at finite temperature, exhibited some interesting features.² The quenching parameter \hat{q} turned out to be proportional to T^3 (which of course provides the correct dimensions) and to $\sqrt{\lambda}$ (thus to $\sqrt{N_c}$), the latter being totally different to the *a priori*

¹After the initial proposal, a host of papers have appeared [17–21] which have shown that this result is not universal. Also the very interesting and related problem of a drag force on the brane has been addressed by several authors [22].

²Although the relation between the results in this framework and real QCD is unclear, the hope is that an understanding of the size and dependences of the jet quenching parameter could provide some kind of upper bound, while the perturbative QCD results should provide a lower bound. In this way, computing the results in less supersymmetric backgrounds might provide an indication (assuming a smooth behavior for the transition from $\mathcal{N} = 4$ SUSY QCD to real QCD) of the expected value for this parameter at strong coupling.

expected dependence on the number of degrees of freedom of the energy or entropy densities, hence $\propto N_c^2$. In this way, the quenching parameter appears not to be a direct measure of the energy density of the system, $\hat{q} \propto \epsilon^{3/4}$ as usually assumed [10], but of the third power of the temperature. Moreover, the numerical values turned out to be astonishingly close to the experimental data: for standard values $\alpha_s = \frac{1}{2}$ and $N_c = 3$, $\hat{q} = 0.94$ (3.16) GeV²/fm for $T = 200$ (300) MeV. Notice, however, that these quantities imply a ‘t Hooft coupling $\lambda = 6\pi$, while the gravity computation is strictly valid in the limit $\lambda \rightarrow \infty$. Thus, a more precise theoretical value of \hat{q} in the strong coupling limit neatly demands the understanding of finite ‘t Hooft coupling corrections to the result in [16]. This is one of the targets in this letter.

The paper is organized as follows. In the next section, trying to keep ourselves as generic as possible, we provide a formula for the jet quenching parameter that can be readily applied to a large class of metrics. As a quick example, we present the results for the thermal deformation of Witten’s D4-brane background [12]. We also comment on the fact that this formula admits a straightforward generalization to encompass quark sources of finite mass, and provide some preliminar numerical analysis. In section 3 we compute the first correction in the inverse ‘t Hooft coupling to the value given by LRW. We show that this correction mildly diminishes the jet quenching parameter. This suggests a smooth interpolation between the strong coupling regime and the perturbative results, in analogy with what has been observed for the free energy and the ratio η/s [23, 24]. In section 4, the effect of turning on chemical potentials³ is thoroughly investigated. The relevant metric corresponds to the background of a stack of rotating D3-branes with maximal number of angular momenta. We explore the evolution of the jet quenching parameter along the space of these three independent charges. We typically find an enhancement within the range of thermodynamical stability. We further compare with recent results which have appeared on the subject [19–21].

2. The jet quenching parameter

In this section we shall provide a formula that allows to readily compute \hat{q} in string theory backgrounds within the class of metrics that are suitable for an AdS/CFT duality, including the case in LRW. We will follow essentially the same steps as in [17]. The family of ten dimensional metrics of interest for us adopt the following form:

$$\begin{aligned}
 ds^2 &= G_{MN} dX^M dX^N \\
 &= -c_T^2 dt^2 + c_X^2 dx^i dx_i + c_R^2 dr^2 + G_{Mn} dX^M dX^n,
 \end{aligned}
 \tag{2.1}$$

where $X^M = (t, x^i, r; X^n)$, $i = 1, \dots, p$, $n = 1, \dots, 8 - p$. This class of metrics encompasses rotating backgrounds which we shall analyze later. We are interested in black brane solutions.

³These chemical potentials are conjugated to R -charge densities of $\mathcal{N} = 4$ SYM theory. They should not be confused with that corresponding to the baryon density in QCD whose implementation in the dual supergravity side is currently an open problem. Indeed, the baryonic charge is not dual to a $U(1)$ isometry of the supergravity background. We thank Krishna Rajagopal for stressing the importance of this point.

Following [16] we will consider the following lightlike Wilson line

$$x^- = \tau, \quad x^2 = \sigma, \quad r = r(\sigma), \quad (2.2)$$

with $x^\pm = (x^0 \pm x^1)/\sqrt{2}$, $\tau \in (0, L^-)$, $\sigma \in (-\frac{L}{2}, \frac{L}{2})$ and $L^- \gg L$ so that isometry along x^- direction holds approximately. Also we shall take a symmetric configuration around $\sigma = 0$, hence $r_0 = r(0)$ is an extremal point, $r'(0) = 0$. The induced metric reads as follows

$$g_{\tau\tau} = \frac{1}{2}(-c_T^2 + c_X^2), \quad g_{\sigma\sigma} = c_X^2 + c_R^2 r'(\sigma)^2. \quad (2.3)$$

From these expressions, the Nambu-Goto action takes the following form

$$S = \frac{L^-}{\sqrt{2\pi\alpha'}} \int_0^{L/2} d\sigma (c_X^2 - c_T^2)^{1/2} (c_X^2 + c_R^2 r'(\sigma)^2)^{1/2}. \quad (2.4)$$

The energy is a first integral of motion, from which the following equation for the profile $r(\sigma)$ can be extracted

$$r'(\sigma)^2 = \frac{c_X^2}{c_R^2} (k c_X^2 (c_X^2 - c_T^2) - 1), \quad (2.5)$$

where k is an integration constant. Let us assume that the r.h.s. of (2.5) does not vanish at any location $r \in (r_H, \infty)$ with r_H the location of the horizon, while, on the other hand, we assume $c_R(r_H) = \infty$. Then, it necessarily holds that $r_0 = r_H$, and the Wilson line extends symmetrically from $r = \infty$ down to r_H . With these assumptions, which we must verify case by case, the profile can be obtained from

$$\sigma(r) = \int_{r_H}^r \frac{c_R}{c_X} \frac{dr}{(k c_X^2 (c_X^2 - c_T^2) - 1)^{1/2}}. \quad (2.6)$$

In particular, the integration constant k is linked with L by the relation $\sigma(\infty) = \frac{L}{2}$. Going to a dimensionless radial coordinate $\rho = r/r_H$, this is

$$L = 2 r_H \int_1^\infty \frac{c_R}{c_X} \frac{d\rho}{(k c_X^2 (c_X^2 - c_T^2) - 1)^{1/2}}. \quad (2.7)$$

The prescription in LRW for \hat{q} calls for the leading behavior with L in the limit $LT \ll 1$. This is clearly related to the limit $k \rightarrow \infty$, i.e.,

$$L = \frac{2 r_H}{\sqrt{k}} \int_1^\infty \frac{c_R d\rho}{c_X^2 (c_X^2 - c_T^2)^{1/2}} + \mathcal{O}(k^{-3/2}). \quad (2.8)$$

Using (2.5) and (2.6), we can write the action as follows

$$S = \frac{r_H L^-}{\sqrt{2\pi\alpha'}} \int_1^\infty \frac{\sqrt{k} (c_X^2 - c_T^2) c_X c_R d\rho}{(k c_X^2 (c_X^2 - c_T^2) - 1)^{1/2}}. \quad (2.9)$$

We must still subtract the contribution corresponding to the self-energy of the quarks. This is given by the Nambu-Goto action for a pair of Wilson lines that stretch straight from the boundary to the horizon,

$$S_0 = \frac{r_H L^-}{\sqrt{2\pi\alpha'}} \int_1^\infty c_R (c_X^2 - c_T^2)^{1/2} d\rho. \quad (2.10)$$

To leading order in k^{-1} , taking into account that L is given by (2.8), $S_I = S - S_0$ reads [17]

$$S_I = \frac{L^-}{\sqrt{2\pi\alpha'}} \frac{L^2}{8r_H} \left(\int_1^\infty \frac{c_R d\rho}{c_X^2 (c_X^2 - c_T^2)^{1/2}} \right)^{-1}. \quad (2.11)$$

From here, let us extract an expression for the jet quenching parameter. We find it convenient to define

$$c_T^2(\rho) = \frac{1}{\Delta_R} \hat{c}_T^2(\rho), \quad c_X^2(\rho) = \frac{1}{\Delta_R} \hat{c}_X^2(\rho), \quad c_R^2(\rho) = \Delta_R \hat{c}_R^2(\rho), \quad (2.12)$$

where the dimensionless quantity Δ_R reads

$$\Delta_R = \left(\frac{(\alpha')^{5-p} \lambda}{r_H^{7-p}} \right)^{1/2}, \quad (2.13)$$

λ being the 't Hooft coupling in the $p + 1$ dimensional dual gauge theory. From these formulas, (1.1) and (1.2), we obtain

$$\hat{q} = \frac{1}{\sqrt{2\pi\lambda}} \left(\frac{r_H}{\alpha'} \right)^{6-p} \left(\int_1^\infty \frac{\hat{c}_R d\rho}{\hat{c}_X^2 (\hat{c}_X^2 - \hat{c}_T^2)^{1/2}} \right)^{-1}. \quad (2.14)$$

As it stands, this formula calls for a translation of r_H in terms of the field theoretical quantities. In the case of non-rotating backgrounds we can provide a more explicit solution. For this class of metrics the Hawking temperature is given by the standard formula

$$T = \frac{1}{4\pi} \frac{c_T'(r)}{\sqrt{c_T^2(r) c_R^2(r)}} \Big|_{r=r_H}. \quad (2.15)$$

Using this definition of the temperature, we can solve for r_H/α' as follows

$$\frac{r_H}{\alpha'} = \left[4\pi\sqrt{\lambda} T \left(\frac{\sqrt{\hat{c}_T^2(\rho) \hat{c}_R^2(\rho)}}{\hat{c}_T'(\rho)} \right) \Big|_{\rho=1} \right]^{\frac{2}{5-p}}, \quad (2.16)$$

and replace it in (2.14) to arrive at the formula

$$\hat{q} = \frac{1}{\sqrt{2\pi}} \left[16\pi^2 \left(\frac{\sqrt{\hat{c}_T^2(1) \hat{c}_R^2(1)}}{\hat{c}_T'(1)} \right)^2 \right]^{\frac{6-p}{5-p}} T^2 (T^2 \lambda)^{\frac{1}{5-p}} \left(\int_1^\infty \frac{\hat{c}_R d\rho}{\hat{c}_X^2 (\hat{c}_X^2 - \hat{c}_T^2)^{1/2}} \right)^{-1}. \quad (2.17)$$

This expression is invariant under reparameterization of the radial coordinate (upon suitable change of the integration limits). Indeed, the dependence on T and λ for generic p also coincides with the discussion in LRW. For example, as it stands, it can be directly used to extract the quenching parameter for the thermal deformation of Witten's D4-brane background. In this metric, the fifth dimension has been compactified to a circle of radius ℓ . Hence, the four dimensional effective coupling is $\tilde{\lambda} = \lambda/\ell \equiv 4\pi\alpha_{\text{SYM}} N_c$. Therefore we may write for the effective quenching parameter the following expression

$$\hat{q} \simeq 14.26 c T^3 \alpha_{\text{SYM}} N_c, \quad (2.18)$$




Figure 1: Evolution of \hat{q} with m for finite mass quarks. The solid curve represents the D3 background and the vertical axis is $\hat{q}/(\sqrt{\alpha_{\text{SYM}} N_c T^3})$ which asymptotes to $18.87\dots$ as in LRW. The dashed curve represents the D4 background. The vertical axis is now $\hat{q}/(c\alpha_{\text{SYM}} N_c T^3)$ which asymptotes to $14.26\dots$. The horizontal axis is $\rho_m = m\alpha'/r_H$. In each case, the quotient α'/r_H can be read off from (2.16).

where $c = \ell T$ is the ratio of radii of the thermal and Kaluza-Klein circles. The particular value $c = 1$ signals the confinement/deconfinement transition temperature $T = T_c = 1/\ell$ [25]. Therefore, strictly speaking, (2.18) is valid for $c \geq 1$, or $T \geq T_c$. The numerical factor in (2.18) is the result of (2.17) with $p = 4$ after inserting $\hat{c}_T^2 = \sqrt{8\pi}\rho^{3/2}(1 - \rho^{-3})$, $\hat{c}_X^2 = \sqrt{8\pi}\rho^{3/2}$ and $\hat{c}_R^2 = (8\pi)^{-1/2}\rho^{-3/2}(1 - \rho^{-3})^{-1}$ [26]. For standard values $\alpha_s = \frac{1}{2}$ and $N_c = 3$, we get $\hat{q} = 0.87(2.93) \text{ GeV}^2/\text{fm}$ for $c = 1$ and $T = 200(300) \text{ MeV}$. These values are just slightly smaller than those in LRW. Yet, the 5d origin of (2.18) is reflected in the linear dependence in the 't Hooft coupling. In the following sections we shall apply the expressions (2.14) and (2.17) to another couple of relevant backgrounds.

Another interesting use of the renormalized expression (2.17) is the possibility to extend the analysis to quark sources of finite mass. Indeed, from the point of view of the derivation, there is nothing peculiar about the integral upper limit being at the boundary, and it could equally well extend up to a finite value $\rho_m = r_m/r_H < \infty$. The physics then is dual to a geometrical setting in which the ends of the fundamental string are attached to a probe brane that is placed in the above background at a fixed distance from the stack, $r_m = m\alpha'$, set up by the mass of the quarks in the fundamental representation [27, 28]. It is quite evident from the analytic form of the formula (2.17) that cutting the value of the integral will decrease the denominator, and hence enhance the value of \hat{q} . Popular scenarios include cases where a probe D7 (alternatively a D6 or a D8) are placed inside D3 (respectively D4) backgrounds. Plotting \hat{q} as a function of the mass of the quarks (figure 1), we observe a very weak dependence until the mass is rather low, where the approximations are questionable. For example, for the case of the D3 background, the horizontal axis is $\rho_m = m/(2\pi^{3/2}\sqrt{\alpha_{\text{SYM}}N_c T}) \simeq m/2.7 \text{ GeV}$ for $\alpha_{\text{SYM}} = 1/2, N_c = 3$ and $T = 200 \text{ MeV}$. In

order to increase \hat{q} by a 10% we must lower ρ_m until a value of $\simeq 10$, hence $m \simeq 27$ GeV.

3. \hat{q} at finite coupling

The AdS/CFT correspondence is a statement that goes beyond the classical limit of string theory. In this limit, it maps classical solutions of supergravity to quantum field theory vacua in the strong coupling limit, $\lambda \rightarrow \infty$. Corrections in λ^{-1} are in direct correspondence with those in powers of α' in the string theory side.⁴ In this paper we shall use the solution given in [23, 30] corresponding to the α' corrected near extremal D3-brane. The relevant pieces of information of this solution can be casted as follows

$$\begin{aligned}\hat{c}_T^2(\rho) &= \rho^2(1 - \rho^{-4})(1 + \gamma T(\rho) + \dots), \\ \hat{c}_X^2(\rho) &= \rho^2(1 + \gamma X(\rho) + \dots), \\ \hat{c}_R^2(\rho) &= \rho^{-2}(1 - \rho^{-4})^{-1}(1 + \gamma R(\rho) + \dots),\end{aligned}$$

to first order in $\gamma = \frac{\zeta(3)}{8} (\alpha'/R^2)^3 \sim 0.15 \lambda^{-3/2}$. Here, following (2.13), we have already extracted $\Delta_R = R^2/r_H^2$ factors, with $R^2 = \sqrt{\lambda} \alpha'$. The intervening functions read

$$\begin{aligned}T(\rho) &= \left(-75\rho^{-4} - \frac{1225}{16}\rho^{-8} + \frac{695}{16}\rho^{-12}\right), \\ X(\rho) &= \left(-\frac{25}{16}\rho^{-8}(1 + \rho^{-4})\right), \\ R(\rho) &= \left(75\rho^{-4} + \frac{1175}{16}\rho^{-8} - \frac{4585}{16}\rho^{-12}\right).\end{aligned}\tag{3.1}$$

Inserting this data in (2.17) with $p = 3$ and expanding to first order in γ gives

$$\hat{q}(\lambda) = \hat{q}(0) \left(1 - \gamma \left[\frac{I}{2a} + 45\right] + \dots\right)\tag{3.2}$$

with $\hat{q}(0)$ as given in [16], and

$$a = \int_1^\infty \frac{d\rho}{\sqrt{\rho^4 - 1}} = \sqrt{\pi} \frac{\Gamma(5/4)}{\Gamma(3/4)},\tag{3.3}$$

$$I = \int_1^\infty \frac{R(\rho) - X(\rho)(\rho^4 + 2) + T(\rho)(\rho^4 - 1)}{\sqrt{\rho^4 - 1}} d\rho = -\frac{30725 \pi^{3/2}}{924\sqrt{2} \Gamma(3/4)^2}.\tag{3.4}$$

Evaluating yields

$$\hat{q}(\lambda) = \hat{q}(0) \left(1 - 1.7652 \lambda^{-3/2} + \dots\right).\tag{3.5}$$

Therefore, we see that finite coupling corrections tend to diminish the value of the quenching parameter. For example, taking $N_c = 3$ and $\alpha_{\text{SYM}} = 1/4, 1/2$ and 1 , we get a reduction factor of 6, 2 and 0.8 percent respectively. Note that the decrease in the jet quenching parameter towards weak coupling is suggestive of a smooth interpolation between the strong coupling regime and the perturbative results. Obviously, the computation of higher order corrections would be necessary to put this conclusion on more solid grounds.

⁴Corrections in α' to timelike or spacelike Wilson lines used to compute the $q\bar{q}$ potential have appeared in [29].

4. \hat{q} at finite chemical potential

The near horizon metric of a rotating black D3-brane with maximal number of angular momenta reads as follows [31, 32], with the conventions of [33]:

$$ds^2 = \sqrt{\Delta} \left(-\mathcal{H}^{-1} f dt^2 + f^{-1} dr^2 + \frac{r^2}{R^2} d\vec{x} \cdot d\vec{x} \right) + \frac{1}{\sqrt{\Delta}} \sum_{i=1}^3 R^2 H_i [d\nu_i^2 + \nu_i^2 (d\phi_i + A^i dt)^2], \quad (4.1)$$

where $\nu_1 = \cos \theta_1$, $\nu_2 = \sin \theta_1 \cos \theta_2$, $\nu_3 = \sin \theta_1 \sin \theta_2$, and $\mathcal{H} = H_1 H_2 H_3$, with

$$A^i = \frac{1}{R} \sqrt{\frac{\mu}{q_i}} (1 - H_i^{-1}), \quad H_i = 1 + \frac{q_i}{r^2}, \quad f = \frac{r^2}{R^2} \mathcal{H} - \frac{\mu}{r^2}, \quad \Delta = \mathcal{H} \sum_{i=1}^3 \frac{\nu_i^2}{H_i}. \quad (4.2)$$

Upon Kaluza-Klein reduction, this becomes a charged AdS black hole solution of $\mathcal{N} = 2$ $U(1)_R^3$ supergravity, where A^i plays the rôle of the gauge field. The holographic gauge theory corresponding to (4.1) is $\mathcal{N} = 4$ $SU(N)$ supersymmetric Yang-Mills at finite temperature and with a chemical potential for the $U(1)_R^3$ symmetry. It will be convenient to trade the nonextremality parameter μ for the horizon radius, $r = r_H$ given as the largest root of $f(r_H) = 0$, i.e.,

$$\mu = \frac{r_H^4}{R^2} \mathcal{H}(r_H), \quad (4.3)$$

and define the adimensional quantities

$$\kappa_i = \frac{q_i}{r_H^2}, \quad \Delta_R = \frac{R^2}{r_H^2}. \quad (4.4)$$

As usual, we go to dimensionless variable $\rho = r/r_H$ and find

$$H_i(\rho) = 1 + \kappa_i \rho^{-2}, \quad f(\rho) = \frac{1}{\Delta_R} (\rho^2 \mathcal{H}(\rho) - \rho^{-2} \mathcal{H}(1)) \equiv \frac{1}{\Delta_R} \hat{f}(\rho). \quad (4.5)$$

Finally, the relevant functions entering the formula (2.14) can be easily extracted from (4.1):

$$\hat{c}_T^2(\rho) = \frac{\sqrt{\Delta} \hat{f}}{\mathcal{H}} - \frac{1}{\sqrt{\Delta}} \sum_{i=1}^3 \frac{\nu_i^2 \mathcal{H}(1)}{\kappa_i H_i} (H_i - 1)^2, \quad \hat{c}_X^2(\rho) = \sqrt{\Delta} \rho^2, \quad \hat{c}_R^2(\rho) = \frac{\sqrt{\Delta}}{\hat{f}}. \quad (4.6)$$

The factors in the metric depend on the internal angles. However, the terms above conspire to give

$$\int_1^\infty \frac{\hat{c}_R d\rho}{\hat{c}_X^2 \sqrt{\hat{c}_X^2 - \hat{c}_T^2}} = \frac{1}{\mathcal{H}(1)} \int_1^\infty d\rho \left(\rho^4 \frac{\mathcal{H}(\rho)}{\mathcal{H}(1)} - 1 \right)^{-1/2}, \quad (4.7)$$

where all information about the internal angular coordinates has dissappeared. Now, given that the Hawking temperature of this solution is given by [31]

$$T = \frac{2 + \sum_{i=1}^3 \kappa_i - \prod_{i=1}^3 \kappa_i}{2\sqrt{\mathcal{H}(1)}} \frac{r_H}{\pi R^2} \quad (4.8)$$




Figure 2: Plot of \hat{q} as function of different combinations of charges κ_i . The charges which are not varied are set equal to zero. The stability bound chops the lower curves, but leaves the upper one unbounded. At the origin the value is $\simeq 18.87$ as found in LRW.

with $R^2 = \sqrt{\lambda} \alpha'$, substituting in (2.14) we find the answer

$$\hat{q}(\kappa_i) = \frac{\pi^2 T^3 \sqrt{\lambda}}{\sqrt{2}} \mathcal{H}(1) \left(\frac{2\sqrt{\mathcal{H}(1)}}{2 + \sum_{i=1}^3 \kappa_i - \prod_{i=1}^3 \kappa_i} \right)^3 \left(\int_1^\infty d\rho \left(\rho^4 \frac{\mathcal{H}(\rho)}{\mathcal{H}(1)} - 1 \right)^{-1/2} \right)^{-1}. \tag{4.9}$$

In order to analyze this result, it must be recalled that the domain of thermodynamical stability of this solution [34, 35] (see also [36]) is bounded by the inequality⁵ $\kappa_1 + \kappa_2 + \kappa_3 - \kappa_1 \kappa_2 \kappa_3 < 2$. Plotting the right hand side of (4.9) numerically, we find the curves that are shown in figure 2. The jet quenching parameter raises its value for nonzero charges. The increase is not monotonous along the whole space of charges. In fact, though hardly noticeable, it changes the sign of the slope along the line κ_1 with $\kappa_2 = \kappa_3 = 0$. To see this better, we have zoomed the interval $\kappa_1 \in [1.4, 2]$ in figure 3, where we can see that the change in slope sign happens around $\kappa_1 = 1.8$.

A comparison with recent results in the literature is in order. We certainly agree and go beyond the perturbative analysis of Cáceres and Guijosa [19]. Also, when restricted to the cases examined by Lin and Matsuo [20] (one charge) and Avramis and Sfetsos [21] (one or two equal charges), we find qualitative agreement within the range of stability. We may also examine less symmetrical configurations on a 3 dimensional plot in figure 4.

Besides analyzing the qualitative behaviour of $\hat{q}(\kappa_i)$ numerically, in order to compare with other approaches, it may be of interest to perform an expansion in powers of quantum field theoretical magnitudes. For this purpose, it is relevant to recall how the thermodynamical magnitudes are related to geometrical quantities. In particular, the density of

⁵A wrong sign in this expression in a previous version of this paper has led to wrong plots that we have corrected in this version.




Figure 3: Zoom of the curve $\hat{q}(\kappa_1)$ showing the value of the turning and end points.




Figure 4: Evolution of \hat{q} in a sector of the plane $(\kappa_1, \kappa_2 = \kappa_3)$. The curve is chopped by the thermodynamical stability bound.

physical charge and chemical potential are given respectively by [35, 37]

$$\rho_i = \frac{\pi N^2 T_0^3}{8} \sqrt{2\kappa_i} \prod_{i=1}^3 (1 + \kappa_i)^{1/2}, \tag{4.10}$$

$$\mu_i \equiv A^i(r)|_{r=r_H} = \frac{\pi T_0 \sqrt{2\kappa_i}}{1 + \kappa_i} \prod_{i=1}^3 (1 + \kappa_i)^{1/2}, \tag{4.11}$$

where $T_0 = r_H/(\pi R^2)$. From these expressions and (4.8), we should invert κ_i in terms of ρ_i and T for the canonical ensemble and in terms of μ_i and T for the grand canonical ensemble. This is difficult in the general case, so we may simplify for equal or vanishing values of κ_i . For example, taking $\kappa_1 = \kappa$ and $\kappa_2 = \kappa_3 = 0$ the two (inverse) expansions

yield

$$\kappa_C = \xi - \xi^2 + \frac{11}{4}\xi^3 + \dots, \tag{4.12}$$

$$\kappa_{GC} = \zeta + \zeta^2 + \frac{5}{4}\zeta^3 + \dots, \tag{4.13}$$

with

$$\xi = \left(\frac{4\sqrt{2}\rho}{\pi N^2 T^3} \right)^2, \quad \zeta = \left(\frac{\mu}{\sqrt{2}\pi T} \right)^2. \tag{4.14}$$

Expanding (4.9) in powers of κ and inserting these series we obtain for the canonical and grand-canonical ensemble respectively the following results

$$\hat{q}_C(\rho) = \hat{q}(0) (1 + 0.63 \xi - 1.08 \xi^2 + 2.83 \xi^3 + \dots), \tag{4.15}$$

$$\hat{q}_{GC}(\mu) = \hat{q}(0) (1 + 0.63 \zeta + 0.18 \zeta^2 + 0.06 \zeta^3 + \dots). \tag{4.16}$$

This expansion fully agrees with the one in [21] upon rescaling⁶ $\xi \rightarrow 2\xi^2$ and $\zeta \rightarrow 2\zeta^2$.

The computation of Wilson loops in thermal AdS backgrounds is a promising line of research, of which the \hat{q} computation is a salient example. The identification of other observables within the AdS/CFT framework (see, for a recent example, [38]) that can be confronted with experimental data is an urgent challenge. It seems clear to us that there are several avenues for further exploration. Among these, the extension of our results to less supersymmetric backgrounds is of neat interest. We hope to report on these issues in the near future.

Acknowledgments

We thank Alex Kovner, Krishna Rajagopal, Carlos Salgado and Urs Wiedemann for useful discussions. Special thanks go to Alfonso V. Ramallo who was involved in the first stages of this work, for his insightful comments and support. NA was supported by Ministerio de Educación y Ciencia of Spain under a contract Ramón y Cajal, by Xunta de Galicia under grant PGIDIT03PXIC20612PN, and by CICYT of Spain under project FPA2005-01963. JDE and JM were supported in part by MCyT, FEDER and Xunta de Galicia under grant FPA2005-00188 and by the EC Commission under grants HPRN-CT-2002-00325 and MRTN-CT-2004-005104. JDE was also supported by the FCT grant POCTI/FNU/38004/2001 and by Ministerio de Educación y Ciencia of Spain under a contract Ramón y Cajal. Institutional support to the Centro de Estudios Científicos (CECS) from Empresas CMPC is gratefully acknowledged. CECS is a Millennium Science Institute and is funded in part by grants from Fundación Andes and the Tinker Foundation.

⁶The rescaling by 2 in ζ is to be traced to the normalization of the gauge fields in [35], which is $\sqrt{2}$ larger than usual. We thank Spyros D. Avramis for pointing this and some calculational errors in a previous version.

References

- [1] PHENIX collaboration, K. Adcox et al., *Formation of dense partonic matter in relativistic nucleus nucleus collisions at rhic: experimental evaluation by the phenix collaboration*, *Nucl. Phys. A* **757** (2005) 184 [nucl-ex/0410003];
 B.B. Back et al., *The phobos perspective on discoveries at RHIC*, *Nucl. Phys. A* **757** (2005) 28 [nucl-ex/0410022];
 BRAHMS collaboration, I. Arsene et al., *Quark gluon plasma and color glass condensate at RHIC? the perspective from the BRAHMS experiment*, *Nucl. Phys. A* **757** (2005) 1 [nucl-ex/0410020];
 STAR collaboration, J. Adams et al., *Experimental and theoretical challenges in the search for the quark gluon plasma: the STAR collaboration's critical assessment of the evidence from RHIC collisions*, *Nucl. Phys. A* **757** (2005) 102 [nucl-ex/0501009].
- [2] R. Baier, Y.L. Dokshitzer, A.H. Mueller, S. Peigne and D. Schiff, *Radiative energy loss and p_T -broadening of high energy partons in nuclei*, *Nucl. Phys. B* **484** (1997) 265 [hep-ph/9608322].
- [3] R. Baier, D. Schiff and B.G. Zakharov, *Energy loss in perturbative QCD*, *Ann. Rev. Nucl. Part. Sci.* **50** (2000) 37 [hep-ph/0002198];
 M. Gyulassy, I. Vitev, X.-N. Wang and B.-W. Zhang, *Jet quenching and radiative energy loss in dense nuclear matter*, nucl-th/0302077.
- [4] A. Kovner and U.A. Wiedemann, *Gluon radiation and parton energy loss*, hep-ph/0304151.
- [5] B.G. Zakharov, *Radiative energy loss of high energy quarks in finite-size nuclear matter and quark-gluon plasma*, *JETP Lett.* **65** (1997) 615 [hep-ph/9704255];
 U.A. Wiedemann, *Gluon radiation off hard quarks in a nuclear environment: opacity expansion*, *Nucl. Phys. B* **588** (2000) 303 [hep-ph/0005129];
 M. Gyulassy, P. Levai and I. Vitev, *Reaction operator approach to non-Abelian energy loss*, *Nucl. Phys. B* **594** (2001) 371 [nucl-th/0006010];
 X.-N. Wang and X.-f. Guo, *Multiple parton scattering in nuclei: Parton energy loss*, *Nucl. Phys. A* **696** (2001) 788 [hep-ph/0102230];
 S. Jeon and G.D. Moore, *Energy loss of leading partons in a thermal QCD medium*, *Phys. Rev. D* **71** (2005) 034901 [hep-ph/0309332].
- [6] I. Vitev, *Testing the theory of QGP-induced energy loss at RHIC and the LHC*, hep-ph/0603010;
 K.J. Eskola, H. Honkanen, C.A. Salgado and U.A. Wiedemann, *The fragility of high- p_T hadron spectra as a hard probe*, *Nucl. Phys. A* **747** (2005) 511 [hep-ph/0406319];
 A. Dainese, C. Loizides and G. Paic, *Leading-particle suppression in high energy nucleus nucleus collisions*, *Eur. Phys. J. C* **38** (2005) 461 [hep-ph/0406201];
 M. Djordjevic, M. Gyulassy, R. Vogt and S. Wicks, *Influence of bottom quark jet quenching on single electron tomography of au + au*, *Phys. Lett. B* **632** (2006) 81 [nucl-th/0507019];
 N. Armesto, M. Cacciari, A. Dainese, C.A. Salgado and U.A. Wiedemann, *How sensitive are high- p_T electron spectra at RHIC to heavy quark energy loss?*, *Phys. Lett. B* **637** (2006) 362 [hep-ph/0511257].
- [7] N. Armesto, C.A. Salgado and U.A. Wiedemann, *Measuring the collective flow with jets*, *Phys. Rev. Lett.* **93** (2004) 242301 [hep-ph/0405301];
 T. Renk and J. Ruppert, *Flow dependence of high p_T parton energy loss in heavy-ion collisions*, *Phys. Rev. C* **72** (2005) 044901 [hep-ph/0507075].

- [8] A. Adil, M. Gyulassy, W.A. Horowitz and S. Wicks, *Collisional energy loss of non asymptotic jets in a QGP*, [nucl-th/0606010](#).
- [9] F. Arleo, *Quenching of hadron spectra in DIS on nuclear targets*, *Eur. Phys. J. C* **30** (2003) 213 [[hep-ph/0306235](#)];
- [10] R. Baier, *Jet quenching*, *Nucl. Phys. A* **715** (2003) 209 [[hep-ph/0209038](#)].
- [11] J.M. Maldacena, *The large- N limit of superconformal field theories and supergravity*, *Adv. Theor. Math. Phys.* **2** (1998) 231, and *Int. J. Theor. Phys.* **38** (1999) 1113 [[hep-th/9711200](#)].
- [12] E. Witten, *Anti-de Sitter space, thermal phase transition and confinement in gauge theories*, *Adv. Theor. Math. Phys.* **2** (1998) 505 [[hep-th/9803131](#)].
- [13] S.-J. Rey and J.-T. Yee, *Macroscopic strings as heavy quarks in large- N gauge theory and anti-de Sitter supergravity*, *Eur. Phys. J. C* **22** (2001) 379 [[hep-th/9803001](#)];
J.M. Maldacena, *Wilson loops in large- N field theories*, *Phys. Rev. Lett.* **80** (1998) 4859 [[hep-th/9803002](#)];
S.-J. Rey, S. Theisen and J.-T. Yee, *Wilson-Polyakov loop at finite temperature in large- N gauge theory and anti-de Sitter supergravity*, *Nucl. Phys. B* **527** (1998) 171 [[hep-th/9803135](#)].
- [14] G. Policastro, D.T. Son and A.O. Starinets, *From AdS/CFT correspondence to hydrodynamics*, *JHEP* **09** (2002) 043 [[hep-th/0205052](#)]; *From AdS/CFT correspondence to hydrodynamics. II: sound waves*, *JHEP* **12** (2002) 054 [[hep-th/0210220](#)].
- [15] G. Policastro, D.T. Son and A.O. Starinets, *The shear viscosity of strongly coupled $N = 4$ supersymmetric Yang-Mills plasma*, *Phys. Rev. Lett.* **87** (2001) 081601 [[hep-th/0104066](#)];
A. Buchel and J.T. Liu, *Universality of the shear viscosity in supergravity*, *Phys. Rev. Lett.* **93** (2004) 090602 [[hep-th/0311175](#)].
- [16] H. Liu, K. Rajagopal and U.A. Wiedemann, *Calculating the jet quenching parameter from AdS/CFT*, [hep-ph/0605178](#).
- [17] A. Buchel, *On jet quenching parameters in strongly coupled non-conformal gauge theories*, *Phys. Rev. D* **74** (2006) 046006 [[hep-th/0605178](#)].
- [18] J.F. Vazquez-Poritz, *Enhancing the jet quenching parameter from marginal deformations*, [hep-th/0605296](#).
- [19] E. Caceres and A. Guijosa, *On drag forces and jet quenching in strongly coupled plasmas*, [hep-th/0606134](#).
- [20] F.-L. Lin and T. Matsuo, *Jet quenching parameter in medium with chemical potential from AdS/CFT*, [hep-th/0606136](#).
- [21] S.D. Avramis and K. Sfetsos, *Supergravity and the jet quenching parameter in the presence of R -charge densities*, [hep-th/0606190](#).
- [22] C.P. Herzog, A. Karch, P. Kovtun, C. Kozcaz and L.G. Yaffe, *Energy loss of a heavy quark moving through $N = 4$ supersymmetric Yang-Mills plasma*, *JHEP* **07** (2006) 013 [[hep-th/0605158](#)];
J. Casalderrey-Solana and D. Teaney, *Heavy quark diffusion in strongly coupled $N = 4$ Yang-Mills*, [hep-ph/0605199](#);
S.S. Gubser, *Drag force in AdS/CFT*, [hep-th/0605182](#);

- C.P. Herzog, *Energy loss of heavy quarks from asymptotically AdS geometries*, hep-th/0605191;
- E. Caceres and A. Guijosa, *Drag force in charged $N = 4$ SYM plasma*, hep-th/0605235;
- J.J. Friess, S.S. Gubser and G. Michalogiorgakis, *Dissipation from a heavy quark moving through $N = 4$ super-Yang-Mills plasma*, hep-th/0605292;
- S.-J. Sin and I. Zahed, *Ampere's law and energy loss in AdS/CFT duality*, hep-ph/0606049.
- [23] S.S. Gubser, I.R. Klebanov and A.A. Tseytlin, *Coupling constant dependence in the thermodynamics of $N = 4$ supersymmetric Yang-Mills theory*, Nucl. Phys. **B 534** (1998) 202 [hep-th/9805156].
- [24] A. Buchel, J.T. Liu and A.O. Starinets, *Coupling constant dependence of the shear viscosity in $N = 4$ supersymmetric Yang-Mills theory*, Nucl. Phys. **B 707** (2005) 56 [hep-th/0406264].
- [25] M. Kruczenski, D. Mateos, R.C. Myers and D.J. Winters, *Towards a holographic dual of large- $N(c)$ QCD*, JHEP **05** (2004) 041 [hep-th/0311270].
- [26] N. Itzhaki, J.M. Maldacena, J. Sonnenschein and S. Yankielowicz, *Supergravity and the large- N limit of theories with sixteen supercharges*, Phys. Rev. **D 58** (1998) 046004 [hep-th/9802042].
- [27] A. Karch and E. Katz, *Adding flavor to AdS/CFT*, JHEP **06** (2002) 043 [hep-th/0205236].
- [28] M. Kruczenski, D. Mateos, R.C. Myers and D.J. Winters, *Meson spectroscopy in AdS/CFT with flavour*, JHEP **07** (2003) 049 [hep-th/0304032].
- [29] H. Dorn and H.J. Otto, *Q anti- q potential from AdS-CFT relation at $t \geq 0$: dependence on orientation in internal space and higher curvature corrections*, JHEP **09** (1998) 021 [hep-th/9807093];
- S. Förste, D. Ghoshal and S. Theisen, *Stringy corrections to the Wilson loop in $N = 4$ super Yang-Mills theory*, JHEP **08** (1999) 013 [hep-th/9903042];
- J. Greensite and P. Olesen, *Worldsheet fluctuations and the heavy quark potential in the AdS/CFT approach*, JHEP **04** (1999) 001 [hep-th/9901057].
- [30] J. Pawelczyk and S. Theisen, *$AdS_5 \times S^5$ black hole metric at $O(\alpha'^3)$* , JHEP **09** (1998) 010 [hep-th/9808126].
- [31] J.G. Russo and K. Sfetsos, *Rotating $D3$ branes and QCD in three dimensions*, Adv. Theor. Math. Phys. **3** (1999) 131 [hep-th/9901056].
- [32] M. Cvetič et al., *Embedding AdS black holes in ten and eleven dimensions*, Nucl. Phys. **B 558** (1999) 96 [hep-th/9903214].
- [33] A. Buchel, *Higher derivative corrections to near-extremal black holes in type-IIB supergravity*, Nucl. Phys. **B 750** (2006) 45 [hep-th/0604167].
- [34] M. Cvetič and S.S. Gubser, *Phases of r -charged black holes, spinning branes and strongly coupled gauge theories*, JHEP **04** (1999) 024 [hep-th/9902195].
- [35] D.T. Son and A.O. Starinets, *Hydrodynamics of r -charged black holes*, JHEP **03** (2006) 052 [hep-th/0601157].
- [36] R.-G. Cai and K.-S. Soh, *Critical behavior in the rotating D -branes*, Mod. Phys. Lett. **A 14** (1999) 1895 [hep-th/9812121];
- T. Harmark and N.A. Obers, *Thermodynamics of spinning branes and their dual field theories*, JHEP **01** (2000) 008 [hep-th/9910036].

- [37] J. Mas, *Shear viscosity from r-charged AdS black holes*, *JHEP* **03** (2006) 016 [[hep-th/0601144](#)].
- [38] K. Peeters, J. Sonnenschein and M. Zamaklar, *Holographic melting and related properties of mesons in a quark gluon plasma*, [hep-th/0606195](#).