

Semiclassical strings in Sasaki-Einstein manifolds and long operators in $\mathcal{N} = 1$ gauge theories

Sergio Benvenuti

*Scuola Normale Superiore
Pisa, Italy, and
INFN, Sezione di Pisa
Pisa, Italy
E-mail: sergio.benvenuti@sns.it*

Martin Kruczenski

*Department of Physics, Brandeis University
Waltham, MA 02454, U.S.A.
E-mail: martink@brandeis.edu*

ABSTRACT: We study the AdS/CFT relation between an infinite class of 5-d $\mathcal{Y}^{p,q}$ Sasaki-Einstein metrics and the corresponding quiver theories. The long BPS operators of the field theories are matched to massless geodesics in the geometries, providing a test of AdS/CFT for these cases. Certain small fluctuations (in the BMN sense) can also be successfully compared.

We then go further and find, using an appropriate limit, a reduced action, first order in time derivatives, which describes strings with large R-charge. In the field theory we consider holomorphic operators with large winding numbers around the quiver and find, interestingly, that, after certain simplifying assumptions, they can be described effectively as strings moving in a particular metric. Although not equal, the metric is similar to the one in the bulk. We find it encouraging that a string picture emerges directly from the field theory and discuss possible ways to improve the agreement.

KEYWORDS: Supersymmetric gauge theory, AdS-CFT Correspondence.

Contents

1. Introduction	1
2. Strings moving in the $\mathcal{Y}^{p,q}$ manifold	4
2.1 Massless geodesics	5
2.2 Reduced action for strings with large R-charge	8
3. The correspondence in the BPS sector	11
3.1 Chiral building blocks	12
3.2 The full mesonic chiral ring	15
3.3 BPS geodesics from the quivers	16
4. The field theory at $\lambda \rightarrow 0$	18
5. Effective action for the spin chain	20
5.1 Long paths in the $\mathcal{Y}^{3,2}$ quiver	21
5.2 Closed paths in $\mathcal{Y}^{p,q}$	25
6. More general operators	29
6.1 Protected building blocks	29
6.2 Near BPS massless geodesics	31
6.2.1 Near BPS massless geodesics from the geometries	32
7. Conclusions	33
A. Useful formulas	34

1. Introduction

The AdS/CFT correspondence [1] gave a precise example of the conjectured relation [2] between the large N limit of gauge theories and string theory. The most studied model is $\mathcal{N} = 4$ SYM, with gauge group $SU(N)$ and coupling g_{YM} . In its simplest form, the correspondence establishes that, in the large N -limit, keeping $\lambda = g_{\text{YM}}^2 N$ fixed, this theory is the same as free IIB strings on $AdS_5 \times S^5$, with N units of RR 5-form flux. The radius R of $AdS_5 \times S^5$ is given by $R/l_s = \lambda^{\frac{1}{4}}$ where l_s is the string length. The effective string tension is therefore $\lambda^{-\frac{1}{2}}$. In the limit $\lambda \rightarrow \infty$ the worldsheet theory becomes classical and can be easily studied. On the other hand, the field theory simplifies in the opposite limit, $\lambda \rightarrow 0$, since then it becomes perturbative. This makes the correspondence very powerful but at the same time difficult to study. In particular, it does not elucidate the point of how

a string description can emerge from a field theory (see [3] for a discussion). An important step in that direction was made in [4] where it was shown how this correspondence can be established for certain ultrarelativistic strings, i.e. strings whose kinetic energy is much larger than their mass. In a related paper [5], the string side of the calculation was understood from a semi-classical point of view which allowed a generalization of [4] to multispin string states in [6, 7] (see [8, 9] for a review and [10] for previous related work). The calculation can also be generalized thanks to the observation of [11] that the one-loop scalar dilatation operator can be interpreted as a Hamiltonian of an integrable $SO(6)$ spin chain. Using a Bethe ansatz method to solve a subsector of the spin chain, in [12, 13] a remarkable agreement was found between energies of various string solutions and eigenvalues of the dilatation operator representing dimensions of particular SYM operators. Moreover, integrable structures appear for certain rigid-shape rotating string configurations [14, 15] and can be mapped [16, 17] to the integrable structure of the spin chain.

Another step was made in [18], where it was shown that one can take the ultrarelativistic limit directly in the string action¹. The resulting, reduced action is a sigma model which turns out to be precisely the semiclassical coherent state action describing the field theory spin chain (in an $SU(2)$ subsector). This makes obvious how a sigma model description of operators can emerge from a field theory as an effective description of very long operators. These ideas can be cast also in the framework of integrable models as later shown in [22]. The results are also useful in understanding higher orders of the semiclassical approximation [23], other subsectors [24–31] including open strings [32], and also quantum corrections [33]. It could be useful also in understanding $1/N$ corrections as the ones discussed in [34].

Moreover, the relation between spin chains and gauge theory is quite generic and in fact it was already noted in QCD [35] implying that these ideas have wide applicability. Therefore, it is natural to wonder if they can be extended to other examples of the AdS/CFT correspondence, for example, with less than four supersymmetries. Examples with at least one supersymmetry are generically² given by IIB backgrounds of the form $AdS_5 \times X^5$, where X^5 is a five dimensional Sasaki-Einstein manifold. The dual superconformal theories are quivers, arising from the low energy excitations of D3-branes at Calabi–Yau singularities. Until recently, the only examples where the metric on X_5 was explicitly known were the homogeneous manifolds S^5 and $T^{1,1}$. The latter case, discussed in detail by Klebanov and Witten [39], gives the paradigmatic example of $\mathcal{N} = 1$ AdS/CFT.

However, one year ago, Gauntlett, Martelli, Sparks and Waldram found an infinite class of inhomogeneous Sasaki–Einstein metrics on $S^2 \times S^3$ [40–42] which are labeled by two integers $0 \leq q \leq p$, and are usually called $\mathcal{Y}^{p,q}$ metrics. The corresponding Calabi–Yau cones, are *toric*, meaning that there is an effectively acting $U(1)^3$ isometry. The toric description of the geometries was given in [43] which allowed [44] to find the superconformal gauge theories dual to Type IIB on $AdS_5 \times \mathcal{Y}^{p,q}$.

¹See [19, 20] for alternative approaches and [21] for a discussion of supersymmetry in the ultrarelativistic limit.

²This does not include the Maldacena–Nuñez solution [36] which uses a different approach. See [37, 38] for more generic solutions.

Once the superconformal field theories are known, it's possible to compute the anomalous dimensions of the chiral fields applying the general a -maximization technique of [45], which relies on general properties of supersymmetric theories [46] and works independently of AdS/CFT. These anomalous dimensions are directly related, in the supergravity dual, to the volume of the dual Sasaki–Einstein manifold, as well as the volumes of supersymmetric submanifolds. In fact, [47] later found the geometric analog of a -maximization, i.e. a general way of computing these volumes for toric Sasaki–Einstein manifolds in any dimension, bypassing the need of an explicit knowledge of the metric.

In a further development [48], a relation was pointed out between toric quivers and dimers, that leads [49] to a general method for obtaining the corresponding brane setups [50]. This 'brane tiling' technique is connected to a correspondence between the statistical mechanics of dimers and topological strings on Calabi-Yau's [51], and significantly generalizes previously known similar constructions [52, 53].

In the present paper similar periodic representations of quivers are considered, in particular, the full mesonic chiral ring of toric gauge theories is naturally encoded in one-cycles of the torus where the quiver itself is drawn.

Another generic feature of quivers associated to toric geometries is that they always admit an exactly marginal deformation analogous to the β -deformation of $\mathcal{N} = 4$ SYM [54], as was shown in [55], using the techniques of [54, 56]. This deformation leaves the toric $U(1)_F^2 \times U(1)_R$ isometry untouched. In [57] a very interesting way of constructing the gravity side of this kind of deformations has been found. The semiclassical sector of the correspondence [58] and integrability properties [59] have been studied.

The knowledge of a general class of geometries should allow the construction of non-conformal examples of the correspondence, in the spirit of the Klebanov-Strassler solution [60]. Progress in this direction was done in [61–65].

In this paper we are interested in improving the understanding of the correspondence in these new examples. We proceed in steps. First we establish a correspondence between massless geodesics and large R-charge chiral primary operators in the field theory. After that, following [4], we consider excited strings whose mass is small compared to their kinetic energy. When the string has few excitations (of certain types) we can find the corresponding operators in the field theory. For a large number of excitations, however, we need a way to obtain a spin chain description of the operators. This is difficult since the theory is not in a perturbative regime. We content ourselves with analyzing the mixing between operators induced by the superpotential and show that they lead to a sigma model action which has similar properties as an action that can be derived directly from the string side of the correspondence. The sigma models are not the same but we suggest that they should have the same infrared limit (in the worldsheet sense). So, we can argue that we indeed were able to find a string action emerging from the field theory. The mapping from one side of the correspondence to the other is that long paths in the quiver correspond to strings. If one draws the paths in a torus then the direction in which the path moves is directly related to the position of the string in the bulk.

The organization of this paper is as follows: In section 2 we analyze strings moving in the $\mathcal{Y}^{p,q}$ manifolds. We find massless geodesics paying special attention to the ones

corresponding to chiral primaries (long) operators in the quivers. We then consider semiclassical fast moving strings, in the limit of [18]. The Sasaki-Einstein geometry seen by these strings is naturally parameterized by a non relativistic effective action, that keeps all the information about the Sasaki-Einstein metric.

In section 3 the BPS sector of the field theory is analyzed. The full mesonic chiral ring is constructed for a general $\mathcal{Y}^{p,q}$ gauge theory, exploiting general features of toric quivers. We reobtain the results on BPS geodesics and find the natural ranges of the coordinates parameterizing the $\mathcal{Y}^{p,q}$ manifolds.

In order to reconstruct the full string background, it is necessary to go beyond the BPS sector and consider chiral non BPS operators. In section 4 we show the existence of a special point on the conformal manifold where some coefficients of the superpotential vanish. At that point the chiral ring is enhanced and a large class of holomorphic operators becomes BPS³. For long operators, non trivially, this class includes the extended semiclassical strings, as can be expected from the string side. The existence of a special point with enhanced chiral ring and symmetry (noticed also in [55] in some particular examples) is an exact result that we expect to be a general feature of superconformal quiver gauge theories and thus of AdS/CFT with critical string theory.

In section 5 we construct a spin chain Hamiltonian for the $\mathcal{Y}^{p,q}$ quivers, considering a simplified approach consisting in studying the mixings between chiral operators induced by the superpotential terms in the Lagrangian. In this way, we are able to reconstruct an $S^2 \times S^3$ geometry from the chiral semiclassical operators in the field theory. Even with the mentioned simplifications, the metric found is very similar to the original Sasaki-Einstein metric. More precisely, we find a Kähler metric on the base but the metric is not Einstein. We suggest that the metric may flow to an Einstein metric in the infrared of the world-sheet.

In section 6, which can be read independently of section 4 and 5, we extend the results of section 3 in a different direction. Instead of considering extended strings we consider non-BPS massless geodesics. We find a class of operators that we conjecture to correspond to a generic non BPS geodesic, and test the idea for massless strings moving along a small perturbation of a BPS geodesic. For short operators, this leads to a proposal for the, generically non protected, operators dual to all supergravity states, i.e. generic Kaluza-Klein harmonics on the transverse Einstein manifold.

Finally we give our conclusions in section 7.

2. Strings moving in the $\mathcal{Y}^{p,q}$ manifold

In this section we study semiclassical strings moving in the $AdS_5 \times \mathcal{Y}^{p,q}$ manifold whose metric is [41]:

$$ds^2 = -dt^2 \cosh^2 \rho + d\rho^2 + \sinh^2 \rho d\Omega_3^2 + ds_{p,q}^2 \tag{2.1}$$

³This is the class of operators that we call "holomorphic sector". In the case of $\mathcal{N} = 4$ this sector is the well known $SU(3)$ subsector. We don't discuss the closure of the sector in the gauge theories, but we expect that for long operators this sector becomes closed, as in $\mathcal{N} = 4$, *Minahan : 2004ds*.

$$ds_{p,q}^2 = \frac{1-y}{6} (d\theta^2 + \sin^2 \theta d\phi^2) + \frac{dy^2}{6p(y)} + \frac{q(y)}{9} (d\psi - \cos \theta d\phi)^2 \quad (2.2)$$

$$+ w(y) [d\alpha + f(y) (d\psi - \cos \theta d\phi)]^2 \quad (2.3)$$

with the functions

$$w(y) = 2 \frac{a-y^2}{1-y}, \quad q(y) = \frac{a-3y^2+2y^3}{a-y^2}, \quad f(y) = \frac{a-2y+y^2}{6(a-y^2)}, \quad (2.4)$$

and

$$p(y) = \frac{w(y)q(y)}{6} = \frac{a-3y^2+2y^3}{3(1-y)} \quad (2.5)$$

The coordinates span the range:

$$0 \leq \theta \leq \pi, \quad 0 \leq \phi \leq 2\pi, \quad 0 \leq \psi \leq 2\pi, \quad 0 \leq \alpha \leq 2\pi\ell, \quad y_1 \leq y \leq y_2 \quad (2.6)$$

The constant a appearing in the metric as well as the constants $y_{1,2}$ and ℓ which determine the range of variation of the coordinates can all be written in terms of the integers p and q that define the manifold:

$$y_{1,2} = \frac{1}{4p} \left(2p \mp 3q - \sqrt{4p^2 - 3q^2} \right) \quad (2.7)$$

$$\ell = \frac{q}{3q^2 - 2p^2 + p\sqrt{4p^2 - 3q^2}} \quad (2.8)$$

$$a = 3y_1^2 - 2y_1^3 \quad (2.9)$$

An important point is that $y_{1,2}$ are zeros of the function $p(y)$ appearing in the metric. It can be seen that, to avoid a conical singularity at $y = y_{1,2}$, we need $p'(y_{1,2}) = \pm 2y_{1,2}$ which is satisfied (choosing a minus sign). We note for further use that there is a third zero of $p(y)$ given by $y_3 = \frac{3}{2} - y_1 - y_2$. Various useful properties of these functions and the metric can be found in the original paper [41] and are collected in an appendix for completeness.

2.1 Massless geodesics

We consider massless geodesic in the reduced metric

$$ds^2 = -dt^2 + ds_{p,q}^2 = -dt^2 + g_{ab} dx^a dx^b \quad (2.10)$$

where $ds_{p,q}^2 = g_{ab} dx^a dx^b$ is the metric of the Sasaki-Einstein manifold and t is the global time in AdS_5 . The massless point-like string is sitting at $\rho = 0$ in the metric (2.1) and the motion is only in the internal manifold.

The action for the motion of a point-like string is

$$S = \frac{\sqrt{\lambda}}{2} \int d\tau \left(-\dot{t}^2 + g_{ij} \dot{x}^i \dot{x}^j \right) \quad (2.11)$$

where $\sqrt{\lambda} = (R/l_s)^2$ is the effective string tension. We include it for completeness but the results do not depend on the tension since the strings are point-like. We need to solve the equations of motion subject to the constraint

$$-\dot{t}^2 + g_{ab} \dot{x}^a \dot{x}^b = 0 \quad (2.12)$$

The equation of motion for t is solved by $t = \kappa\tau$ and therefore the action reduces

$$S = \int d\tau \mathcal{L} = \frac{\sqrt{\lambda}}{2} \int d\tau \left(g_{ab} \dot{x}^a \dot{x}^b \right) \quad (2.13)$$

$$= \frac{\sqrt{\lambda}}{2} \int d\tau \left\{ \frac{1-y}{6} \left(\dot{\theta}^2 + \sin^2 \theta \dot{\phi}^2 \right) + \frac{1}{w(y)q(y)} \dot{y}^2 + \frac{q(y)}{9} \left(\dot{\psi} - \cos \theta \dot{\phi} \right)^2 \right. \quad (2.14)$$

$$\left. + w(y) \left[\dot{\alpha} + f(y) \left(\dot{\psi} - \cos \theta \dot{\phi} \right) \right]^2 \right\} \quad (2.15)$$

namely free motion in the Sasaki-Einstein manifold. The momentum P_t conjugate to t is the energy of the string and therefore is equal to the conformal dimension Δ of the dual operator:

$$\Delta = P_t = \sqrt{\lambda} \kappa \quad (2.16)$$

We can also introduce the other conjugate momenta as

$$p_a = \frac{\partial \mathcal{L}}{\partial \dot{x}^a} \quad (2.17)$$

and the Hamiltonian which is given by

$$H = \frac{1}{2} g^{ab} p_a p_b \quad (2.18)$$

From the action we see immediately that the momenta P_ϕ , P_ψ and P_α are conserved quantities. This is a consequence of the $SU(2) \times U(1) \times U(1)$ isometry since P_ϕ is the third component of the $SU(2)$ angular momentum and P_ψ , P_α are associated to the $U(1)$ factors. There is a further conserved quantity corresponding to the total $SU(2)$ angular momentum given by:

$$J^2 = P_\theta^2 + \frac{1}{\sin^2 \theta} (P_\phi + \cos \theta P_\psi)^2 + P_\psi^2 \quad (2.19)$$

The momenta in terms of the velocities are given by

$$\frac{1}{\sqrt{\lambda}} P_y = \frac{1}{6p(y)} \dot{y} \quad (2.20)$$

$$\frac{1}{\sqrt{\lambda}} P_\theta = \frac{1-y}{6} \dot{\theta} \quad (2.21)$$

$$\frac{1}{\sqrt{\lambda}} (P_\phi + \cos \theta P_\psi) = \frac{1-y}{6} \sin^2 \theta \dot{\phi} \quad (2.22)$$

$$\frac{1}{\sqrt{\lambda}} (P_\psi - f(y) P_\alpha) = \frac{q(y)}{9} \left(\dot{\psi} - \cos \theta \dot{\phi} \right) \quad (2.23)$$

$$\frac{1}{\sqrt{\lambda}} P_\alpha = w(y) \left(\dot{\alpha} + f(y) \left(\dot{\psi} - \cos \theta \dot{\phi} \right) \right) \quad (2.24)$$

In terms of the momenta the Hamiltonian can be written as

$$\sqrt{\lambda} H = \frac{\lambda}{2} \kappa^2 = \frac{1}{2} \Delta^2 = \frac{1}{2} \left\{ 6p(y) P_y^2 + \frac{6}{1-y} (J^2 - P_\psi^2) + \frac{1-y}{2(a-y^2)} P_\alpha^2 \right. \quad (2.25)$$

$$\left. + \frac{9(a-y^2)}{a-3y^2+2y^3} \left(P_\psi - \frac{a-2y+y^2}{6(a-y^2)} P_\alpha \right)^2 \right\} \quad (2.26)$$

where we also used the constraint (2.12) to relate H to κ and further used (2.16) to relate κ and Δ . As expected the relation between the conformal dimension Δ and the momenta does not involve the tension $\sqrt{\lambda}$.

The only non trivial equation of motion we need to solve now is that of $y(\tau)$ which is simply a one dimensional motion in a potential as follows from the conservation of H and the fact that $P_y \propto \dot{y}/p(y)$. Before proceeding it is useful to introduce the R-charge:

$$Q_R = 2P_\psi - \frac{1}{3}P_\alpha \tag{2.27}$$

which gives, after some algebra,

$$\begin{aligned} \Delta^2 = & \left(\frac{3}{2}Q_R\right)^2 + \frac{1}{6p(y)}(P_\alpha + 3yQ_R)^2 \\ & + 6p(y)P_y^2 + \frac{6}{1-y}(J^2 - P_\psi^2) \end{aligned} \tag{2.28}$$

where we used the function $p(y)$ that was defined in (2.5). As we said this last equation should be understood as an equation of motion for $y(\tau)$.

The full set of geodesics moving only in the transverse SE manifold is completely described by eq. (2.28). We note that the set of geodesics on a five dimensional manifold is itself a manifold with eight dimensions; in the case of S^5 , for instance, this set is the manifold $SO(6)/(SO(2) \times SO(4))$. Since from (2.19) $J^2 \geq P_\psi^2$, all solutions have $\Delta \geq \frac{3}{2}Q_R$. We want now to restrict to solutions where this bound is saturated. These geodesics correspond to chiral primary, or BPS, operators that will be analyzed in the next section. From (2.28) it is clear than in order to have $\Delta = \frac{3}{2}Q_R$ we must require

$$P_y = 0, \quad J^2 = P_\psi^2, \tag{2.29}$$

The first equation implies $y = y_0$ is constant. The constant y_0 should be set to the minimum of Δ^2 , namely

$$y_0 = -\frac{P_\alpha}{3Q_R} \tag{2.30}$$

This ensures that the equation of motion for y is satisfied and at the same time implies $\Delta = \frac{3}{2}Q_R$. The restriction however is that, to obtain a geodesic, we need:

$$y_1 \leq -\frac{P_\alpha}{3Q_R} \leq y_2 \tag{2.31}$$

In this sense, $y_{1,2}$ can be thought as defining the range of variation of P_α/Q_R .

To summarize, for all these BPS geodesics we obtain:

$$P_\alpha = -3y_0Q_R, \quad J = \frac{1}{2}(1 - y_0)Q_R \tag{2.32}$$

and the y_0 independent relations

$$\Delta = \frac{3}{2}Q_R, \quad Q_R = 2J - \frac{1}{3}P_\alpha. \tag{2.33}$$

The last equality follows from the definition of Q_R , namely eq. (2.27), and the fact that $J = P_\psi$ for these geodesics. Together with the first relation in (2.32) it implies the second one, namely $J = \frac{1}{2}(1 - y_0)Q_R$.

Using the definitions of the momenta in terms of the velocities one can see that the geodesics in question are simply given by

$$y = y_0, \quad \theta = \theta_0, \quad \phi = \phi_0, \quad \dot{\alpha} + \frac{1}{6}\dot{\psi} = 0 \quad (2.34)$$

which suggests introducing a new angle β through

$$\beta = 6\alpha + \psi, \quad \tilde{\psi} = \psi \quad (2.35)$$

This implies

$$P_\beta = \frac{1}{6}P_\alpha \quad (2.36)$$

$$P_{\tilde{\psi}} = P_\psi - \frac{1}{6}P_\alpha = \frac{1}{2}Q_R \quad (2.37)$$

Now, the geodesics are such that $\dot{\beta} = 0$. Note also that $\tilde{\psi} = \psi$ is now conjugate to the R-charge.

Concluding, a four dimensional subset of geodesic is BPS, and corresponds, as expected, to point-like strings moving only along the R-charge direction, or $\tilde{\psi}$ direction. In section 3 we reconstruct these BPS geodesics from chiral primaries in the quivers. In section 6 we will study small deviations from the BPS case, corresponding to $\Delta > \frac{3}{2}Q_R$.

2.2 Reduced action for strings with large R-charge

In this section we consider classical strings which move with large angular momentum corresponding to field theory operators with large R-charge. Such strings move fast in the ψ direction as in the previous section but now we do not use the approximation that the string is small. Any five-dimensional Sasaki–Einstein metric can be written in the following form

$$ds^2 = -dt^2 + \frac{1}{6}g_{ij}dx^i dx^j + \frac{1}{9}(d\psi + A_i dx^i)^2 \quad (2.38)$$

where g_{ij} is a local Kähler–Einstein metric on the base, parameterized by the coordinates x_i . Both g_{ij} and A_i depend only on the four coordinates x_i . The external derivative of the one form $A_i dx^i$ is proportional to the Kähler form of the four dimensional base. It also completely specifies the Kähler form of the Calabi-Yau cone over the SE manifold.

Now we introduce a coordinate $\psi_1 = \psi - 3t$. The metric becomes

$$ds^2 = \frac{2}{3}dt(d\psi_1 + A_i dx^i) + \frac{1}{9}(d\psi_1 + A_i dx^i)^2 + \frac{1}{6}g_{ij}dx^i dx^j \quad (2.39)$$

If we choose $t = \kappa\tau$ we can write the Polyakov action

$$S = \frac{1}{2} \int \frac{2}{3} \kappa (\partial_\tau \psi_1 + A_i \partial_\tau x^i) + \frac{1}{9} (\partial_\tau \psi_1 + A_i \partial_\tau x^i)^2 + \frac{1}{6} g_{ij} \partial_\tau x^i \partial_\tau x^j \quad (2.40)$$

$$- \frac{1}{9} (\partial_\sigma \psi_1 + A_i \partial_\sigma x^i)^2 - \frac{1}{6} g_{ij} \partial_\sigma x^i \partial_\sigma x^j \quad (2.41)$$

and the conformal constraints

$$0 = \frac{1}{3}\kappa(\partial_\tau\psi_1 + A_i\partial_\tau x^i) + \frac{1}{9}(\partial_\tau\psi_1 + A_i\partial_\tau x^i)^2 + \frac{1}{6}g_{ij}\partial_\tau x^i\partial_\tau x^j \quad (2.42)$$

$$+ \frac{1}{9}(\partial_\sigma\psi_1 + A_i\partial_\sigma x^i)^2 + \frac{1}{6}g_{ij}\partial_\sigma x^i\partial_\sigma x^j \quad (2.43)$$

$$0 = \frac{2}{3}\kappa(\partial_\sigma\psi_1 + A_i\partial_\sigma x^i) + \frac{1}{9}(\partial_\tau\psi_1 + A_i\partial_\tau x^i)(\partial_\sigma\psi_1 + A_i\partial_\sigma x^i) + \frac{1}{6}g_{ij}\partial_\tau x^i\partial_\sigma x^j \quad (2.44)$$

In this system of coordinates the string moves slowly (which means it moves almost at the speed of light in the original ones). We therefore consider the limit [18]

$$\partial_\tau X \rightarrow 0, \quad \kappa \rightarrow \infty, \quad \kappa \partial_\tau X \text{ fixed} \quad (2.45)$$

where X denotes all coordinates, ψ_1 and x^i . What we did is to focus on strings that move very fast in the original coordinate ψ and therefore carry a large R -charge. The reason is that we want to match those strings with large R -charge field theory operators.

Going back to our problem, in that limit the conformal constraints reduce to

$$0 = \kappa(\partial_\tau\psi_1 + A_i\partial_\tau x^i) + \frac{1}{2}g_{ij}\partial_\sigma x^i\partial_\sigma x^j \quad (2.46)$$

$$0 = \partial_\sigma\psi_1 + A_i\partial_\sigma x^i \quad (2.47)$$

Taking the limit in the action and using the constraint (2.47) we get

$$S = \int \frac{1}{3}\kappa(\partial_\tau\psi_1 + A_i\partial_\tau x^i) - \frac{1}{12}g_{ij}\partial_\sigma x^i\partial_\sigma x^j \quad (2.48)$$

which is the final form of the reduced action describing strings with large R -charge.

We can now specify this general derivation to our case of interest, the $\mathcal{Y}^{p,q}$ metrics. Using the coordinates discussed at the end of the previous subsection, (θ, ϕ, y, β) , the local Kähler–Einstein metric g_{ij} and the $U(1)$ -fibration A_i are

$$g_{ij}dx^i dx^j = (1-y)(d\theta^2 + \sin^2\theta d\phi^2) + \frac{dy^2}{p(y)} + p(y)(d\beta - \cos\theta d\phi)^2 \quad (2.49)$$

$$A_i dx^i = -y d\beta - (1-y)\cos\theta d\phi \quad (2.50)$$

It should be noticed that the metric (2.49) is valid only locally, and has orbifold singularities at the zeros of $p(y)$.⁴ The constraint (2.47) becomes

$$\partial_\sigma\psi_1 - \cos\theta\partial_\sigma\phi = y(\partial_\sigma\beta - \cos\theta\partial_\sigma\phi) \quad (2.51)$$

and the effective action (2.48) takes the explicit form

$$S = \sqrt{\lambda} \int \frac{1}{3}\kappa(\partial_\tau\psi_1 - y\partial_\tau\beta - (1-y)\cos\theta\partial_\tau\phi) \quad (2.52)$$

$$- \frac{1}{12} \left[(1-y) \left((\partial_\sigma\theta)^2 + \sin^2\theta(\partial_\sigma\phi)^2 \right) + \frac{(\partial_\sigma y)^2}{p(y)} + p(y)(\partial_\sigma\beta - \cos\theta\partial_\sigma\phi)^2 \right] \quad (2.53)$$

⁴We note also that the one form $A_i dx^i$ does not depend on $p(y)$; as a consequence the Kähler forms of the four dimensional base and of the metric cone over the five-manifolds do not depend on the precise form of $p(y)$. In section 5 we derive a metric from the spin chain which differs from (2.49) only in the form of $p(y)$ and for A_i gives the same result as (2.50).

where we restored the factor $(R/l_s)^2 = \sqrt{\lambda}$ in front of the action. We can immediately identify the following conserved quantities

$$\mathcal{P}_{\psi_1} = \frac{1}{3}\sqrt{\lambda}\kappa \int d\sigma = \frac{2\pi}{3}\sqrt{\lambda}\kappa \quad (2.54)$$

$$\mathcal{P}_\beta = -\frac{1}{3}\sqrt{\lambda}\kappa \int d\sigma y \quad (2.55)$$

$$\mathcal{P}_\phi = -\frac{1}{3}\sqrt{\lambda}\kappa \int d\sigma (1-y) \cos \theta \quad (2.56)$$

$$\mathcal{H} = \frac{\sqrt{\lambda}}{12} \int d\sigma \left[(1-y) \left((\partial_\sigma \theta)^2 + \sin^2 \theta (\partial_\sigma \phi)^2 \right) + \right. \quad (2.57)$$

$$\left. \frac{(\partial_\sigma y)^2}{p(y)} + p(y) (\partial_\sigma \beta - \cos \theta \partial_\sigma \phi)^2 \right] \quad (2.58)$$

where \mathcal{P}_{ψ_1} is (half) the R-charge, \mathcal{P}_β , \mathcal{P}_ϕ , are the $U(1)_F$ and the third component of the $SU(2)$ charges respectively. Finally \mathcal{H} is the Hamiltonian which corresponds to $\Delta - \frac{3}{2}Q_R$ in the field theory. Furthermore, if we remember that $P_{\psi_1} = P_{\tilde{\psi}}$, we see that the relations (2.32), (2.33) are satisfied at each value of σ implying that each point of the string moves approximately along a BPS geodesic. As expected, \mathcal{H} vanishes precisely when all the four local coordinates do not depend on σ . In this case one recovers the results of the previous subsection. It is useful to note that if we use the coordinate $t = \tau/\kappa$ and replace κ by the R-charge $Q_R = \frac{1}{2}\mathcal{P}_{\psi_1}$ the reduced action can be written as

$$S = \frac{Q_R}{4\pi} \left\{ \int (\partial_t \psi_1 - y \partial_t \beta - (1-y) \cos \theta \partial_t \phi) \right. \quad (2.59)$$

$$\left. - \frac{4\pi^2}{9} \frac{\lambda}{Q_R^2} \left[(1-y) \left((\partial_\sigma \theta)^2 + \sin^2 \theta (\partial_\sigma \phi)^2 \right) + \frac{(\partial_\sigma y)^2}{p(y)} + p(y) (\partial_\sigma \beta - \cos \theta \partial_\sigma \phi)^2 \right] \right\}$$

We see that the corrections introduced by a sigma dependence of the coordinates, namely for an extended string, are small for large R-charge as expected. The result is valid for large λ but a naive extrapolation to $\lambda = 0$ suggests that the corrections vanish in that point, a fact that we use later.

We conclude this subsection by noting that one can write the reduced action (2.53) in the following general form:

$$S = -i\kappa \int \left(\dot{z}^a \partial_a K - \dot{\bar{z}}^{\bar{a}} \partial_{\bar{a}} K \right) - \frac{1}{2} \int \partial_{\bar{a}\bar{b}} K \partial_\sigma z^a \partial_\sigma \bar{z}^{\bar{b}} \quad (2.60)$$

where we introduced two complex variables $z^{a=1,2}$ and a Kähler potential $K(z^1 \bar{z}^{\bar{1}} + z^2 \bar{z}^{\bar{2}})$. In terms of the original variables they are:

$$z_1 = \sin\left(\frac{\theta}{2}\right) e^{-i\frac{1}{2}(\beta-\phi)} \prod_{i=1}^3 |y - y_i|^{\frac{1}{4y_i}} \quad (2.61)$$

$$z_2 = \cos\left(\frac{\theta}{2}\right) e^{-i\frac{1}{2}(\beta+\phi)} \prod_{i=1}^3 |y - y_i|^{\frac{1}{4y_i}} \quad (2.62)$$

$$(2.63)$$

where $y_1 < y_2 < y_3$ are the three roots of $p(y) = 0$ already introduced in (2.9). This relation defines complex coordinates only locally since for example the periodicity of β is not 2π . We see that $\rho = z^1 \bar{z}^1 + z^2 \bar{z}^2$ is a function of y only. This means that the (local) Kähler potential is also a function of y and turns out to be given by

$$K = \frac{1}{6} \sum_{i=1}^3 \frac{1-y_i}{y_i} \ln |y - y_i| \tag{2.64}$$

With these definitions it is easy to check that (2.60) is equivalent to (2.53). In doing so it is useful to note that

$$\frac{\partial K}{\partial y} = -\frac{(1-y)}{3p(y)} \tag{2.65}$$

$$\frac{\partial \rho}{\partial y} = -\frac{\rho}{p(y)} \tag{2.66}$$

The form of the action (2.60) means simply that the base of the Sasaki-Einstein manifold is locally Kähler with complex coordinates $z_{1,2}$ and Kähler potential K . The fact that K depends only on $\rho = z^1 \bar{z}^1 + z^2 \bar{z}^2$ means that there is an $U(2) = SU(2) \times U(1)$ isometry. Actually this fact supplemented by the condition that the metric is Einstein completely determines the reduced action (up to the constants $y_{1,2}$ and the couplings).

3. The correspondence in the BPS sector

We now want to study the various operators dual to the semiclassical strings moving on the $\mathcal{Y}^{p,q}$ manifolds. Since we consider strings without AdS angular momenta, the operators are scalars constructed only with the matter bifundamental fields. Moreover, if the string is moving fast only along the ψ direction, as is the case for strings described by the effective action of section 2.2, the operators will be holomorphic, or chiral, i.e. products of chiral bifundamentals.

In this section we restrict to chiral primaries, or BPS operators. We will at first focus on the generators of the mesonic chiral ring. 'Mesonic' means that these operators are constructed taking the trace of products of bifundamental fields; in order to be gauge invariant each of these operators has to correspond to a loop in the quiver. Then we describe a generic BPS operator. This study reobtains our previous geometric results on BPS geodesics, and constitutes a first step towards the description of the holomorphic operators dual to semiclassical extended strings. Even if we consider the $\mathcal{Y}^{p,q}$ models, we will uncover generic features of toric quivers.

We refer to [44] for the description of how the superconformal field theories are constructed. Table 1 gives the values of the R-charges Q_R , the $SU(2)$ -spin J and the $U(1)$ flavor charges Q_F for all the bifundamental fields present in a generic $\mathcal{Y}^{p,q}$ quiver⁵. As is well known, for chiral operators there is a simple relation between the R-charge Q_R and

⁵In the case of $\mathcal{Y}^{2,1}$, for which the superconformal field theory was constructed in [66], the values of the R-charges were computed in [67].

Field	number	Q_R	$U(1)_B$	Q_F
Y	$p + q$	$(3q^2 - 4p^2 + 2pq + (2p - q)\sqrt{4p^2 - 3q^2})/3q^2$	$p - q$	-1
Z	$p - q$	$(3q^2 - 4p^2 - 2pq + (2p + q)\sqrt{4p^2 - 3q^2})/3q^2$	$p + q$	$+1$
U^α	p	$(4p^2 - 2p\sqrt{4p^2 - 3q^2})/3q^2$	$-p$	0
V^α	q	$(3q^2 - 2pq + q\sqrt{4p^2 - 3q^2})/3q^2$	q	$+1$

Table 1: Charges of the bifundamental fields present in the $\mathcal{Y}^{p,q}$ quivers found in [44].

the scaling dimension Δ : $\Delta = 3/2Q_R$. Since the R-charges add under multiplication of chiral operators, the knowledge of the R-charge of the four types of bifundamental fields, Y, Z, U^α and V^α , suffices to determine the R-charge of all holomorphic operators and the scaling dimension of all chiral primaries.

3.1 Chiral building blocks

The basic chiral operators correspond to some loops in the quiver that have been considered in [44] in order to give a field theoretical computation of the topology of the supersymmetric 3-cycles. Moreover, the analysis of the short loops (R-charge 2 chiral ring) has been given in [55] in order to determine the conformal manifold. We will, for the sake of clarity, describe nevertheless in detail the various operators also here.

Let us start from some simple examples of mesonic chiral operators. The simplest chiral single trace operators are of the form⁶

$$tr(ZUY_qU) \quad \text{or} \quad tr(UVY_c) \tag{3.1}$$

for 'short' loops of the quivers. For 'long loops' of the quivers of table 2 one finds

$$tr(ZUVUVUZU) \tag{3.2}$$

for counter-clockwise loops, and

$$tr(Y_qUY_qUY_cY_c) \quad \text{or} \quad tr(Y_qUY_qY_cY_cU) \quad \text{or} \quad tr(Y_qUY_qY_cUY_c) \tag{3.3}$$

for clockwise loops. These examples are valid for $Y^{4,2}$, in general there are operators like (3.1) of length 3 and 4, operators like (3.2) of length $2p$ and (3.3) of length $2p - q$. These three types of operators constitute the basic building blocks for any scalar chiral operator. Notice that operators corresponding to 'long loops' carry a non zero winding number around the quivers; this winding number counts the value of the charge associated to the $U(1)$ flavor symmetry. Another thing that can be observed immediately is that the baryonic charge is always vanishing for any mesonic operator (this gives the constraints on the topology of the SUSY 3-cycles of the Sasaki-Einstein manifolds [44]). As a consequence, we do not consider the baryonic charge in the remaining part of this paper.

⁶With the hope of helping in visualize the operators on the quivers, we denote by Y_c the Y -fields entering a cubic superpotential term, and by Y_q the Y -fields entering a quartic superpotential term.




Table 2: Example of the recursive construction of the $\mathcal{Y}^{p,q}$ quivers, as from [44].

Because of \mathcal{F} -terms relations only a subset of these holomorphic operators are chiral primaries, or BPS, and have dimension $\Delta = 3/2R$ on the whole IR conformal surface.

Let us consider the short loops, having R-charge 2. There are $2p$ such loops: $2q$ of length 3 and $p-q$ of length 4. Moreover, since the fields U and V transform in the spin-1/2 of the global $SU(2)$, all the short loops are in the $1/2 \otimes 1/2 = 0 \oplus 1$. We are thus dealing with $4(p+q)$ operators. The \mathcal{F} -term relations will imply that only 3 of them are chiral. The explicit superpotential [44] is

$$W = \sum_{i=1}^q \epsilon_{\alpha\beta} (U_i^\alpha V_i^\beta Y_{2i-1} + V_i^\alpha U_{i+1}^\beta Y_{2i}) + \sum_{j=q+1}^p \epsilon_{\alpha\beta} Z_j U_{j+1}^\alpha Y_{2j-1} U_j^\beta. \quad (3.4)$$

It is important to remember that this writing is schematic, the precise coefficients multiplying every $SU(2)$ -invariant term depend on the position in the conformal surface, similarly to the gauge couplings. The equations of motion of the Y -fields

$$U_i^1 V_i^2 = U_i^2 V_i^1 \quad (3.5)$$

$$V_i^1 U_{i+1}^2 = V_i^2 U_{i+1}^1 \quad (3.6)$$

$$U_j^1 Z_j U_{j+1}^2 = U_j^2 Z_j U_{j+1}^1 \quad (3.7)$$

immediately say that the spin-0 parts are zero in the chiral ring. The equations of motion for the 'external' U , V , Z fields enable to 'move' the short loops around the quiver. All these short loops are thus equal in the chiral ring. The superconformal BPS operator is a symmetrization over the quiver of all these short operators:

$$\mathcal{S}^I = \sum_{i=1}^q \sigma_{\alpha\beta}^I (U_i^\alpha V_i^\beta Y_{2i-1} + V_i^\alpha U_{i+1}^\beta Y_{2i}) + \sum_{j=q+1}^p \sigma_{\alpha\beta}^I Z_j U_{j+1}^\alpha Y_{2j-1} U_j^\beta, \quad (3.8)$$

where σ^I are the 3 Pauli matrices. Also in (3.8) the precise coefficients in front of every $SU(2)$ -covariant term depend on the position in the conformal surface. In conclusion there are only 3 operators, \mathcal{S}^\pm and \mathcal{S}^0 , with R-charge 2 and scaling dimension $\Delta = 3$ over the whole conformal surface, and they transform in the spin-1 representation of $SU(2)$. Note that the Q_F charge of \mathcal{S} is 0. The chiral operator with vanishing spin- z , \mathcal{S}^0 , lies in the

Meson	spin J	Q_R	Q_F
\mathcal{S}	1	2	0
\mathcal{L}_+	$\frac{p+q}{2}$	$p+q - \frac{1}{3\ell}$	$+p$
\mathcal{L}_-	$\frac{p-q}{2}$	$p-q + \frac{1}{3\ell}$	$-p$

Table 3: Charge assignments for the three basic mesonic fields. Notice that $2J - Q_R$ is proportional to Q_F for all the operators.

chiral ring for any toric superconformal quiver and drives the exactly marginal deformation called β -deformation [55]. For more general toric quivers, without an “accidental” $SU(2)$ global symmetry, BPS building blocks with non vanishing $U(1) \times U(1)$ flavor charge, like \mathcal{S}^\pm , are not short: their analog are similar to the ‘long’ chiral operators we are going to study now.

Considering the winding operators we distinguish between clockwise and counter-clockwise loops. The length- $2p$ loop (counter-clockwise, of the form (3.2)) is made of p U -fields, q V -fields and $p - q$ Z -fields. The set of operators (in total 2^{p+q}) corresponding to this loop transform in the $SU(2)$ -representation with spin

$$\left(\otimes^p \frac{1}{2}\right) \otimes \left(\otimes^q \frac{1}{2}\right) = \frac{p+q}{2} \oplus \frac{p+q-2}{2} \oplus \dots \quad (3.9)$$

The \dots represent lower dimensional $SU(2)$ -representations. All the non-maximal $SU(2)$ -representations in the chiral ring vanish, due to the Y -fields \mathcal{F} -term relations. We denote this operator \mathcal{L}_+ . One thus finds that it carries spin $\frac{p+q}{2}$ and, from table 1 a non vanishing positive $U(1)_F$ charge.

The other type of winding operators, clockwise loops of the form (3.2), are a little bit more difficult to visualize. They have length $2p - q$ and are made of p Y -fields and $p - q$ U -fields (including U - or Z -fields in such a loop is equivalent to multiply by an \mathcal{S} operators, and would not be a building block). The Z - and Y -fields \mathcal{F} -term relations imply that the $SU(2)$ indices have to be completely symmetrized, i.e. the BPS operators transform in the spin- $\frac{p-q}{2}$ representation. On the other hand, V -field \mathcal{F} -terms enable one to move the position of the various U - and Y_c -fields present in the operators of the form (3.2). This implies that there is only one BPS clockwise loop, with $J = \frac{p-q}{2}$, that we call \mathcal{L}_- .⁷

The R-charge of the long loops is computed using table 1

$$Q_R[\mathcal{L}_\pm] = p \pm (p(2p - \sqrt{4p^2 - 3q^2})/3q) = p \pm (q - \frac{1}{3\ell}), \quad (3.10)$$

where in the last equality the relation (2.8) has been used. The final results are summarized in table 3.

⁷In the case of $\mathcal{Y}^{p,p}$ there are two different \mathcal{L}_- operators. This fact does not lead to a strong enhancement of the chiral ring, since in this case $J = 0$, in particular it is not possible to use this two operators to construct long operators dual to non point-like semiclassical strings.

The natural way to think of the mesonic operator is in terms of the quiver diagram drawn on a two-torus, as suggested by the connection between dimers and toric geometry [48, 49]. The point is that any toric quiver can be drawn on a torus in such a way to provide a polygonalization of the torus. The quiver diagram is precisely the dual diagram of the dimer model. Each face is surrounded by bifundamental fields going either in the clockwise or in the counterclockwise direction, and precisely corresponds to a superpotential term. For the toric phases of $\mathcal{Y}^{p,q}$ quivers there are only cubic and quartic superpotential terms, so all the faces are triangles or squares. The torus has one 'short' homology cycle and one 'long' homology cycle.

In this picture, the generators of the $\mathcal{Y}^{p,q}$ chiral ring are as follows:

- \mathcal{S}^0 is the only chiral operator that does not wind around any homology cycle of the torus.
- \mathcal{S}^\pm wind around the short homology cycle, in opposite directions.
- \mathcal{L}_\pm wind around the long homology cycle. The value of the z -spin counts the winding number around the short homology cycle.

In all cases we analyzed, we find that the values of the two commuting $U(1)_F$ charges (that are always present) are counted precisely by the two winding numbers of the operator. This is actually valid for any mesonic operator, not just BPS ones. We expect this to be a generic property of toric quivers. A consistency check is that, since all terms in the superpotential (which correspond to the faces of the quiver diagram) have zero flavor charge, the total charge of an operator corresponding to a given path in the periodic quiver, depend only on the initial and final points of the path and not on the particular path we take. Clearly, the winding numbers have the same property.

Of course the generators of the chiral ring we found satisfy various non linear relations. Studying these relations, it should be possible to reconstruct the algebro-geometric description of the $\mathcal{Y}^{p,q}$ Calabi-Yau cones. Instead of doing this, we will reconstruct the transverse geometry through the analysis of semiclassical holomorphic operators, in section 5. This will give also information about the metric.

3.2 The full mesonic chiral ring

We now want to consider 'multiloop' operators. First of all we see what happens multiplying two \mathcal{S} . An operator like

$$tr(U_i V_i Y_{2i+2} U_i V_i Y_{2i+2}) \sim tr(U_i V_i Y_{2i+2} U_i Y_{2i+3} V_{i+1}) \tag{3.11}$$

transforms in the $\otimes^4 1/2 = 2 \oplus \dots$ and can be seen simply as the product of two short mesons \mathcal{S} . It is easy to convince that this is general: any chiral operator that does not wind all the way around the quiver is of the form \mathcal{S}^n and transforms in the spin- n of $SU(2)$. In other words, BPS operators do not carry a position on the quiver, and always transform in the maximal possible $SU(2)$ -irrep.

Consider now the product of \mathcal{L}_+ and \mathcal{L}_- . This product, which winds 0 times around the quiver, can be expressed in terms of the \mathcal{S} operators.

$$\mathcal{L}_+\mathcal{L}_- \sim \text{tr}(\dots UVU YYU \dots) \sim \text{tr}(\dots UVY UYU \dots) \quad (3.12)$$

where we used the equation of motion for V . We thus see that the resulting operator is the product of various \mathcal{S} s. More precisely

$$\mathcal{L}_+\mathcal{L}_- \sim \mathcal{S}^p \quad (3.13)$$

Note that this relation is consistent with the charge assignments of table 3.

We are now in the position of giving the classification of the mesonic chiral BPS operators of the $Y^{p,q}$ quivers. A general operator \mathcal{O} can be seen as the product of \mathcal{S} and \mathcal{L} :

$$\mathcal{O}_{s,l} = \mathcal{S}^s \mathcal{L}^l \quad (3.14)$$

Where we denote $\mathcal{L} = \mathcal{L}_+$ and $\mathcal{L}^{-1} = \mathcal{L}_-$. s is a non-negative integer, while the integer l can be positive or negative. The R-charge of $\mathcal{O}_{s,l}$ is given by

$$Q_R[\mathcal{O}_{s,l}] = 2s + p|l| + l \left(q - \frac{1}{3\ell} \right) \quad (3.15)$$

while the flavor charge

$$Q_F[\mathcal{O}_{s,l}] = pl \quad (3.16)$$

Finally, $\mathcal{O}_{s,l}$ transforms in the irreducible $SU(2)$ -representation with spin J

$$J[\mathcal{O}_{s,l}] = s + |l| \frac{p}{2} + l \frac{q}{2} \quad (3.17)$$

Again, the precise form of these operators can be obtained by a complete symmetrization over the quiver (imposed by U - and V -fields \mathcal{F} -terms) and over the $SU(2)$ indices (imposed by Y - and Z -fields \mathcal{F} -terms). A complete symmetrization over the trace is also to be performed.

3.3 BPS geodesics from the quivers

The operators corresponding to point-like strings moving along a null BPS geodesic are chiral primaries and therefore should be among the ones we just described. In this section we make the mapping precise and compare with the results of section 2.1.

Before doing so note that, heuristically, we can understand that these operators correspond to point-like strings because, due to the complete symmetrization imposed by \mathcal{F} -term relations, the three values of the $U(1)^3$ charges are constant along the operator, for long operators and in a sense made precise later when we study a coherent state representation of the operator (see eqs. (5.6) and (5.7)). There we also see that we can get non-BPS operators by tuning continuously, along the operator, the ratio l/s , corresponding to the value of the $U(1)_F$ charge and the z -component of the $SU(2)$, corresponding to the difference between the number of A^1 -fields and the number of A^2 -fields (A^α stands for U^α or V^α .) The R-charge is determined in terms of l and s by the relation 3.15.

Going back to our main problem in this section, the first task is to reobtain, from the field theory, the quantities $y_{1,2}$ and ℓ (defined in (2.9)) that play an important role in the supergravity background.

We start by writing the charges of a chiral operator made out of n_+ operators \mathcal{L}_+ , n_- \mathcal{L}_- 's and s operators \mathcal{S} composed to maximum $SU(2)$ spin J . The result is

$$Q_F = pn_+ - pn_- = pn_\alpha \quad (3.18)$$

$$J = n_+ \frac{p+q}{2} + n_- \frac{p-q}{2} + s = \frac{1}{2}(p\bar{n} + 2s) + \frac{1}{2}qn_\alpha \quad (3.19)$$

$$Q_R = n_+ \left(p + q - \frac{1}{3\ell} \right) + n_- \left(p - q + \frac{1}{3\ell} \right) + 2s = (p\bar{n} + 2s) + \left(q - \frac{1}{3\ell} \right) n_\alpha \quad (3.20)$$

where we introduced $\bar{n} = n_+ + n_-$ and $n_\alpha = n_+ - n_-$. We see that we can use n_α instead of Q_F . Furthermore, \bar{n} and s appear only in the combination $p\bar{n} + 2s$ which follows from the fact that actually $\mathcal{L}_+\mathcal{L}_- \sim \mathcal{S}^p$ in the chiral ring. This means that there are only two independent numbers and therefore from (3.19) and (3.20) a relation between the charges follows

$$Q_R - 2J = -\frac{1}{3\ell}n_\alpha \quad (3.21)$$

So, ℓ^{-1} has appeared as a natural unit for $U(1)_F$ charge. We can define two new variables:

$$P_\alpha = \frac{n_\alpha}{\ell}, \quad \text{and} \quad y_0 = -\frac{P_\alpha}{3Q_R} \quad (3.22)$$

Therefore, in the field theory, y_0 is the relation between $U(1)_F$ and R-charges for a given operator. The range of y_0 is determined by noticing that its minimum and maximum values correspond to \mathcal{L}_+ and \mathcal{L}_- respectively, namely for $n_- = s = 0$ and $n_+ = s = 0$:

$$y_0(\mathcal{L}_+) = -\frac{n_\alpha(\mathcal{L}_+)}{3\ell Q_R(\mathcal{L}_+)} = y_1 \quad (3.23)$$

$$y_0(\mathcal{L}_-) = -\frac{n_\alpha(\mathcal{L}_-)}{3\ell Q_R(\mathcal{L}_-)} = y_2 \quad (3.24)$$

where we used the Q_R charges of \mathcal{L}_+ and \mathcal{L}_- from table 3. In this way we recover, from the field theory, that

$$y_1 \leq y \leq y_2 \quad (3.25)$$

If we rewrite now the charges (3.20) in terms of y_0 we get perfect agreement with (2.32) and (2.33):

$$J = \frac{1}{2}(1 - y_0)Q_R, \quad P_\alpha = -3y_0Q_R \quad (3.26)$$

We have therefore identified the chiral operators with massless geodesics. Particular examples are: an operator made out only of \mathcal{L}_+ with the geodesic at $y_0 = y_1$, one made out only of \mathcal{L}_- with the geodesic at $y_0 = y_2$ and one made out of equal number of \mathcal{L}_+ and \mathcal{L}_- with the geodesic at $y_0 = 0$.

4. The field theory at $\lambda \rightarrow 0$

Since BPS operators are protected, the matching between chiral primary operators and massless geodesics is valid at generic points on the conformal surface of the $\mathcal{Y}^{p,q}$ quivers. The $SU(2)$ invariant points on the conformal manifolds are parametrized by two complex parameters⁸. On the string side [39, 61, 68], these parameters are the complex dilaton (g_s) and the vev of the B -fields (RR and NSNS) on the 2-cycle present in the manifold $S^2 \times S^3$.

On the string side we further considered a set of string states described by the effective action of section 2.2. Since these states are not BPS the effective action is valid only in the regime $g_s \ll 1$ (to ignore string loops) and small curvature, namely $\lambda = (R/l_s)^{\frac{1}{4}} = g_s N \gg 1$.

In the case of $\mathcal{N} = 4$ this effective action can be compared to a similar action derived from the field theory in the opposite regime $\lambda \ll 1$ (which can also be interpreted as taking g_s to zero keeping N fixed and large). In our case, since the effective action is proportional to λ , a naive extrapolation to λ small suggests that, in that regime, the result might be interpreted as a small perturbation around a point with $\lambda = 0$. If such a point exists (actually should be a line) it is special since all the semiclassical operators described by the reduced action would satisfy $\Delta = 3/2Q_R$.

In the case of $\mathcal{N} = 4$ SYM (and orbifolds thereof), this point is the free theory. In the case of the conifold [55] there is a line of conformal fixed points with vanishing superpotential which is part of the conformal manifold. Having $\mathcal{W} = 0$ implies that *all* chiral operators are chiral primaries, or BPS, operators; in other words the chiral ring is much bigger on this line, depicted in figure 1. This means that this particular line should be identified with the $\lambda = 0$ point as suggested in [69] where the BMN limit of the conifold was studied. Notice that the gauge couplings are not zero which precludes doing standard perturbation theory. Rather, one should do conformal perturbation theory on a marginal perturbation given by the quartic superpotential. As a last preliminary example, for the $\mathcal{Y}^{2,0}$ quiver in the “single impurities phase”, this $\mathcal{W} = 0$ conformal line has an enhanced chiral ring and a global symmetry $SU(2)^4$. What is interesting here is that one can Seiberg dualize [70] this phase with $\mathcal{W} = 0$, obtaining a special line on the “double impurity phase” [68]; in this dual description there are the standard Seiberg $\mathcal{M}q\tilde{q}$ superpotential terms, so the superpotential does not vanish. We conclude that $\lambda = 0$ on the string side is not, in general, equivalent to $\mathcal{W} = 0$ on the gauge side.

We can now consider a generic $\mathcal{Y}^{p,q}$ quiver. In this case the superpotential has cubic and quartic terms. Since there are several terms in the superpotential we do not expect to be able to cancel all of them since we can only vary two parameters on the conformal surface. If only gauge couplings are turned on, the conformal dimension decreases and we can never get to the conformal manifold where for example the V -fields have R-charge




Figure 1: Space of couplings for the conifold field theory with no superpotential. The conformal manifold is indicated.

⁸In [55] a description of the full conformal manifold is given.

larger than $2/3$ ⁹. We conclude that, at least for the toric phases, there is no point on the conformal manifold with vanishing superpotential.

In fact this problem was studied in [71], where it is pointed out that, if a IR conformal manifold exists¹⁰, it is sufficient that *one* gauge group is asymptotically free in order to be able to flow from the free theory to the conformal manifold but, in order to reach this point, the superpotential couplings are crucial. For the $\mathcal{Y}^{p,q}$ models, the flow has been qualitatively described in [55]: one flows at first the $N_F = 2N_C$ nodes, then some cubic superpotential terms and so on.

We can actually go further by using an idea of Kutasov [72], that can be thought of as extending the a -maximization procedure of Intriligator and Wecht [45] away from the conformal manifold. In our case this consists in introducing $2p$ Lagrange multipliers $\lambda_{i=1\dots 2p}$, one for each gauge group, and $\mu_{k=1\dots p+q}$, one for each term in the superpotential. We can define a as:

$$a = 3\text{tr}(R - 1)^3 - \text{tr}(R - 1) - \sum_i \lambda_i (\text{tr}(\epsilon_i(R - 1)) + 2) + \sum_k \mu_k (\text{tr}(\nu_k R) - 2) \quad (4.1)$$

where ϵ_i is a diagonal matrix in the space of fields which is 1 or 0 if the corresponding field is charged or not with respect to the i -gauge group. The same for ν_k which is 1 or 0 if the field appears or not in the k -th term in the superpotential. If we maximize a with respect to the R-charges they become functions of the Lagrange multipliers (λ_i, μ_k) . It was further argued in [72] that the Lagrange multipliers can be used to parameterize the space of couplings. Although the relation between the couplings and the multipliers is still somewhat conjectural, one thing is clear: a coupling is zero if and only if the corresponding Lagrange multipliers is zero, since the corresponding constraint has not to be imposed. We want to see now if, on the conformal surface, we can make some of the Lagrange multipliers to vanish. Since we are on the conformal surfaces we have to impose anomaly cancellation conditions. After that one can see, working in specific examples, that one can put some quartic terms in the superpotential to zero. (Here we are also using that all Lagrange multipliers are positive in the physical region of the couplings).

We can now understand the superpotential corresponding to $\lambda = 0$. Decreasing λ , all the couplings (taken in their absolute value) decrease. At some point a coupling becomes zero. This coupling has thus to correspond to a quartic term in the superpotential.

Having a vanishing superpotential coupling generates a change in the chiral ring of the theory but it is not obvious that the holomorphic long operators corresponding to the semiclassical strings of section 2.2 become protected. To see that we focus on an example, $\mathcal{Y}^{4,3}$, which should clarify the general structure.

Consider the operator \mathcal{L}_+ in the $\mathcal{Y}^{4,3}$ quiver.

$$\mathcal{L}_+ = \text{tr}(ZUVUVUVU) \quad (4.2)$$

To be chiral primary, namely not a descendant, \mathcal{L}_+ has to transform in the spin-7/2 representation of $SU(2)$, as follows from using the \mathcal{F} -term relation of the Y -field entering in the cubic terms. If we multiply two \mathcal{L}_+ operators, for $\mathcal{W}_4 \neq 0$, we saw in section 3.2

⁹The presence of fields with $Q_R \geq 2/3$ follows, directly, from the presence of nodes with $N_F = 3N_C$.

¹⁰This is not the case for a generic $\mathcal{N} = 1$ quiver theory.

that we get an operator that transform in the spin-7 representation (with 15 states), of the form

$$\text{tr}(ZUVUVUVU ZUVUVUVU) \quad (4.3)$$

If $\mathcal{W}_4 = 0$, however, there are more than 15 states. The reason is that one cannot use the \mathcal{F} -term relations coming from the quartic terms to move the $SU(2)$ spins from the first $UVUVUVU$ block to the second $UVUVUVU$ block. More generally, operators of the form $(\mathcal{L}_+)^n$ contain a number of states that grows as 8^n if $\mathcal{W}_4 = 0$ and as $8n$ if $\mathcal{W}_4 \neq 0$. It is also possible to see that the spin-1 \mathcal{S} operator gets enhanced to spin- $(1 \oplus 0)$. This \mathcal{S} operator with $J = 0$ generates an exactly marginal deformation, which is precisely the conformal line parameterized by g_s .

What happens is thus that, at $\mathcal{W}_4 = 0$, the chiral ring is much larger which leads us to identify this point with $\lambda = 0$ although it is not true that all chiral operators are in the chiral ring.

We consider now semiclassical operators at the $\mathcal{W}_4 = 0$ point. Let us focus for simplicity on operators of the form $(\mathcal{L}_+)^n$. These correspond to semiclassical strings moving only on the round two-sphere, satisfying $y(\sigma, \tau) = y_1$. A class of operators of the form $(\mathcal{L}_+)^n$ is as follows:

$$\text{tr} \left(\prod_{i=1}^n R(\theta_i, \phi_i) \mathcal{L}_+ \right) \quad (4.4)$$

where $R(\theta_i, \phi_i)$ is an $SU(2)$ rotation applied to \mathcal{L}_+ (which has maximum z -spin P_ϕ). Taking n to be large, and the angles (θ_i, ϕ_i) which parameterize the rotation to vary smoothly with i , we see that we are constructing a semiclassical string extended along the S^2 sphere parameterized by (θ, ϕ) . This is similar to what happens in the $SU(2)$ sector of $\mathcal{N} = 4$ operators [18]. Reconstructing the directions y and β is more involved and can be recovered from the results the next section (naively, the absence of the \mathcal{F} -term relations coming from \mathcal{W}_4 implies that one cannot exchange \mathcal{L}_+ with \mathcal{L}_-). We just emphasize that the important point here is that a generic operator like (4.4), at $\mathcal{W}_4 = 0$, is BPS, and satisfies the relation $\Delta = 3/2Q_R$.

5. Effective action for the spin chain

In the previous sections we studied conformal primaries and compared them to the massless geodesics in the metric. As we just discussed, going further is difficult since the theories are strongly coupled and we cannot use a perturbative expansion to compute anomalous dimensions. In principle, as argued in the previous section, we should use conformal perturbation theory around a conformal point where some terms in the superpotential vanish. Instead of doing that, to simplify the problem, we are going to consider all terms of the superpotential on equal footing and extract a simple spin chain model that captures the generic features of the operator mixing that the superpotential produces. Even then we are




Figure 2: Quiver diagram corresponding to $\mathcal{Y}^{4,3}$.

going to simplify the problem further. From the point of view of the resulting spin chain what we are doing is trying to obtain the correct long distance physics so we expect that the microscopic details should not be important.

Using coherent states we obtain an effective action for the spin chain which is similar to the one we derived from string theory, namely eq. (2.53) albeit with a different function $p(y)$.

We analyze first the case of $\mathcal{Y}^{3,2}$ which should make clear the generic case we discuss afterwards.

5.1 Long paths in the $\mathcal{Y}^{3,2}$ quiver

The quiver corresponding to $\mathcal{Y}^{3,2}$ is depicted in figure 3. Gauge invariant operators correspond to closed paths in the quiver. An example is the outer counterclockwise loop that we called \mathcal{L}_+ . Other important example is the operator \mathcal{L}_- . It is a linear combination (with equal coefficients) of the three paths depicted in the figure and can be written as:

$$\mathcal{L}_-^{(3,2)} = \frac{1}{\sqrt{3}} [\text{tr}(Y_q U Y Y) + \text{tr}(Y_q Y U Y) + \text{tr}(Y_q Y Y U)] \quad (5.1)$$

Here, for clarity, we denoted as Y_q the operator Y that appears in the quartic superpotential term $\text{tr}(U Y_q U Z)$. We see that \mathcal{L}_- is a mix of three operators where the operator U moves between three possible positions among the Y 's. This mixing comes from the cubic vertices of the superpotential as can be seen in the example of figure 4. The mixing matrix induced by these vertices is proportional to

$$H = h \begin{pmatrix} -1 & 1 & 0 \\ 1 & -2 & 1 \\ 0 & 1 & -1 \end{pmatrix} \quad (5.2)$$

The off-diagonal terms correspond to the mixing. The diagonal terms come also from the superpotential. They have opposite sign due to the relative sign between different terms in the superpotential and one is double of the others since, for that state, the U has two neighboring Y 's. Instead, when the U is between Y_q and Y only the Y counts since there is no term in the superpotential involving U and Y_q (and no Z). We wrote the mixing matrix as H since one can think of it as a Hamiltonian whose eigenvalues are the conformal dimensions. In this case we find the eigenvector $\frac{1}{\sqrt{3}}(1, 1, 1)$ with eigenvalue zero which is precisely \mathcal{L}_- . The constant h denotes the superpotential couplings (and other factors appearing in the computation) and therefore cannot be taken to be small in general. This implies that to obtain the correct spin chain Hamiltonian one should use non perturbative techniques that sum all the diagrams. What is clear is that $\frac{1}{\sqrt{3}}(1, 1, 1)$ is always a ground state, since it is a protected operator.

Now we should investigate what happens for more generic operators, namely to all possible closed loops in the quiver. These loops form a basis in a Hilbert space. In such space we can define a Hamiltonian that converts a given path in a linear combination of all paths that can be obtained from it by using the “moves” of the type described in figure 4.




Figure 3: Quiver diagram corresponding to $\mathcal{Y}^{3,2}$. We show, on the top left, the path \mathcal{L}_+ . The other three paths are those whose symmetric linear combination is \mathcal{L}_- .




Figure 4: The superpotential generates mixing among the chiral operators, namely the closed paths in the quiver. Diagrams as the one shown on the right give mixing between paths that differ by the “move” shown on the left: $324 \leftrightarrow 342$. Here F_V denotes the F component of the chiral field V . The other moves are: $423 \leftrightarrow 453$, $534 \leftrightarrow 564$, $6125 \leftrightarrow 645$, $231 \leftrightarrow 2561$.

In fact, to understand the dynamics of the paths it is better to plot the quiver in a plane where the two axis give the angular momentum J and $U(1)_F$ charge Q_F . In figure 5 we can see such a plot. Each point is a vertex of the quiver and the operators U, V, Y and




Figure 5: It is convenient to draw the quiver on a plane. The horizontal axis corresponds to the P_α charge in ℓ^{-1} units and the vertical axis to the total J (assuming that we compose the operators to maximum $SU(2)$ spin). The “moves” that convert one path into another are now very simple as exemplified in the figure where the solid path can take alternative routes depicted with dashed lines according to the “moves” $564 \rightarrow 534$ and $645 \rightarrow 6125$.

Z are the arrows plotted according to the J and Q_F of each operator (see table 1). The diagram is infinite but periodic as is clear from the labels of the vertices. A closed path in the quiver is given here by an open path where the initial and end points should have the same label. The difference in the coordinates of the initial and final point determine the charges of the operator. This representation is similar to the doubly periodic representation of toric quivers dual to the dimer picture. However it is adapted to the fact that here we have an $SU(2)$ global symmetry. In the general case, with only the toric $U(1) \times U(1)$ flavor symmetry, the $U(1)$ charges of each bifundamental fields in the torus representation are in correspondence with the direction of the field.

After trying different paths it is easy to see that although individual jumps can be done in several directions, in average, the slope of a path lies between the one corresponding to \mathcal{L}_- and \mathcal{L}_+ . Also, for chiral primaries, this can be seen by parameterizing the slope in terms of y as in 3.26,

$$\frac{J}{P_\alpha} = -\frac{1}{6y} + \frac{1}{6} \quad (5.3)$$

We know, from the analysis of the previous section that, in the field theory (as in the string side) one has $y_1 < y < y_2$. So, in this case, the limit in the slopes that we mentioned is the same as (3.25).

A simplification appears when we consider how the moves that determine the Hamiltonian are represented in this diagram. It is easy to see (as exemplified in figure 5) that they simply correspond to moving the path across the polygons or faces in the diagram. For example we can convert $\dots 564\dots$ into $\dots 534\dots$ etc. In this way we can get from a given path all paths that join the two given vertices. In such moves, the number of operators is not conserved but the R -charge is and therefore we can use the R -charge as a measure of the length.

One other thing to take into account is that not all moves have same weight, since they correspond to different terms in the superpotential. In particular, moving the path across a shadowed region in figure 5 requires the use of the quartic superpotential and therefore it is suppressed at the points with $\mathcal{W}_4 \simeq 0$. This also shows that at this point there are semiclassical operators with a non trivial $y(\sigma)$ satisfying $\Delta = 3/2Q_R$, similar to eq. (4.4). However, if we want to study very long paths, namely very large R -charge, we can take a limit where the paths become continuous and the details of the diagram are irrelevant. What remains is the fact that there is a maximum and minimum slope for the paths. The Hamiltonian acting on a path produces infinitesimal deformations weighted by an effective coupling that vanishes at the special points with $\mathcal{W}_4 = 0$. Each path can be described (up to reparametrizations) by the slope as a function of σ , the coordinate along the path. We associate the slope to the variable y . Since this is configuration space, in the classical limit we need also a momentum conjugate to y that turns out to be the angle β . Furthermore, each portion of path has an angular momentum $\Delta J = \frac{1}{2}(1 - y)\Delta Q_R$ which can be oriented in a direction parameterized by two angles (θ, ϕ) . In this way we see that each path is determined by four variables function of σ . Therefore, the path itself becomes the string that we found on the string side!

To be more precise we have to compute the action for these paths as determined by the Hamiltonian.

One can consider a related, discrete model, where the paths have the same properties and therefore should be described by the same long distance physics (long distance in the sense of the paths). The model is depicted in figure 6. We consider the lattice formed by the dashed lines which are parallel to the directions determined by \mathcal{L}_+ and \mathcal{L}_- . The parallel lines are one unit of R -charge from each other. Consider the points A , B and C lying in a line of equal R -charge. From the origin to the points A (or B) there is only one path corresponding to a chiral primary operator. However, to a point such as C there are many path that should be entangled. The Hamiltonian is taken to be the one that moves a path across one parallelogram.

More precisely, if we describe the paths as a succession of two “effective operators” with charges

$$\begin{aligned} \mathcal{L}_1 : Q_R = 1, \quad J_1 = \frac{1}{2}(1 - y_1), \quad P_\alpha^{(1)} = -3y_1 \\ \mathcal{L}_2 : Q_R = 1, \quad J_2 = \frac{1}{2}(1 - y_2), \quad P_\alpha^{(2)} = -3y_2 \end{aligned} \tag{5.4}$$

the Hamiltonian can be written as

$$H = h_{\text{eff.}} \sum_{i=1}^L (1 - P_{ii+1}) \tag{5.5}$$




Figure 6: Effective description of the lattice in figure 5. We tile the wedge where paths are contained by parallelograms whose sides carry one unit of R -charge. In this way lines of constant R -charge are such as ACB . The moves are similar as in the other case. For example the two paths shown are connected by applying four moves and therefore mix under renormalization.

where P_{i+1} is the permutation operator between neighboring sites¹¹. The identity is included so that we do not get corrections to operators made out only of \mathcal{L}_1 's (or \mathcal{L}_2 's). The coefficient h_{eff} is an effective coupling that should be computed by matching to the description in the quiver. The Hamiltonian permutes $\mathcal{L}_1\mathcal{L}_2$ into $\mathcal{L}_2\mathcal{L}_1$ which moves the path across the lattice in a similar way as happens in the quiver. h_{eff} is small when $\mathcal{W}_4 \simeq 0$, since without one quartic coupling one cannot permute \mathcal{L}_1 and \mathcal{L}_2 .

In the continuum limit the paths in this lattice are continuous paths such that the slope is contained between the ones of \mathcal{L}_1 and \mathcal{L}_2 and the Hamiltonian moves such paths around. Since the continuum description is the same we expect that this simplified model is described by the same effective action as the one in the quiver.

Perhaps a more detailed analysis can be desirable but we do not expect that changes this simple picture.

We can now analyze the operators constructed out of \mathcal{L}_1 and \mathcal{L}_2 . However at this point it is clear that we can repeat the discussion for any $\mathcal{Y}^{p,q}$ quiver and the result will be the same except with different values of y_1 and y_2 . So we proceed now to the generic case.

5.2 Closed paths in $\mathcal{Y}^{p,q}$

In the previous subsection we argued that long operators in the quiver can be modeled by operators constructed out of the two effective operators defined in eq. (5.4) with Hamiltonian (5.5). We now want to derive a classical action that describe the dynamics of these paths in the limit in which they are very long.

¹¹Similar expressions are familiar in the $\mathcal{N} = 4$ case [11, 73]

At first sight, such paths seem equivalent to a Heisenberg model if we associate *e.g.* \mathcal{L}_1 to spin up and \mathcal{L}_2 to spin down. However we should remember that \mathcal{L}_1 and \mathcal{L}_2 also carry $SU(2)$ spin (given by J_1 and J_2) and therefore we need two angles to describe their orientation. Nevertheless in deriving the classical action it is clear that we can apply the same coherent state techniques that are useful in the Heisenberg model.

Following the method of coherent states, we are going to consider operators of the type

$$|\mathcal{O}\rangle = \prod_{i=1}^{Q_R} |\mathcal{O}_i\rangle \tag{5.6}$$

where we used $|\mathcal{O}\rangle$ to denote an operator to emphasize that we also consider it as a state of a physical system. Also, Q_R is the length, namely R -charge of the operator since we chose elementary jumps each with a unit R -charge. The operators \mathcal{O}_i are defined as a linear combination of \mathcal{L}_1 and \mathcal{L}_2 :

$$\mathcal{O}_i = \sum_{a=1}^2 \rho_{ia} e^{iP_\alpha^{(a)}\alpha_i} U(\theta_i, \phi_i, \psi_i) \mathcal{L}_a \tag{5.7}$$

Here we took \mathcal{L}_1 and \mathcal{L}_2 to have maximum projection in the direction S_3 of $SU(2)$ and then applied a rotation $U(\theta_i, \phi_i, \psi_i)$ parameterized by three Euler angles θ_i, ϕ_i, ψ_i . These angles are slowly varying with i which implies that the total $SU(2)$ spin is not maximal and therefore the operator is not primary. One other point is that since P_α as defined in (3.22) is quantized in units of ℓ^{-1} the range of variation of α is $0 \leq \alpha \leq 2\pi\ell$ in agreement with (2.6).

The coefficients ρ_a determine the relative weight between \mathcal{L}_1 and \mathcal{L}_2 and therefore correspond to the slope of a particular piece of the path. They have to satisfy the conditions

$$\rho_1^2 + \rho_2^2 = 1, \quad y_1\rho_1^2 + y_2\rho_2^2 = y \tag{5.8}$$

In fact this can be taken as the definition of y as the mean value of the slopes y_1 and y_2 . It implies that J and P_α are given by

$$J = J_1\rho_1^2 + J_2\rho_2^2 = \frac{1}{2}(1 - y) \tag{5.9}$$

$$P_\alpha = P_\alpha^{(1)}\rho_1^2 + P_\alpha^{(2)}\rho_2^2 = -3y \tag{5.10}$$

which are by now familiar expressions (if we remember that $Q_R = 1$ for this effective operators). In terms of y we can write ρ_1 and ρ_2 as

$$\rho_1^2 = \frac{y_2 - y}{y_2 - y_1}, \quad \rho_2^2 = \frac{y - y_1}{y_2 - y_1} \tag{5.11}$$

Here y should also be consider to be a slowly varying function of i .

Finally the angle α_i determines the relative phase in the linear combination and, in the coherent state action turns out to be the canonical conjugate of the variable y_i .

Since a global phase is irrelevant there is a redundancy between ψ and α that we are going to resolve later by an appropriate “gauge choice”.

The classical action for the coherent states is given by

$$S = \int d\tau \text{Im} \langle \mathcal{O} | \frac{\partial}{\partial \tau} | \mathcal{O} \rangle - \int d\tau \langle \mathcal{O} | H | \mathcal{O} \rangle \quad (5.12)$$

where the Hamiltonian was given in eq. (5.5). Its mean value is:

$$\langle \mathcal{O} | H | \mathcal{O} \rangle = \sum_{i=1}^{L=Q_R} \left(1 - |\langle \mathcal{O}_i | \mathcal{O}_{i+1} \rangle|^2 \right) \quad (5.13)$$

where we identified the length L of the chain with the total R-charge Q_R . It can be computed in the continuum limit in an expansion in derivatives. Up to second order it gives, after a somewhat lengthy but simple computation:

$$\langle \mathcal{O} | H | \mathcal{O} \rangle = \frac{1}{2} \int_0^{Q_R} d\sigma \left\{ \sum_{a=1}^2 (\partial_\sigma \rho_a)^2 + \langle (D_\sigma^{(a)} \alpha)^2 \rangle - \langle D_\sigma^{(a)} \alpha \rangle^2 \right. \quad (5.14)$$

$$\left. + \sum_{a=1}^2 \rho_a^2 J_a \left((\partial_\sigma \theta)^2 + \sin^2 \theta \partial_\sigma \phi^2 \right) \right\} \quad (5.15)$$

where, for brevity we defined

$$D_\sigma^{(a)} \alpha = P_\alpha^{(a)} \partial_\sigma \alpha + J_a (\partial_\sigma \psi + \cos \theta \partial_\sigma \phi) \quad (5.16)$$

and $\langle \dots \rangle$ denotes average in the sense $\langle \xi_a \rangle = \rho_1^2 \xi_1 + \rho_2^2 \xi_2$.

Doing the change of variables

$$\alpha = \frac{1}{6\ell} (\beta - \psi) \quad (5.17)$$

we obtain that

$$D_\sigma^{(a)} \alpha = \frac{P_\alpha^{(a)}}{6\ell} (\partial_\sigma \beta + \cos \theta \partial_\sigma \phi) + \frac{1}{2} (\partial_\sigma \psi + \cos \theta \partial_\sigma \phi) \quad (5.18)$$

We see now explicitly the redundancy between ψ and α (or β and ψ now). We can fix this ambiguity by choosing

$$\partial_\sigma \psi + \cos \theta \partial_\sigma \phi = y (\partial_\sigma \beta + \cos \theta \partial_\sigma \phi) \quad (5.19)$$

to agree with eq. (2.51). We can therefore write

$$D_\sigma^{(a)} \alpha = \left(\frac{P_\alpha^{(a)}}{6} + \frac{1}{2} y \right) (\partial_\sigma \beta + \cos \theta \partial_\sigma \phi) \quad (5.20)$$

We can now compute

$$\langle D_\sigma^{(a)} \alpha \rangle = \sum_{a=1}^2 \rho_a^2 \left(\frac{P_\alpha^{(a)}}{6} + \frac{1}{2} y \right) (\partial_\sigma \beta + \cos \theta \partial_\sigma \phi) = 0 \quad (5.21)$$

and

$$\langle (D_\sigma^{(a)} \alpha)^2 \rangle = \sum_{a=1}^2 \rho_a^2 \left(\frac{P_\alpha^{(a)}}{6} + \frac{1}{2} y \right)^2 (\partial_\sigma \beta + \cos \theta \partial_\sigma \phi)^2 = \frac{1}{4} (y - y_1)(y_2 - y) (\partial_\sigma \beta + \cos \theta \partial_\sigma \phi)^2 \quad (5.22)$$

which together with

$$\sum_{a=1}^2 \rho_a^2 J_a = \frac{1}{2} (1 - y) \quad (5.23)$$

completes the evaluation of $\langle \mathcal{O} | H | \mathcal{O} \rangle$. Replacing in the action we get

$$S = \frac{Q_R}{2\pi} \int d\tau d\sigma \left[\frac{1}{2} \partial_\tau \psi - \frac{1}{2} y \partial_\tau \beta + \frac{1}{2} (1 - y) \cos \theta \partial_\tau \phi \right] \quad (5.24)$$

$$+ \frac{\pi h_{\text{eff}}}{2Q_R} \int d\tau d\sigma \left\{ (1 - y) [(\partial_\sigma \theta)^2 + \sin^2 \theta (\partial_\sigma \phi)^2] \right. \quad (5.25)$$

$$\left. - \frac{(\partial_\sigma y)^2}{(y_2 - y)(y - y_1)} + (y - y_1)(y_2 - y) (\partial_\sigma \beta + \cos \theta \partial_\sigma \phi)^2 \right\} \quad (5.26)$$

where we also computed the Wess-Zumino term using similar methods and $0 \leq \sigma \leq 2\pi$.

Comparing with (2.53) one has to identify h_{eff} with λ . We see that there is agreement, except that the function $p(y)$ is different. The function $p(y)$ that we obtained also vanishes at $y = y_1$ and $y = y_2$ and can be considered as a first approximation to the actual $p(y)$. However, it should be noted that the metric of the five dimensional manifold is singular since, as observed in section 2, to avoid a conical singularity, we need $p'(y_{1,2}) = \pm 2y_{1,2}$ which this $p(y)$ does not satisfy.

It is clear that the rest, namely the dependence in the angles, is largely determined by symmetry so the partial agreement does not seem like a big accomplishment. However the purpose here was to derive this action directly from the field theory without reference to the AdS/CFT correspondence. From that point of view it is not even clear that such action should exist and the mere fact that one can find a string representation for these operators in the field theory should be considered as a check of the relation between strings and gauge theories. Moreover it is plausible that in the infrared of the world sheet this model flows to the one derived from the string side. We leave this problem for future work.

It would also be nice to apply this procedure to other examples, as the ones discussed in [74].

One final point is that we can find again a local Kähler potential for this model of the form

$$K = - \left\{ \frac{1 - y_1}{y_2 - y_1} \ln |y - y_1| + \frac{1 - y_2}{y_1 - y_2} \ln |y_2 - y| \right\} \quad (5.27)$$

with complex coordinates

$$z_1 = \sin\left(\frac{\theta}{2}\right) e^{-i\frac{1}{2}(\beta - \phi)} \left(\frac{y_2 - y}{y - y_1} \right)^{\frac{1}{2}(y_2 - y_1)} \quad (5.28)$$

$$z_2 = \cos\left(\frac{\theta}{2}\right) e^{-i\frac{1}{2}(\beta + \phi)} \left(\frac{y_2 - y}{y - y_1} \right)^{\frac{1}{2}(y_2 - y_1)} \quad (5.29)$$

6. More general operators

In the string analysis we found massless BPS geodesics which we mapped to chiral primary operators. After that we extended the result to certain excited strings which gave chiral operators which for large R-charge Q_R have anomalous dimensions $(\Delta - 3/2Q_R)$ of order $\frac{\lambda}{Q_R^2}$, where λ is the string tension.

Now we want to extend the discussion to massless strings moving along non-BPS geodesics. As seen in section 2, in that case the conformal dimensions do not depend on λ at least in the region of large λ in which the results are valid. This suggests that these operators might be protected, namely, their conformal dimension do not depend on the point of the conformal manifold where they are computed. As a particular case one can consider geodesics which move close to a BPS geodesic or large R-charge Q_R . As we show below, the conformal dimension Δ of the corresponding near BPS operators behaves as $\Delta = \frac{3}{2}Q_R + \delta\Delta + \mathcal{O}\left(\frac{1}{Q_R}\right)$ where $\delta\Delta$ is of order 1 in an expansion for large R-charge. In the limit of large λ , $\delta\Delta$ is independent of λ since Δ is. A more conservative point of view is to suggest that only the first correction $\delta\Delta$ is protected. In the rest of the section we find a description of the corresponding operators and leave further consideration about the dependence on λ for future study.

Notice that this problem is absent in the $\mathcal{N} = 4$ case since there all massless geodesics in the S^5 are protected. The discussion is therefore closer to what was discussed for the $T^{1,1}$ background in [75, 76] through an analysis of the Laplacian and in [69] in terms of the Penrose limit.

Before starting, however, let us recall that there are more protected short operators than the chiral primaries (namely those annihilated by $\bar{\mathcal{D}}$). These are the conserved currents, which are annihilated by $\bar{\mathcal{D}}^2$ and \mathcal{D}^2 and thus satisfy shortening conditions as well. Their conformal dimension is independent of the coupling, but for them $\Delta \neq \frac{3}{2}Q_R$.

In general, our analysis leads to a proposal for the structure of the generic scalar operators (built out of the bifundamental fields) dual to supergravity states. The scaling dimension of these operators should thus be independent of the conformal couplings, at least in the large N limit. For these operators we are able to provide the 3 Abelian charges, but not the precise scaling dimension. It would be interesting to match the counting of these states from the gauge theory and the gravity point of view, performing an analysis of the Laplacian spectrum on the $\mathcal{Y}^{p,q}$.¹²

6.1 Protected building blocks

We consider in this subsection the building blocks, or 'minimal' operators. Let us start from a simple, well known, example. In the special case of the conifold, which is also $\mathcal{Y}^{1,0}$, minimal operators are quadratic in the bifundamental fields. More precisely, all bilinear gauge invariant operators (except the Konishi operator) of the conifold field theory are protected, and can be recognized as 4 chiral operators of the form $tr(AB)$, 4 antichiral

¹²Note added: after this work appeared, some properties of the general Laplacian spectrum for the $\mathcal{Y}^{p,q}$ s have been studied in [77].

operators $tr(\bar{B}\bar{A})$, and 7 real operators, $tr(A\bar{A} + \bar{B}B)$, that are part of supermultiplets containing the conserved currents of the global non-R symmetry $SU(2) \times SU(2) \times U(1)_B$.

For general p and q , the simplest protected operators satisfying $\Delta > 3/2Q_R$ are, as above, conserved currents, that are easy to describe. The global symmetries of the $\mathcal{Y}^{p,q}$ quivers are $SU(2) \times U(1)_F \times U(1)_B \times U(1)_R$, so there are six conserved currents, whose dimension on the full conformal manifold is 3. For the non-R symmetries, these currents are part of real multiplets \mathcal{K} , quadratic in the bifundamental fields, satisfying the condition

$$\mathcal{D}^2\mathcal{K} = \bar{\mathcal{D}}^2\mathcal{K} = 0 \tag{6.1}$$

and can be easily written down explicitly, using table 1

$$\mathcal{K}_{SU(2)}^I = \sum_{i=1} \sigma_{\alpha\beta}^I (U_i^\alpha \bar{U}_i^\beta + V_i^\alpha \bar{V}_{i+1}^\beta) \tag{6.2}$$

$$\mathcal{K}_F = \sum_{i=1} (Z_i \bar{Z}_i - Y_i \bar{Y}_i + V_i^1 \bar{V}_{i+1}^1 + V_i^2 \bar{V}_{i+1}^2) \tag{6.3}$$

(The baryonic current has a very similar structure). These protected operators have vanishing values for Q_R and Q_F , their scaling dimension Δ is 2. The $SU(2)$ -current has $J = 1$, so there is one operator with vanishing spin- z : $P_\varphi = 0$.

Also here we see a generic feature of toric superconformal quivers: there are always two uncharged flavor currents, corresponding to the two non-R $U(1)$ isometries of the toric Sasaki–Einstein manifold. In the case of the $\mathcal{Y}^{p,q}$ this generic isometry is enhanced to $SU(2) \times U(1)_F$, and there are two more length-2 protected operators. This is precisely analog to the situation of section 3.1, note indeed that the two currents \mathcal{K}^\pm wind around the short homology cycle of the torus.

Up to now we exhibited a class of operators satisfying a shortening condition, that are thus protected by the superconformal algebra. Their BPS conditions are very well known in 4D superconformal field theories. Now we propose an extension of this class.

Let us start from the long chiral operator \mathcal{L}_+ , of the form $UVUVUVUZUZU \dots$. Now substitute a piece UV , or VU , or UZU , with the 'nearby' antichiral operator \bar{Y} and symmetrize this 'impurity' all over the quiver. To be explicit, in the case of $\mathcal{Y}^{4,3}$, one passes from

$$\mathcal{L}_+ = tr(UVUVUVUZ) \tag{6.4}$$

to

$$\bar{Y}UVUVUZ + U\bar{Y}VUVUZ + UV\bar{Y}UVUZ + \tag{6.5}$$

$$UVU\bar{Y}VUZ + UVUV\bar{Y}UZ + UVUVU\bar{Y}Z + VUVUV\bar{Y} \tag{6.6}$$

Notice that this new operator is not BPS. It is clear that it is minimal, the only way to have a gauge invariant operator is to take one single trace.

Our proposal is that, if the position of the impurity, or 'shortcut', is symmetrized over the quiver and the $SU(2)$ spin J is taken to be the largest possible, these are precisely the operators that correspond to single particle AdS_5 supergravity states, and should thus be

Meson	spin	J	Q_F	Q_R	Δ
\mathcal{S}	1	0	0	2	3
\mathcal{K}_F	0	0	0	0	2
$\mathcal{K}_{SU(2)}$	1	0	0	0	2
$\mathcal{L}_{+,n}$	$\frac{p+q}{2} - n$	$+p$	$p + q - \frac{1}{3\ell} - 2n$?	?

Table 4: Charge assignments for some more general building blocks. The first three lines satisfy shortening conditions.

protected at least in the large N limit. It should be possible to see their duals by studying the scalar Laplacian on the $\mathcal{Y}^{p,q}$ manifolds, as has been done for $\mathcal{Y}^{1,0}$ in [75, 76].

This new operator can be thought of as \mathcal{L}_+ 'divided' by the short chiral loop \mathcal{S}^I and multiplied by the conserved current \mathcal{K}_F . So the R-charge is $Q_R[\mathcal{L}_+] - 2$, the total spin is $J[\mathcal{L}_+] - 1$, while Q_F does not change.

It should be clear now how to add more 'shortcuts' to our original BPS operator \mathcal{L}_+ , and generalize it to $\mathcal{L}_{+,n}$, where n is the number of shortcuts. In order to have a protected operator one has to fully symmetrize over the positions of the impurities and take the maximal $SU(2)$ spin. The values of the 3 commuting $U(1)$ charges simply adds, while the scaling dimensions, that we are not able to determine, should depend non linearly on n (for $q < p$).

In the case of \mathcal{L}_- the length of the operators, the values of J and Q_R can increase or decrease: one can replace a piece UY_c with a \bar{V} , or a Y_q with a piece $\bar{U}\bar{Z}\bar{U}$.

A similar procedure can be applied to the antichiral versions of \mathcal{L}_\pm . It is non trivial that in this way one can interpolate between chiral and antichiral operators.

In table 4 we give a list of the operators discussed.

Let us emphasize again that we don't have a field theoretical proof of the fact that the operators we discussed are protected. We can however check that in the case of $q = p$ and $q = 0$ this is actually the case.

For $p = q$ (the quiver becomes an orbifold of the $\mathcal{N} = 4$ SYM), it is easy to verify that our set of operators are precisely the orbifold descendant of the well known 1/2 BPS chiral primaries of $\mathcal{N} = 4$ SYM (notice indeed the 1/2 BPS operators in $\mathcal{N} = 4$ SYM are more than the chiral operator of a chosen $\mathcal{N} = 1$ parameterization).

Also in the case of $p = 0$ (the quiver becomes an orbifold of the conifold) the set of operators we proposed fits the set of protected operators of the mother theory, that are known from the spectrum of the scalar Laplacian on the conifold [75, 76].

6.2 Near BPS massless geodesics

In the previous subsection we proposed a set of minimal protected operators significantly larger than the set of minimal BPS operators. With these building blocks one can construct a lot of long operators. As for the BPS case, one has to symmetrize the impurities all over the quiver and all over the trace, and take the maximal $SU(2)$ spin. We can use all the

minimal operators, both the conserved currents and the $\mathcal{L}_{\pm,n}$ operators. Notice that these symmetrizations imply that the operators are not localized in a particular point of the quiver. This has to be the case if one wants to compare with the geometry: for instance, taking orbifolds the quivers become bigger, while the number of supergravity states does not increase at all.

Taking the limit of long operators one finds operators with constant densities of the three $U(1)$ charges (because of the symmetrizations), so the operators corresponds to non BPS geodesics.

From the gauge theory side, it is clear how to find the values of the 3 commuting $U(1)$ charges, while we do not know what is the precise value of the scaling dimensions Δ . We can however give a quantitative treatment in the case of a small number of excitations around a long BPS operator. Let us consider a BPS geodesic with $Q_F > 0$ and add one, symmetrized, impurity. We want to understand the change in the scaling dimension Δ . For $p = q$ it is obvious that, since the length of the operator decreases by 1 unit, $\delta\Delta = -1$. For $p = 0$ the length of the operator changes by 2 units, and we know from [69] that, in the limit of infinite length, $\delta\Delta = -1$. This can be obtained from the formulas for the Laplacian on $T^{1,1}$ [75], taking the limit of large R-charge with a fixed number of 'impurities' or 'shortcuts' [69]. Imposing monotonicity in q for $\delta\Delta$, one concludes that for any q the change in the scaling dimension induced by one symmetrized shortcut is precisely -1 .

In the limit of operators of infinite length, satisfying a near BPS condition, we can thus find the scaling dimensions of our operators. (Note that this is similar to the BMN limit, but much simpler, since we are sticking to operators symmetrized over the trace.)

We can thus proceed and consider all the various oscillations leading from a BPS geodesic to a near BPS geodesic. All these impurities do not wind around the quiver, so the value of Q_F does not change.

- Adding or removing a chiral \mathcal{S} operator simply changes a little bit the position of the geodesic (y_0 and θ_0 values). This gives $\delta\Delta = \pm 3$, $\delta Q_R = \pm 2$, $\delta J = \pm 1$.
- Adding a \mathcal{K}_F current gives $\delta\Delta = 2$, $\delta Q_R = 0$, $\delta J = 0$.
- Adding a $\mathcal{K}_{SU(2)}$ current gives $\delta\Delta = 2$, $\delta Q_R = 0$, $\delta J = 1$.
- Adding a 'shortcut' gives $\delta\Delta = \pm 1$, $\delta Q_R = \pm 2$, $\delta J = \pm 1, 0$.

Notice that the addition of a shortcut can be thought of as a combination of a removal of a \mathcal{S} and an addition of a \mathcal{K} , or viceversa.

6.2.1 Near BPS massless geodesics from the geometries

Now we want to study deviations from the BPS geodesics, namely when $\Delta > \frac{3}{2} Q_R$. We recall eq. (2.28):

$$\Delta^2 = \left(\frac{3}{2} Q_R\right)^2 + \frac{1}{6p(y)} (P_\alpha + 3y Q_R)^2 + 6p(y) P_y^2 + \frac{6}{1-y} (J^2 - P_\psi^2) \quad (6.7)$$

Appropriately quantizing this Hamiltonian is equivalent to solving the Laplacian operator on the $\mathcal{Y}^{p,q}$ manifolds. Here we study small perturbations around a BPS geodesic. There are two non trivial perturbations, one leading to $J > P_\psi$ and one to $P_y \neq 0$.

In the first case ($J > P_\psi$) we have

$$\Delta\delta\Delta = \frac{6}{1-y_0} J\delta J = 3Q_R\delta J = 2\Delta\delta J, \Rightarrow \delta\Delta = 2\delta J \quad (6.8)$$

where we used the relations (2.32) and (2.33) valid for the unperturbed geodesic.

In the second case ($P_y \neq 0$) we perturb y away from the minimum $y = y_0 + \delta y$ and get a Hamiltonian for the perturbation

$$H = \frac{1}{2}\Delta^2 = \frac{1}{2}\left(\frac{3}{2}Q_R\right)^2 + \frac{1}{2}\left[6p(y_0)P_{\delta y}^2 + \frac{3Q_R^2}{2p(y_0)}\delta y^2\right] \quad (6.9)$$

This is a standard harmonic oscillator with mass $m = \frac{1}{6p(y_0)}$ and angular frequency $\omega = 3Q_R$. This means that there are classical geodesics that oscillate in the y direction around the BPS one. From the worldsheet point of view, these oscillations should be quantized. This leads to

$$\Delta^2 = \left(\frac{3}{2}Q_R\right)^2 + 2n\omega \quad (6.10)$$

For a small variation we therefore get

$$\Delta\delta\Delta = n\omega, \Rightarrow \delta\Delta = \frac{\omega}{\Delta}n = \frac{3Q_R}{\Delta}n = 2n \quad (6.11)$$

To summarize, we found two non trivial types of perturbations characterized by:

$$\begin{aligned} I) \quad & \delta\Delta = 2n \quad \delta Q_R = 0 \quad \delta P_\alpha = 0 \quad \delta J = n \\ II) \quad & \delta\Delta = 2n \quad \delta Q_R = 0 \quad \delta P_\alpha = 0 \quad \delta J = 0 \end{aligned} \quad (6.12)$$

It is straightforward to see that combining these fluctuations with BPS fluctuations, that do not change $\Delta - 3/2Q_R$, one gets precisely the fluctuations found on the quiver side.

7. Conclusions

We have described the computation of a set of chiral primary operators in the $\mathcal{Y}^{p,q}$ quiver gauge theories. Those operators were successfully matched to massless geodesics in the corresponding supergravity backgrounds. The matching gives the interpretation of the coordinate y in the bulk as the ratio between the $U(1)_F$ charge and the R -charge of an operator (precisely $P_\alpha = -3yQ_R$). From the analysis of the operators one can find the maximum and minimum values of such ratio. They agree precisely with y_2 and y_1 as expected from the bulk. Small fluctuation around the BPS geodesics were identified with the insertion of conserved currents associated with the global charges.

After that we analyzed very long operators. Such operators correspond to long loops in the quiver. The matrix of anomalous dimensions, induced by the superpotential has a simple description in term of moves that convert one path into another. Diagonalizing the

matrix of anomalous dimensions reduces to the study of the dynamics of such paths. We constructed a simple model which we argued has the same behavior for long paths, namely in the continuum limit. Using the coherent state method we obtained a classical action which is similar but not the same as the one obtained from a limit of the string action. We suggest that in the infrared limit (in the sense of the spin chain) the action we found flows to the one from the bulk but we leave that point for future investigation.

In any case it is encouraging that in these more complicated cases the string action can be reproduced at least in part by an analysis of the operators in the gauge theory.

Acknowledgments

We are very grateful to Guido Festuccia and especially Carlos Nuñez for collaboration in the initial stages of this project as well as for numerous suggestions and discussions. We thank Amihay Hanany, Igor Klebanov, Albion Lawrence, Antonello Scardicchio, Matt Strassler and Arkady Tseytlin for useful comments.

S. B. is grateful to MIT for kind hospitality during the last stages of this work.

The work of M. K. is supported in part by NSF through grants PHY-0331516, PHY99-73935 and DOE under grant DE-FG02-92ER40706.

A. Useful formulas

Throughout the paper we used various relations that do not belong to any specific section. We decide to collect them here in the hope that can be useful to reproduce some of the calculations. The definitions of the functions and constant involved can be found in the main text.

Relating f, q, w :

$$f(y) - \frac{1}{6} = \frac{2}{3} \frac{y}{w(y)} \tag{A.1}$$

$$1 - y + 6yf(y) = q(y) \tag{A.2}$$

Relating $y_{1,2,3}$ to p, q :

$$y_2 - y_1 = \frac{3q}{2p} \tag{A.3}$$

$$p\ell = \frac{y_1 - y_2}{6y_1y_2} = -\frac{q}{4py_1y_2} \tag{A.4}$$

$$Q_R(\mathcal{L}_+) = -\frac{1}{3y_1\ell} = \frac{2py_2}{y_2 - y_1} \tag{A.5}$$

$$Q_R(\mathcal{L}_-) = \frac{1}{3y_2\ell} = -\frac{2py_1}{y_2 - y_1} \tag{A.6}$$

$$y_1 + y_2 + y_3 = \frac{3}{2} \tag{A.7}$$

$$y_1y_2 + y_1y_3 + y_2y_3 = 0 \Rightarrow \frac{1}{y_1} + \frac{1}{y_2} + \frac{1}{y_3} = 0 \tag{A.8}$$

References

- [1] J.M. Maldacena, *The large- N limit of superconformal field theories and supergravity*, *Adv. Theor. Math. Phys.* **2** (1998) 231 [[hep-th/9711200](#)];
S.S. Gubser, I.R. Klebanov and A.M. Polyakov, *Gauge theory correlators from non-critical string theory*, *Phys. Lett.* **B 428** (1998) 105 [[hep-th/9802109](#)];
E. Witten, *Anti-de Sitter space and holography*, *Adv. Theor. Math. Phys.* **2** (1998) 253 [[hep-th/9802150](#)];
O. Aharony, S.S. Gubser, J.M. Maldacena, H. Ooguri and Y. Oz, *Large- N field theories, string theory and gravity*, *Phys. Rept.* **323** (2000) 183 [[hep-th/9905111](#)].
- [2] G. 't Hooft, *A planar diagram theory for strong interactions*, *Nucl. Phys.* **B 72** (1974) 461; *A two-dimensional model for mesons*, *Nucl. Phys.* **B 75** (1974) 461.
- [3] A.M. Polyakov, *Gauge fields and space-time*, *Int. J. Mod. Phys.* **A17S1** (2002) 119–136 [[hep-th/0110196](#)].
- [4] D. Berenstein, J.M. Maldacena and H. Nastase, *Strings in flat space and pp waves from $N = 4$ super Yang-Mills*, *JHEP* **04** (2002) 013 [[hep-th/0202021](#)].
- [5] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, *A semi-classical limit of the gauge/string correspondence*, *Nucl. Phys.* **B 636** (2002) 99 [[hep-th/0204051](#)].
- [6] S. Frolov and A.A. Tseytlin, *Semiclassical quantization of rotating superstring in $AdS_5 \times S^5$* , *JHEP* **06** (2002) 007 [[hep-th/0204226](#)].
- [7] S. Frolov and A.A. Tseytlin, *Multi-spin string solutions in $AdS_5 \times S^5$* , *Nucl. Phys.* **B 668** (2003) 77 [[hep-th/0304255](#)].
- [8] A.A. Tseytlin, *Semiclassical quantization of superstrings: $AdS_5 \times S^5$ and beyond*, *Int. J. Mod. Phys.* **A 18** (2003) 981 [[hep-th/0209116](#)].
- [9] A.A. Tseytlin, *Spinning strings and AdS/CFT duality*, [hep-th/0311139](#).
- [10] H.J. de Vega, A.L. Larsen and N.G. Sanchez, *Semiclassical quantization of circular strings in de Sitter and anti-de Sitter space-times*, *Phys. Rev.* **D 51** (1995) 6917 [[hep-th/9410219](#)].
- [11] J.A. Minahan and K. Zarembo, *The bethe-ansatz for $N = 4$ super Yang-Mills*, *JHEP* **03** (2003) 013 [[hep-th/0212208](#)].
- [12] N. Beisert, J.A. Minahan, M. Staudacher and K. Zarembo, *Stringing spins and spinning strings*, *JHEP* **09** (2003) 010 [[hep-th/0306139](#)].
- [13] N. Beisert, S. Frolov, M. Staudacher and A.A. Tseytlin, *Precision spectroscopy of AdS/CFT*, *JHEP* **10** (2003) 037 [[hep-th/0308117](#)].
- [14] G. Arutyunov, S. Frolov, J. Russo and A.A. Tseytlin, *Spinning strings in $AdS_5 \times S^5$ and integrable systems*, *Nucl. Phys.* **B 671** (2003) 3 [[hep-th/0307191](#)].
- [15] G. Arutyunov, J. Russo and A.A. Tseytlin, *Spinning strings in $AdS_5 \times S^5$: new integrable system relations*, *Phys. Rev.* **D 69** (2004) 086009 [[hep-th/0311004](#)].
- [16] G. Arutyunov and M. Staudacher, *Matching higher conserved charges for strings and spins*, *JHEP* **03** (2004) 004 [[hep-th/0310182](#)]; *Two-loop commuting charges and the string /gauge duality*, [hep-th/0403077](#).
- [17] J. Engquist, J.A. Minahan and K. Zarembo, *Yang-Mills duals for semiclassical strings on $AdS_5 \times S^5$* , *JHEP* **11** (2003) 063 [[hep-th/0310188](#)].

- [18] M. Kruczenski, *Spin chains and string theory*, *Phys. Rev. Lett.* **93** (2004) 161602 [[hep-th/0311203](#)].
- [19] A. Mikhailov, *Speeding strings*, *JHEP* **12** (2003) 058 [[hep-th/0311019](#)]; *Slow evolution of nearly-degenerate extremal surfaces*, *J. Geom. Phys.* **54** (2005) 228 [[hep-th/0402067](#)]; *Supersymmetric null-surfaces*, *JHEP* **09** (2004) 068 [[hep-th/0404173](#)].
- [20] A. Gorsky, *Spin chains and gauge/string duality*, *Theor. Math. Phys.* **142** (2005) 153 [[hep-th/0308182](#)].
- [21] D. Mateos, T. Mateos and P.K. Townsend, *Supersymmetry of tensionless rotating strings in $AdS_5 \times S^5$ and nearly-BPS operators*, *JHEP* **12** (2003) 017 [[hep-th/0309114](#)]; *More on supersymmetric tensionless rotating strings in $AdS_5 \times S^5$* , [hep-th/0401058](#).
- [22] V.A. Kazakov, A. Marshakov, J.A. Minahan and K. Zarembo, *Classical/quantum integrability in AdS/CFT* , *JHEP* **05** (2004) 024 [[hep-th/0402207](#)].
- [23] M. Kruczenski, A.V. Ryzhov and A.A. Tseytlin, *Large spin limit of $AdS_5 \times S^5$ string theory and low energy expansion of ferromagnetic spin chains*, *Nucl. Phys. B* **692** (2004) 3 [[hep-th/0403120](#)].
- [24] R. Hernandez and E. Lopez, *The $SU(3)$ spin chain sigma model and string theory*, *JHEP* **04** (2004) 052 [[hep-th/0403139](#)].
- [25] J. Stefanski, B. and A.A. Tseytlin, *Large spin limits of AdS/CFT and generalized Landau-Lifshitz equations*, *JHEP* **05** (2004) 042 [[hep-th/0404133](#)].
- [26] H. Dimov and R.C. Rashkov, *A note on spin chain/string duality*, *Int. J. Mod. Phys. A* **20** (2005) 4337 [[hep-th/0403121](#)].
- [27] M. Kruczenski and A.A. Tseytlin, *Semiclassical relativistic strings in S^5 and long coherent operators in $N = 4$ SYM theory*, *JHEP* **09** (2004) 038 [[hep-th/0406189](#)].
- [28] R. Hernandez and E. Lopez, *Spin chain sigma models with fermions*, *JHEP* **11** (2004) 079 [[hep-th/0410022](#)].
- [29] M. Kruczenski, *Spiky strings and single trace operators in gauge theories*, *JHEP* **08** (2005) 014 [[hep-th/0410226](#)].
- [30] J. Stefanski, B. and A.A. Tseytlin, *Super spin chain coherent state actions and $AdS_5 \times S^5$ superstring*, *Nucl. Phys. B* **718** (2005) 83 [[hep-th/0503185](#)].
- [31] S. Bellucci, P.Y. Casteill and J.F. Morales, *Superstring sigma models from spin chains: the $SU(1,1|1)$ case*, *Nucl. Phys. B* **729** (2005) 163 [[hep-th/0503159](#)].
- [32] D. Berenstein, D.H. Correa and S.E. Vazquez, *Quantizing open spin chains with variable length: an example from giant gravitons*, *Phys. Rev. Lett.* **95** (2005) 191601 [[hep-th/0502172](#)];
Y. Susaki, Y. Takayama and K. Yoshida, *Open semiclassical strings and long defect operators in $AdS/dCFT$ correspondence*, *Phys. Rev. D* **71** (2005) 126006 [[hep-th/0410139](#)].
- [33] N. Beisert, A.A. Tseytlin and K. Zarembo, *Matching quantum strings to quantum spins: one-loop vs. finite-size corrections*, *Nucl. Phys. B* **715** (2005) 190 [[hep-th/0502173](#)];
R. Hernandez, E. Lopez, A. Perianez and G. Sierra, *Finite size effects in ferromagnetic spin chains and quantum corrections to classical strings*, *JHEP* **06** (2005) 011 [[hep-th/0502188](#)];
I.Y. Park, A. Tirziu and A.A. Tseytlin, *Spinning strings in $AdS_5 \times S^5$: one-loop correction to energy in $SL(2)$ sector*, *JHEP* **03** (2005) 013 [[hep-th/0501203](#)];

- S.A. Frolov, I.Y. Park and A.A. Tseytlin, *On one-loop correction to energy of spinning strings in S^5* , *Phys. Rev. D* **71** (2005) 026006 [[hep-th/0408187](#)];
H. Fuji and Y. Satoh, *Quantum fluctuations of rotating strings in $AdS_5 \times S^5$* , *Int. J. Mod. Phys. A* **21** (2006) 3673 [[hep-th/0504123](#)].
- [34] K. Peeters, J. Plefka and M. Zamaklar, *Splitting strings and chains*, *Fortschr. Phys.* **53** (2005) 640 [[hep-th/0501165](#)]; *Splitting spinning strings in AdS/CFT* , *JHEP* **11** (2004) 054 [[hep-th/0410275](#)].
- [35] V.M. Braun, S.E. Derkachov and A.N. Manashov, *Integrability of three-particle evolution equations in QCD*, *Phys. Rev. Lett.* **81** (1998) 2020 [[hep-ph/9805225](#)];
V.M. Braun, S.E. Derkachov, G.P. Korchemsky and A.N. Manashov, *Baryon distribution amplitudes in QCD*, *Nucl. Phys. B* **553** (1999) 355 [[hep-ph/9902375](#)];
A.V. Belitsky, *Integrability and WKB solution of twist-three evolution equations*, *Nucl. Phys. B* **558** (1999) 259 [[hep-ph/9903512](#)]; *Fine structure of spectrum of twist-three operators in QCD*, *Phys. Lett. B* **453** (1999) 59 [[hep-ph/9902361](#)].
- [36] J.M. Maldacena and C. Núñez, *Towards the large- N limit of pure $N = 1$ super Yang-Mills*, *Phys. Rev. Lett.* **86** (2001) 588 [[hep-th/0008001](#)].
- [37] A. Butti, M. Grana, R. Minasian, M. Petrini and A. Zaffaroni, *The baryonic branch of Klebanov-Strassler solution: a supersymmetric family of $SU(3)$ structure backgrounds*, *JHEP* **03** (2005) 069 [[hep-th/0412187](#)].
- [38] S.S. Gubser, C.P. Herzog and I.R. Klebanov, *Variations on the warped deformed conifold*, *Comptes Rendus Physique* **5** (2004) 1031–1038 [[hep-th/0409186](#)].
- [39] I.R. Klebanov and E. Witten, *Superconformal field theory on threebranes at a Calabi-Yau singularity*, *Nucl. Phys. B* **536** (1998) 199 [[hep-th/9807080](#)].
- [40] J.P. Gauntlett, D. Martelli, J. Sparks and D. Waldram, *Supersymmetric AdS_5 solutions of M-theory*, *Class. and Quant. Grav.* **21** (2004) 4335 [[hep-th/0402153](#)].
- [41] J.P. Gauntlett, D. Martelli, J. Sparks and D. Waldram, *Sasaki-einstein metrics on $S^2 \times S^3$* , *Adv. Theor. Math. Phys.* **8** (2004) 711 [[hep-th/0403002](#)].
- [42] J.P. Gauntlett, D. Martelli, J.F. Sparks and D. Waldram, *A new infinite class of Sasaki-Einstein manifolds*, *Adv. Theor. Math. Phys.* **8** (2006) 987 [[hep-th/0403038](#)].
- [43] D. Martelli and J. Sparks, *Toric geometry, Sasaki-Einstein manifolds and a new infinite class of AdS/CFT duals*, *Commun. Math. Phys.* **262** (2006) 51 [[hep-th/0411238](#)].
- [44] S. Benvenuti, S. Franco, A. Hanany, D. Martelli and J. Sparks, *An infinite family of superconformal quiver gauge theories with Sasaki-Einstein duals*, *JHEP* **06** (2005) 064 [[hep-th/0411264](#)].
- [45] K. Intriligator and B. Wecht, *The exact superconformal R -symmetry maximizes a* , *Nucl. Phys. B* **667** (2003) 183 [[hep-th/0304128](#)].
- [46] D. Anselmi, D.Z. Freedman, M.T. Grisaru and A.A. Johansen, *Nonperturbative formulas for central functions of supersymmetric gauge theories*, *Nucl. Phys. B* **526** (1998) 543 [[hep-th/9708042](#)]; *Positivity constraints on anomalies in supersymmetric gauge theories*, *Phys. Rev. D* **57** (1998) 7570 [[hep-th/9711035](#)].
- [47] D. Martelli, J. Sparks and S.-T. Yau, *The geometric dual of a -maximisation for toric sasaki-Einstein manifolds*, [hep-th/0503183](#).

- [48] A. Hanany and K.D. Kennaway, *Dimer models and toric diagrams*, hep-th/0503149.
- [49] S. Franco, A. Hanany, K.D. Kennaway, D. Vegh and B. Wecht, *Brane dimers and quiver gauge theories*, *JHEP* **01** (2006) 096 [hep-th/0504110].
- [50] A. Hanany and E. Witten, *Type IIB superstrings, BPS monopoles and three-dimensional gauge dynamics*, *Nucl. Phys. B* **492** (1997) 152 [hep-th/9611230].
- [51] A. Okounkov, N. Reshetikhin and C. Vafa, *Quantum Calabi-Yau and classical crystals*, hep-th/0309208.
- [52] O. Aharony and A. Hanany, *Branes, superpotentials and superconformal fixed points*, *Nucl. Phys. B* **504** (1997) 239 [hep-th/9704170];
O. Aharony, A. Hanany and B. Kol, *Webs of (p,q) 5-branes, five dimensional field theories and grid diagrams*, *JHEP* **01** (1998) 002 [hep-th/9710116].
- [53] A. Hanany and A. Zaffaroni, *On the realization of chiral four-dimensional gauge theories using branes*, *JHEP* **05** (1998) 001 [hep-th/9801134];
A. Hanany and A.M. Uranga, *Brane boxes and branes on singularities*, *JHEP* **05** (1998) 013 [hep-th/9805139];
M. Aganagic, A. Karch, D. Lüüst and A. Miemiec, *Mirror symmetries for brane configurations and branes at singularities*, *Nucl. Phys. B* **569** (2000) 277 [hep-th/9903093];
B. Feng, A. Hanany, Y.-H. He and A.M. Uranga, *Toric duality as Seiberg duality and brane diamonds*, *JHEP* **12** (2001) 035 [hep-th/0109063].
- [54] R.G. Leigh and M.J. Strassler, *Exactly marginal operators and duality in four-dimensional $N = 1$ supersymmetric gauge theory*, *Nucl. Phys. B* **447** (1995) 95 [hep-th/9503121].
- [55] S. Benvenuti and A. Hanany, *Conformal manifolds for the conifold and other toric field theories*, *JHEP* **08** (2005) 024 [hep-th/0502043].
- [56] B. Kol, *On conformal deformations*, *JHEP* **09** (2002) 046 [hep-th/0205141].
- [57] O. Lunin and J.M. Maldacena, *Deforming field theories with $U(1) \times U(1)$ global symmetry and their gravity duals*, *JHEP* **05** (2005) 033 [hep-th/0502086].
- [58] S.A. Frolov, R. Roiban and A.A. Tseytlin, *Gauge - string duality for superconformal deformations of $N = 4$ super Yang-Mills theory*, *JHEP* **07** (2005) 045 [hep-th/0503192].
- [59] S. Frolov, *Lax pair for strings in Lunin-Maldacena background*, *JHEP* **05** (2005) 069 [hep-th/0503201].
- [60] I.R. Klebanov and N.A. Nekrasov, *Gravity duals of fractional branes and logarithmic RG flow*, *Nucl. Phys. B* **574** (2000) 263 [hep-th/9911096];
I.R. Klebanov and A.A. Tseytlin, *Gravity duals of supersymmetric $SU(N) \times SU(N + M)$ gauge theories*, *Nucl. Phys. B* **578** (2000) 123 [hep-th/0002159];
I.R. Klebanov and M.J. Strassler, *Supergravity and a confining gauge theory: duality cascades and χ SB-resolution of naked singularities*, *JHEP* **08** (2000) 052 [hep-th/0007191].
- [61] C.P. Herzog, Q.J. Ejaz and I.R. Klebanov, *Cascading RG flows from new sasaki-einstein manifolds*, *JHEP* **02** (2005) 009 [hep-th/0412193].
- [62] S.S. Pal, *A new Ricci flat geometry*, *Phys. Lett. B* **614** (2005) 201 [hep-th/0501012].
- [63] B.A. Burington, J.T. Liu, M. Mahato and L.A. Pando Zayas, *Towards supergravity duals of chiral symmetry breaking in Sasaki-Einstein cascading quiver theories*, *JHEP* **07** (2005) 019 [hep-th/0504155].

- [64] D. Berenstein, C.P. Herzog, P. Ouyang and S. Pinansky, *Supersymmetry breaking from a Calabi-Yau singularity*, *JHEP* **09** (2005) 084 [[hep-th/0505029](#)].
- [65] S. Franco, A. Hanany, F. Saad and A.M. Uranga, *Fractional branes and dynamical supersymmetry breaking*, *JHEP* **01** (2006) 011 [[hep-th/0505040](#)].
- [66] B. Feng, A. Hanany and Y.-H. He, *D-brane gauge theories from toric singularities and toric duality*, *Nucl. Phys.* **B 595** (2001) 165 [[hep-th/0003085](#)].
- [67] M. Bertolini, F. Bigazzi and A.L. Cotrone, *New checks and subtleties for AdS/CFT and a-maximization*, *JHEP* **12** (2004) 024 [[hep-th/0411249](#)].
- [68] S. Benvenuti, A. Hanany and P. Kazakopoulos, *The toric phases of the $Y(p,q)$ quivers*, *JHEP* **07** (2005) 021 [[hep-th/0412279](#)].
- [69] J. Gomis and H. Ooguri, *Penrose limit of $N = 1$ gauge theories*, *Nucl. Phys.* **B 635** (2002) 106 [[hep-th/0202157](#)];
N. Itzhaki, I.R. Klebanov and S. Mukhi, *Pp wave limit and enhanced supersymmetry in gauge theories*, *JHEP* **03** (2002) 048 [[hep-th/0202153](#)];
L.A. Pando Zayas and J. Sonnenschein, *On Penrose limits and gauge theories*, *JHEP* **05** (2002) 010 [[hep-th/0202186](#)].
- [70] N. Seiberg, *Electric-magnetic duality in supersymmetric nonAbelian gauge theories*, *Nucl. Phys.* **B 435** (1995) 129 [[hep-th/9411149](#)].
- [71] S. Benvenuti and A. Hanany, *New results on superconformal quivers*, [hep-th/0411262](#).
- [72] D. Kutasov, *New results on the 'a-theorem' in four dimensional supersymmetric field theory*, [hep-th/0312098](#).
- [73] N. Beisert, C. Kristjansen and M. Staudacher, *The dilatation operator of $N = 4$ super Yang-Mills theory*, *Nucl. Phys.* **B 664** (2003) 131 [[hep-th/0303060](#)].
- [74] A. Hanany, P. Kazakopoulos and B. Wecht, *A new infinite class of quiver gauge theories*, [hep-th/0503177](#).
- [75] S.S. Gubser, *Einstein manifolds and conformal field theories*, *Phys. Rev.* **D 59** (1999) 025006 [[hep-th/9807164](#)].
- [76] A. Ceresole, G. Dall'Agata, R. D'Auria and S. Ferrara, *Spectrum of type IIB supergravity on $AdS_5 \times T_{11}$: Predictions on $N = 1$ SCFT's*, *Phys. Rev.* **D 61** (2000) 066001 [[hep-th/9905226](#)];
A. Ceresole, G. Dall'Agata and R. D'Auria, *PKK spectroscopy of type IIB supergravity on $AdS_5 \times T_{11}$* , *JHEP* **11** (1999) 009 [[hep-th/9907216](#)].
- [77] H. Kihara, M. Sakaguchi and Y. Yasui, *Scalar Laplacian on Sasaki-Einstein manifolds $Y(p,q)$* , [hep-th/0505259](#).
- [78] B. Feng, S. Franco, A. Hanany and Y. H. He, *Symmetries of toric duality*, *JHEP* **12** (2002) 076 [[hep-th/0205144](#)].
- [79] J. A. Minahan, *Higher loops beyond the $SU(2)$ sector*, *JHEP* **10** (2004) 053 [[hep-th/0405243](#)].