

Heavy quark effective theory computation of the mass of the bottom quark

Michele Della Morte,^a Nicolas Garron,^b Mauro Papinutto^c and Rainer Sommer^b

^a*Institut für Physik, Humboldt Universität
Newtonstr. 15, 12489 Berlin, Germany*

^b*DESY
Platanenallee 6, 15738 Zeuthen, Germany*

^c*INFN Sezione di Roma Tre
Via della Vasca Navale 84, I-00146, Rome, Italy*

*E-mail: dellamor@mail.cern.ch, garron@ifh.de, Mauro.Papinutto@desy.de,
sommer@ifh.de*

ABSTRACT: We present a fully non-perturbative computation of the mass of the b-quark in the quenched approximation. Our strategy starts from the matching of HQET to QCD in a finite volume and finally relates the quark mass to the spin averaged mass of the B_s meson in HQET. All steps include the terms of order Λ^2/m_b . Expanding on [1], we discuss the computation and renormalization of correlation functions at order $1/m_b$. With the strange quark mass fixed from the Kaon mass and the QCD scale set through $r_0 = 0.5$ fm, we obtain a renormalization group invariant mass $M_b = 6.758(86)$ GeV or $\bar{m}_b(\bar{m}_b) = 4.347(48)$ GeV in the $\overline{\text{MS}}$ scheme. The uncertainty in the computed Λ^2/m_b terms contributes little to the total error and Λ^3/m_b^2 terms are negligible. The strategy is promising for full QCD as well as for other B-physics observables.

KEYWORDS: Quark Masses and SM Parameters, Heavy Quark Physics, Lattice QCD, Nonperturbative Effects.

Contents


1. Introduction	1
2. Heavy quark effective theory on the lattice	3
2.1 Formulation	3
2.2 $1/m_b$ expansion in a geometry without boundaries	5
2.3 Schrödinger functional correlation functions	7
3. Finite volume observables and step scaling functions	9
3.1 Observables	9
3.2 Step scaling functions	10
4. M_b including $1/m_b$ corrections	11
5. Results	12
5.1 QCD observables	13
5.2 HQET step scaling functions	14
5.3 Large volume matrix elements and M_b	16
5.4 Comparison to results from an alternative strategy	18
6. Conclusions	19
A. Finite volume simulations	21
B. Large volume simulations and extraction of matrix elements	25
C. Alternative strategy	28
D. Propagating uncertainties in L_i/r_0 and $\bar{g}^2(L_i)$	29

1. Introduction

The mass of the b-quark, M_b , is a relevant input parameter for phenomenology analysis based on perturbation theory. Let us just mention the extraction of V_{ub} from inclusive b-decays [2, 3]. M_b is a fundamental parameter of the Standard Model of particle physics. Thus it should be determined precisely. One may of course turn the very first observation around. For instance, applying high order perturbation theory to sufficiently well integrated cross sections, the quark mass can be determined [4–15]. Still, the achievable precision is limited by the intrinsic uncertainty of perturbation theory and maybe more by experimental difficulties. On the other hand, the use of lattice QCD offers a strategy to compute the

fundamental renormalization group invariant (RGI) parameters of QCD with very precisely known experimental input, e.g. N_f (= number of quarks flavours) meson masses as well as the nucleon mass; see [16] for a basic introduction. However, for the b-quark mass, such a computation is more involved than for the light quarks because the achievable inverse lattice spacings are below the mass of the quark. Effective theories have to be employed in a numerical treatment of the bound states. The most serious problem that arises is that a power law divergent ($\sim g_0^2/a$) additive renormalization of the mass is present due to the absence of a chiral symmetry in the effective theories (even in the continuum). Although at the lowest order in Heavy Quark Effective Theory (HQET) the subtraction is known to order g_0^6/a [17–19], an (in the continuum limit) divergent remainder is unavoidable and the total uncertainty is difficult to estimate as long as the renormalization is carried out perturbatively.

In [1] a general strategy was described which allows HQET at zero velocity to be implemented non-perturbatively on the lattice, including all renormalizations.



The basic idea is illustrated in the above diagram. It is founded on the knowledge of the relation between the RGI mass and the bare mass in QCD [20, 21]. In a finite volume of extent $L_1 \approx 0.4$ fm, one chooses lattice spacings a sufficiently smaller than $1/m_b$, such that the b-quark propagates correctly up to controllable discretization errors of order a^2 . Finite volume observables $\Phi_i(L_1, M_b)$ may then be computed as a function of the RGI mass M_b including an extrapolation to the continuum limit. The resulting values are equated to their representation in HQET — a step called matching, indicated by the r.h.s. of the diagram. Choosing now $L_1/a = O(10)$, with the same physical value of L_1 , one uses the knowledge of $\Phi_i(L_1, M_b)$ to determine the bare parameters in the effective theory for a -values of about 0.025 fm to 0.05 fm. At these lattice spacings one then computes the same observables in a larger volume $L_2 = 2L_1$. Again these observables can be extrapolated to the continuum limit. Next, the knowledge of $\Phi_i(L_2, M_b)$ and the choice $L_2/a = O(10)$ yields the bare parameters of the effective theory for a around 0.05 fm to 0.1 fm. One then has full control over the effective theory at lattice spacings where large volume observables, such as the B-meson mass, can be computed. Perturbation theory is completely avoided with the power divergent subtractions being taken care of non-perturbatively.

We return to the specific application of computing M_b . The whole chain allows to express m_B in terms of $\Phi_i(L_1, M_b)$ and thus as a function of M_b . This function naturally splits into various pieces which may be computed individually as they separately have a continuum limit. In particular, the step scaling functions σ relate $\Phi_i(L_1)$ to $\Phi_i(L_2)$. As we will see below, at first order in $1/m_b$, two matching observables Φ_1, Φ_2 are sufficient if we consider the spin averaged B-meson mass.

The strategy requires all considered observables to be accurately described by the $1/m_b$ expansion. Naive counting estimates the accuracy of the quark mass as $\Lambda \times \frac{\Lambda^2}{m_b^2}$ and $\Lambda \times \frac{1}{L_1^2 m_b^2}$. For a typical QCD scale $\Lambda \approx 400 \text{ MeV}$ both these terms yield the same, very small, estimate. In [22] the $1/m_b$ expansion was tested for an even smaller $L = L_0 = L_1/2$ and found to be well behaved, as it is also the case in perturbation theory [23]. Here we will have additional cross checks by choosing different quantities Φ_i in the matching step.

In section 2 we will go through the definition of the effective theory in order to fix some notations and give rules how the $1/m_b$ expansion is implemented in practice. We also discuss correlation functions in the Schrödinger functional [24, 25], which defines our finite volume geometry. These correlation functions are then used in section 3 to form suitable dimensionless observables Φ_i , followed by a section which lists the step scaling functions. Section 4 discusses the final formula for the RGI b-quark mass M_b . Numerical results for all quantities in the quenched approximation are discussed in section 5. This includes also results from an alternative strategy as a check on the smallness of the $1/m_b^2$ terms.

2. Heavy quark effective theory on the lattice

We start from the Eichten Hill static quark Lagrangian [26], using the notation of [27], but setting the mass counter term δm to zero. Its effect is taken into account in the overall energy shift m_{bare} between the effective theory and QCD. Thus m_{bare} is regularization dependent with a $\sim g_0^2/a$ divergence. For the sake of a light notation, we also drop the superscript W [27] for the different lattice discretizations of the static Lagrangian, but in the numerical computations these different versions will be used and referred to exactly as in that reference. We remind the reader that they differ only by the choice of the covariant derivative D_0 .

The terms of first order in $1/m_b$ are introduced exactly as in [1], but we use a slightly different notation which is convenient when one does not go beyond that order.

2.1 Formulation

The lowest order (static) Lagrangian,

$$\mathcal{L}_{\text{stat}}(x) = \bar{\psi}_h(x) D_0 \psi_h(x), \tag{2.1}$$

is written in terms of the backward covariant derivative D_0 as in [27] and the 4-component heavy quark field subject to the constraints $P_+ \psi_h = \psi_h$, $\bar{\psi}_h P_+ = \bar{\psi}_h$ with $P_+ = (1 + \gamma_0)/2$. At the first order we write the HQET Lagrangian

$$\mathcal{L}_{\text{HQET}}(x) = \mathcal{L}_{\text{stat}}(x) + \mathcal{L}^{(1)}(x), \tag{2.2}$$

$$\mathcal{L}^{(1)}(x) = -\omega_{\text{spin}} \mathcal{O}_{\text{spin}}(x) - \omega_{\text{kin}} \mathcal{O}_{\text{kin}}(x), \tag{2.3}$$

$$\mathcal{O}_{\text{spin}}(x) = \bar{\psi}_h(x) \boldsymbol{\sigma} \cdot \mathbf{B} \psi_h(x) \quad \mathcal{O}_{\text{kin}} = \bar{\psi}_h(x) \mathbf{D}^2 \psi_h(x), \tag{2.4}$$

such that the *classical* values for the coefficients are $\omega_{\text{kin}} = \omega_{\text{spin}} = 1/(2m_b)$. We use the discretized version $\boldsymbol{\sigma} \cdot \mathbf{B} = \sum_{k,j} \sigma_{kj} \hat{F}_{kj}/(2i)$, with σ_{kj} and the lattice field tensor \hat{F} defined

in [28]. The kinetic term \mathbf{D}^2 is represented by the nearest neighbor covariant 3-d Laplacian. The effective theory is expected to be renormalizable at each (fixed) order in $1/m_b$ if (and only if) path integral expectation values are defined by expanding the path integral weight as [1]

$$\begin{aligned} \exp(-a^4 \sum_x [\mathcal{L}_{\text{HQET}}(x) + \mathcal{L}_{\text{light}}(x)]) &= \exp(-a^4 \sum_x [\mathcal{L}_{\text{stat}}(x) + \mathcal{L}_{\text{light}}(x)]) \\ &\times \left(1 + a^4 \sum_x [\omega_{\text{spin}} \mathcal{O}_{\text{spin}}(x) + \omega_{\text{kin}} \mathcal{O}_{\text{kin}}(x)] \right). \end{aligned} \quad (2.5)$$

For correlation functions of some multilocal fields \mathcal{O} this means

$$\langle \mathcal{O} \rangle = \langle \mathcal{O} \rangle_{\text{stat}} + \omega_{\text{kin}} a^4 \sum_x \langle \mathcal{O} \mathcal{O}_{\text{kin}}(x) \rangle_{\text{stat}} + \omega_{\text{spin}} a^4 \sum_x \langle \mathcal{O} \mathcal{O}_{\text{spin}}(x) \rangle_{\text{stat}} \quad (2.6)$$

$$\equiv \langle \mathcal{O} \rangle_{\text{stat}} + \omega_{\text{kin}} \langle \mathcal{O} \rangle_{\text{kin}} + \omega_{\text{spin}} \langle \mathcal{O} \rangle_{\text{spin}}, \quad (2.7)$$

where $\langle \mathcal{O} \rangle_{\text{stat}}$ denotes the static expectation value with Lagrangian $\mathcal{L}_{\text{stat}}(x) + \mathcal{L}_{\text{light}}(x)$. All terms composed of just the relativistic quarks and the gauge fields are summarized in $\mathcal{L}_{\text{light}}(x)$. Note that as one performs the Wick contractions of the heavy quark field, the $1/m_b$ terms $\mathcal{O}_{\text{kin}}(x), \mathcal{O}_{\text{spin}}(x)$ leave behind insertions in the static heavy quark propagators. From the point of view of renormalization all terms in eq. (2.6) are correlation functions in the static effective theory, which is power counting renormalizable.

The above form assumes that \mathcal{O} contains all $1/m_b$ terms needed to represent the local fields in the effective theory. A relevant example is the time component of the heavy light axial current. In the effective theory it is represented as

$$A_0^{\text{HQET}}(x) = Z_A^{\text{HQET}} [A_0^{\text{stat}}(x) + c_A^{\text{HQET}} \delta A_0^{\text{stat}}(x)], \quad (2.8)$$

$$A_0^{\text{stat}}(x) = \bar{\psi}_1(x) \gamma_0 \gamma_5 \psi_h(x), \quad (2.9)$$

$$\delta A_0^{\text{stat}}(x) = \bar{\psi}_1(x) \frac{1}{2} (\overleftarrow{\nabla}_i + \overleftarrow{\nabla}_i^*) \gamma_i \gamma_5 \psi_h(x). \quad (2.10)$$

Later we will also use the space components of the vector current represented by

$$V_k^{\text{HQET}}(x) = Z_V^{\text{HQET}} [V_k^{\text{stat}}(x) + c_V^{\text{HQET}} \delta V_k^{\text{stat}}(x)], \quad (2.11)$$

$$V_k^{\text{stat}}(x) = \bar{\psi}_1(x) \gamma_k \psi_h(x), \quad (2.12)$$

$$\delta V_k^{\text{stat}}(x) = -\bar{\psi}_1(x) \frac{1}{2} (\overleftarrow{\nabla}_i + \overleftarrow{\nabla}_i^*) \gamma_i \gamma_k \psi_h(x). \quad (2.13)$$

We have chosen $V_k^{\text{stat}}, \delta V_k^{\text{stat}}$ such that they are exactly related to $A_0^{\text{stat}}, \delta A_0^{\text{stat}}$ by a spin rotation.

The coefficients $\omega_{\text{kin}}, \omega_{\text{spin}}, Z_A^{\text{HQET}}, c_A^{\text{HQET}}, Z_V^{\text{HQET}}, c_V^{\text{HQET}}$ are functions of the bare coupling g_0 and of the heavy quark mass in lattice units. They represent bare parameters of the effective theory, which are to be fixed by matching to QCD. Just like $\omega_{\text{kin}}, \omega_{\text{spin}}$, the coefficients $c_A^{\text{HQET}}, c_V^{\text{HQET}}$ are of order $1/m_b$, while we may write

$$Z_A^{\text{HQET}} = Z_A^{\text{stat}} + Z_A^{(1)}, \quad \text{with } Z_A^{(1)} = \mathcal{O}(1/m_b), \quad (2.14)$$

and similarly for Z_V^{HQET} ¹. Note that in the expansion to first order, terms such as $\omega_{\text{kin}} c_A^{\text{HQET}} \propto 1/m_b^2$ are to be dropped.

Below we will consider an example and discuss that indeed the bare parameters $\omega_{\text{kin}}, \omega_{\text{spin}}, Z_A^{\text{HQET}}, c_A^{\text{HQET}}, Z_V^{\text{HQET}}, c_V^{\text{HQET}}$ and m_{bare} are sufficient to absorb all divergences in the effective theory at this order in $1/m_b$.

2.2 $1/m_b$ expansion in a geometry without boundaries

In order to illustrate further how the expansion works, we consider a two-point function of a composite field in a space-time without boundaries, i.e. with periodic boundary conditions or in infinite volume. We choose the example

$$C_{AA}(x_0) = Z_A^2 a^3 \sum_{\mathbf{x}} \left\langle A_0(x) (A_0)^\dagger(0) \right\rangle \quad (2.15)$$

with the heavy-light axial current in QCD, $A_\mu = \bar{\psi}_1 \gamma_\mu \gamma_5 \psi_b$, and Z_A ensuring the natural normalization of the current consistent with current algebra [29, 30]. The $1/m_b$ expansion reads

$$\begin{aligned} C_{AA}(x_0) &= e^{-m_{\text{bare}} x_0} (Z_A^{\text{HQET}})^2 a^3 \sum_{\mathbf{x}} \left[\left\langle A_0^{\text{stat}}(x) (A_0^{\text{stat}}(0))^\dagger \right\rangle_{\text{stat}} \right. \\ &\quad + \omega_{\text{kin}} \left\langle A_0^{\text{stat}}(x) (A_0^{\text{stat}}(0))^\dagger \right\rangle_{\text{kin}} + \omega_{\text{spin}} \left\langle A_0^{\text{stat}}(x) (A_0^{\text{stat}}(0))^\dagger \right\rangle_{\text{spin}} \\ &\quad \left. + c_A^{\text{HQET}} \left\langle A_0^{\text{stat}}(x) (\delta A_0^{\text{stat}}(0))^\dagger \right\rangle_{\text{stat}} + c_A^{\text{HQET}} \left\langle \delta A_0^{\text{stat}}(x) (A_0^{\text{stat}}(0))^\dagger \right\rangle_{\text{stat}} \right] \\ &\equiv e^{-m_{\text{bare}} x_0} (Z_A^{\text{HQET}})^2 \left[C_{AA}^{\text{stat}}(x_0) + \omega_{\text{kin}} C_{AA}^{\text{kin}}(x_0) + \omega_{\text{spin}} C_{AA}^{\text{spin}}(x_0) \right. \\ &\quad \left. + c_A^{\text{HQET}} [C_{\delta AA}^{\text{stat}}(x_0) + C_{A\delta A}^{\text{stat}}(x_0)] \right] \end{aligned} \quad (2.16)$$

up to terms of order $1/m_b^2$. As mentioned in the introduction, the mass shift $m_{\text{bare}} = \mathcal{O}(m_b)$ includes an additive mass renormalization. It is also split up as

$$m_{\text{bare}} = m_{\text{bare}}^{\text{stat}} + m_{\text{bare}}^{(1)}, \quad \text{with } m_{\text{bare}}^{(1)} = \mathcal{O}(1/m_b), \quad (2.17)$$

and the expansion $e^{-m_{\text{bare}} x_0} \equiv e^{-m_{\text{bare}}^{\text{stat}} x_0} (1 - x_0 m_{\text{bare}}^{(1)})$ is understood.

For illustration we check the self consistency of eq. (2.17). The relevant question concerns renormalization, namely: are the “free” parameters $m_{\text{bare}} \dots c_A^{\text{HQET}}$ sufficient to absorb all divergences on the r.h.s.? We consider the most difficult term, $C_{AA}^{\text{kin}}(x_0)$. According to the standard rules of renormalization of composite operators, it is renormalized as

$$\left(C_{AA}^{\text{kin}} \right)_R(x_0) = e^{-m_{\text{bare}}^{\text{stat}} x_0} (Z_A^{\text{stat}})^2 a^7 \sum_{\mathbf{x}, z} \left\langle A_0^{\text{stat}}(x) (A_0^{\text{stat}}(0))^\dagger (\mathcal{O}_{\text{kin}})_R(z) \right\rangle_{\text{stat}} + \text{C.T.}, \quad (2.18)$$

¹If $\mathcal{O}(a)$ improvement is desired in the static approximation, there are also $a \delta A_0^{\text{stat}}$, $a \delta V_k^{\text{stat}}$ corrections to the currents. They are not relevant in the present discussion but will be taken into account when necessary.

where C.T. denotes contact terms to be discussed shortly. The renormalized operator $(\mathcal{O}_{\text{kin}})_R(z)$ involves a subtraction of lower dimensional ones,

$$(\mathcal{O}_{\text{kin}})_R(z) = Z_{\mathcal{O}_{\text{kin}}}(\mathcal{O}_{\text{kin}}(z) + \frac{c_1}{a} \bar{\psi}_h(z) D_0 \psi_h(z) + \frac{c_2}{a^2} \bar{\psi}_h(z) \psi_h(z)), \quad (2.20)$$

written here in terms of dimensionless c_i . Since we are interested in on-shell observables ($x_0 > 0$ in eq. (2.19)), we may use the equation of motion $D_0 \psi_h(z) = 0$ to eliminate the second term. The third one, $\frac{c_2}{a^2} \bar{\psi}_h(z) \psi_h(z)$, is equivalent to a mass shift and only changes $m_{\text{bare}}^{(1)}$, which is hence quadratically divergent². Thus all terms which are needed for the renormalization of \mathcal{O}_{kin} are present in eq. (2.17). It remains to consider the contact terms in eq. (2.19). They originate from singularities in the operator products $\mathcal{O}_{\text{kin}}(z) A_0^{\text{stat}}(x)$ as $z \rightarrow x$ (and $\mathcal{O}_{\text{kin}}(z) (A_0^{\text{stat}})^\dagger(0)$ as $z \rightarrow 0$). Using the operator product expansion they can be represented as linear combinations of $A_0^{\text{stat}}(x)$ and $\delta A_0^{\text{stat}}(x)$. Such terms are contained in eq. (2.17) in the form of $C_{AA}^{\text{stat}}, C_{\delta AA}^{\text{stat}}$ and $C_{A\delta A}^{\text{stat}}$ ³.

We conclude that all terms which are needed for the renormalization of $C_{AA}^{\text{kin}}(x_0)$ are present in eq. (2.17); the parameters may thus be adjusted to absorb all infinities and with properly chosen coefficients the continuum limit of the r.h.s. is expected to exist. The basic assumption of the effective field theory is that once the finite parts of the coefficients have been determined by matching a set of observables to QCD, these coefficients are applicable to any other observables.

The B-meson mass is given by $C_{AA}(x_0)$ in large volume via

$$m_B = - \lim_{x_0 \rightarrow \infty} \frac{\partial_0 + \partial_0^*}{2} \log C_{AA}(x_0), \quad (2.21)$$

with

$$\partial_0 f(x_0) = \frac{1}{a} [f(x_0 + a) - f(x_0)], \quad \partial_0^* f(x_0) = \frac{1}{a} [f(x_0) - f(x_0 - a)]. \quad (2.22)$$

Inserting the HQET expansion we derive

$$m_B = m_B^{\text{stat}} + m_B^{(1)}, \quad (2.23)$$

with

$$m_B^{\text{stat}} = m_{\text{bare}}^{\text{stat}} + E_{\text{stat}}, \quad E_{\text{stat}} = - \lim_{x_0 \rightarrow \infty} \frac{\partial_0 + \partial_0^*}{2} \log C_{AA}^{\text{stat}}(x_0), \quad (2.24)$$

$$m_B^{(1)} = m_{\text{bare}}^{(1)} + \omega_{\text{kin}} E_{\text{kin}} + \omega_{\text{spin}} E_{\text{spin}}, \quad (2.25)$$

$$E_{\text{kin}} = - \lim_{x_0 \rightarrow \infty} \frac{\partial_0 + \partial_0^*}{2} [C_{AA}^{\text{kin}}(x_0) / C_{AA}^{\text{stat}}(x_0)], \quad (2.26)$$

²Using the explicit form of the static propagator, eq. (2.4) of reference [27], one can check that indeed $a^3 \sum_{\mathbf{x}} \langle A_0^{\text{stat}}(x) (A_0^{\text{stat}}(0))^\dagger a^4 \sum_z \bar{\psi}_h(z) \psi_h(z) \rangle_{\text{stat}} = x_0 C_{AA}^{\text{stat}}(x_0)$.

³ $A_0^{\text{stat}}(x)$ and $\delta A_0^{\text{stat}}(x)$ are the only operators of dimension 3 and 4 with the correct quantum numbers. Higher dimensional operators contribute only terms of order a . Note that the $A_0^{\text{stat}}(x)$ term is power divergent $\sim 1/(am_b)$. This divergence is absorbed by a power divergent contribution to Z_A^{HQET} (at order $1/m_b$).

$$E_{\text{spin}} = - \lim_{x_0 \rightarrow \infty} \frac{\partial_0 + \partial_0^*}{2} [C_{\text{AA}}^{\text{spin}}(x_0)/C_{\text{AA}}^{\text{stat}}(x_0)]. \quad (2.27)$$

Here the terms $\propto c_A^{\text{HQET}}$ of eq. (2.17) do not contribute. They are proportional to the derivative of ratios $C_{\delta\text{AA}}^{\text{stat}}(x_0)/C_{\text{AA}}^{\text{stat}}(x_0)$. At large x_0 these ratios approach a constant since δA_0^{stat} has the same quantum numbers as A_0^{stat} . Using the transfer matrix formalism (with normalization $\langle B|B \rangle = 1$), one further observes that

$$E_{\text{kin}} = -\langle B|a^3 \sum_{\mathbf{z}} \mathcal{O}_{\text{kin}}(0, \mathbf{z})|B \rangle_{\text{stat}}, \quad E_{\text{spin}} = -\langle B|a^3 \sum_{\mathbf{z}} \mathcal{O}_{\text{spin}}(0, \mathbf{z})|B \rangle_{\text{stat}}. \quad (2.28)$$

As expected, only the parameters of the action are relevant in the expansion of a hadron mass. In the above relations $m_{\text{bare}}^{\text{stat}}$ absorbs a linear divergence of E_{stat} and $m_{\text{bare}}^{(1)}$ a quadratic divergence of E_{kin} .

Going through the same steps in the vector channel and using the spin symmetry of the static action is one way to see that the combination

$$m_{\text{B}}^{\text{av}} \equiv \frac{1}{4}[m_{\text{B}} + 3m_{\text{B}^*}] = m_{\text{bare}} + E_{\text{stat}} + \omega_{\text{kin}} E_{\text{kin}} \quad (2.29)$$

is independent of ω_{spin} . It is instructive to represent this equation in a different way, subtracting the $1/a$ (and $1/a^2$) divergences of E_{stat} (and E_{kin}). In this way we have

$$m_{\text{B}}^{\text{av}} = m_{\text{B}}^{(0a)} + m_{\text{B}}^{(0b)} + m_{\text{B}}^{(1a)} + m_{\text{B}}^{(1b)}, \quad (2.30)$$

$$m_{\text{B}}^{(0a)} = m_{\text{bare}}^{\text{stat}} + E_{\text{stat}}^{\text{sub}}, \quad (2.31)$$

$$m_{\text{B}}^{(0b)} = E_{\text{stat}} - E_{\text{stat}}^{\text{sub}}, \quad (2.32)$$

$$m_{\text{B}}^{(1a)} = m_{\text{bare}}^{(1)} + \omega_{\text{kin}} E_{\text{kin}}^{\text{sub}}, \quad (2.33)$$

$$m_{\text{B}}^{(1b)} = \omega_{\text{kin}} [E_{\text{kin}} - E_{\text{kin}}^{\text{sub}}], \quad (2.34)$$

with finite terms $m_{\text{B}}^{(0a)}, m_{\text{B}}^{(0b)}, m_{\text{B}}^{(1a)}, m_{\text{B}}^{(1b)}$. Our strategy, described in the introduction can be seen as a way of determining the coefficient ω_{kin} as well as the subtractions $E_{\text{stat}}^{\text{sub}}, E_{\text{kin}}^{\text{sub}}$ from finite volume computations in QCD and HQET. Finite parts in the subtraction terms do of course depend on the detailed choice of kinematical parameters such as the matching volume, but the end result is unique up to terms of order $1/m_{\text{b}}^2$. Note that by the same logics, the order $1/m_{\text{b}}$ term, $m_{\text{B}}^{(1a)} + m_{\text{B}}^{(1b)}$, is *not unique* but depends on the details of the strategy.

Since the prediction eq. (2.29) requires only the knowledge of two parameters, we also need only two finite volume observables to perform the matching with QCD. The Schrödinger functional is particularly useful to find suitable observables [1, 31, 22]. We proceed to discuss the $1/m_{\text{b}}$ expansion in this situation.

2.3 Schrödinger functional correlation functions

The pure gauge Schrödinger functional has thoroughly been discussed in [24], relativistic and static quarks were introduced in [25] and [28]. In particular in the last reference also Symanzik $O(a)$ -improvement was discussed. The improvement of the Schrödinger

functional requires the addition of dimension four local composite fields localized at the boundaries [32]. However, it turns out that there are no dimension four composite fields which involve static quarks fields and which are compatible with the symmetries of the static action and the Schrödinger functional boundary conditions and which do not vanish by the equations of motion. Thus there are no $O(a)$ boundary counter terms with static quark fields. For the same reason there are also no $O(1/m_b)$ boundary terms in HQET. This then means the HQET expansion of the boundary quark fields $\zeta, \bar{\zeta}$ is trivial up to and including $1/m_b$ terms.

For details of the boundary conditions as well as the definition of the fields $\zeta, \bar{\zeta}$ we refer to [28], where also our notation is explained. For a general understanding it is, however, sufficient to note a few facts. In space the fermion fields are taken to be periodic up to a phase,

$$\psi(x + \hat{k}L) = e^{i\theta}\psi(x), \quad \bar{\psi}(x + \hat{k}L) = e^{-i\theta}\bar{\psi}(x), \quad (2.35)$$

with the same phase θ for all quark fields, whether relativistic or described by HQET. In time we take homogeneous Dirichlet boundary conditions at $x_0 = 0$ and $x_0 = T$ [28]. Correlation functions can be formed from composite fields in the bulk, $0 < x_0 < T$, and boundary quark fields $\zeta, \bar{\zeta}$. In QCD, correlation functions in the pseudoscalar and vector channel are

$$f_A(x_0, \theta) = -\frac{a^6}{2} \sum_{\mathbf{y}, \mathbf{z}} \langle (A_I)_0(x) \bar{\zeta}_b(\mathbf{y}) \gamma_5 \zeta_1(\mathbf{z}) \rangle, \quad (2.36)$$

$$k_V(x_0, \theta) = -\frac{a^6}{6} \sum_{\mathbf{y}, \mathbf{z}, k} \langle (V_I)_k(x) \bar{\zeta}_b(\mathbf{y}) \gamma_k \zeta_1(\mathbf{z}) \rangle. \quad (2.37)$$

The $O(a)$ improved currents A_I, V_I can be found in [1]. Furthermore we consider boundary to boundary correlation functions

$$f_1(\theta) = -\frac{a^{12}}{2L^6} \sum_{\mathbf{u}, \mathbf{v}, \mathbf{y}, \mathbf{z}} \langle \bar{\zeta}_1'(\mathbf{u}) \gamma_5 \zeta_b'(\mathbf{v}) \bar{\zeta}_b(\mathbf{y}) \gamma_5 \zeta_1(\mathbf{z}) \rangle, \quad (2.38)$$

$$k_1(\theta) = -\frac{a^{12}}{6L^6} \sum_{\mathbf{u}, \mathbf{v}, \mathbf{y}, \mathbf{z}, k} \langle \bar{\zeta}_1'(\mathbf{u}) \gamma_k \zeta_b'(\mathbf{v}) \bar{\zeta}_b(\mathbf{y}) \gamma_k \zeta_1(\mathbf{z}) \rangle. \quad (2.39)$$

Their renormalization is standard [33], for example

$$[f_A]_R(x_0, \theta) = Z_A Z_\zeta^2 f_A(x_0, \theta), \quad [f_1]_R(\theta) = Z_\zeta^4 f_1(\theta), \quad (2.40)$$

with Z_ζ a renormalization factor of the relativistic boundary quark fields.

In complete analogy to the case of a manifold without boundary we can write down the expansions to first order in $1/m_b$. They read

$$[f_A]_R = Z_A^{\text{HQET}} Z_{\zeta_b} Z_\zeta e^{-m_{\text{bare}} x_0} \left\{ f_A^{\text{stat}} + c_A^{\text{HQET}} f_{\delta A}^{\text{stat}} + \omega_{\text{kin}} f_A^{\text{kin}} + \omega_{\text{spin}} f_A^{\text{spin}} \right\}, \quad (2.41)$$

$$[k_V]_R = Z_V^{\text{HQET}} Z_{\zeta_b} Z_\zeta e^{-m_{\text{bare}} x_0} \left\{ k_V^{\text{stat}} + c_V^{\text{HQET}} k_{\delta V}^{\text{stat}} + \omega_{\text{kin}} k_V^{\text{kin}} + \omega_{\text{spin}} k_V^{\text{spin}} \right\}, \quad (2.42)$$

$$= -Z_V^{\text{HQET}} Z_{\zeta_h} Z_{\zeta} e^{-m_{\text{bare}} x_0} \left\{ f_A^{\text{stat}} + c_V^{\text{HQET}} f_{\delta A}^{\text{stat}} + \omega_{\text{kin}} f_A^{\text{kin}} - \frac{1}{3} \omega_{\text{spin}} f_A^{\text{spin}} \right\},$$

$$[f_1]_{\text{R}} = Z_{\zeta_h}^2 Z_{\zeta}^2 e^{-m_{\text{bare}} T} \left\{ f_1^{\text{stat}} + \omega_{\text{kin}} f_1^{\text{kin}} + \omega_{\text{spin}} f_1^{\text{spin}} \right\}, \quad (2.43)$$

$$[k_1]_{\text{R}} = Z_{\zeta_h}^2 Z_{\zeta}^2 e^{-m_{\text{bare}} T} \left\{ f_1^{\text{stat}} + \omega_{\text{kin}} f_1^{\text{kin}} - \frac{1}{3} \omega_{\text{spin}} f_1^{\text{spin}} \right\}. \quad (2.44)$$

Apart from

$$f_{\delta A}^{\text{stat}}(x_0, \theta) = -\frac{a^6}{2} \sum_{\mathbf{y}, \mathbf{z}} \langle \delta A_0^{\text{stat}}(x) \bar{\zeta}_h(\mathbf{y}) \gamma_5 \zeta_1(\mathbf{z}) \rangle \quad (2.45)$$

the labeling of the different terms follows directly the one introduced in eq. (2.7). We have used identities such as $f_A^{\text{kin}} = -k_V^{\text{kin}}$, $f_A^{\text{spin}} = 3k_V^{\text{spin}}$. As a simple consequence of the spin symmetry of the static action, these are valid at any lattice spacing.

3. Finite volume observables and step scaling functions

3.1 Observables

We concentrate on a strategy based on the correlation functions f_1, k_1 alone. This is advantageous, since the additional coefficients $c_A^{\text{HQET}}, c_V^{\text{HQET}}$ in eq. (2.41), eq. (2.42) are avoided. Apart from the b-quark, we set the masses of all quarks to zero.

In terms of the spin-averaged combination,

$$F_1(L, \theta) = \frac{1}{4} [\log f_1(\theta) + 3 \log k_1(\theta)], \quad (3.1)$$

we form

$$R_1(L, \theta_1, \theta_2) = F_1(L, \theta_1) - F_1(L, \theta_2) \text{ at } T = L/2 \quad (3.2)$$

$$\Gamma_1(L, \theta_0) = -\frac{\partial_T + \partial_T^*}{2} F_1(L, \theta_0) \text{ at } T = L/2. \quad (3.3)$$

Note that the boundary quark wave function renormalization cancels in R_1 and in Γ_1 . They are thus finite after renormalization of the parameters of the Lagrangian.

The dimensionless observables,

$$\Phi_1(L, M_b) = R_1(L, \theta_1, \theta_2) - R_1^{\text{stat}}(L, \theta_1, \theta_2), \quad (3.4)$$

$$\Phi_2(L, M_b) = L \Gamma_1(L, \theta_0), \quad (3.5)$$

$$R_1^{\text{stat}}(L, \theta_1, \theta_2) = \log [f_1^{\text{stat}}(L, \theta_1) / f_1^{\text{stat}}(L, \theta_2)] \text{ at } T = L/2 \quad (3.6)$$

are parametrized in terms of the RGI mass of the b-quark, M_b . They have a particularly simple $1/m_b$ expansion

$$\Phi_1(L, M_b) = \omega_{\text{kin}} R_1^{\text{kin}}(L, \theta_1, \theta_2), \quad (3.7)$$

$$\Phi_2(L, M_b) = L [m_{\text{bare}} + \Gamma_1^{\text{stat}}(L, \theta_0) + \omega_{\text{kin}} \Gamma_1^{\text{kin}}(L, \theta_0)], \quad (3.8)$$

which involves

$$R_1^{\text{kin}}(L, \theta_1, \theta_2) = \frac{f_1^{\text{kin}}(L, \theta_1)}{f_1^{\text{stat}}(L, \theta_1)} - \frac{f_1^{\text{kin}}(L, \theta_2)}{f_1^{\text{stat}}(L, \theta_2)} \text{ at } T = L/2, \quad (3.9)$$

$$\Gamma_1^{\text{stat}}(L, \theta_0) = -\frac{\partial_T + \partial_T^*}{2} \log f_1^{\text{stat}}(\theta_0) \text{ at } T = L/2, \quad (3.10)$$

$$\Gamma_1^{\text{kin}}(L, \theta_0) = -\frac{\partial_T + \partial_T^*}{2} [f_1^{\text{kin}}(\theta_0)/f_1^{\text{stat}}(\theta_0)] \text{ at } T = L/2. \quad (3.11)$$

The $\theta_0, \theta_1, \theta_2$ dependence of Φ_i is not explicitly written, but will of course be relevant in the numerical results. For the reader familiar with [34, 1], we point out that Γ_1 differs from Γ which was used in those references. Note that in eq. (3.4) we subtract the static term. This simplifies subsequent formulae. In fact, whenever such a lowest order contribution is universal (in the sense of having a universal continuum limit) and independent of an HQET parameter, it will be convenient to subtract it. Despite this subtraction, we refer to Φ_1 as an observable in QCD.

The reader may be surprised that we introduce the quantity Γ_1 which contains a (discretized) derivative with respect to the time extent, T . Its MC evaluation requires two separate simulations⁴. However, obviously a quantity of order m_b is needed and this is obtained from some logarithmic derivative of a correlation function. Boundary-to-boundary correlation functions are then very convenient since one does not have to deal with the $1/m_b$ corrections to the currents. It is a useful feature of the Schrödinger functional that such gauge invariant correlation functions are available.

3.2 Step scaling functions

We turn to the relations between $\Phi_i(L, M_b)$ and $\Phi_i(2L, M_b)$ in the effective theory. The dimensionful variable L is replaced by the Schrödinger functional renormalized coupling $\bar{g}^2(L)$ [35] over which we have good control in numerical computations [20]. Straightforward substitution yields

$$\Phi_1(2L, M_b) = \sigma_1^{\text{kin}}(u) \Phi_1(L, M_b), \quad (3.12)$$

$$\Phi_2(2L, M_b) = 2\Phi_2(L, M_b) + \sigma_m(u) + \sigma_2^{\text{kin}}(u) \Phi_1(L, M_b), \quad (3.13)$$

where always $u = \bar{g}^2(L)$. Our continuum step scaling functions σ (with any subscripts or superscripts) are defined in terms of those at finite lattice spacing as

$$\sigma(u) = \lim_{a/L \rightarrow 0} \Sigma(u, a/L). \quad (3.14)$$

At finite lattice spacing we have

$$\Sigma_1^{\text{kin}}(u, a/L) = \frac{R_1^{\text{kin}}(2L, \theta_1, \theta_2)}{R_1^{\text{kin}}(L, \theta_1, \theta_2)} \Big|_{u=\bar{g}^2(L)}, \quad (3.15)$$

$$\Sigma_2^{\text{kin}}(u, a/L) = \frac{2L [\Gamma_1^{\text{kin}}(2L, \theta_0) - \Gamma_1^{\text{kin}}(L, \theta_0)]}{R_1^{\text{kin}}(L, \theta_1, \theta_2)} \Big|_{u=\bar{g}^2(L)}, \quad (3.16)$$

$$\Sigma_m(u, a/L) = 2L [\Gamma_1^{\text{stat}}(2L, \theta_0) - \Gamma_1^{\text{stat}}(L, \theta_0)] \Big|_{u=\bar{g}^2(L)}. \quad (3.17)$$

⁴In appendix C we discuss a different strategy, which is based on the x_0 -derivative of f_A and thus requires less simulations. Note, however, that these additional simulations do not represent a significant effort.

The above equations are easily derived. In a first step, just from the $1/m_b$ expansions of Φ_i , one obtains them at a given resolution a/L or equivalently at fixed bare coupling, g_0 . One then uses that $\Phi_i(L, M_b)$ are dimensionless physical observables with a continuum limit. Since the self energy of a static quark cancels in σ_m , also that quantity has a finite continuum limit. Thus the continuum limit of the step scaling functions $\Sigma_m, \Sigma_i^{\text{kin}}$ exists and eqs. (3.12,3.13) can be written in terms of continuum quantities, as we have done.

4. M_b including $1/m_b$ corrections

Before giving the equation for M_b , we recall the overall strategy. For $L_1 \approx 0.4$ fm we compute $\Phi_1(L_1, M_b), \Phi_2(L_1, M_b)$ for a few quark masses around the physical one in quenched QCD. It is understood that the continuum limit is reached by an extrapolation and with a suitable interpolation of Φ_i in M_b , these quantities can be considered to be known as a function of M_b . With the step scaling functions described in the previous section and computed in the effective theory, we then arrive at $\Phi_1(L_2, M_b), \Phi_2(L_2, M_b)$, where $L_2 = 2L_1$. It remains to express the spin averaged B-meson mass m_B^{av} in terms of $\Phi_1(L_2, M_b), \Phi_2(L_2, M_b)$.

To this end, we straightforwardly combine eqs. (3.7,3.8) with eq. (2.29) and obtain

$$Lm_B^{\text{av}} = \Phi_2(L, M_b) + L[E^{\text{stat}} - \Gamma_1^{\text{stat}}(L, \theta_0)] + \frac{L[E^{\text{kin}} - \Gamma_1^{\text{kin}}(L, \theta_0)]}{R_1^{\text{kin}}(L, \theta_1, \theta_2)} \Phi_1(L, M_b). \quad (4.1)$$

We now set $L = L_2$ in this equation and insert eq. (3.13). In the form of eq. (2.30) we then have

$$L_2 m_B^{(0a)}(M_b) = \sigma_m(u_1) + 2 \Phi_2(L_1, M_b) \quad (4.2)$$

$$L_2 m_B^{(0b)} = L_2 [E^{\text{stat}} - \Gamma_1^{\text{stat}}(L_2, \theta_0)], \quad (4.3)$$

$$L_2 m_B^{(1a)}(M_b) = \sigma_2^{\text{kin}}(u_1) \Phi_1(L_1, M_b), \quad (4.4)$$

$$L_2 m_B^{(1b)}(M_b) = L_2 \frac{E^{\text{kin}} - \Gamma_1^{\text{kin}}(L_2, \theta_0)}{R_1^{\text{kin}}(L_2, \theta_1, \theta_2)} \sigma_1^{\text{kin}}(u_1) \Phi_1(L_1, M_b), \quad (4.5)$$

where

$$u_1 = \bar{g}^2(L_1), \quad L_2 = 2L_1. \quad (4.6)$$

The subtraction of power divergences in eq. (2.32), eq. (2.34) are $E_{\text{stat}}^{\text{sub}} = \Gamma_1^{\text{stat}}(L_2, \theta_0)$, $E_{\text{kin}}^{\text{sub}} = \Gamma_1^{\text{kin}}(L_2, \theta_0)$ and $\sigma_1^{\text{kin}}(u_1) \Phi_1(L_1, M_b) / R_1^{\text{kin}}(L_2, \theta_1, \theta_2)$ is a representation of the bare parameter ω_{kin} in eq. (2.34). The other parts, $m_B^{(0a)}, m_B^{(1a)}$, are computable entirely in finite volume.

The step scaling functions σ have been discussed before. They can be computed with lattice spacings such that a/L_1 is reasonably small, say below $1/6$. Of course they should be extrapolated to the continuum. We work with lattice spacings $a \leq 0.07$ fm in this step. The relativistic observables $\Phi_i(L_1, M_b)$, $i = 1, 2$ are computed for $a \leq 0.02$ fm, where a relativistic b-quark can be described by the $O(a)$ -improved Wilson action with controlled

a^2 -effects. Finally, the combinations $L_2[E^{\text{stat}} - \Gamma_1^{\text{stat}}(L_2, \theta_0)]$ and $L_2 \frac{E^{\text{kin}} - \Gamma_1^{\text{kin}}(L_2, \theta_0)}{R_1^{\text{kin}}(L_2, \theta_1, \theta_2)}$ are computed for lattice spacings of $a \leq 0.1$ fm such that finite size effects in E_{stat} and E_{kin} are negligible on lattices with an affordable number of points.

The mass of the b-quark is obtained from eq. (2.30) by expanding

$$M_b = M_b^{(0)} + M_b^{(1)}, \quad (4.7)$$

where $M_b^{(0)}$ is the solution of the static equation

$$m_B^{\text{av}} = m_B^{(0a)}(M_b^{(0)}) + m_B^{(0b)}(M_b^{(0)}) \quad (4.8)$$

and the $1/m_b$ correction is

$$M_b^{(1)} = -\frac{1}{S} [m_B^{(1a)}(M_b^{(0)}) + m_B^{(1b)}(M_b^{(0)})] \quad (4.9)$$

with

$$S = \frac{d}{dM_b} [m_B^{(0a)}(M_b) + m_B^{(0b)}(M_b)] = \frac{d}{dM_b} [m_B^{(0a)}(M_b)]. \quad (4.10)$$

We finish the discussion of the strategy with a remark on the dependence on the mass of the light quarks. This is relevant because it is of course better to consider the spin-averaged B_s quark mass in eq. (2.29); the necessary large volume computations are easier than for the B_d meson. In the quenched approximation the parameters in the action $m_{\text{bare}}, \omega_{\text{kin}}$ are independent of the light quark mass.⁵ Since our strategy determines them through finite volume computations, it follows that in all these computations the light quark mass may be set to zero, a convenient choice. Only E_{kin} and E_{stat} are then to be computed at the mass of the light quark of the meson whose (spin averaged) mass is considered.

5. Results

We have performed a numerical computation in the quenched approximation, using the $O(a)$ improved Wilson action [41, 32, 42]. The box size L_2 is chosen as $L_2 = 1.436r_0$, where r_0 , defined in terms of the static quark potential [43], has a phenomenological value of $r_0 \approx 0.5$ fm. From [20] we know the Schrödinger functional coupling $\bar{g}^2(L_1) = \bar{g}^2(L_2/2) \approx 3.48$. Given the knowledge of r_0/a as a function of g_0 of ref. [44] and that of the renormalized coupling [20], it is then convenient to fix g_0 in different ways for the different steps of the calculation. The differences are of course only a -effects which disappear in the continuum extrapolations. We give more details below. We will take the uncertainties in the relations $\bar{g}^2(L_1) \approx 3.48$ and $\bar{g}^2(L_1/4) \approx 1.8811$ (which we need later) into account in the very end.

In order to complete our definitions, we further choose $\theta_0 = 0$ and $\theta_1, \theta_2 \in \{0, 1/2, 1\}$. The different values of θ_1, θ_2 offer the possibility to check whether our final results are independent of these arbitrary parameters as they should be up to small $1/m_b^2$ terms. Simulation parameters as well as raw results are listed in tables in appendix A and B.

⁵In general, δm (and hence also m_{bare}) will contain a term like $b(g_0)m_1$, where for simplicity the light quarks are assumed to be degenerate with mass m_1 . Obviously, $b(g_0) = O(g_0^4)$ does, however, vanish for $N_f = 0$. As a renormalization term odd in m_1 , it is also absent for twisted mass lattice QCD [36] and QCD with exact chiral symmetry [37–40].




Figure 1: Continuum extrapolation of $\Phi_2(L_1, M_b)$, for $z = 10.4, 12.1, 13.3$ from bottom to top. The errors in the relation between bare quark mass \tilde{m}_q and the RGI mass M are translated into errors in Φ_2 . The g_0 -independent part of that error is included *after* [31] the continuum extrapolation (left side error bar). On the right, the equivalent in the alternative strategy is shown for $\theta_0 = 1/2$ (see appendix C).

5.1 QCD observables

For this part of the computation, we determined the bare parameters as in [31]: g_0 is fixed by requiring $\bar{g}^2(L_1/4) = 1.8811$ for given resolutions a/L . The PCAC mass of the light quark, defined exactly as in that reference, is set to zero. Our heavy quark masses are chosen such that $z = M_b L_1 \approx 10 - 13$. The bare parameters are listed in table 3.

We focus our attention directly on the continuum extrapolations. As an example we show $\Phi_1(M_b, L_1)$ and $\Phi_2(M_b, L_1)$ in Fig. 2 and Fig. 1. Note that for the static subtraction $R_1^{\text{stat}}(L_1, 1/2, 1)$, displayed on the right of Fig. 2, our lattice spacings are roughly a factor three larger, since in the effective theory we only have to respect $a/L_1 \ll 1$, not $aM_b \ll 1$ (for details see appendix A). Data have been obtained for two static actions, HYP1 and HYP2 [27]. In fitting them to the expected a -dependence, their continuum limit value is constrained to be independent of the action, but the a^2 slopes are of course different. The data for the different actions are highly correlated. As in all such cases, the errors of the continuum limit are computed from jackknife samples.

For values of θ_i which differ from the choice made in the figures, the a -dependence is very similar. In all these cases we find that extrapolations linear in a^2 using all four available lattice spacings are compatible with the ones where the data point at largest lattice spacing is ignored. We take the extrapolations with three points as our results for further analysis, since they have the more conservative error bars. The continuum limits are listed together with the raw numbers in tables 6 and 4. From a fit of the continuum $\Phi_2(z)$ to a linear function, we then extract the slope

$$S = \frac{d}{dz} \Phi_2 = 0.61(5) \tag{5.1}$$




Figure 2: Continuum extrapolation of $\Phi_1(L_1, M_b)$, separately for $R_1(L_1, 1/2, 1)$ in QCD (left) and for $R_1^{\text{stat}}(L_1, 1/2, 1)$ in the static approximation (right). Circles denote results with action HYP1 and squares, displaced slightly for visibility, are from action HYP2. The corresponding continuum extrapolation lines are slightly displaced as well.

and we are done with the matching. The rest of the numerical computations is carried out in the effective theory.

5.2 HQET step scaling functions

Next we discuss the connection of $\Phi_i(L_1, M_b)$ to $\Phi_i(L_2, M_b)$, $L_2 = 2L_1$. It is given by the step scaling functions of section 3.2. The bare parameters used in their computation are described in appendix A, and the values at finite resolution a/L_1 are given in tables 8-10⁶.

At lowest order in $1/m_b$, only σ_m contributes. In its continuum extrapolation (Fig. 3, table 8) we allow for a slope in a^2 , although the data are compatible with a vanishing slope. Note that the *absolute* error of σ_m is negligible in comparison to twice the one of Φ_2 (see Fig. 1) to which it is added in eq. (4.2). In fact the uncertainty in σ_m corresponds to an error of only 5 MeV in the b-quark mass, illustrating the possible precision in the static effective theory with these actions [27, 45].

A relevant question is how the precision deteriorates when one includes the first order corrections in $1/m_b$. Then two more step scaling functions contribute. In Fig. 4, we illustrate how the continuum limit of σ_1^{kin} is obtained. Here we have to allow for a *linear* dependence on the lattice spacing, since the theory is not $O(a)$ improved at the level of the $1/m_b$ contributions [1]. Taking the more conservative fit with only three points, we arrive at the continuum limit listed in table 9 for all combinations θ_1, θ_2 . In eq. (4.5), σ_1^{kin} is multiplied by small numbers (of order $1/m_b$). This means that its error will be negligible in the overall error budget.

⁶For $\Sigma_m, \Sigma_2^{\text{kin}}$ the coarsest resolution considered is $a/L_1 = 8$. Due to the derivative ∂_T at $T = L/2$, smaller values of L_1/a would involve a very short time separation.




Figure 3: Continuum extrapolation of Σ_m and $\widetilde{\Sigma}_m$.




Figure 4: Continuum extrapolation of Σ_1^{kin} for $\theta_1 = 1/2, \theta_2 = 1$.

Instead of σ_2^{kin} we show directly the continuum extrapolation of $m_B^{(1a)}$, eq. (4.4). As for Σ_m , the data shows no significant a -dependence. Nevertheless, in order to have a realistic error estimate, we allow for a linear slope in a (Fig. 5). In table 10 we list the raw numbers for Σ_2^{kin} as well as the extracted continuum limit for further analysis.




Figure 5: Lattice spacing dependence of $L_2m_B^{(1a)}$ for $M_b = M_b^{(0)}$. On the left we show $m_B^{(1a)}$ as introduced in section 4, with $\theta_1 = 1/2$, $\theta_2 = 1$. We insert Φ_1 in the continuum limit, such that the lattice spacing dependence is just due to Σ_2^{kin} . On the right the corresponding quantity is shown for the alternative strategy of appendix C, again with continuum values for $\tilde{\Phi}_i$. There we set $\theta_0 = 1/2$, $\theta_1 = 1/2$, $\theta_2 = 1$.




Figure 6: Lattice spacing dependence of $L_2m_B^{(0b)}$, details as in Fig. 5.

5.3 Large volume matrix elements and M_b

The last missing pieces in eq. (2.30) are the large volume static energy E_{stat} , eq. (2.24), and the matrix element of the kinetic operator E_{kin} , eq. (2.28). Here, in contrast to the rest of our numerical evaluations, the light quark mass is set to the mass of the strange quark in order to avoid a chiral extrapolation. The spin averaged mass of the B_s system is then to be inserted into eq. (2.29).

Although E_{stat} and E_{kin} can be computed with periodic boundary conditions we here follow [46] and evaluate also these quantities with Schrödinger functional boundary conditions in a large volume of about $T \times (1.5 \text{ fm})^3$, with $1.5 \text{ fm} \leq T \leq 3 \text{ fm}$ (also a check for finite size effects is carried out). The extraction is fairly standard, but still care has to be




Figure 7: Lattice spacing dependence of $L_2m_B^{(1b)}$, details as in Fig. 5.

taken to make sure that the ground state contribution is obtained. This is a particularly relevant issue for B-physics, because the gap to the first excited state is rather small. We relegate details to appendix B and discuss immediately the universal combinations $L_2m_B^{(0b)}$, $L_2m_B^{(1b)}$ which enter in eq. (2.30). The static contribution, shown in Fig. 6, is known with very good precision ⁷. In contrast, the $1/m_b$ correction $L_2m_B^{(1b)}$ does have a noticeable total uncertainty (Fig. 7, table 1). Still, this error is only about 50% of the one on $2\Phi_2$. Note also that this error is almost entirely due to E_{kin} which may possibly be computed more precisely by other techniques [47].

We now have all pieces necessary to solve the equations for M_b . The static one, eq. (4.8), is illustrated in Fig. 8. The horizontal error band is given by subtracting the static pieces $\sigma_m + L_2m_B^{(0b)}$ from the experimental number

$$m_B^{\text{av}} = 5.405 \text{ GeV} . \quad (5.2)$$

The figure demonstrates again that the main source of error is contained in the QCD computation of Φ_2 . Finally, by interpolating $\Phi_i(L_1, M_b)$ to $M_b = M_b^{(0)}$ we obtain ($\theta_1 = 1/2, \theta_2 = 1$)

$$r_0 M_b^{(0)} = 17.25(20), \quad M_b^{(0)} = 6.806(79) \text{ GeV} \quad \text{for } r_0 = 0.5 \text{ fm} \quad (5.3)$$

$$r_0 M_b^{(1)} = -0.12(9), \quad M_b^{(1)} = -0.049(39) \text{ GeV} \quad \text{for } r_0 = 0.5 \text{ fm} \quad (5.4)$$

$$r_0 M_b = 17.12(22), \quad M_b = 6.758(86) \text{ GeV} \quad \text{for } r_0 = 0.5 \text{ fm} . \quad (5.5)$$

Here the small difference $\bar{g}^2(L_1/4) - 1.8811$ as well as the statistical uncertainties in $\bar{g}^2(L_1)$ and L_1/r_0 have been taken into account, as explained in appendix D. Moreover, one can see in table 1 that the θ_i dependence of the $1/m_b$ contribution is absorbed. With

⁷We show the results given for the static action HYP2. The continuum extrapolation with action HYP1 looks very similar, but the fit has a smaller χ^2 .

θ_1	θ_2	$r_0 M_b^{(1a)}$	$r_0 M_b^{(1b)}$
0	1/2	-0.06(3)	-0.06(8)
1/2	1	-0.06(3)	-0.06(8)
1	0	-0.06(3)	-0.06(8)

Table 1: RGI results of $1/m_b$ correction of the b-quark mass, in units of r_0 .




Figure 8: Graphical solution of eq. (4.8). On the left hand side, data points are $2\Phi_2$ and the horizontal error band is $L_2 m_B^{\text{av}} - \sigma_m - L_2[E^{\text{stat}} - \Gamma_1^{\text{stat}}(L_2, \theta_0)]$. On the right hand side, the analogous terms are shown for the alternative strategy ($\theta_0 = 1/2$, see section 5.4).

$\Lambda_{\overline{\text{MS}}} r_0 = 0.602(48)$ [35, 48], the 4-loop β function and the mass anomalous dimension [49–52], we translate $M_b = M_b^{(0)} + M_b^{(1)}$ to the mass in the $\overline{\text{MS}}$ scheme at its own scale,

$$\overline{m}_b(\overline{m}_b) = 4.347(48)\text{GeV} ; \tag{5.6}$$

the associated perturbative uncertainty can safely be neglected. For completeness we note that in the $\overline{\text{MS}}$ scheme the $1/m_b$ term amounts to $-27(22)\text{MeV}$.

5.4 Comparison to results from an alternative strategy

As mentioned earlier, at first sight it appears more natural to base the computation of M_b on the logarithmic derivative of the spin average of f_A and k_V as the prime finite volume quantity. We have not chosen this option as our standard strategy since then three observables are needed for matching. However, it is useful to consider also that alternative strategy in order to perform an explicit check that $1/m_b^2$ terms are as small as expected. The results can be appreciated without detailed definitions of the observables and step scaling functions, the interested reader can find them in appendix C. Here we note that within this alternative strategy we actually have nine different sets of $\{\tilde{\Phi}_1, \tilde{\Phi}_2, \tilde{\Phi}_3\}$. Only one observable, $\tilde{\Phi}_1 = \Phi_1$, is in common to the two strategies. For our graphs we have selected (arbitrarily) one choice of parameters.

θ_1	θ_2	$r_0(M_b^{(0)} + M_b^{(1a)} + M_b^{(1b)})$			
		Main strategy	Alternative strategy		
			$\theta_0 = 0$	$\theta_0 = 1/2$	$\theta_0 = 1$
0	1/2	17.12(22)	17.25(28)	17.23(28)	17.17(32)
1/2	1	17.12(22)	17.23(27)	17.21(27)	17.14(30)
1	0	17.12(22)	17.24(27)	17.22(28)	17.15(30)

Table 2: RGI results of M_b including the $1/m_b$ correction, and comparison of the two strategies.

First, let us summarize what kind of differences one expects in such a comparison apart from a -effects. In the order of magnitude counting, we take $L_1^{-1} \sim \Lambda \sim 0.5 \text{ GeV}$ and of course $L_2 = 2L_1$. The matching observables $\Gamma_1, \Gamma^{\text{av}}$ are constructed to be equal to the quark mass at the leading order in the HQET expansion. They start to differ at the next to leading order, which means by terms of order Λ . Also their dependence on θ_i is of that magnitude. Since $\Phi_2(L_1, M_b)$ and $\tilde{\Phi}_3(L_1, M_b)$ have been made dimensionless by multiplication with L_1 and L_1 happens to be around Λ^{-1} , the differences of $\Phi_2(L_1, M_b)$ and $\tilde{\Phi}_3(L_1, M_b)$ are order one. The step scaling functions $\sigma_m, \tilde{\sigma}_m$ as well as $L_2 m_B^{(0b)}$ are added to Φ_2 (or $\tilde{\Phi}_3$) to obtain $L_2 m_B$ in static approximation. Thus they depend on the details at the same level, apart from a trivial $L_2/L_1 = 2$ factor. Of course, in the total static estimate $r_0 M_b^{(0)}$ this dependence is reduced to $r_0 M_b \times (\Lambda/m_b)^2 \sim 1/5$. In the same way, the $1/m_b$ corrections $L_2 m_B^{(1a)}, L_2 m_B^{(1b)}$ themselves have a dependence on the matching conditions which is $L_2 \times \Lambda^2/m_b \sim 1/5$ but the final result $r_0 M_b$ including these terms is accurate and unique up to $r_0 \times (\Lambda^3/m_b)^2 \sim 1/50$ corrections.

We leave it to the reader to check in figure 1 to 7 that these expectations are fully satisfied by our results ⁸. In fact it appears that our estimate for the expansion parameter, $\Lambda/m_b \sim 1/10$ is quite realistic. Of course, to find this out requires an explicit computation of the correction terms as presented here. In some cases, such as $m_B^{(1b)}$, our precision is not good enough to resolve a dependence on the matching conditions.

In the b-quark mass in the static approximation, $r_0 M_b^{(0)}$ (eq. (5.3) and table 12), the maximum difference is 0.5(2), which is of the predicted order of magnitude. Finally, when we add all contributions together, the results from the alternative strategy, table 2, are fully in agreement with eq. (5.5). As expected $1/m_b^2$ terms are not visible with our precision. They can safely be neglected.

6. Conclusions

The main conclusion of this work is that fully non-perturbative computations in lattice HQET, as they have been suggested in [1], are possible in practice. In particular, the uncertainties in the $1/m_b$ corrections are smaller than those in the static approximation,

⁸We note in passing that $\tilde{\Sigma}_m$, in contrast to Σ_m , does in principle require an improvement coefficient, c_A^{stat} [28], for $O(a)$ -improvement. It has been set to the 1-loop values from [27], but the results are rather insensitive to c_A^{stat} , so its uncertainty can be neglected.

despite the fact that we numerically cancel large a^{-2} divergences in the $1/m_b$ terms. The final error in the mass of the b-quark is dominated by the uncertainty in the renormalization in QCD. Errors due to simulations in the effective theory can almost be neglected in comparison.

A very nice result is the independence of the final numbers for M_b of the matching condition: table 2 shows that within our reasonably small uncertainties, we get the same results for the quark mass for altogether twelve different matching conditions. This is expected up to very small terms of order $r_0 M_b \times (\Lambda/m_b)^3 \sim 0.02$, which should be compared to our result $r_0 M_b = 17.12(22) - 17.25(28)$. Here the quoted range is due to the different matching conditions. In the order of magnitude estimates we have made a guess for the typical scale of $\Lambda \sim 0.5 \text{ GeV}$. In the static approximation, some of the matching conditions yield slightly differing results for the quark mass in agreement with the expectation for such variations of $r_0 M_b \times (\Lambda/m_b)^2 \sim 0.2$.

Both this explicit test of the magnitude of the different orders in the expansion and the naive order of magnitude estimate say that $1/m_b^2$ corrections are completely negligible.

Still, in aspects of the computation, considerable improvement can be envisaged. For example, return to the $1/m_b$ contribution to the B-meson mass Fig. 7. The statistical errors grow rapidly as one decreases the lattice spacing. The by far dominating uncertainty in the shown combination is the one of the large volume matrix E_{kin} . It appears worth while to look for improvements, maybe along the line of [53]. Due to these errors, and of course the missing $O(a)$ -improvement of the theory at order $1/m_b$ [1], the continuum extrapolation is not easy. Fortunately it is still precise enough for the present case. It will be very interesting to see cases where the $1/m_b$ corrections are larger, as it is expected, for example, for F_B .

Let us now turn to the computed value of \bar{m}_b , eq. (5.6). Starting from a precisely specified input, namely r_0 , m_K and $(m_{B_s} + 3m_{B_s^*})/4$, the value of M_b is unambiguous in the quenched approximation, because these inputs fix the bare coupling, strange and beauty quark masses. We have used the experimental meson masses and $r_0 = 0.5 \text{ fm}$. Our numbers for M_b or \bar{m}_b may then be used as a benchmark result for other methods. Indeed, a comparison shows agreement with [54] and the recent extension of that work [55] $\bar{m}_b = 4.42(7) \text{ GeV}$.

Earlier, the review [56] quoted $\bar{m}_b = 4.30(5)(5) \text{ GeV}$ and $\bar{m}_b = 4.34(3)(6) \text{ GeV}$, based on static computations [57] and an extrapolation of NRQCD results to the static limit [58] respectively. A perturbative subtraction [18, 59, 60] of the linear divergence δm was carried out in these static estimates and, of course, a continuum extrapolation could not be done.

However, if other inputs are used, the result may change because r_0 is only approximately known and because the quenched approximation is not real QCD. A rough idea on the possible changes can be obtained by varying r_0 by $\pm 0.05 \text{ fm}$. This changes $\bar{m}_b(\bar{m}_b)$ by roughly $\pm 80 \text{ MeV}$.

These remarks just serve to stress the obvious necessity of performing computations with $N_f > 0$. The ALPHA-collaboration is presently starting with $N_f = 2$, where the renormalization of the quark mass in QCD is known [21]. The necessary HQET computations are *not* expected to be a big numerical challenge, apart from the large volume B-meson matrix elements: simulations of the Schrödinger functional for $L \leq 1 \text{ fm}$ are not

$\frac{L}{a}$	β	κ_1	$\bar{g}^2(\frac{L}{4})$	$Z_P(g_0, \frac{L}{2})$	b_m	Z	κ_h
20	7.2611	0.134145	1.8811(19)	0.6826(3)	-0.621	1.0955	0.124195
							0.122119
							0.120483
24	7.4082	0.133961	1.8811(22)	0.6764(6)	-0.622	1.0941	0.126055
							0.124528
							0.123383
32	7.6547	0.133632	1.8811(28)	0.6713(8)	-0.622	1.0916	0.127991
							0.126967
							0.126222
40	7.8439	0.133373	1.8811(22)	0.6679(8)	-0.623	1.0900	0.128989
							0.128214
							0.127656

Table 3: Bare parameters used in the computation of the QCD observables for $L = \tilde{L}_1$.

very demanding with nowadays computing capabilities [61]. Altogether the extension of the present work to full QCD is feasible and should be carried out, since presently no better method is known to compute the b-quark mass from lattice QCD.

Acknowledgments

We thank Stephan Dürr for collaboration in the early stages of this work [62]. We thank NIC for allocating computer time on the APEmille computers at DESY Zeuthen to this project and the APE group for its help. This work is supported by the Deutsche Forschungsgemeinschaft in the SFB/TR 09.

A. Finite volume simulations

For the matching in a finite volume, we performed one set of simulations of (quenched) QCD and one of HQET. In the case of the relativistic theory, we used $L = \tilde{L}_1$, defined by $\bar{g}^2(\tilde{L}_1/4) = 1.8811$.⁹ The parameters of these simulations have been taken from [31] (see table 3). The difference is that here $L = \tilde{L}_1 = 2\tilde{L}_0$ (and $T = L/2$ and $T = L/2 \pm a$ in addition to $T = L$) compared to $L = \tilde{L}_0$ in [31].

The parameters for the resolution $\tilde{L}_1/a = 20$ cannot be found in the mentioned reference. For that point, the gauge coupling β has been chosen such that $\bar{g}^2(\tilde{L}_1/4) = 1.8811$ for $\tilde{L}_1/4a = 5$, see [20]. The renormalization constant Z_P and $\kappa_1 = \kappa_c$ have been computed here, while b_m and Z have been extrapolated from the values in table 2 of [31]. These factors are put into the relationship between the bare mass $m_{q,h}$ and the RGI mass [20, 63],

$$M = h Z_m m_{q,h} (1 + b_m a m_{q,h}) , \tag{A.1}$$

⁹ \tilde{L}_1 differs slightly from L_1 defined in the main text by $L_1 = 0.718r_0$. This mismatch is however corrected, as explained later in this appendix and in appendix D.

L/a	z	R_1			Φ_2
		$\theta_1 = 0$ $\theta_2 = 1/2$	$\theta_1 = 1/2$ $\theta_2 = 1$	$\theta_1 = 1$ $\theta_2 = 0$	$\theta_0 = 0$
20	10.4	0.09795(13)	0.27426(30)	-0.37221(42)	7.847(40)
20	12.1	0.09512(12)	0.26588(30)	-0.36100(43)	9.108(46)
20	13.3	0.09336(12)	0.26068(30)	-0.35404(43)	10.068(50)
24	10.4	0.09958(18)	0.27904(37)	-0.37862(52)	7.697(44)
24	12.1	0.09689(17)	0.27110(37)	-0.36799(52)	8.866(50)
24	13.3	0.09528(17)	0.26632(36)	-0.36159(50)	9.716(54)
32	10.4	0.10157(30)	0.28481(71)	-0.38638(93)	7.512(53)
32	12.1	0.09897(30)	0.27717(71)	-0.37614(92)	8.623(58)
32	13.3	0.09744(30)	0.27265(71)	-0.37008(92)	9.411(62)
40	10.4	0.10283(30)	0.28806(52)	-0.39089(76)	7.484(51)
40	12.1	0.10027(30)	0.28052(52)	-0.38079(75)	8.575(56)
40	13.3	0.09876(29)	0.27608(52)	-0.37484(74)	9.344(60)
<i>CL</i>	10.4	0.10450(44)	0.29297(89)	-0.39748(125)	7.341(96)
<i>CL</i>	12.1	0.10202(44)	0.28567(90)	-0.38769(124)	8.386(102)
<i>CL</i>	13.3	0.10058(44)	0.28143(91)	-0.38202(124)	9.106(107)

Table 4: Simulation results of the finite volume ($L = \tilde{L}_1$) relativistic observables needed in our main strategy. The continuum limits, obtained by linear extrapolation in $(a/L)^2$ of the results for $L/a \geq 24$, are indicated by *CL*.

where

$$Z_m = \frac{Z Z_A}{Z_P}, \quad \text{and} \quad am_{q,h} = \frac{1}{2} \left(\frac{1}{\kappa_h} - \frac{1}{\kappa_c} \right). \quad (\text{A.2})$$

The renormalization constant $Z_A(g_0^2)$ is known non-perturbatively from [30], while

$$h = \frac{M}{\bar{m}(\mu_0)} = 1.544(14), \quad \mu_0 = 2/\tilde{L}_1, \quad (\text{A.3})$$

relates the running quark mass in the Schrödinger functional scheme [20] at the scale μ_0 , to the renormalization group invariant quark mass M ¹⁰.

For all values of \tilde{L}_1/a three hopping parameters κ_h have then been fixed in order to achieve

$$z = \tilde{L}_1 M = 10.4, 12.1, 13.3. \quad (\text{A.4})$$

We collect these parameters in table 3, whereas the results for the quantities needed in the matching step are summarized in tables 4 and 5. The errors there include systematics due to the uncertainties in the Z -factors, in particular, the error on the universal factor h has been propagated only *after* performing the continuum limit extrapolations. Ensembles of roughly 2000 (for $\tilde{L}_1/a = 20$) to few hundreds (for $\tilde{L}_1/a = 40$) gauge configurations

¹⁰In $h = M/\bar{m}(\mu_0)$ we take the small difference between the above defined \tilde{L}_0 and the value $L_0 = L_1/2$ into account. It causes a change of less than 1% of the value of h used in [31].

L/a	z	R_{av}			$\tilde{\Phi}_3$		
		$\theta_1 = 0$	$\theta_1 = 1/2$	$\theta_1 = 1$	$\theta_1 = 0$	$\theta_1 = 1/2$	$\theta_1 = 1$
		$\theta_2 = 1/2$	$\theta_2 = 1$	$\theta_2 = 0$			
20	10.4	0.1699(9)	0.4299(12)	-0.5998(20)	8.059(37)	8.293(37)	8.993(37)
20	12.1	0.1668(9)	0.4198(11)	-0.5867(20)	9.315(37)	9.545(37)	10.234(37)
20	13.3	0.1649(9)	0.4137(11)	-0.5787(19)	10.271(37)	10.500(37)	11.180(37)
24	10.4	0.1739(23)	0.4391(31)	-0.6130(54)	7.864(39)	8.102(38)	8.822(39)
24	12.1	0.1710(23)	0.4295(30)	-0.6005(52)	9.027(39)	9.263(38)	9.971(39)
24	13.3	0.1693(22)	0.4239(29)	-0.5931(51)	9.874(39)	10.109(38)	10.809(38)
32	10.4	0.1760(41)	0.4494(48)	-0.6254(90)	7.713(43)	7.941(41)	8.661(42)
32	12.1	0.1733(40)	0.4403(46)	-0.6135(87)	8.818(42)	9.045(41)	9.753(42)
32	13.3	0.1717(40)	0.4349(45)	-0.6066(85)	9.603(42)	9.828(41)	10.531(42)
40	10.4	0.1790(70)	0.4493(72)	-0.6283(142)	7.656(45)	7.894(42)	8.624(44)
40	12.1	0.1763(68)	0.4403(70)	-0.6166(138)	8.743(45)	8.979(42)	9.698(44)
40	13.3	0.1747(67)	0.4352(68)	-0.6099(136)	9.509(45)	9.744(42)	10.456(44)
<i>CL</i>	10.4	0.1801(75)	0.4587(84)	-0.6392(159)	7.533(89)	7.765(86)	8.496(88)
<i>CL</i>	12.1	0.1776(73)	0.4502(81)	-0.6280(154)	8.573(91)	8.805(88)	9.524(89)
<i>CL</i>	13.3	0.1761(72)	0.4452(79)	-0.6218(151)	9.289(93)	9.519(91)	10.234(92)

Table 5: Same as table 4 in the case of the alternative strategy.

L/a	R_1^{stat}					
	$\theta_1 = 0$		$\theta_1 = 1/2$		$\theta_1 = 1$	
	$\theta_2 = 1/2$		$\theta_2 = 1$		$\theta_2 = 0$	
	HYP1	HYP2	HYP1	HYP2	HYP1	HYP2
6	0.06936(5)	0.06939(4)	0.18583(7)	0.18591(7)	-0.25519(12)	-0.25530(11)
8	0.07572(6)	0.07574(6)	0.20452(11)	0.20457(11)	-0.28024(17)	-0.28031(17)
10	0.07821(5)	0.07822(5)	0.21246(8)	0.21249(8)	-0.29067(13)	-0.29071(13)
12	0.07934(8)	0.07935(8)	0.21622(13)	0.21625(13)	-0.29556(21)	-0.29559(20)
<i>CL</i>	0.08238(12)		0.22596(21)		-0.30835(32)	

Table 6: Lattice results of R_1^{stat} for $L = L_1$. The continuum limits are obtained by a linear extrapolation in $(a/L)^2$ of the results for $L/a \geq 8$.

have been generated for this part of the computation. The lattice $\tilde{L}_1/a = 20$ is not used in the extrapolations but rather to check for the smallness of higher order cutoff effects for $\tilde{L}_1/a \geq 24$.

Concerning the simulation of HQET, we have computed the various quantities in the two required volumes. The first one, where we match the effective theory with QCD, has a space extent L_1 . The second one is such that $L_2 = 2L_1$. The value of the Schrödinger functional renormalized coupling is fixed at $\bar{g}^2(L_1) = 3.48$, and we have used the resolutions $L_1/a = 6, 8, 10, 12$. The corresponding values of β as well as $\kappa = \kappa_c$ can be found in table A.1 of [1]. All these quantities are computed with two different actions, HYP1 and HYP2. The continuum values are then obtained by constraining the fits to give the same values for these actions. We note that the results for HYP1 and HYP2 are statistically correlated.

L/a	$R_{\text{av}}^{\text{stat}}$					
	$\theta_1 = 0$		$\theta_1 = 1/2$		$\theta_1 = 1$	
	$\theta_2 = 1/2$		$\theta_2 = 1$		$\theta_2 = 0$	
	HYP1	HYP2	HYP1	HYP2	HYP1	HYP2
6	0.1502(3)	0.1543(3)	0.3562(3)	0.3688(3)	-0.5231(6)	-0.5231(6)
8	0.1544(4)	0.1575(4)	0.3672(4)	0.3765(4)	-0.5216(7)	-0.5340(8)
10	0.1571(5)	0.1595(5)	0.3724(6)	0.3710(6)	-0.5295(10)	-0.5391(10)
12	0.1561(8)	0.1579(8)	0.3729(8)	0.3786(9)	-0.5289(15)	-0.5365(16)
<i>CL</i>	0.08238(12)		0.22596(21)		-0.30835(32)	

Table 7: Lattice results of $R_{\text{av}}^{\text{stat}}$. The details are the same as in table 6.

L/a	$\Sigma_{\text{m}}(3.48, a/L)$	
	HYP1	HYP2
8	0.431(11)	0.411(11)
10	0.437(11)	0.424(10)
12	0.422(16)	0.418(16)
<i>CL</i>	0.430(25)	

Table 8: Lattice results of the step scaling function Σ_{m} . The bare parameters are described in the text. The continuum limit is obtained by a linear extrapolation in $(a/L)^2$ of the results for $L/a \geq 8$.

L/a	$\Sigma_1^{\text{kin}}(3.48, a/L)$					
	$\theta_1 = 0$		$\theta_1 = 1/2$		$\theta_1 = 1$	
	$\theta_2 = 1/2$		$\theta_2 = 1$		$\theta_2 = 0$	
	HYP1	HYP2	HYP1	HYP2	HYP1	HYP2
6	0.6241(17)	0.6245(11)	0.6219(60)	0.6223(5)	0.6225(8)	0.6228(6)
8	0.5790(20)	0.5797(13)	0.5789(65)	0.5793(5)	0.5789(10)	0.5794(7)
10	0.5587(47)	0.5586(22)	0.5585(14)	0.5588(9)	0.5586(22)	0.5590(14)
12	0.5364(66)	0.5342(39)	0.5424(19)	0.5417(12)	0.5409(30)	0.5398(18)
<i>CL</i>	0.457(10)		0.471(3)		0.467(5)	

Table 9: Lattice results of the step scaling function Σ_1^{kin} . The continuum limits are obtained by a linear extrapolation in a/L of the results for $L/a \geq 8$.

For the computation of the step scaling functions one uses the same β, κ and $L_2/a = 2L_1/a$. All these computations are done with several thousand gauge configurations. Note that, even if L_1 is the same in QCD and in HQET, the typical lattice spacings are much larger in the effective theory. The results of R_1^{stat} and R_{av} can be found in tables 6 and 7. The values of the step scaling functions are collected in tables 8, 9 and 10.

Finally there are simulations in small volume to obtain the subtractions $\Gamma_1^{\text{stat}}(L_2)$ and $\Gamma_1^{\text{kin}}(L_2)$. These are done with $L_2 = 1.436 r_0$ and β determined from the knowledge of

L/a	$\Sigma_2^{\text{kin}}(3.48, a/L)$					
	$\theta_1 = 0$		$\theta_1 = 1/2$		$\theta_1 = 1$	
	$\theta_2 = 1/2$		$\theta_2 = 1$		$\theta_2 = 0$	
	HYP1	HYP2	HYP1	HYP2	HYP1	HYP2
8	4.81(44)	4.72(32)	1.58(15)	1.55(10)	-1.19(11)	-1.17(8)
10	4.34(58)	4.20(39)	1.43(19)	1.39(13)	-1.08(15)	-1.04(10)
12	4.79(86)	3.98(58)	1.58(28)	1.31(19)	-1.19(21)	-0.99(14)
CL	2.9(1.5)		0.96(50)		-0.71(38)	

Table 10: Same as table 9 for Σ_2^{kin} .

β	κ_s	$L^3 \times T$	aE_{stat}		a^2E_{kin}	
			HYP1	HYP2	HYP1	HYP2
6.0219	0.133849	$16^3 \times 24$ [32]	0.4345(21)	0.4029(32)	0.750(4)	0.774(3)
6.0219	0.133849	$24^3 \times 32$	0.4378(25)	0.4034(20)	0.746(7)	0.776(5)
6.2885	0.1349798	$24^3 \times 48$	0.3295(21)	0.3034(29)	0.643(7)	0.676(5)
6.4956	0.1350299	$32^3 \times 64$	0.2724(20)	0.2461(14)	0.599(10)	0.620(11)

Table 11: Parameters of the large volume simulations. Where present, the numbers in brackets refer to a second dataset at the same (β, κ) values.

r_0/a [48]. The parameters, including $\kappa = \kappa_c$, are listed in table 6 of [27]. The values of β do of course agree with the ones employed in the large volume, which we describe in the next appendix.

B. Large volume simulations and extraction of matrix elements

The parameters for the simulations in large volume are collected in table 11 together with the results for E_{stat} and E_{kin} . The lattice extension L/a and β are such that $L = 4L_1 \approx 3/2$ fm except for the second lattice where we have $L = 6L_1 \approx 2$ fm. This lattice is used only to check for the absence of finite size effects. We see from table 11 that finite size effects are indeed very small, the difference between the results from the $L/a = 16$ and the $L/a = 24$ lattices at $\beta = 6.0219$ is consistent with zero within at most one standard deviation (aE_{stat} from HYP1). The number of configurations generated ranges from 4300 at $\beta = 6.0219$ to 2200 at $\beta = 6.4956$ (for the larger volume at $\beta = 6.0219$ we had 1300 configurations). Since our phenomenological input is the mass of the (spin averaged) B_s meson, we set κ to κ_s in order to reproduce the quenched value of the strange quark mass from ref. [64], i.e.

$$M_s r_0 = 0.35(1) , \tag{B.1}$$

with M_s the renormalization group invariant strange quark mass defined as in appendix A after replacing κ_h by κ_s .

The numbers for E_{stat} and E_{kin} have been obtained by applying two different fitting procedures to two independent datasets (where available). The quoted errors are such that both the results are covered and they therefore provide a reasonable estimate of the systematics associated with the fits. We now sketch these procedures.

Let us consider in QCD the effective “mass” $\Gamma(x_0)$ obtained from the correlation function $F_{\text{av}}(x_0)$ in eq. (C.1) and its quantum-mechanical decomposition

$$\Gamma(x_0) = -\frac{\partial_0 + \partial_0^*}{2} F_{\text{av}} = E_0 + Ae^{-\Delta x_0} + \dots \quad (\text{B.2})$$

where E_0 is the energy of the ground state, Δ is the gap between the ground and the first excited states and the dots refer to contributions from higher states. The $1/m_b$ expansion reads

$$\begin{aligned} \Gamma(x_0) &= E_{\text{stat}} + \omega_{\text{kin}} E_{\text{kin}} + (A^{\text{stat}} + \omega_{\text{kin}} A^{\text{kin}}) e^{-\Delta^{\text{stat}} x_0} (1 - \omega_{\text{kin}} x_0 \Delta^{\text{kin}}) + \dots \\ &= \Gamma^{\text{stat}}(x_0) + \omega_{\text{kin}} \Gamma^{\text{kin}}(x_0) + \dots \end{aligned} \quad (\text{B.3})$$

where Γ^{stat} and Γ^{kin} are defined in analogy to eqs. (3.10, 3.11) in terms of the correlators $f_{\text{A}}^{\text{stat}}(x_0)$ and $f_{\text{A}}^{\text{kin}}(x_0)$.

In the correlation function $f_{\delta\text{A}}(x_0)$ the same states contribute as in $f_{\text{A}}(x_0)$. Performing again first the quantum-mechanical decomposition and then the $1/m_b$ expansion of these correlators, it is easy to see that the ratios

$$P_{\text{A}}^{\text{stat}}(x_0) = \frac{f_{\text{A}}^{\text{stat}}(x_0)}{f_{\delta\text{A}}^{\text{stat}}(x_0)} \quad \text{and} \quad P_{\text{A}}^{\text{kin}}(x_0) = P_{\text{A}}^{\text{stat}}(x_0) \left[\frac{f_{\text{A}}^{\text{kin}}(x_0)}{f_{\text{A}}^{\text{stat}}(x_0)} - \frac{f_{\delta\text{A}}^{\text{kin}}(x_0)}{f_{\delta\text{A}}^{\text{stat}}(x_0)} \right] \quad (\text{B.4})$$

have the following form

$$P_{\text{A}}^{\text{stat}} = b_1 + b_2 e^{-\Delta^{\text{stat}} x_0}, \quad (\text{B.5})$$

$$P_{\text{A}}^{\text{kin}} = b_3 + b_4 e^{-\Delta^{\text{stat}} x_0} - b_2 \Delta^{\text{kin}} x_0 e^{-\Delta^{\text{stat}} x_0}. \quad (\text{B.6})$$

They can therefore be used to further constrain Δ^{stat} and Δ^{kin} . We are thus lead to perform a combined fit

$$\Gamma^{\text{stat}} = b_5 + b_6 e^{-\Delta^{\text{stat}} x_0}, \quad (\text{B.7})$$

$$\Gamma^{\text{kin}} = b_7 + b_8 e^{-\Delta^{\text{stat}} x_0} - b_6 \Delta^{\text{kin}} x_0 e^{-\Delta^{\text{stat}} x_0}, \quad (\text{B.8})$$

together with eq. (B.5) and (B.6), with non-linear parameters $a_1 = \Delta^{\text{stat}}$ and $a_2 = \Delta^{\text{kin}}$ and the linear parameters b_i , which contain the desired $b_5 = E_{\text{stat}}$ and $b_7 = E_{\text{kin}}$.

Since the correction terms are nevertheless not so easy to compute at the smaller lattice spacings, we perform the above fit first at $\beta = 6.0219$ and extract $a\Delta^{\text{stat}}$ and $a^2\Delta^{\text{kin}}$. We then use that these quantities scale *roughly* (i.e. $r_0 \Delta^{\text{stat}} \approx \text{constant}$ and $r_0^2 \Delta^{\text{kin}} \approx \text{constant}$). To implement this, we input the scaled means as priors [65] in a second step where we add

$$\chi_{\text{prior}}^2 = \sum_{i=1,2} \frac{(a_i - a_i^{\text{prior}})^2}{(\delta a_i^{\text{prior}})^2}, \quad (\text{B.9})$$




Figure 9: Results for P_A^{stat} , P_A^{kin} , Γ^{stat} and Γ^{kin} at $\beta = 6.2885$ (HYP2) with the corresponding functions obtained by the fit.

to the standard χ^2 . The uncertainty $\delta a_i^{\text{prior}}$ is taken from the fit result at $\beta = 6.0219$. However, in order to remain on the safe side, it is not scaled but kept constant at the smaller lattice spacing. Thus $\delta a_2^{\text{prior}}/a_2^{\text{prior}} \propto 1/a^2$ for example. The constraint due to the priors becomes weaker as we approach the continuum.

Here and in the following procedure the fit range is chosen to keep a minimum physical distance from the boundaries, namely $x_0 \geq t_{\text{min}} \approx 2r_0$. The stability of the results is checked by varying t_{min} to $t_{\text{min}} - r_0/2$. As an example we show in figure 9 the results for P_A^{stat} , P_A^{kin} , Γ^{stat} and Γ^{kin} at $\beta = 6.2885$. One observes that P_A^{stat} , Γ^{stat} provide very good constraints of the parameters Δ^{stat} , b_2 , b_6 . The remaining ones are then effectively linear fit parameters. Nevertheless, the error band of E_{kin} (dashed line) resulting from the fit is not that small.

An alternative strategy is used to get a second estimate of E_{stat} at the two coarser lattice spacings, where we have two independent datasets. Exploiting again the remark before eq. (B.4) we construct an effective mass $\Gamma_{\delta A}$ from the correlator $f_{\delta A}(x_0)$ in the very same way as Γ^{stat} is obtained from $f_A^{\text{stat}}(x_0)$. The idea is to combine the two effective masses in order to eliminate the contribution from the first excited state and then perform

a fit to a constant (in the mentioned fit range). In practice we minimize the quantity

$$Q = \sum_{x_0=t_{\min}}^{t_{\max}} [\Gamma^{\text{stat}}(x_0) + \alpha\Gamma_{\delta A}(x_0) - B]^2, \quad (\text{B.10})$$

with respect to α , B . Finally the weighted average of $[\Gamma^{\text{stat}}(x_0) + \alpha\Gamma_{\delta A}(x_0)]/(1+\alpha)$ yields the estimate of E^{stat} . The quality of the result is comparable to that obtained in the first approach.

C. Alternative strategy

We briefly introduce an alternative strategy, based on the correlation functions f_A, k_V in addition to f_1, k_1 . With

$$F_{\text{av}}(x_0, \theta) = \frac{1}{4} \log \left(-[f_A]_{\text{R}}(x_0, \theta) \times [k_V]_{\text{R}}^3(x_0, \theta) \right) \quad (\text{C.1})$$

we introduce

$$R_{\text{av}}(L, \theta_1, \theta_2) = F_{\text{av}}(x_0, \theta_1) - F_{\text{av}}(x_0, \theta_2) \text{ at } x_0 = L/2, T = L \quad (\text{C.2})$$

$$\Gamma^{\text{av}}(L, \theta_0) = -\frac{\partial_0 + \partial_0^*}{2} F_{\text{av}}(x_0, \theta_0) \text{ at } x_0 = L/2, T = L. \quad (\text{C.3})$$

Keeping Φ_1 , from the standard strategy, we define the set of observables

$$\tilde{\Phi}_1(L, M_b) = \Phi_1(L, M_b), \quad (\text{C.4})$$

$$\tilde{\Phi}_2(L, M_b) = R_{\text{av}}(L, \theta_1, \theta_2) - R_{\text{av}}^{\text{stat}}(L, \theta_1, \theta_2), \quad (\text{C.5})$$

$$\tilde{\Phi}_3(L, M_b) = L\Gamma^{\text{av}}(L, \theta_0), \quad (\text{C.6})$$

with the $1/m_b$ expansion

$$\tilde{\Phi}_2(L, M_b) = \omega_{\text{kin}} R_{\text{A}}^{\text{kin}}(L, \theta_1, \theta_2) + c_{\text{av}}^{\text{HQET}} R_{\delta A}(L, \theta_1, \theta_2) \quad (\text{C.7})$$

$$\tilde{\Phi}_3(L, M_b) = L [m_{\text{bare}} + \Gamma^{\text{stat}}(L, \theta_0) + \omega_{\text{kin}} \Gamma^{\text{kin}}(L, \theta_0) + c_{\text{av}}^{\text{HQET}} \Gamma_{\delta A}(L, \theta_0)], \quad (\text{C.8})$$

where due to the spin average the combination

$$c_{\text{av}}^{\text{HQET}} = \frac{1}{4} [c_{\text{A}}^{\text{HQET}} + 3c_{\text{V}}^{\text{HQET}}] \quad (\text{C.9})$$

is present. The so far undefined terms $R_{\text{av}}^{\text{stat}}, R_{\text{A}}^{\text{kin}}, \Gamma^{\text{kin}}, R_{\delta A}, \Gamma_{\delta A}$ are straightforwardly obtained from our definitions.

The alternative observables change from L to $2L$ via

$$\tilde{\Phi}_i(2L, M_b) = \sum_{j \leq i} \sigma_{ij}(u) \tilde{\Phi}_j(L, M_b) + \delta_{i3} \tilde{\sigma}_m(u), \quad (\text{C.10})$$

$$\sigma_{ij}(u) = \lim_{a/L \rightarrow 0} \Sigma_{ij}(u, a/L) \quad (\text{C.11})$$

with the step scaling functions (we drop arguments θ_1, θ_2 and $u = \bar{g}^2(L)$ is understood)

θ_0	$r_0 M_b^{(0)}$	$r_0 M_b^{(1a)}$			$r_0 M_b^{(1b)}$		
		$\theta_1 = 0$	$\theta_1 = 1/2$	$\theta_1 = 1$	$\theta_1 = 0$	$\theta_1 = 1/2$	$\theta_1 = 1$
		$\theta_2 = 1/2$	$\theta_2 = 1$	$\theta_2 = 0$	$\theta_2 = 1/2$	$\theta_2 = 1$	$\theta_2 = 0$
0	17.05(25)	0.17(6)	0.17(6)	0.17(6)	0.02(9)	0.02(8)	0.02(9)
1/2	17.01(22)	0.20(7)	0.18(6)	0.19(7)	0.02(10)	0.02(9)	0.02(9)
1	16.78(28)	0.34(11)	0.30(7)	0.32(8)	0.06(12)	0.06(9)	0.06(10)

Table 12: RGI results of M_b in the static approximation and of the $1/m_b$ correction for the alternative strategy.

$$\Sigma_{11}(u, a/L) = R_1^{\text{kin}}(2L)/R_1^{\text{kin}}(L) = \Sigma_1^{\text{kin}}(u, a/L) \quad (\text{C.12})$$

$$\Sigma_{21}(u, a/L) = \frac{1}{R_1^{\text{kin}}(L)} \{R_A^{\text{kin}}(2L) - R_A^{\text{kin}}(L) \Sigma_{22}(u, a/L)\} \quad (\text{C.13})$$

$$\Sigma_{22}(u, a/L) = R_{\delta A}(2L)/R_{\delta A}(L) \quad (\text{C.14})$$

$$\Sigma_{31}(u, a/L) = \frac{2L\{\Gamma^{\text{kin}}(2L) - \Gamma^{\text{kin}}(L)\}}{R_1^{\text{kin}}(L)} - \Sigma_{32}(u, a/L) \frac{R_A^{\text{kin}}(L)}{R_1^{\text{kin}}(L)} \quad (\text{C.15})$$

$$\Sigma_{32}(u, a/L) = 2L \frac{\Gamma_{\delta A}(2L) - \Gamma_{\delta A}(L)}{R_{\delta A}(L)} \quad (\text{C.16})$$

$$\Sigma_{33}(u, a/L) = 2 \quad (\text{C.17})$$

$$\widetilde{\sigma}_m(u) = \lim_{a/L \rightarrow 0} 2L [\Gamma^{\text{stat}}(2L) - \Gamma^{\text{stat}}(L)]. \quad (\text{C.18})$$

The final relation for the B-meson mass is eq. (2.30) with

$$L_2 m_B^{(0a)}(M_b) = \widetilde{\sigma}_m(u_1) + 2\widetilde{\Phi}_3(L_1, M_b), \quad (\text{C.19})$$

$$L_2 m_B^{(0b)}(M_b) = L_2 [E^{\text{stat}} - \Gamma^{\text{stat}}(L_2)], \quad (\text{C.20})$$

$$L_2 m_B^{(1a)}(M_b) = \sigma_{31}(u_1) \widetilde{\Phi}_1(L_1, M_b) + \sigma_{32}(u_1) \widetilde{\Phi}_2(L_1, M_b), \quad (\text{C.21})$$

$$L_2 m_B^{(1b)}(M_b) = L_2 \left[\frac{E^{\text{kin}} - \Gamma^{\text{kin}}(L_2)}{R_1^{\text{kin}}(L_2)} + \frac{\Gamma_{\delta A}(L_2) R_A^{\text{kin}}(L_2)}{R_{\delta A}(L_2) R_1^{\text{kin}}(L_2)} \right] \sigma_1^{\text{kin}}(u_1) \widetilde{\Phi}_1(L_1, M_b) \\ - L_2 \frac{\Gamma_{\delta A}(L_2)}{R_{\delta A}(L_2)} \left[\sigma_{21}(u_1) \widetilde{\Phi}_1(L_1, M_b) + \sigma_{22}(u_1) \widetilde{\Phi}_2(L_1, M_b) \right]. \quad (\text{C.22})$$

Although the results have been already given in table 2, the reader will find more details in table 12.

D. Propagating uncertainties in L_i/r_0 and $\bar{g}^2(L_i)$

In our simulations we have fixed \tilde{L}_1 by $\bar{g}^2(\tilde{L}_1/4) = 1.8811$, because the corresponding bare parameters β, κ are available in the literature. We here give the estimate of the small effect caused by $\tilde{L}_1 \neq L_1$ in the static approximation. From the polynomial interpolations of the

step scaling function of the coupling, $\sigma(u)$ [20], we estimate the corresponding mismatch in couplings as

$$\tilde{u} - u = \bar{g}^2(\tilde{L}_1) - \bar{g}^2(L_1) = \sigma(\sigma(1.8811)) - 3.48 = -0.17(5). \quad (\text{D.1})$$

Let us write

$$\frac{M_b}{m_B} = \rho(\tilde{u}, z) [1 + K(u)] \quad \text{at} \quad \tilde{u} = u \quad (\text{D.2})$$

with

$$K(u) = \frac{\Gamma_1^{\text{stat}}(L_1) - E_{\text{stat}}}{m_B}, \quad \rho(u, z) = \frac{z}{\Phi_2(u, z)}. \quad (\text{D.3})$$

The relation $\frac{d}{du} \frac{M_b}{m_B} = 0$ gives

$$-\frac{1 + K(u)}{\rho(u, z)} \frac{d}{du} \rho(u, z) = K'(u) = \frac{1}{m_B} \frac{d}{du} \Gamma_1^{\text{stat}}. \quad (\text{D.4})$$

Denoting by ΔM_b the correction we have to add to M_b when it is computed with $\tilde{u} \neq u$ (as we did), we get from the above equations

$$\frac{1}{m_B} \Delta M_b = [\tilde{u} - u] \times \rho(u) K'(u), \quad (\text{D.5})$$

where $K'(u)$ is easily estimated by taking a numerical derivative of Γ_1^{stat} . From the difference of $L/a = 12$ and $L/a = 10$ at fixed g_0^2 (with $\bar{g}^2|_{L/a=12} = 3.48$) and with $\rho(u, z) \approx 1.44$ we arrive at the small shift

$$r_0 \Delta M_b = -0.055(17). \quad (\text{D.6})$$

A similar error is be taken into account due to the 2% uncertainty in the relation $L_2 = 1.436r_0$ [48]. In the same way it leads to a *statistical error* of

$$r_0 \Delta M_b = 0.016. \quad (\text{D.7})$$

The two contributions eq. (D.6), eq. (D.7) are combined to

$$r_0 \Delta M_b = -0.055(23), \quad (\text{D.8})$$

which we have taken into account in section 5.3. Because of the smallness of these effects, they can be neglected in the $1/m_b$ -corrections.

In the case of our alternative strategy, the shift depends on the value of θ_0 . We find

$$\theta_0 = 0 \quad r_0 \Delta M_b = -0.042(20), \quad (\text{D.9})$$

$$\theta_0 = 1/2 \quad r_0 \Delta M_b = 0.009(11), \quad (\text{D.10})$$

$$\theta_0 = 1 \quad r_0 \Delta M_b = 0.150(45). \quad (\text{D.11})$$

References

- [1] ALPHA collaboration, J. Heitger and R. Sommer, *Non-perturbative heavy quark effective theory*, *JHEP* **02** (2004) 022 [[hep-lat/0310035](#)].
- [2] I.I.Y. Bigi, N.G. Uraltsev and A.I. Vainshtein, *Nonperturbative corrections to inclusive beauty and charm decays: QCD versus phenomenological models*, *Phys. Lett.* **B 293** (1992) 430 [[hep-ph/9207214](#)].
- [3] M. Battaglia et al., *The CKM matrix and the unitarity triangle*, [hep-ph/0304132](#).
- [4] K. Melnikov and A. Yelkhovsky, *The B quark low-scale running mass from epsilon sum rules*, *Phys. Rev.* **D 59** (1999) 114009 [[hep-ph/9805270](#)].
- [5] M. Beneke and A. Signer, *The bottom $\bar{M}S$ quark mass from sum rules at next-to-next-to-leading order*, *Phys. Lett.* **B 471** (1999) 233 [[hep-ph/9906475](#)].
- [6] A.H. Hoang, *1S and $\bar{M}S$ bottom quark masses from epsilon sum rules*, *Phys. Rev.* **D 61** (2000) 034005 [[hep-ph/9905550](#)].
- [7] J.H. Kühn and M. Steinhauser, *Determination of α_s and heavy quark masses from recent measurements of R_s* , *Nucl. Phys.* **B 619** (2001) 588 [[hep-ph/0109084](#)].
- [8] J. Erler and M.X. Luo, *Precision determination of heavy quark masses and the strong coupling constant*, *Phys. Lett.* **B 558** (2003) 125 [[hep-ph/0207114](#)].
- [9] M. Eidemuller, *QCD moment sum rules for coulomb systems: the charm and bottom quark masses*, *Phys. Rev.* **D 67** (2003) 113002 [[hep-ph/0207237](#)].
- [10] G. Corcella and A.H. Hoang, *Uncertainties in the $\bar{M}S$ bottom quark mass from relativistic sum rules*, *Phys. Lett.* **B 554** (2003) 133 [[hep-ph/0212297](#)].
- [11] J. Bordes, J. Penarrocha and K. Schilcher, *Bottom quark mass and QCD duality*, *Phys. Lett.* **B 562** (2003) 81 [[hep-ph/0212083](#)].
- [12] A. Pineda and A. Signer, *Renormalization group improved sum rule analysis for the bottom quark mass*, *Phys. Rev.* **D 73** (2006) 111501 [[hep-ph/0601185](#)].
- [13] R. Boughezal, M. Czakon and T. Schutzmeier, *Charm and bottom quark masses from perturbative QCD*, *Phys. Rev.* **D 74** (2006) 074006 [[hep-ph/0605023](#)].
- [14] A.A. Penin and A.A. Pivovarov, *Next-to-next-to-leading order vacuum polarization function of heavy quark near threshold and sum rules for B anti-B system*, *Phys. Lett.* **B 435** (1998) 413 [[hep-ph/9803363](#)].
- [15] A.A. Penin and A.A. Pivovarov, *Bottom quark pole mass and $-v(cb)$ matrix element from $R(e^+e^- \rightarrow B\bar{B})$ and $\gamma(SL)(B \rightarrow CL\nu/l)$ in the next-to-next-to-leading order*, *Nucl. Phys.* **B 549** (1999) 217 [[hep-ph/9807421](#)].
- [16] ALPHA collaboration, R. Sommer and H. Wittig, 2002, [physics/0204015](#).
- [17] E. Eichten and B. Hill, *Static effective field theory: $1/M$ corrections*, *Phys. Lett.* **B 243** (1990) 427.
- [18] G. Martinelli and C.T. Sachrajda, *Computation of the b-quark mass with perturbative matching at the next-to-next-to-leading order*, *Nucl. Phys.* **B 559** (1999) 429 [[hep-lat/9812001](#)].

- [19] F. Di Renzo and L. Scorzato, *The residual mass in lattice heavy quark effective theory to α^3 order*, *JHEP* **02** (2001) 020 [[hep-lat/0012011](#)].
- [20] ALPHA collaboration, S. Capitani, M. Lüscher, R. Sommer and H. Wittig, *Non-perturbative quark mass renormalization in quenched lattice QCD*, *Nucl. Phys.* **B 544** (1999) 669 [[hep-lat/9810063](#)].
- [21] ALPHA collaboration, M. Della Morte et al., *Non-perturbative quark mass renormalization in two-flavor QCD*, *Nucl. Phys.* **B 729** (2005) 117 [[hep-lat/0507035](#)].
- [22] ALPHA collaboration, J. Heitger, A. Jüttner, R. Sommer and J. Wennekers, *Non-perturbative tests of heavy quark effective theory*, *JHEP* **11** (2004) 048 [[hep-ph/0407227](#)].
- [23] ALPHA collaboration, M. Kurth and R. Sommer, *Heavy quark effective theory at one-loop order: an explicit example*, *Nucl. Phys.* **B 623** (2002) 271 [[hep-lat/0108018](#)].
- [24] M. Lüscher, R. Narayanan, P. Weisz and U. Wolff, *The Schrödinger functional: a renormalizable probe for nonabelian gauge theories*, *Nucl. Phys.* **B 384** (1992) 168 [[hep-lat/9207009](#)].
- [25] S. Sint, *On the Schrödinger functional in QCD*, *Nucl. Phys.* **B 421** (1994) 135 [[hep-lat/9312079](#)].
- [26] E. Eichten and B. Hill, *An effective field theory for the calculation of matrix elements involving heavy quarks*, *Phys. Lett.* **B 234** (1990) 511.
- [27] M. Della Morte, A. Shindler and R. Sommer, *On lattice actions for static quarks*, *JHEP* **08** (2005) 051 [[hep-lat/0506008](#)].
- [28] ALPHA collaboration, M. Kurth and R. Sommer, *Renormalization and $O(a)$ -improvement of the static axial current*, *Nucl. Phys.* **B 597** (2001) 488 [[hep-lat/0007002](#)].
- [29] M. Bochicchio, L. Maiani, G. Martinelli, G.C. Rossi and M. Testa, *Chiral symmetry on the lattice with Wilson fermions*, *Nucl. Phys.* **B 262** (1985) 331.
- [30] M. Lüscher, S. Sint, R. Sommer and H. Wittig, *Non-perturbative determination of the axial current normalization constant in $O(a)$ improved lattice QCD*, *Nucl. Phys.* **B 491** (1997) 344 [[hep-lat/9611015](#)].
- [31] ALPHA collaboration, J. Heitger and J. Wennekers, *Effective heavy-light meson energies in small-volume quenched QCD*, *JHEP* **02** (2004) 064 [[hep-lat/0312016](#)].
- [32] M. Lüscher, S. Sint, R. Sommer and P. Weisz, *Chiral symmetry and $O(a)$ improvement in lattice QCD*, *Nucl. Phys.* **B 478** (1996) 365 [[hep-lat/9605038](#)].
- [33] S. Sint and P. Weisz, *Further results on $o(a)$ improved lattice QCD to one-loop order of perturbation theory*, *Nucl. Phys.* **B 502** (1997) 251 [[hep-lat/9704001](#)].
- [34] ALPHA collaboration, J. Heitger and R. Sommer, *A strategy to compute the b quark mass with non-perturbative accuracy*, *Nucl. Phys.* **106** (*Proc. Suppl.*) (2002) 358 [[hep-lat/0110016](#)].
- [35] M. Lüscher, R. Sommer, P. Weisz and U. Wolff, *A precise determination of the running coupling in the $SU(3)$ Yang-Mills theory*, *Nucl. Phys.* **B 413** (1994) 481 [[hep-lat/9309005](#)].
- [36] ALPHA collaboration, R. Frezzotti, P.A. Grassi, S. Sint and P. Weisz, *Lattice QCD with a chirally twisted mass term*, *JHEP* **08** (2001) 058 [[hep-lat/0101001](#)].

- [37] H. Neuberger, *Exactly massless quarks on the lattice*, *Phys. Lett.* **B 417** (1998) 141 [hep-lat/9707022].
- [38] M. Lüscher, *Exact chiral symmetry on the lattice and the Ginsparg-Wilson relation*, *Phys. Lett.* **B 428** (1998) 342 [hep-lat/9802011].
- [39] P. Hasenfratz, V. Laliena and F. Niedermayer, *The index theorem in QCD with a finite cut-off*, *Phys. Lett.* **B 427** (1998) 125 [hep-lat/9801021].
- [40] Y. Shamir, *Chiral fermions from lattice boundaries*, *Nucl. Phys.* **B 406** (1993) 90 [hep-lat/9303005].
- [41] B. Sheikholeslami and R. Wohlert, *Improved continuum limit lattice action for QCD with Wilson fermions*, *Nucl. Phys.* **B 259** (1985) 572.
- [42] M. Lüscher, S. Sint, R. Sommer, P. Weisz and U. Wolff, *Non-perturbative $O(a)$ improvement of lattice QCD*, *Nucl. Phys.* **B 491** (1997) 323 [hep-lat/9609035].
- [43] R. Sommer, *A new way to set the energy scale in lattice gauge theories and its applications to the static force and α_s in SU(2) Yang-Mills theory*, *Nucl. Phys.* **B 411** (1994) 839 [hep-lat/9310022].
- [44] ALPHA collaboration, M. Guagnelli, R. Sommer and H. Wittig, *Precision computation of a low-energy reference scale in quenched lattice QCD*, *Nucl. Phys.* **B 535** (1998) 389 [hep-lat/9806005].
- [45] ALPHA collaboration, M. Della Morte et al., *Lattice HQET with exponentially improved statistical precision*, *Phys. Lett.* **B 581** (2004) 93 [hep-lat/0307021].
- [46] ALPHA collaboration, M. Guagnelli, J. Heitger, R. Sommer and H. Wittig, *Hadron masses and matrix elements from the QCD Schrödinger functional*, *Nucl. Phys.* **B 560** (1999) 465 [hep-lat/9903040].
- [47] UKQCD collaboration, C. Michael and J. Peisa, *Maximal variance reduction for stochastic propagators with applications to the static quark spectrum*, *Phys. Rev.* **D 58** (1998) 034506 [hep-lat/9802015].
- [48] S. Necco and R. Sommer, *The $N_f = 0$ heavy quark potential from short to intermediate distances*, *Nucl. Phys.* **B 622** (2002) 328 [hep-lat/0108008].
- [49] T. van Ritbergen, J.A.M. Vermaseren and S.A. Larin, *The four-loop β function in quantum chromodynamics*, *Phys. Lett.* **B 400** (1997) 379 [hep-ph/9701390].
- [50] K.G. Chetyrkin, *Quark mass anomalous dimension to $O(\alpha_s^4)$* , *Phys. Lett.* **B 404** (1997) 161 [hep-ph/9703278].
- [51] J.A.M. Vermaseren, S.A. Larin and T. van Ritbergen, *The 4-loop quark mass anomalous dimension and the invariant quark mass*, *Phys. Lett.* **B 405** (1997) 327 [hep-ph/9703284].
- [52] M. Czakon, *The four-loop QCD beta-function and anomalous dimensions*, *Nucl. Phys.* **B 710** (2005) 485 [hep-ph/0411261].
- [53] J. Foley et al., *Practical all-to-all propagators for lattice QCD*, *Comput. Phys. Commun.* **172** (2005) 145 [hep-lat/0505023].
- [54] G.M. de Divitiis, M. Guagnelli, R. Petronzio, N. Tantalo and F. Palombi, *Heavy quark masses in the continuum limit of lattice QCD*, *Nucl. Phys.* **B 675** (2003) 309 [hep-lat/0305018].

- [55] D. Guazzini, R. Sommer and N. Tantalo, *PoS LAT 2006* 084.
- [56] V. Lubicz, *Quark masses on the lattice: light and heavy*, *Nucl. Phys.* **94** (*Proc. Suppl.*) (2001) 116 [[hep-lat/0012003](#)].
- [57] V. Gimenez, L. Giusti, G. Martinelli and F. Rapuano, *Nnlo unquenched calculation of the B quark mass*, *JHEP* **03** (2000) 018 [[hep-lat/0002007](#)].
- [58] S. Collins, *The mass of the B quark from lattice nrqcd*, [hep-lat/0009040](#).
- [59] F. Di Renzo and L. Scorzato, *The residual mass in lattice heavy quark effective theory to α^3 order*, *JHEP* **02** (2001) 020 [[hep-lat/0012011](#)].
- [60] G.P. Lepage, P.B. Mackenzie, N.H. Shakespeare and H.D. Trottier, *Perturbative two- and three-loop coefficients from large beta Monte Carlo*, *Nucl. Phys.* **83** (*Proc. Suppl.*) (2000) 866 [[hep-lat/9910018](#)].
- [61] ALPHA collaboration, M. Della Morte et al., *Simulating the schroedinger functional with two pseudo-fermions*, *Comput. Phys. Commun.* **156** (2003) 62 [[hep-lat/0307008](#)].
- [62] S. Dürr, A. Jüttner, J. Rolf and R. Sommer, *Signal at subleading order in lattice Hqet*, *Nucl. Phys.* **140** (*Proc. Suppl.*) (2005) 488 [[hep-lat/0409058](#)].
- [63] ALPHA collaboration, M. Guagnelli et al., *Non-perturbative results for the coefficients $B(m)$ and $B(a)$ - $B(p)$ in $O(a)$ improved lattice QCD*, *Nucl. Phys.* **B 595** (2001) 44 [[hep-lat/0009021](#)].
- [64] ALPHA collaboration, J. Garden, J. Heitger, R. Sommer and H. Wittig, *Precision computation of the strange quark's mass in quenched QCD*, *Nucl. Phys.* **B 571** (2000) 237 [[hep-lat/9906013](#)].
- [65] G.P. Lepage et al., *Constrained curve fitting*, *Nucl. Phys.* **106** (*Proc. Suppl.*) (2002) 12 [[hep-lat/0110175](#)].