

Gravitino dark matter in R-parity breaking vacua

Wilfried Buchmüller,^a Laura Covi,^a Koichi Hamaguchi,^b Alejandro Ibarra,^a Tsutomu T. Yanagida^b

^a *Deutsches Elektronen-Synchrotron DESY
Hamburg, Germany*

^b *Department of Physics, University of Tokyo
Tokyo, Japan*

E-mail: buchmuwi@mail.desy.de, laura.covi@desy.de,
hama@hep-th.phys.s.u-tokyo.ac.jp, alejandro.ibarra@desy.de

ABSTRACT: We show that in the case of small R-parity and lepton number breaking couplings, primordial nucleosynthesis, thermal leptogenesis and gravitino dark matter are naturally consistent for gravitino masses $m_{3/2} \gtrsim 5$ GeV. We present a model where R-parity breaking is tied to B-L breaking, which predicts the needed small couplings. The metastable next-to-lightest superparticle has a decay length that is typically larger than a few centimeters, with characteristic signatures at the LHC. The photon flux produced by relic gravitino decays may be part of the apparent excess in the extragalactic diffuse gamma-ray flux obtained from the EGRET data for a gravitino mass $m_{3/2} \sim 10$ GeV. In this case, a clear signal can be expected from GLAST in the near future.

KEYWORDS: Supersymmetry Phenomenology, Cosmology of Theories beyond the SM, Supergravity Models.

Contents

1.	Introduction	1
2.	R-parity breaking and B-L breaking	3
2.1	A model of R-parity breaking	3
2.2	Scale of B-L breaking and thermal leptogenesis	5
3.	Neutrino masses	6
4.	Cosmology and collider physics	8
4.1	Gravitino decay	8
4.2	Extragalactic diffuse gamma-ray emission	9
4.3	Collider signatures	11
4.4	Microscopic determination of the Planck mass	12
5.	Conclusions	13

1. Introduction

Most supersymmetric extensions of the standard model impose R-parity [1] as an exact symmetry of the supergravity Lagrangian. In this way, one forbids renormalizable baryon and lepton number violating interactions which might cause too rapid proton decay [2]. On theoretical grounds, however, theories with and without R-parity are on the same footing, and in low-energy effective theories obtained from string compactifications R-parity plays no preferred role.

One can also construct supersymmetric extensions of the standard model without R-parity [3], and the phenomenological constraints on these theories have been studied in great detail [4]. Without R-parity conservation, the lightest superparticle (LSP) is no longer stable and, in general, it does not contribute to dark matter.

Stringent constraints on the lepton number and R-parity violating interactions

$$W_{\Delta L=1} = \lambda_{ikj} l_i e_j^c l_k + \lambda'_{kji} d_i^c q_j l_k \tag{1.1}$$

are imposed by baryogenesis. Both operators contain lepton doublets. Together with sphaleron processes they therefore influence the baryon asymmetry at high temperature in the early universe. The requirement that an existing baryon asymmetry is not erased before the electroweak transition typically implies [5]

$$\lambda, \lambda' < 10^{-7} . \tag{1.2}$$

It is very remarkable that for such a small breaking of R-parity a gravitino LSP has a lifetime much longer than the age of the universe [6]. This is due to the double suppression by the inverse Planck mass and the R-parity breaking coupling, $\Gamma_{3/2} \propto \lambda^2 m_{3/2}^3 / M_{\text{P}}^2$. We find for the gravitino lifetime

$$\tau_{3/2} \sim 10^{26} \text{s} \left(\frac{\lambda}{10^{-7}} \right)^{-2} \left(\frac{m_{3/2}}{10 \text{ GeV}} \right)^{-3}, \quad (1.3)$$

which is consistent with gravitino dark matter.

For a gravitino LSP, the properties of the next-to-lightest superparticle (NLSP) are strongly constrained by primordial nucleosynthesis (BBN). In the particularly interesting case of a charged NLSP, like a scalar τ -lepton, its lifetime has to be relatively short, $\tau_{\text{NLSP}} \lesssim 10^3 - 10^4 \text{ s}$ [7]¹, which typically requires $m_{3/2} < 1 \text{ GeV}$. Even for neutral particles, BBN excludes a neutralino NLSP for lifetimes longer than 10^2 s due to the strong constraints from hadronic showers [9]. Only a sneutrino NLSP could be marginally acceptable also with longer lifetimes, and therefore larger gravitino mass, in the region where the hadronic branching ratio of the decay is below 10^{-3} [11].

On the other hand, standard thermal leptogenesis [12], an attractive model for baryogenesis, needs a large reheating temperature in the early universe, $T_R \gtrsim 10^9 \text{ GeV}$ (cf. [13, 14]). This reheating temperature implies $m_{3/2} \gtrsim 5 \text{ GeV}$ for a gluino mass of $m_{\tilde{g}} = 500 \text{ GeV}$ in order to avoid overclosure of the universe due to thermal gravitino production [15, 16]². The lower bound on the gravitino mass scales as $m_{3/2}^{\text{min}} \sim T_R m_{\tilde{g}}^2$.

All these cosmological problems are automatically solved without any fine tuning of parameters in the case of a small breaking of R-parity, as given in eq. (1.2), with a gravitino LSP. The NLSP lifetime becomes sufficiently short for $\lambda, \lambda' > 10^{-14}$,

$$\tau_{\text{NLSP}} \simeq 10^3 \text{s} \left(\frac{\lambda}{10^{-14}} \right)^{-2} \left(\frac{m_{\text{NLSP}}}{100 \text{ GeV}} \right)^{-1}. \quad (1.4)$$

Therefore, primordial nucleosynthesis, thermal leptogenesis and gravitino dark matter are naturally consistent for $10^{-14} < \lambda, \lambda' < 10^{-7}$ and $m_{3/2} \gtrsim 5 \text{ GeV}$. This is the main point of this paper.

The paper is organised as follows. In section 2 we present a model where R-parity breaking is tied to B-L breaking, yielding the needed small R-parity breaking couplings. Section 3 deals with constraints from neutrino masses. Section 4 deals with implications for cosmology and collider physics. The results are discussed in section 5.

¹See also [8]. Here, we consider $m_{\text{NLSP}} = \mathcal{O}(100 \text{ GeV})$. For a heavier charged NLSP, $m_{\text{NLSP}} > \mathcal{O}(1 \text{ TeV})$, the bound on the lifetime becomes even more stringent (cf. [9]). We do not consider a late time entropy production in this paper, which is another possible way to avoid these BBN constraints [10].

²We use the perturbative result for the gravitino production rate to leading order in the strong gauge coupling g . Since g and also the thermal gluon mass are large, the perturbative expansion is problematic [15]. The uncertainty due to higher orders in g and nonperturbative effects is $\mathcal{O}(1)$. Possible effects due to thermal masses are also $\mathcal{O}(1)$ [17].

2. R-parity breaking and B-L breaking

2.1 A model of R-parity breaking

We consider a supersymmetric extension of the standard model whose symmetry group G includes $U(1)_{B-L}$ and R-invariance,

$$G = SU(3) \times SU(2) \times U(1)_Y \times U(1)_{B-L} \times U(1)_R . \quad (2.1)$$

Three quark-lepton generations can be grouped into the $SU(5)$ representations $\mathbf{10}_i = (q, u^c, e^c)_i$, $\bar{\mathbf{5}}_i = (d^c, l)_i$ and $\mathbf{1} = \nu_i^c$, which together form 16-plets of $SO(10)$. In addition, we have two Higgs doublets H_u and H_d , two standard model singlets N^c and N , and three $SO(10)$ singlets X , Φ and Z . The two Higgs doublets are contained in $\mathbf{5}$ - and $\bar{\mathbf{5}}$ -plets of $SU(5)$, which we shall also denote as H_u and H_d , respectively. N^c and N are contained in $\mathbf{16}$ and $\bar{\mathbf{16}}$ of $SO(10)$, which fixes their B-L charge to be +1 and -1, respectively. X , Φ and Z have B-L charge zero. This set of fields is familiar from $SO(10)$ orbifold GUTs (cf. [18]): matter fields form complete $SO(10)$ representations, whereas fields which break $SU(2) \times U(1)_Y$ and $U(1)_{B-L}$ appear as ‘split’ multiplets. For simplicity, we shall use in the following often $SU(5)$ notation also for the Higgs multiplets.

The matter sector of the superpotential has the usual form

$$W_M = h_{ij}^{(u)} \mathbf{10}_i \mathbf{10}_j H_u + h_{ij}^{(d)} \bar{\mathbf{5}}_i \mathbf{10}_j H_d + h_{ij}^{(\nu)} \bar{\mathbf{5}}_i \mathbf{1}_j H_d + \frac{1}{M_P} h_{ij}^{(n)} \mathbf{1}_i \mathbf{1}_j N^2 , \quad (2.2)$$

where $M_P = 2.4 \times 10^{18}$ GeV is the Planck mass. The expectation values of the Higgs multiplets H_u and H_d generate Dirac masses of quarks and leptons, whereas the expectation value of the singlet Higgs field N generates the Majorana mass matrix of the right-handed neutrinos $\mathbf{1}_i$. The superpotential responsible for B-L breaking is chosen as

$$W_{B-L} = X(NN^c - \Phi^2) , \quad (2.3)$$

where unknown Yukawa couplings have been set equal to one. Φ plays the role of a spectator field, which will finally be replaced by its expectation value, $\langle \Phi \rangle = v_{B-L}$. Similarly, Z is a spectator field³, which breaks supersymmetry and $U(1)_R$, $\langle Z \rangle = F_Z \theta \theta$. The superpotential in eqs. (2.2) and (2.3) is the most general one consistent with the R-charges listed in table 1, up to higher order terms which we will discuss later. Note that the choice of a negative R-charge for N^c forbids the dangerous superpotential terms

$$\bar{\mathbf{5}}_i H_d N^c , \quad \bar{\mathbf{5}}_i \bar{\mathbf{5}}_j \mathbf{10}_k N^c , \quad (2.4)$$

which would yield too large bilinear mixings and too rapid proton decay, respectively.

The expectation value of Φ leads to the breaking of $B - L$,

$$\langle N \rangle = \langle N^c \rangle = \langle \Phi \rangle = v_{B-L} , \quad (2.5)$$

³For simplicity, we use a spectator chiral superfield to describe supersymmetry breaking. The field Z is not essential for the connection between R-parity breaking and B-L breaking discussed in this section.

	$\mathbf{10}_i$	$\mathbf{5}_i^*$	$\mathbf{1}_i$	H_u	H_d	N	N^c	Φ	X	Z
R	1	1	1	0	0	0	-2	-1	4	0

Table 1: R-charges of matter fields, Higgs fields and $SO(10)$ singlets.

where the first equality is a consequence of the $U(1)_{B-L}$ D-term. This generates a Majorana mass matrix M for the right-handed neutrinos with three large eigenvalues, with $M_1 < M_2 < M_3$. If the largest eigenvalue of $h^{(n)}$ is $\mathcal{O}(1)$, one has $M_3 \simeq v_{B-L}^2/M_P$. The heavy Majorana neutrinos can be integrated out yielding for the matter part of the superpotential

$$W_M = h_{ij}^{(u)} \mathbf{10}_i \mathbf{10}_j H_u + h_{ij}^{(d)} \bar{\mathbf{5}}_i \mathbf{10}_j H_d - \frac{1}{2} (h^{(\nu)} \frac{1}{M} h^{(\nu)T})_{ij} (\bar{\mathbf{5}}_i H_u) (\bar{\mathbf{5}}_j H_u), \quad (2.6)$$

with the familiar dimension-5 seesaw operator for light neutrino masses.

Since the field Φ carries R-charge -1 , the VEV $\langle \Phi \rangle$ breaks R-parity, which is conserved by the VEV $\langle Z \rangle$. Thus, the breaking of B-L is tied to the breaking of R-parity. This is the key feature of the mechanism for R-parity breaking presented in this paper⁴. The breaking of R-parity is transmitted to the low-energy degrees of freedom via higher-dimensional operators in the superpotential and the Kähler potential. The leading correction to the Kähler potential is

$$\delta K_1 = \frac{1}{M_P^3} (a_i Z^\dagger + a'_i Z) \Phi^\dagger N^c \bar{\mathbf{5}}_i H_u + \frac{1}{M_P^3} (c_i Z^\dagger + c'_i Z) \Phi N^\dagger \bar{\mathbf{5}}_i H_u + h.c. \quad (2.7)$$

Replacing the spectator fields Z and Φ , as well as N^c by their expectation values, one obtains the correction to the superpotential

$$\delta W_1 = \mu_i \Theta \bar{\mathbf{5}}_i H_u, \quad (2.8)$$

with

$$\mu_i = \mathcal{O}(m_{3/2}), \quad \Theta = \frac{v_{B-L}^2}{M_P^2} \simeq \frac{M_3}{M_P}, \quad (2.9)$$

where $m_{3/2} = F_Z/(\sqrt{3}M_P)$ is the gravitino mass. Note that Θ can be increased or decreased by an appropriate choice of Yukawa couplings in eqs. (2.2) and (2.3). Eq. (2.8) is the familiar bilinear R-parity breaking term [3]. The correction to the Kähler potential

$$\delta K_0 = \frac{k}{M_P} Z^\dagger H_d H_u + h.c. \quad (2.10)$$

yields the corresponding R-parity conserving term [21]

$$\delta W_0 = \mu H_d H_u, \quad \mu = \mathcal{O}(m_{3/2}). \quad (2.11)$$

Note that μ and μ_i are generated by operators of different mass dimension. Hence, their values may easily differ by one or two orders of magnitude, allowing for $\mu > \mu_i, m_{3/2}$ and a gravitino LSP.

⁴For a recent discussion of the connection between B-L breaking and R-parity breaking in the context of string compactifications, see [19, 20].

To analyse the complete superpotential including the R-symmetry breaking terms, it is convenient to perform a rotation of the Higgs and lepton superfields,

$$H_d = H'_d - \epsilon_i l'_i, \quad l_i = l'_i + \epsilon_i H'_d, \quad (2.12)$$

where $\epsilon_i = \mu_i \Theta / \mu$. In terms of the new fields the superpotential reads

$$\begin{aligned} W &= W_M + \delta W_0 + \delta W_1 \\ &= \mu H'_d H_u + h_{ij}^{(u)} q_i u_j^c H_u + h_{ij}^{(d)} d_i^c q_j H'_d + h_{ij}^{(e)} l'_i e_j^c H'_d \\ &\quad - \epsilon_k h_{ij}^{(d)} d_i^c q_j l'_k - \epsilon_k h_{ij}^{(e)} l'_i e_j^c l'_k - \frac{1}{2} (h^{(\nu)} \frac{1}{M} h^{(\nu)T})_{ij} (l'_i H_u) (l'_j H_u) + \mathcal{O}(\epsilon^2, \epsilon m_\nu). \end{aligned} \quad (2.13)$$

The mixing of Higgs and lepton superfields has induced trilinear R-parity breaking terms $\mathcal{O}(\epsilon)$. As we will discuss in section 3, the mixing terms yield vacuum expectation values for the scalar neutrinos that in turn induce mixing terms $\mathcal{O}(\epsilon)$ of neutrinos with the neutralinos, and neutrino masses suppressed by $\mathcal{O}(\epsilon^2)$.

It is remarkable that the potentially dangerous operator leading to proton decay is strongly suppressed compared to the trilinear terms $\mathcal{O}(\epsilon)$ in eq. (2.13). The leading operator is

$$\delta W_2 = \frac{1}{M_{\text{P}}^5} u^c d^c d^c N^c \Phi^3 X. \quad (2.14)$$

For global supersymmetry one has $\langle X \rangle = 0$, which in supergravity is modified to $\langle X \rangle = \mathcal{O}(m_{3/2})^5$. One then obtains

$$\delta W_2 \propto \frac{m_{3/2} v_{B-L}^4}{M_{\text{P}}^5} u^c d^c d^c + \dots \quad (2.15)$$

For λ, λ' satisfying eq. (1.2), the coefficient of the dangerous dimension-4 $\Delta B = 1$ operator is much smaller than the upper bound from the proton lifetime [4].

2.2 Scale of B-L breaking and thermal leptogenesis

The phenomenological viability of the model depends on the size of R-parity breaking mixings ϵ_i and therefore on the scale v_{B-L} of R-parity breaking. An important constraint comes from baryogenesis. As already discussed in the introduction, the potential washout of a baryon asymmetry before the electroweak phase transition is avoided if the R-parity violating Yukawa couplings satisfy $\lambda_{ijk}, \lambda'_{ijk} \lesssim 10^{-7}$, which in turn implies:

$$\left(\frac{\epsilon_i}{10^{-6}} \right) \left(\frac{\tan \beta}{10} \right) \lesssim 1. \quad (2.16)$$

This is a sufficient condition, which can be relaxed for some flavour structures [5].

As an illustration for possible scales of B-L breaking we use a model [22] for quark and lepton mass hierarchies based on a Froggatt-Nielsen $U(1)$ flavour symmetry. The mass hierarchy is generated by the expectation value of a singlet field ϕ with charge $Q_\phi = -1$

⁵The VEV $\langle X \rangle = \mathcal{O}(m_{3/2})$ also causes an additional contribution to the bilinear term via $\delta W = (1/M_{\text{P}}^3) X \Phi N^c \bar{5}_i H_u$, which is comparable to those from δK_1 .

ψ_i	$\mathbf{10}_3$	$\mathbf{10}_2$	$\mathbf{10}_1$	$\mathbf{5}_3^*$	$\mathbf{5}_2^*$	$\mathbf{5}_1^*$	$\mathbf{1}_3$	$\mathbf{1}_2$	$\mathbf{1}_1$	H_u	H_d	Φ	X	Z
Q_i	0	1	2	a	a	a+1	b	c	d	0	0	0	0	0

Table 2: Chiral charges: $a = 0$ or 1 , and $0 \leq b \leq c \leq d$.

via nonrenormalizable interactions with a scale $\Lambda = \langle \phi \rangle / \eta > \Lambda_{GUT}$, $\eta \simeq 0.06$. Yukawa couplings and bilinear terms for $SU(5)$ multiplets ψ_i with charge Q_i scale like

$$h_{ij} \propto \eta^{Q_i+Q_j}, \quad \mu_i \propto \eta^{Q_i}. \quad (2.17)$$

Charges Q_i describing qualitatively the observed quark and lepton masses and mixings are listed in table 2. The model also predicts the observed baryon asymmetry via leptogenesis for the cases where $a + d = 2$. There are two, at low energies indistinguishable, consistent scales of B-L breaking: $M_3 \sim 10^{15}$ GeV ($a = b = 0, c = 1, d = 2$) and $M_3 \sim 10^{12}$ GeV ($b = c = 0, a = d = 1$). For $\mu_i/\mu = 1.0 \dots 0.01$ these two cases lead to the R-parity breaking mixing parameters (cf. eq. (2.9))

$$(I) \quad \frac{\epsilon_i}{\eta^{Q_i}} = 10^{-3} \dots 10^{-5}, \quad (II) \quad \frac{\epsilon_i}{\eta^{Q_i}} = 10^{-6} \dots 10^{-8}. \quad (2.18)$$

In the extreme case $M_3 \sim M_2 \sim M_1 \sim 10^{10}$ GeV without Froggatt-Nielsen symmetry, where leptogenesis may still work for an appropriate enhancement of the CP asymmetry, one has

$$(III) \quad \epsilon_i = 10^{-8} \dots 10^{-10}. \quad (2.19)$$

In the flavour models (I) and (II) the RPV mixings ϵ_i are suppressed by η^{Q_i} . As we shall see in the following section, model (I) is inconsistent with the constraints from neutrino masses and baryogenesis washout; the models (II) and (III) are consistent with both constraints.

The expected mass scale of right-handed neutrinos depends on the mechanism which breaks B-L. The expectation value of a field with lepton number $L = 2$ can generate heavy Majorana masses via renormalizable Yukawa couplings. With B-L broken at the GUT scale, and for Yukawa coupling $\mathcal{O}(1)$ for the third family, one then obtains the canonical result $M_3 \sim v_{B-L} \sim 10^{15}$ GeV. On the other hand, if right-handed neutrino masses are generated via a nonrenormalizable dimension-5 operator and the expectation value of a field with $L = 1$, as in eq. (2.2), one has instead $M_3 \sim v_{B-L}^2/M_P \sim 10^{12}$ GeV. This illustrates how the two mass scales for M_3 , which correspond to the two cases (I) and (II), respectively, might be obtained.

3. Neutrino masses

The model we are considering generates after supersymmetry breaking bilinear R-parity violating terms, eq. (2.8), and tiny R-parity violating Yukawa couplings, eq. (2.15), that we neglect in what follows. Scenarios with just bilinear R-parity violation have been thoroughly studied in the literature [23]. Here, we will limit ourselves to estimate the size of neutrino masses, following closely [24].

At the high-energy scale, the soft SUSY breaking Lagrangian reads

$$-\mathcal{L}_{soft} = m_{H_d}^2 |H_d|^2 + m_{H_u}^2 |H_u|^2 + m_{\tilde{l}_i}^2 |\tilde{l}_i|^2 + (B H_d H_u + B_i \tilde{l}_i H_u + m_{\tilde{l}_i H_d}^2 \tilde{l}_i H_d^* + h.c.) + \dots \quad (3.1)$$

For the computation of neutrino masses we find convenient to work in the basis where the R-parity violating bilinear couplings in the superpotential are rotated away, $\mu_i = 0$, through the field redefinition eq. (2.12). This choice of basis has the advantage that once the basis has been fixed at the high energy scale, the condition $\mu_i = 0$ holds at any scale, and it is not necessary to redefine the basis again at low energies. We also choose the phases of the lepton doublets such that the ϵ_i are real. In this basis the soft SUSY breaking Lagrangian is given by

$$-\mathcal{L}_{soft} = m_{H'_d}^2 |H'_d|^2 + m_{H_u}^2 |H_u|^2 + m_{\tilde{l}'_i}^2 |\tilde{l}'_i|^2 + (B' H'_d H_u + B'_i \tilde{l}'_i H_u + m_{\tilde{l}'_i H'_d}^2 \tilde{l}'_i H'^*_d + h.c.) + \dots \quad (3.2)$$

where

$$\begin{aligned} m_{H'_d}^2 &= m_{H_d}^2 + \epsilon_i \text{Re}(m_{\tilde{l}_i H_d}^2) + \mathcal{O}(|\epsilon_i|^2), \\ m_{\tilde{l}'_i}^2 &= m_{\tilde{l}_i}^2 - \epsilon_i \text{Re}(m_{\tilde{l}_i H_d}^2) + \mathcal{O}(|\epsilon_i|^2), \\ B' &= B + B_i \epsilon_i, \\ B'_i &= B_i - B \epsilon_i, \\ m_{\tilde{l}'_i H'_d}^2 &= m_{\tilde{l}_i H_d}^2 + \epsilon_i (m_{\tilde{l}_i}^2 - m_{H_d}^2) + \mathcal{O}(|\epsilon_i|^2). \end{aligned} \quad (3.3)$$

Minimisation of the scalar potential yields non-vanishing vacuum expectation values for the neutral components of the Higgs doublets, as well as for the sneutrinos,

$$\langle \tilde{\nu}'_i \rangle = \frac{B'_i \tan \beta + m_{\tilde{l}'_i H'_d}^2}{m_{\tilde{l}'_i}^2 - \frac{1}{2} M_Z^2 \cos 2\beta} \langle H'_d \rangle, \quad v_\nu^2 = \sum_{i=1}^3 \langle \tilde{\nu}'_i \rangle^2. \quad (3.4)$$

These vacuum expectation values induce mixings between neutrinos and gauginos, giving rise to one non-vanishing neutrino mass through an “electroweak” seesaw mechanism. The resulting neutrino mass is

$$m_\nu^{Rp} = \frac{1}{2} g_Z^2 v_\nu^2 \sum_{\alpha=1}^4 \frac{|c_{\tilde{z}\alpha}|^2}{m_{\chi_\alpha^0}}, \quad (3.5)$$

where g_Z is the Z-boson gauge coupling, $m_{\chi_\alpha^0}$ are the neutralino masses and $c_{\tilde{z}\alpha}$ is the Zino component of the neutralino χ_α^0 . In the following estimates we will replace the sum over inverse neutralino masses by the inverse of the characteristic SUSY breaking scale, $1/\tilde{m}$.

The size of the neutrino masses depends crucially on the mechanism for supersymmetry breaking. Generically, one expects

$$B'_i \sim \mu_i \tilde{m} \sim \epsilon_i \tilde{m}^2, \quad m_{\tilde{l}'_i H'_d}^2 = \epsilon_i \tilde{m}^2. \quad (3.6)$$

Then, using eqs. (3.4) and (3.5) we find for $\epsilon_1, \epsilon_2 \leq \epsilon_3$,

$$m_\nu^{Rp} \sim 10^{-4} \text{eV} \left(\frac{\epsilon_3}{10^{-7}} \right)^2 \left(\frac{\tilde{m}}{200 \text{GeV}} \right)^{-1}. \quad (3.7)$$

We can now insert the values of ϵ_3 for the different flavour models considered in the previous section. Clearly, model (I), which has a high scale of B-L breaking, is excluded. In models (II) and (III) one has $\epsilon_3 < 10^{-7}$ and $\epsilon_3 < 10^{-8}$, respectively. Here the neutrino mass terms induced by R-parity breaking are negligible and the baryogenesis constraint, eq. (2.16), is fulfilled for all values of $\tan \beta$.

4. Cosmology and collider physics

4.1 Gravitino decay

Since R-parity is broken, the gravitino is no more stable in our setting, but it still has a lifetime much longer than the age of the universe, since it is suppressed both by the Planck mass and the small R-parity breaking parameters.

The two-body decay is determined by the mixing of the neutralinos with the neutrinos. Neglecting the small neutrino masses, one has [6]

$$\Gamma(\psi_{3/2} \rightarrow \gamma\nu) = \frac{1}{32\pi} |U_{\tilde{\gamma}\nu}|^2 \frac{m_{3/2}^3}{M_{\text{P}}^2}. \quad (4.1)$$

The photino-neutrino mixing can be approximated by (cf. eq. (3.4))

$$|U_{\tilde{\gamma}\nu}| \simeq g_z \left| \sum_{\alpha=1}^4 c_{\tilde{\gamma}\alpha} c_{z\alpha}^* \frac{v_\nu}{m_{\chi_\alpha^0}} \right| \sim 10^{-8} \left(\frac{\epsilon_3}{10^{-7}} \right) \left(\frac{\tilde{m}}{200 \text{ GeV}} \right)^{-1}, \quad (4.2)$$

for $\epsilon_1, \epsilon_2 \leq \epsilon_3$, where we made the rough estimate $0.1/\tilde{m}$ for the weighted sum of neutralino masses and the coupling, taking into account that not all mixings can be maximal. Using $M_{\text{P}} = 2.4 \times 10^{18} \text{ GeV}$, one obtains for the gravitino lifetime [6]

$$\tau_{3/2}^{2\text{-body}} \simeq 4 \times 10^{27} \text{ s} \left(\frac{\epsilon_3}{10^{-7}} \right)^{-2} \left(\frac{\tilde{m}}{200 \text{ GeV}} \right)^2 \left(\frac{m_{3/2}}{10 \text{ GeV}} \right)^{-3}. \quad (4.3)$$

The three-body decay is usually subdominant due to the phase-space and intermediate heavy particle suppression. For the decay with intermediate heavy $\tilde{\tau}_R$, neglecting all external masses in the phase space factor, we find

$$\Gamma(\psi_{3/2} \rightarrow \tau_R l_i l_j) = \frac{|\lambda_{ij3}|^2}{3(32)^2 \pi^3} \frac{m_{3/2}^3}{M_{\text{P}}^2} F \left(\frac{m_{\tilde{\tau}_R}}{m_{3/2}} \right), \quad (4.4)$$

where

$$F(\alpha) = \int_0^1 dx \frac{x^3(1-x)^2}{(1-x-\alpha^2)^2} \simeq \frac{1}{60\alpha^4}. \quad (4.5)$$

The full expression has been obtained in [25].

In the case where only bilinear R-parity breaking is present, the λ_{ijk} couplings are generated from the Yukawa couplings as

$$\lambda_{ijk} = \epsilon_i h_{jk}^{(e)}. \quad (4.6)$$

Then the inverse partial width for the three-body decay,

$$\Gamma(\psi_{3/2} \rightarrow \tau_R l_i l_j)^{-1} \simeq 4 \times 10^{37} s \left(\frac{\epsilon_3}{10^{-7}} \right)^{-2} \left(\frac{\tan \beta}{10} \right)^{-2} \left(\frac{\tilde{m}}{200 \text{ GeV}} \right)^4 \left(\frac{m_{3/2}}{10 \text{ GeV}} \right)^{-7}, \quad (4.7)$$

is much larger than the lifetime determined from the two-body decay, eq. (4.3), as long as the mixing between photino and neutrino is not unnaturally suppressed.

4.2 Extragalactic diffuse gamma-ray emission

A stringent astrophysical constraint for decaying gravitino dark matter is the measured gamma-ray flux [6]. Assuming that the gravitino constitutes the dominant component of dark matter, its decay into neutrino and photon gives rise to an extragalactic diffuse gamma-ray flux with a characteristic energy spectrum, corresponding to a red shifted monochromatic line. A photon with measured energy $E = m_{3/2}/(2(1+z))$ has been emitted at the comoving distance $\chi(z)$, with $d\chi/dz = (1+z)^{-3/2}/(a_0 H_0 \sqrt{\Omega_M(1+\kappa(1+z)^{-3})})$. Here a_0 and H_0 are the present scale factor and Hubble parameter, respectively, and $\kappa = \Omega_\Lambda/\Omega_M \simeq 3$, with $\Omega_\Lambda + \Omega_M = 1$, assuming a flat universe. For the photon flux one obtains, for $\tau_{3/2} \gg H_0^{-1}$,

$$E^2 \frac{dJ_{eg}}{dE} = C_\gamma \left(1 + \kappa \left(\frac{2E}{m_{3/2}} \right)^3 \right)^{-1/2} \left(\frac{2E}{m_{3/2}} \right)^{5/2} \theta \left(1 - \frac{2E}{m_{3/2}} \right), \quad (4.8)$$

with

$$C_\gamma = \frac{\Omega_{3/2} \rho_c}{8\pi \tau_{3/2} H_0 \Omega_M^{1/2}} = 10^{-6} (\text{cm}^2 \text{str s})^{-1} \text{GeV} \left(\frac{\tau_{3/2}}{10^{28} \text{s}} \right)^{-1}; \quad (4.9)$$

here $\tau_{3/2}$ is given by eq. (4.3), and we have taken the gravitino density equal to the Cold Dark Matter density as $\Omega_{3/2} h^2 = 0.1$, $\rho_c = 1.05 h^2 \times 10^{-5} \text{GeV cm}^{-3}$, $\Omega_M = 0.25$ and $H_0 = h 100 \text{ km s}^{-1} \text{Mpc}^{-1}$ with $h = 0.73$ [26].

In addition to the extragalactic signal one also expects a sharp line from the halo of our galaxy with an intensity comparable to the extragalactic signal and strong anisotropy [27]. We have in fact from the decay of halo gravitinos

$$E^2 \frac{dJ_{halo}}{dE} = D_\gamma \delta \left(1 - \frac{2E}{m_{3/2}} \right), \quad (4.10)$$

where

$$D_\gamma = C_\gamma \frac{H_0 \Omega_M^{1/2}}{\Omega_{3/2} \rho_c} \int_{l.o.s.} \rho_{halo}(\vec{l}) d\vec{l}. \quad (4.11)$$

The ratio D_γ/C_γ is given only by cosmological constants and the halo dark matter density integrated along the line of sight, so the intensity and angular distribution of the halo signal is very sensitive to the distribution of the dark matter in the Milky Way. It is surprising that for typical halo models, such number is of order unity [27] and shows moderate angular dependence if one excludes the galactic centre region. The anisotropic part of the halo signal may be partially hidden in the diffuse galactic γ -ray emission due to conventional

astrophysical processes. We expect therefore the isotropic signal in the extragalactic γ -ray flux to be a combination of both the continuum spectrum in eq. (4.8) and part of the halo line in eq. (4.10).

Assuming that one understands the diffuse galactic γ -ray flux, one can extract from the EGRET data the extragalactic diffuse component. The first analysis of Sreekumar *et al.* [28] gave an extragalactic flux described by the power law

$$E^2 \frac{dJ}{dE} = 1.37 \times 10^{-6} \left(\frac{E}{1 \text{ GeV}} \right)^{-0.1} (\text{cm}^2 \text{str s})^{-1} \text{GeV} \quad (4.12)$$

in the energy range 50 MeV–10 GeV. A non-observation of a γ -ray line can then be used to constrain the allowed gravitino mass and lifetime [6]. Assuming the gravitinos to make up all the Cold Dark Matter density, and taking a 3σ upper bound on the flux above 100 MeV corresponding to $2.23 \times 10^{-6} (\text{cm}^2 \text{str s})^{-1} \text{GeV}$ [28], we can have directly a rough bound on the gravitino lifetime from C_γ as

$$\tau_{3/2} \gtrsim 4 \times 10^{27} \text{s} . \quad (4.13)$$

The more recent analysis of the EGRET data [29] shows in the 50 MeV – 2 GeV range a power law behaviour, but a clear excess between 2 GeV and 10 GeV. The maximal flux allowed by the data taking into account the model dependence and systematic errors is not very far from the one obtained in the old analysis, in fact the integrated flux between 0.1-10 GeV is given as $(11.1 \pm 0.1) \times 10^{-6} \text{cm}^{-2} \text{str}^{-1} \text{s}^{-1}$ compared to $(14.5 \pm 0.5) \times 10^{-6} \text{cm}^{-2} \text{str}^{-1} \text{s}^{-1}$ [29].

This is precisely the energy range where, based on our lower bound on the gravitino mass of 5 GeV, one may expect a gravitino signal. It is very remarkable that also the measured flux corresponds to the expectation of the model for R-parity and B-L breaking discussed in section 2 as can be seen from the bound eq. (4.13). On the other hand we would expect also an anisotropic flux from the halo component that EGRET does not resolve probably due to the galactic background, which is an order of magnitude larger than the extracted extragalactic signal.

The excess in the extragalactic γ -ray flux above 2 GeV from the EGRET data [29] has also been related to the annihilation of heavy neutralinos in the galactic halo [30]. Due to the current limitations in the determination of the diffuse galactic γ -ray emission [31] theoretical interpretations of the EGRET excess remain uncertain at present. Clarification can be expected from the Gamma Ray Large Area Space Telescope (GLAST) [32], to be launched this fall, that aims to improve by a factor 30–50 the sensitivity of the EGRET satellite to the diffuse gamma ray flux in the range 20 MeV–10 GeV.

Another constraint comes from the neutrino flux. In the energy range of interest, from about 1 GeV to 1 TeV, the extraterrestrial neutrino flux is constrained by the flux of upward-going muons measured by the IMB experiment, that does not show any discrepancy with respect to the expected neutrino flux from cosmic ray interactions in the atmosphere. The requirement that the neutrino flux from gravitino decay does not exceed the observed flux, translates into a lower bound on the gravitino lifetime, which is roughly $\tau_{3/2} \gtrsim 6 \times 10^{24} \text{s}$

for $m_{3/2} = 1$ TeV, and becomes weaker for smaller gravitino masses [33]. This bound is clearly consistent with a signal in the EGRET data, as discussed above.

4.3 Collider signatures

The collider signatures depend on the nature of the NLSP. Here we consider the cases that the NLSP is the lightest stau or the lightest neutralino.

The lightest stau, that we assume mainly right-handed, decays through $\tilde{\tau}_R \rightarrow \tau\nu_\mu, \mu\nu_\tau$. On the other hand, the small left-handed component of the stau mass eigenstate can trigger a decay into two jets through $\tilde{\tau}_L \rightarrow b^c t$, provided the process is kinematically open. The hadronic decays are enhanced compared to the leptonic decays by the larger bottom Yukawa coupling and by the colour factor, but are usually suppressed by the small left-right mixing.

If the leptonic decay channel is the dominant mode, the decay length can be approximated by

$$c\tau_{\tilde{\tau}}^{lep} \sim 30 \text{ cm} \left(\frac{m_{\tilde{\tau}}}{200\text{GeV}} \right)^{-1} \left(\frac{\epsilon_2}{10^{-7}} \right)^{-2} \left(\frac{\tan \beta}{10} \right)^{-2}. \quad (4.14)$$

It is intriguing that the sufficient condition to avoid the erasure of the baryon asymmetry, eq. (2.16), implies the observation of a displaced stau vertex at future colliders, more than 3mm away from the beam axis for $\epsilon_2 < 10^{-6}$. In the particular case of the flavour model (II) discussed in section 2.2, $\epsilon_2 \sim 6 \times 10^{-8}$, one has a spectacular signal consisting on a heavily ionising charged track of length ~ 0.8 m, followed by a muon track or a jet and missing energy, corresponding to $\tilde{\tau} \rightarrow \mu\nu_\tau$ or $\tilde{\tau} \rightarrow \tau\nu_\mu$, respectively.

If the hadronic channel $\tilde{\tau}_L \rightarrow b^c t$ is the dominant mode, the decay length is given by

$$c\tau_{\tilde{\tau}}^{had} \sim 1.4 \text{ m} \left(\frac{m_{\tilde{\tau}}}{200\text{GeV}} \right)^{-1} \left(\frac{\epsilon_3}{10^{-7}} \right)^{-2} \left(\frac{\tan \beta}{10} \right)^{-2} \left(\frac{\cos \theta_\tau}{0.1} \right)^{-2}, \quad (4.15)$$

where θ_τ denotes the mixing angle of the staus. This channel also yields a very unique signature at colliders, consisting of a heavily ionising charged track followed by two jets.

These characteristic signatures would allow to distinguish at colliders our scenario from the case with conserved R-parity where the decay $\tilde{\tau} \rightarrow \psi_{3/2}\tau$ leads to (cf. [34])

$$c\tau_{\tilde{\tau}}^{3/2} \sim 40 \text{ cm} \left(\frac{m_{3/2}}{1 \text{ keV}} \right)^2 \left(\frac{m_{\tilde{\tau}}}{200 \text{ GeV}} \right)^{-5}. \quad (4.16)$$

Hence, for a gravitino mass $m_{3/2} \lesssim \mathcal{O}(10 \text{ keV})$, the decay length of the lightest stau is shorter than $\mathcal{O}(10 \text{ m})$, and would therefore decay inside the detector into tau and gravitino. The experimental signature for this process would be identical to the decay $\tilde{\tau} \rightarrow \tau\nu_\mu$. However, the scenario with R-parity violation also predicts the decay $\tilde{\tau} \rightarrow \mu\nu_\tau$, with very similar branching ratio due to $SU(2)$ invariance. Although this signature could be mimicked by a scenario with conserved R-parity if lepton flavour is violated, through the decay $\tilde{\tau} \rightarrow \mu\psi_{3/2}$, large branching ratios are precluded from present bounds on flavour violation [35]. In consequence, the observation of a comparable number of tau and muon events in stau decays would constitute a signature for the scenario with R-parity violation. Also, the observation of a stau decaying into two jets would undoubtedly point to the scenario with R-parity violation.

On the other hand, if the lightest neutralino is the NLSP, it decays through $\chi_1^0 \rightarrow \tau^\pm W^\mp$ [36], or through $\chi_1^0 \rightarrow b b^c \nu$ [37] if the former decay channel is kinematically closed. The corresponding decay lengths can be approximated by

$$c\tau_{\chi_1^0}^{2\text{-body}} \sim 20 \text{ cm} \left(\frac{m_{\chi_1^0}}{200 \text{ GeV}} \right)^{-3} \left(\frac{\epsilon_3}{10^{-7}} \right)^{-2} \left(\frac{\tan \beta}{10} \right)^2, \quad (4.17)$$

$$c\tau_{\chi_1^0}^{3\text{-body}} \sim 600 \text{ m} \left(\frac{m_{\tilde{\nu}_L}}{300 \text{ GeV}} \right)^4 \left(\frac{m_{\chi_1^0}}{200 \text{ GeV}} \right)^{-5} \left(\frac{\epsilon_3}{10^{-7}} \right)^{-2} \left(\frac{\tan \beta}{10} \right)^{-2}. \quad (4.18)$$

Again, this scenario can be easily discriminated at future colliders from the scenario with conserved R-parity. In this case, the neutralino decays into gravitino and photon [38] with decay length

$$c\tau_{\chi_1^0}^{3/2} \sim 80 \text{ cm} \left(\frac{m_{3/2}}{1 \text{ keV}} \right)^2 \left(\frac{m_{\chi_1^0}}{200 \text{ GeV}} \right)^{-5}. \quad (4.19)$$

For a gravitino mass $m_{3/2} \lesssim \mathcal{O}(10 \text{ keV})$ the neutralino would decay inside the detector producing an energetic photon and missing energy, which is clearly distinguishable from the signals in the R-parity violating scenario that in general involve jets.

4.4 Microscopic determination of the Planck mass

Recently, a method has been proposed for the microscopic determination of the Planck mass in collider experiments [34], providing a direct test of supergravity. The method requires a very long lived stau NLSP which decays mostly into tau and gravitino, which is difficult to reconcile with recent constraints from BBN [7, 9], unless there is a late-time entropy production [10]. In the picture proposed in this letter, where primordial nucleosynthesis, thermal leptogenesis and dark matter are naturally consistent, this method cannot be pursued, as the stau decays predominantly in the R-parity violating channel into charged lepton and neutrino.

Nevertheless, from a gravitino signal in the diffuse γ -ray flux and the width for the stau decay into two jets, one can still obtain a microscopic estimate of the Planck mass. The gravitino mass is given by the maximal energy of the photon, $m_{3/2} = 2E_\gamma$, and the gravitino lifetime can be determined from the photon flux, eqs. (4.8), (4.10). Then, using the expression for the gravitino decay rate, eq. (4.1), one can rewrite the Planck mass in terms of the gravitino mass, lifetime and photino-neutrino mixing as

$$\begin{aligned} M_{\text{P}} &= \left(\frac{m_{3/2}^3 \tau_{3/2}}{32\pi} \right)^{1/2} |U_{\tilde{\gamma}\nu}| \\ &= 2.5 \times 10^{18} \text{ GeV} \left(\frac{m_{3/2}}{10 \text{ GeV}} \right)^{3/2} \left(\frac{\tau_{3/2}}{4 \times 10^{27} \text{ s}} \right)^{1/2} \left(\frac{|U_{\tilde{\gamma}\nu}|}{10^{-8}} \right) \end{aligned} \quad (4.20)$$

where $|U_{\tilde{\gamma}\nu}|$ is related to the decay rate of the stau into two jets⁶. We can cast the dependence on the decay rate as a dependence on the decay length of the stau in this

⁶Note that the decay rate of the stau into leptons depends on ϵ_2 whereas $|U_{\tilde{\gamma}\nu}|$ depends on the sneutrino VEV and therefore mainly on ϵ_3 for the hierarchical case.

channel, yielding

$$|U_{\tilde{\gamma}\nu}|^2 \simeq 10^{-16} \left(\frac{c\tau_{\tilde{\tau}}^{had}}{1.4\text{m}} \right)^{-1} \left(\frac{\tilde{m}}{200\text{GeV}} \right)^{-3} \left(\frac{\tan\beta}{10} \right)^2 \left(\frac{\cos\theta_{\tau}}{0.1} \right)^2. \quad (4.21)$$

The measurement of the decay length of the stau in the hadronic channel, complemented with additional information about supersymmetry breaking parameters, can provide a determination of $|U_{\tilde{\gamma}\nu}|$. The measurement of the photon energy and the photon flux in the diffuse γ -ray background then gives the gravitino mass and lifetime and, using eq. (4.20), an estimate of the Planck mass.

5. Conclusions

On theoretical grounds, theories with and without R-parity are on equal footing. In this paper we have presented a simple model where R-parity is not conserved and its violation is connected to the scale of B-L breaking. One can then have R-parity violating couplings that are small enough to be consistent with baryogenesis and gravitino dark matter, yet large enough to allow for the NLSP decay before nucleosynthesis. For gravitino masses above 5 GeV one obtains a cosmological history consistent with thermal leptogenesis, thermally produced gravitino dark matter and primordial nucleosynthesis.

Relic gravitino decays into neutrino and photon yield a diffuse halo and extragalactic γ -ray flux which depends on the R-parity violating Yukawa couplings. It is remarkable that for a gravitino mass $m_{3/2} = \mathcal{O}(10)$ GeV, the predicted photon flux could be part of the apparent excess in the extragalactic diffuse γ -ray flux obtained from the EGRET data. However, given the current uncertainties in the determination of the diffuse galactic γ -ray emission, this consistency may be accidental. Unequivocal evidence for decaying gravitino dark matter could come from the results of GLAST.

The flavour dependent pattern of R-parity breaking can give striking signatures at the LHC, in particular a vertex of the NLSP, that is significantly displaced from the beam axis. Together with the measurement of supersymmetry breaking parameters at the LHC, the observation of a redshifted photon spectral line from gravitino decay by GLAST can allow a microscopic determination of the Planck mass. In the less optimistic case where the R-parity breaking Yukawa couplings are near to their lower bound, astrophysical detection will be very challenging whereas signals hinting at R-parity breaking and gravitino dark matter could still come from stau decays, as in the case of R-parity conservation.

Acknowledgments

We are grateful to G. Heinzlmann and J. Ripken for valuable discussions on the diffuse γ -ray background. L.C. acknowledges the support of the ‘‘Impuls- und Vernetzungsfond’’ of the Helmholtz Association, contract number VH-NG-006.

References

- [1] P. Fayet, *Supersymmetric theories of particles*, *Orbis Scientiae* **0413** (1978) QCD161:C6:1978.

- [2] N. Sakai and T. Yanagida, *Proton decay in a class of supersymmetric grand unified models*, *Nucl. Phys. B* **197** (1982) 533;
S. Weinberg, *Supersymmetry at ordinary energies, 1. Masses and conservation laws*, *Phys. Rev. D* **26** (1982) 287.
- [3] L.J. Hall and M. Suzuki, *Explicit R-parity breaking in supersymmetric models*, *Nucl. Phys. B* **231** (1984) 419.
- [4] For recent discussions and references, see B.C. Allanach, A. Dedes and H.K. Dreiner, *The R parity violating minimal supergravity model*, *Phys. Rev. D* **69** (2004) 115002 [[hep-ph/0309196](#)];
R. Barbier et al., *R-parity violating supersymmetry*, *Phys. Rept.* **420** (2005) 1 [[hep-ph/0406039](#)].
- [5] B.A. Campbell, S. Davidson, J.R. Ellis and K.A. Olive, *Cosmological baryon asymmetry constraints on extensions of the standard model*, *Phys. Lett. B* **256** (1991) 457;
W. Fischler, G.F. Giudice, R.G. Leigh and S. Paban, *Constraints on the baryogenesis scale from neutrino masses*, *Phys. Lett. B* **258** (1991) 45;
H.K. Dreiner and G.G. Ross, *Sphaleron erasure of primordial baryogenesis*, *Nucl. Phys. B* **410** (1993) 188 [[hep-ph/9207221](#)].
- [6] F. Takayama and M. Yamaguchi, *Gravitino dark matter without R-parity*, *Phys. Lett. B* **485** (2000) 388 [[hep-ph/0005214](#)].
- [7] M. Pospelov, *Particle physics catalysis of thermal big bang nucleosynthesis*, [hep-ph/0605215](#);
R.H. Cyburt, J. Ellis, B.D. Fields, K.A. Olive and V.C. Spanos, *Bound-state effects on light-element abundances in gravitino dark matter scenarios*, *JCAP* **11** (2006) 014 [[astro-ph/0608562](#)].
- [8] K. Kohri and F. Takayama, *Big bang nucleosynthesis with long lived charged massive particles*, [hep-ph/0605243](#);
M. Kaplinghat and A. Rajaraman, *Big bang nucleosynthesis with bound states of long-lived charged particles*, *Phys. Rev. D* **74** (2006) 103004 [[astro-ph/0606209](#)].
- [9] M. Kawasaki, K. Kohri and T. Moroi, *Big-bang nucleosynthesis and hadronic decay of long-lived massive particles*, *Phys. Rev. D* **71** (2005) 083502 [[astro-ph/0408426](#)];
J.L. Feng, S. Su and F. Takayama, *Supergravity with a gravitino LSP*, *Phys. Rev. D* **70** (2004) 075019 [[hep-ph/0404231](#)];
F.D. Steffen, *Gravitino dark matter and cosmological constraints*, *JCAP* **09** (2006) 001 [[hep-ph/0605306](#)].
- [10] W. Buchmüller, K. Hamaguchi, M. Ibe and T.T. Yanagida, *Eluding the BBN constraints on the stable gravitino*, [hep-ph/0605164](#).
- [11] T. Kanzaki, M. Kawasaki, K. Kohri and T. Moroi, *Cosmological constraints on gravitino lsp scenario with sneutrino NLSP*, *Phys. Rev. D* **75** (2007) 025011 [[hep-ph/0609246](#)].
- [12] M. Fukugita and T. Yanagida, *Phys. Lett. B* **174** (1986) 45.
- [13] S. Davidson and A. Ibarra, *A lower bound on the right-handed neutrino mass from leptogenesis*, *Phys. Lett. B* **535** (2002) 25 [[hep-ph/0202239](#)].
- [14] W. Buchmüller, P. Di Bari and M. Plümacher, *Leptogenesis for pedestrians*, *Ann. Phys. (NY)* **315** (2005) 305 [[hep-ph/0401240](#)].

- [15] M. Bolz, A. Brandenburg and W. Buchmüller, *Thermal production of gravitinos*, *Nucl. Phys.* **B 606** (2001) 518 [[hep-ph/0012052](#)].
- [16] J. Pradler and F.D. Steffen, *Thermal gravitino production and collider tests of leptogenesis*, *Phys. Rev.* **D 75** (2007) 023509 [[hep-ph/0608344](#)].
- [17] V.S. Rychkov and A. Strumia, *Thermal production of gravitinos*, [hep-ph/0701104](#).
- [18] T. Asaka, W. Buchmüller and L. Covi, *Quarks and leptons between branes and bulk*, *Phys. Lett.* **B 563** (2003) 209 [[hep-ph/0304142](#)].
- [19] R. Tatar and T. Watari, *Proton decay, Yukawa couplings and underlying gauge symmetry in string theory*, *Nucl. Phys.* **B 747** (2006) 212 [[hep-th/0602238](#)]; *A stable proton without R-parity: implications for the lsp* , [hep-ph/0605315](#).
- [20] W. Buchmüller, K. Hamaguchi, O. Lebedev and M. Ratz, *Supersymmetric standard model from the heterotic string. ii*, [hep-th/0606187](#).
- [21] G.F. Giudice and A. Masiero, *A natural solution to the μ problem in supergravity theories*, *Phys. Lett.* **B 206** (1988) 480.
- [22] W. Buchmüller and T. Yanagida, *Quark lepton mass hierarchies and the baryon asymmetry*, *Phys. Lett.* **B 445** (1999) 399 [[hep-ph/9810308](#)].
- [23] S. Roy and B. Mukhopadhyaya, *Some implications of a supersymmetric model with r-parity breaking bilinear interactions*, *Phys. Rev.* **D 55** (1997) 7020 [[hep-ph/9612447](#)];
M. Hirsch, M.A. Diaz, W. Porod, J.C. Romao and J.W.F. Valle, *Neutrino masses and mixings from supersymmetry with bilinear R-parity violation: a theory for solar and atmospheric neutrino oscillations*, *Phys. Rev.* **D 62** (2000) 113008 [[hep-ph/0004115](#)];
A. Abada, S. Davidson and M. Losada, *Neutrino masses and mixings in the MSSM with soft bilinear R_p violation*, *Phys. Rev.* **D 65** (2002) 075010 [[hep-ph/0111332](#)];
E.J. Chun, D.-W. Jung and J.D. Park, *Bi-large neutrino mixing from bilinear r-parity violation with non-universality*, *Phys. Lett.* **B 557** (2003) 233 [[hep-ph/0211310](#)].
- [24] F. Takayama and M. Yamaguchi, *Pattern of neutrino oscillations in supersymmetry with bilinear R-parity violation*, *Phys. Lett.* **B 476** (2000) 116 [[hep-ph/9910320](#)].
- [25] G. Moreau and M. Chemtob, *R-parity violation and the cosmological gravitino problem*, *Phys. Rev.* **D 65** (2002) 024033 [[hep-ph/0107286](#)].
- [26] W.-M. Yao et al., *Journal of Physics G* 33 (2006) 1.
- [27] T. Asaka, J. Hashiba, M. Kawasaki and T. Yanagida, *Spectrum of background x-rays from moduli dark matter*, *Phys. Rev.* **D 58** (1998) 023507 [[hep-ph/9802271](#)].
- [28] EGRET collaboration, P. Sreekumar et al., *EGRET observations of the extragalactic gamma ray emission*, *Astrophys. J.* **494** (1998) 523 [[astro-ph/9709257](#)].
- [29] A.W. Strong, I.V. Moskalenko and O. Reimer, *A new determination of the diffuse galactic and extragalactic gamma-ray emission*, [astro-ph/0506359](#); *Diffuse galactic continuum gamma rays. A model compatible with egret data and cosmic-ray measurements*, *Astrophys. J.* **613** (2004) 962 [[astro-ph/0406254](#)]; *A new determination of the extragalactic diffuse gamma-ray background from EGRET data*, *Astrophys. J.* **613** (2004) 956 [[astro-ph/0405441](#)].
- [30] D. Elsässer and K. Mannheim, *Supersymmetric dark matter and the extragalactic gamma ray background*, *Phys. Rev. Lett.* **94** (2005) 171302 [[astro-ph/0405235](#)].

- [31] I.V. Moskalenko, S.W. Digel, T.A. Porter, O. Reimer and A.W. Strong, *Understanding limitations in the determination of the diffuse galactic gamma-ray emission*, astro-ph/0609768.
- [32] <http://glast.gsfc.nasa.gov>.
- [33] P. Gondolo, G. Gelmini and S. Sarkar, *Cosmic neutrinos from unstable relic particles*, *Nucl. Phys. B* **392** (1993) 111 [hep-ph/9209236].
- [34] W. Buchmüller, K. Hamaguchi, M. Ratz and T. Yanagida, *Supergravity at colliders*, *Phys. Lett. B* **588** (2004) 90 [hep-ph/0402179].
- [35] K. Hamaguchi and A. Ibarra, *Probing lepton flavour violation in slepton NLSP scenarios*, *JHEP* **02** (2005) 028 [hep-ph/0412229].
- [36] B. Mukhopadhyaya, S. Roy and F. Vissani, *Correlation between neutrino oscillations and collider signals of supersymmetry in an R-parity violating model*, *Phys. Lett. B* **443** (1998) 191 [hep-ph/9808265];
E.J. Chun and J.S. Lee, *Implication of super-KamioKande data on R-parity violation*, *Phys. Rev. D* **60** (1999) 075006 [hep-ph/9811201].
- [37] H.K. Dreiner and G.G. Ross, *R-parity violation at hadron colliders*, *Nucl. Phys. B* **365** (1991) 597.
- [38] S. Dimopoulos, M. Dine, S. Raby and S.D. Thomas, *Experimental signatures of low energy gauge mediated supersymmetry breaking*, *Phys. Rev. Lett.* **76** (1996) 3494 [hep-ph/9601367].