

Precision study of the $SU(3)$ topological susceptibility in the continuum

Stephan Dürr

*Universität Bern, Institut für theoretische Physik
Sidlerstr. 5, CH-3012 Bern, Switzerland*

Zoltan Fodor,^{abc} Christian Hoelbling^a and Thorsten Kurth^a

^a*Bergische Universität Wuppertal
Gaussstr. 20, D-42119 Wuppertal, Germany*

^b*Eötvös University, Physics Department
Pázmány 1, H-1117 Budapest, Hungary*

^c*University of California at San Diego
9500 Gilman Drive, La Jolla, CA 92093-0319, U.S.A.*

*E-mail: fodor@bodri.elte.hu, christian.hoelbling@cern.ch,
thorsten.kurth@uni-wuppertal.de*

ABSTRACT: We determine the topological susceptibility in the $SU(3)$ pure gauge theory. We perform a series of high-statistics lattice studies and take the combined continuum and infinite volume limit. We find $\chi_{\text{top}} r_0^4 = 0.0524(7)(6)$, which translates into $\chi_{\text{top}}^{1/4} = 193(1)(8)$ MeV with the second error exclusively due to the intrinsic scale ambiguity.

KEYWORDS: Lattice Gauge Field Theories, 1/N Expansion.

Contents

1.	Introduction	1
2.	Topological charge definition	2
3.	Lattice simulations	4
4.	Continuum and infinite volume extrapolations	7
5.	Conversion to physical units	10
6.	Summary	10
A.	Parametrization of r_0/a	11

1. Introduction

In QCD with N_c colors and N_f light dynamical quarks there are two notions of the topological susceptibility, defined as the second moment of the global topological charge distribution.

On the one hand, the *actual* topological susceptibility $\chi_{\text{top}}^{\text{QCD}}$ shows a clear sensitivity on the dynamical (sea) quark masses [1]. This property renders it an ideal vacuum diagnostics tool, as emphasized in [2] and exploited in a number of recent studies [3–10].

On the other hand, the *quenched* topological susceptibility $\chi_{\text{top}}^{\text{YM}}$ is the quantity which is linked via the famous Witten-Veneziano formula [11, 12]

$$\chi_{\text{top}}^{\text{YM}} \doteq \frac{F^2}{2N_f} (M_{\eta'}^2 + M_{\eta}^2 - 2M_K^2) \tag{1.1}$$

to the excess of the η' mass over the pseudoscalar octet masses, with a proportionality factor which contains the pseudoscalar decay constant¹ in the chiral limit. This relation is supposed to hold at the leading order in an expansion in $1/N_c$ (for a different viewpoint see [13–15]). To be precise, this quenched susceptibility is the susceptibility of the underlying $SU(N_c)$ Yang-Mills (YM) theory, and this is what makes (1.1) appealing from a theorist’s viewpoint — it relates two different theories. Specifically, on the lattice one can measure the l.h.s. of (1.1) for several N_c and evaluate the r.h.s. for various (N_c, N_f) combinations, and finally check whether the agreement is parametrically controlled by $1/N_c$ or N_f/N_c .

¹We use the Bern normalization where $F_{\pi}^{\text{phys}} = 92.4(3)$ MeV and $F = 86.2(5)$ MeV in the chiral limit.

In this paper we elaborate on the first step in this program — we determine with unprecedented precision the quenched topological susceptibility for the case $N_c = 3$. We begin with an exposition of the main avenues towards defining a topological charge on the lattice and how one extracts the topological susceptibility. The next two sections contain details of our lattice simulations and of our combined continuum and infinite volume extrapolation. Having a result in terms of the Sommer radius r_0 [16], the latter needs to be identified with a length-scale in fm, and we discuss both the input that goes into such an identification and the remaining ambiguity. In the concluding section we compare our result to other recent determinations of $\chi_{\text{top}} \equiv \chi_{\text{top}}^{\text{YM}}$. Details of a new parameterization of $r_0(\beta)$ have been arranged in an appendix.

2. Topological charge definition

In the continuum the topological charge of a given gauge background is defined as

$$q = \frac{1}{16\pi^2} \int dx \operatorname{tr}(F_{\mu\nu}(x)\tilde{F}_{\mu\nu}(x)) = \frac{1}{32\pi^2} \int dx \operatorname{tr}(\epsilon_{\mu\nu\sigma\rho}F_{\mu\nu}(x)F_{\sigma\rho}(x)) \quad (2.1)$$

where $F_{\mu\nu} = F_{\mu\nu}^a \lambda^a / 2$ is the field strength tensor. For toroidal space-time geometry (2.1) is integer and linked to the index of the Dirac operator D via the Atiyah-Singer theorem [17]

$$q = n_+ - n_- \quad (2.2)$$

where n_{\pm} denotes the number of zero modes of D with positive or negative chirality.

On the lattice the definition of the topological charge is not unique. Aiming for the gluonic side of (2.2), one may choose any discretization of $F_{\mu\nu}$ which has the correct perturbative continuum limit, and form the so-called “naive” (or unrenormalized field-theoretic) charge

$$\begin{aligned} q_{\text{nai}}[U] &= \frac{1}{32\pi^2} \sum_x \operatorname{tr}(\epsilon_{\mu\nu\sigma\rho}F_{\mu\nu}(x)F_{\sigma\rho}(x)) \\ &= \frac{1}{4\pi^2} \sum_x \operatorname{tr}(F_{12}F_{34} + F_{13}F_{42} + F_{14}F_{23}) . \end{aligned} \quad (2.3)$$

In general this definition does not lead to an integer² charge.

On the other hand, the fermionic side of (2.2) always yields an integer answer. This is most straightforward with the massless overlap³ operator $D = \rho[1 + D_{\text{W},-\rho}(D_{\text{W},-\rho}^\dagger D_{\text{W},-\rho})^{-1/2}]$ [22] or with any other Dirac operator D which satisfies the Ginsparg-Wilson relation [23]. In this case the index can be written in the closed form [24, 25]

$$q_{\text{fer}}[U] = -\frac{1}{2\rho} \operatorname{tr}(\gamma_5 D) . \quad (2.4)$$

With a non-chiral Dirac operator (e.g. D_{W}) explicit mode counting prescriptions may be set up. Below, the only point which matters is that $q_{\text{fer}}[U]$ is necessarily an integer.

²There are two known exceptions, the Lüscher [18] and the Phillips-Stone [19] definitions of $q_{\text{th}}[U]$.

³It does not matter whether one uses the Wilson operator D_{W} or another doubler-free kernel [20, 21].




Figure 1: Histogram of the topological charge q_{nai} with 3 HYP steps for the 12^4 lattices at $\beta = 6.0$, before and after rescaling with the renormalization factors defined in (2.6).

The continuum topological susceptibility at zero virtuality is defined as

$$\chi_{\text{top}} = \lim_{V \rightarrow \infty} \frac{\langle q^2 \rangle}{V} \tag{2.5}$$

and this shows that a finite volume is mandatory for the definition. Still, contributions to $\langle q^2 \rangle$ which grow less than linearly in the 4D volume V create finite volume effects in χ_{top} .

On the lattice the details of the topological charge definition reflect themselves in the precise form of the latticized version of (2.5) [26]. In case one starts with the gluonic definition (2.3), the traditional approach has been to form the bare susceptibility $\langle q_{\text{nai}}^2 \rangle / V$ which is then subject to both additive and multiplicative renormalization [27–29]. On the other hand, starting from the fermionic definition (2.4), one just forms $\langle q_{\text{fer}}^2 \rangle / V$, since q_{fer} is already a renormalized charge. Likewise, if we first compute a renormalized (integer) field-theoretic charge, the susceptibility based on it will not require any further renormalization⁴.

In this paper we investigate the utility of such a renormalized gluonic charge definition for a precision measurement of the topological susceptibility in the pure SU(3) gauge theory. We start from the standard “clover-leaf” definition of $F_{\mu\nu}(x)$ (it uses the average of the antihermitean part of the plaquette $U_{\mu\nu}$ in $x, x-\hat{\mu}, x-\hat{\nu}, x-\hat{\mu}-\hat{\nu}$) based on HYP smeared [34] gauge links. Plugging this into (2.3) we have the bare charge $q_{\text{nai}}[U]$ which is a real number. One of our q_{nai} distributions ($\beta=6.0, 12^4, 3$ HYP steps) is shown in the left panel of figure 1. Thanks to CP symmetry only a multiplicative renormalization applies, and we opt for a non-perturbatively defined Z -factor. Still, there are various possibilities, and we choose a

⁴Note that the round-to-integer operation (2.7) brings in a global element, and our q_{ren} is *not* given by the integral over a local charge density. Accordingly, there is no conflict with the result by Stamatescu and Seiler that in general $q(x)$ mixes with the identity and the correlator $\int q(x)q(0) dx$ has a contact term [30]. Any overlap based charge definition sticks out in the sense that it is an integral of a local density *and* avoids the mixing with the identity [31–33]. The argument for the absence of additive mass renormalization is then standard — in the zero-charge sector the susceptibility vanishes exactly. Note that it is essential to define the sectors with the same operator that is used in the χ_{top} estimator — if one uses one overlap charge to define the sector and another overlap charge to measure $\langle q^2 \rangle$, then even the overlap based susceptibility has an additive renormalization.

strategy which makes use of the fact that on fine enough lattices the overall distribution of q_{nai} tends to cluster near integer values (cf. figure 1). We find Z as the solution⁵ of

$$\min_{Z>1}(\chi^2) \quad \text{where} \quad \chi^2 = \sum_U \left(Z q_{\text{nai}}[U] - \text{round}(Z q_{\text{nai}}[U]) \right)^2 \quad (2.6)$$

and use it to define, via rounding to the nearest integer, the renormalized field-theoretic charge

$$q_{\text{ren}}[U] = \text{round}(Z q_{\text{nai}}[U]) \quad (2.7)$$

which, by construction, is an integer. This charge definition has already been used in [35].

The alert reader might be surprised by our frequent use of expressions like “we choose” or “we opt for” in this passage. Indeed, there is a huge amount of freedom in how one attributes an integer charge to a lattice configuration. However, according to the standard scaling hypothesis by Symanzik this ambiguity reflects itself in different $O(a^2)$ cut-off effects of observables built from the topological charge. It has been checked that the “disagreement rate” between any pair of charge definitions quickly vanishes with $\beta \rightarrow \infty$ [37] and that the gluonic or fermionic charge definition yields the same continuum limit for the topological susceptibility [29, 38, 39]. Clearly, this is not a mathematical proof, but it is worth emphasizing that this is exactly the behavior that one expects to see, if one is in the Symanzik scaling regime.

3. Lattice simulations

Our goal is to perform a series of simulations in a fixed physical volume that will allow us to determine the topological charge distribution in the continuum limit (in that volume). This will be complemented by a second series of simulations (at a fixed lattice spacing) to assess possible finite volume effects.

We use the MILC code [40] to produce the SU(3) gauge ensembles. We choose the Wilson gauge action and run parameters as detailed in table 1 (for the scaling series) and table 2 (for the volume extrapolation series). The scale is set via the Sommer radius r_0 [16], based on a new parameterization of data from [41] as specified in the appendix. The physical value is a separate topic that will be discussed along with the presentation of the final result.

We start with a closer look at table 1. Throughout, our runs are designed to yield $\sim 10^5$ measurements. Anticipating that the autocorrelation time of the topological charge rapidly grows with β , we have increased the number of updates, n_{sepa} , between adjacent measurements. Still, it turns out that the measured integrated autocorrelation times are somewhat larger on the finer lattices. The renormalization factor Z for q_{nai} with 3HYP steps, as defined in (2.6), seems to go monotonically towards 1 with increasing β , as expected. Based on it we determine $q = q_{\text{ren}}$ defined in (2.7). As a first check, we measure $|\langle q \rangle|$, and this moment is consistent with zero on all lattices. The observable of interest, $\langle q^2 \rangle$, is measured

⁵The restriction $Z > 1$ is a technical aspect of the minimization procedure to avoid the global minimum $\chi^2 = 0$ at $Z = 0$ [35]. In perturbation theory one finds $Z = 1 + \text{const } g_0^2$ with $\text{const} > 0$ [36].

β	5.8980	6.0000	6.0938	6.1802	6.2602	6.3344
L/a	10	12	14	16	18	20
n_{sepa}	10	10	10	50	50	100
n_{conf}	100010	101600	103705	101710	112222	105314
Z	1.2902	1.2353	1.1993	1.1742	1.1570	1.1430
$\tau_{\text{int}}(q)$	0.93	2.18	4.83	2.27	4.65	4.61
$\tau_{\text{int}}(q^2)$	0.61	1.09	2.44	1.19	2.45	2.52
$\tau_{\text{int}}(\text{sign}(q))$	0.89	2.05	4.42	2.10	4.25	4.27
$ \langle q \rangle $	0.004(5)	0.007(8)	0.007(11)	0.007(8)	0.021(10)	0.010(10)
$\langle q^2 \rangle$	1.695(9)	1.592(11)	1.490(15)	1.465(10)	1.427(14)	1.381(14)
$\langle q^4 \rangle / \langle q^2 \rangle^2 - 3$	0.214(21)	0.238(24)	0.244(27)	0.227(21)	0.241(30)	0.204(32)
$\langle q^4 \rangle / \langle q^2 \rangle - 3\langle q^2 \rangle$	0.363(36)	0.378(38)	0.363(40)	0.333(32)	0.344(43)	0.282(44)
$\langle q^4 \rangle - 3\langle q^2 \rangle^2$	0.616(61)	0.602(62)	0.541(60)	0.487(46)	0.490(63)	0.389(61)
$\langle q^2 \rangle_{q \neq 0}$	1.702(12)	1.610(13)	1.526(22)	1.486(13)	1.461(21)	1.400(19)

Table 1: Summary of the scaling series of runs. Unless stated otherwise, q is the renormalized (i.e. integer) gluonic topological charge q_{ren} with 3HYP steps. The subscript “ $q \neq 0$ ” means that the distribution of $|q|$ with $q \neq 0$ has been fitted to a half-Gaussian. Statistical errors have been estimated via a jackknife with the blocklength set to $\text{round}(10\tau_{\text{int}}(q))$.

β	6.0000	6.0000	6.0000	6.0000	6.0000
L/a	10	12	14	16	18
n_{sepa}	10	10	10	10	10
n_{conf}	103600	101600	99000	100000	105000
Z	[1.2441]	1.2353	[1.2333]	[1.2332]	[1.2237]
$\tau_{\text{int}}(q)$	1.80	2.18	2.25	2.09	2.13
$\tau_{\text{int}}(q^2)$	1.34	1.09	1.10	1.11	1.13
$\tau_{\text{int}}(\text{sign}(q))$	1.74	2.05	1.96	1.74	1.71
$ \langle q \rangle $	0.004(4)	0.007(8)	0.003(11)	0.020(14)	0.023(17)
$\langle q^2 \rangle$	0.586(5)	1.592(11)	3.012(20)	5.198(34)	8.233(53)
$\langle q^4 \rangle / \langle q^2 \rangle^2 - 3$	1.277(41)	0.238(24)	0.095(20)	0.065(19)	0.008(16)
$\langle q^4 \rangle / \langle q^2 \rangle - 3\langle q^2 \rangle$	0.748(24)	0.378(38)	0.287(59)	0.338(97)	0.069(139)
$\langle q^4 \rangle - 3\langle q^2 \rangle^2$	0.439(15)	0.602(62)	0.86 (18)	1.76 (50)	0.57 (114)
$\langle q^2 \rangle_{q \neq 0}$	0.780(10)	1.610(13)	3.009(26)	5.202(51)	8.242(54)

Table 2: Summary of the volume extrapolation series of runs. Throughout, the charge renormalization factors of the $(6.0, 12^4)$ lattices have been used. For details see caption of table 1.

with 1% statistical accuracy (or better), throughout. We have also checked that using a 2HYP or 4HYP charge would change the susceptibility by an amount which is an order of magnitude smaller than the statistical error of the $\langle q^2 \rangle$ given. Our data are precise enough to evaluate the fourth moment of the distribution, and the kurtosis $\langle q^4 \rangle / \langle q^2 \rangle^2 - 3$ turns out to deviate from zero, for most lattices in the scaling series, by about 10σ . It seems




Figure 2: Distribution of q_{ren} for the $(6.0, 12^4)$ [left] and $(6.3344, 20^4)$ lattices [right]. The fit to a Gaussian form excludes the $q=0$ sector; the excess gives rise to the kurtosis $\langle q^4 \rangle / \langle q^2 \rangle^2 - 3$ reported in table 1, an effect that goes away in the infinite volume limit, as evident from table 2.

independent of the lattice spacing and it is thus natural to ask whether a non-vanishing $\langle q^4 \rangle / \langle q^2 \rangle^2 - 3$ is a finite volume effect.

This brings us to discuss the second series of runs, as detailed in table 2. Potential finite volume effects are commonly attributed to infrared physics; we thus restrict ourselves to a single coupling, $\beta = 6.0$. Again, we aim for $O(10^5)$ measurements per run. For larger volumes the “crest and valley” structure in figure 1 becomes less pronounced,⁶ and we pursue the analysis with the Z -factor determined on the $(6.0, 12^4)$ lattices. Now one expects the autocorrelation times to be independent of the volume. This happens to be true, except in the $L/a=10$ case, where we find severe finite volume effects in $\langle q^2 \rangle$, too (see below). Again $\langle q \rangle$ is basically consistent with zero. The second moment $\langle q^2 \rangle$ is measured, as before, with 1% statistical accuracy (or better) on all lattices. The main point of this series is clear evidence that the kurtosis $\langle q^4 \rangle / \langle q^2 \rangle^2 - 3$, which quantifies the deviation from a Gaussian distribution, tends to zero with $V \rightarrow \infty$.

This observation lets us plot the charge histograms for two lattices from the scaling series in figure 2. It turns out that the relative weight of the topologically non-trivial sectors is almost consistent with a Gaussian form, while the $q=0$ sector shows a clear excess. It is thus tempting to define a new $\langle q^2 \rangle_{q \neq 0}$ as the width of this Gaussian, where the fit takes only the nontrivial sectors into account. In fact, due to CP symmetry the distribution is even, and we produce a histogram of $|q|$. The $|q| \geq 1$ sectors are then fitted to a half-Gaussian. This procedure stabilizes the contribution from the tails (for this reason a similar “improved estimator” was used in [42]), but it also introduces a model dependence.

β	6.0000	6.0938	6.1802	6.2602	6.3344
L/a	12	14	16	18	20
$\langle q^2 \rangle (r_0/L)^4$	6330(60)	5926(71)	5826(57)	5673(67)	5490(66)
$\langle q^2 \rangle_{q \neq 0} (r_0/L)^4$	6401(69)	6066(95)	5907(66)	5808(92)	5568(84)

Table 3: Summary of the data from the scaling series of runs, in units of 10^{-5} . Here, an extra 0.68% error has been added w.r.t. what is reported in table 1.

L/a ($\beta=6.0$)	10	12	14	16	18
$\langle q^2 \rangle (r_0/L)^4$	4834(42)	6330(42)	6464(43)	6540(43)	6468(42)
$\langle q^2 \rangle_{q \neq 0} (r_0/L)^4$	6428(79)	6401(53)	6458(56)	6545(64)	6474(42)

Table 4: Summary of the data from the infinite volume extrapolation series, in units of 10^{-5} . Here, no extra error has been added w.r.t. what is reported in table 2.

4. Continuum and infinite volume extrapolations

For the continuum and infinite volume extrapolation of the quenched topological susceptibility we convert our data to physical units by multiplying them with the appropriate power of r_0/a . In the scaling series the uncertainty of L/r_0 reflects itself independently for each datapoint. Assuming that L/r_0 is known (at each β) with 0.17% precision (see appendix) an extra 0.68% error needs to be included in table 3. By contrast, in the volume extrapolation series the uncertainty of L/r_0 is correlated, and no extra error has been included in table 4.

For the continuum extrapolation in a fixed physical volume we plot our data against $(a/r_0)^2$. This power of the lattice spacing is motivated by a Symanzik analysis (both the Wilson action and our topological charge operator contain dimension 6, but no dimension 5 operators). In principle, the uncertainty of r_0/a also leads to a horizontal error bar, but its effect is negligible, and we shall omit it. The resulting fits are shown in figure 3. The observables $\langle q^2 \rangle (r_0/L)^4$ and $\langle q^2 \rangle_{q \neq 0} (r_0/L)^4$ extrapolate in a parallel manner, giving the continuum values 0.05092(71) and 0.05205(71), respectively. The difference, if evaluated inside a jackknife, is significant.

This brings us to the discussion of finite volume effects. Standard reasoning suggests that the dominant finite volume corrections in the YM theory come from glueball states travelling around the box. Since euclidean Green's functions fall off exponentially, this would imply

$$Z_\nu(L) = Z_\nu(\infty) \left(1 + \text{const } e^{-M_G L} + \dots \right) \tag{4.1}$$

in close analogy to relative finite volume effects in QCD being proportional to $\exp(-M_\pi L)$. Simple models for the YM vacuum (random population of unit cells by instantons or anti-instantons) suggest no finite volume corrections for the susceptibility and a $1/V$ type

⁶Here it is understood that the total statistics is kept fixed, so that the number of configurations in a given sector drops. Via appropriately increasing the statistics, Z can be determined in a volume as large as desired.




Figure 3: Continuum extrapolation of $\chi_{\text{top}} r_0^4$ in a fixed physical volume, $V = (2.2394 r_0)^4$.




Figure 4: Infinite volume extrapolation of $\chi_{\text{top}} r_0^4$, versus r_0/L [cf. (4.1)] and $(r_0/L)^4$ [cf. (4.2)].

kurtosis.⁷ As a result, one may consider the alternative form of the partition function

$$Z_\nu(L) = Z_\nu(\infty) \left(1 + \text{const}/L^4 + \dots \right). \quad (4.2)$$

The data of table 4 are shown in figure 4. If we assume exponential finite-volume effects, the data for the first estimator may be fitted all the way out to $L/a=10$. The difference of

⁷With *independent* fluctuations in unit cells V_1, \dots, V_N this is generic: $\langle \nu^2 \rangle = \langle (\nu_1 + \dots + \nu_N)^2 \rangle = N \langle \nu_1^2 \rangle$ and $\langle \nu^4 \rangle = \sum_i^N \langle \nu_i^4 \rangle + 6 \sum_{i < j}^N \langle \nu_i^2 \rangle \langle \nu_j^2 \rangle = N \langle \nu_1^4 \rangle + 3N(N-1) \langle \nu_1^2 \rangle \langle \nu_2^2 \rangle$, where we use that all odd moments vanish.




Figure 5: Continuum extrapolation of $\langle q^4 \rangle / \langle q^2 \rangle^2 - 3$ in a fixed physical volume, $V = (2.2394r_0)^4$.




Figure 6: Left: Infinite volume extrapolation of the kurtosis $\langle q^4 \rangle / \langle q^2 \rangle^2 - 3$ in log-log form; this quantity vanishes with $V \rightarrow \infty$. Right: The reduced moment ratio $\langle q^4 \rangle / \langle q^2 \rangle - 3\langle q^2 \rangle$ versus r_0^4/V . This plot includes constant fits to the last 2, 3 or 4 datapoints and a linear fit to all 5 points. We cannot discriminate a finite from a vanishing limit of this quantity with $V \rightarrow \infty$.

the fit curve at $L/a = 12$ and at $L/a = \infty$ suggests that the continuum result for $\langle q^2 \rangle (r_0/L)^4$ is suppressed by the amount 0.00175, and this gives a corrected $\langle q^2 \rangle (r_0/L)^4 = 0.05267(71)$. With the other estimator the fitted value for “const” in (4.1) is consistent with zero, and we stay with $\langle q^2 \rangle_{q \neq 0} (r_0/L)^4 = 0.05205(71)$, as quoted above. If we assume $1/V$ -type effects,

finite-volume corrections are hard to quantify (at least the volume-scaling regime with the first estimator is smaller). Therefore, we decide to stay with the extrapolation via (4.1). Taking the average of both estimators but keeping the full difference as a theoretical uncertainty, we consider

$$\chi_{\text{top}} r_0^4 = 0.05236(71)(62) \tag{4.3}$$

our final result, where the first error is statistical and the second systematic.

Our data for the kurtosis $\langle q^4 \rangle / \langle q^2 \rangle^2 - 3$ scale perfectly (figure 5), and the infinite volume extrapolation is consistent with both an exponential and a power law behavior. We find that in the YM theory a non-zero kurtosis is a pure finite volume effect and has nothing to do with the discretization (figure 6). Since our “independent cell” model of footnote 7 and large N_c arguments suggest that $\langle q^4 \rangle / \langle q^2 \rangle - 3 \langle q^2 \rangle$ has a finite limit with $V \rightarrow \infty$, we plot this quantity versus $1/V$ in figure 6, too. It turns out that our data are consistent with the value 0.276(84) given in [43], but we cannot rule out the possibility that the infinite volume limit might actually vanish.

5. Conversion to physical units

Having an unambiguous result for the topological susceptibility in the combined continuum and infinite volume limit in units of the Sommer scale r_0^{-1} [16], we are left with the question which value [in MeV] the latter should be identified with.

In QCD r_0 is a well-defined quantity. In other words, one could measure aM_p , aM_π and aM_K in a 2+1 flavor simulations in which their ratios are always adjusted to the respective experimental ratios. By considering the continuum limit of $r_0 M_p$, one would have the correct physical value of r_0 , and there is nothing wrong with using this value also in pure YM theory. The original estimate $r_0 = 0.5$ fm [16] for the outcome of this procedure has been superseded, more recently, by values like $r_0 = 0.467(6)$ fm [44, 45]. Alternatively, one could set the scale in the quenched theory via $r_0 f_K = 0.4146(94)$ [46], using $f_K = 160(2)$ MeV from experiment. This has the same effect as attributing $r_0 = 0.512(12)$ fm. To encompass this spread we set the scale via $r_0 = 0.49$ fm and add a 4% error to reflect the intrinsic scale ambiguity in pure YM theory.

With this choice for r_0 our value $\chi_{\text{top}} r_0^4$, as quoted in the previous section, amounts to

$$\chi_{\text{top}}^{1/4} = 193(1)(8) \text{ MeV} \tag{5.1}$$

where the first error bar contains all statistical and systematic uncertainties of our calculation, and the second one reflects the scale setting ambiguity in a theory which is not full QCD.

6. Summary

In this paper we have performed a precision study of the topological susceptibility in pure SU(3) Yang Mills theory in the combined continuum and infinite volume limit using a field-theoretic definition of the topological charge.

	$\chi_{\text{top}} r_0^4$	$\chi_{\text{top}}^{1/4}$ [MeV]	$\chi_{\text{top}}^{1/4}$ [MeV]
Ref. [49] (2001)	0.072(7)	209(5)(8)	—
Ref. [43] (2002)	0.057(3)	197(3)(8)	180(2)(8)
Ref. [39] (2003)	0.055(10)	195(9)(8)	188(12)(8)
Ref. [50] (2003)	0.059(5)	198(4)(8)	—
Ref. [42] (2004)	0.059(3)	198(3)(8)	191(5)(8)
this work	0.0524(9)	193(1)(8)	

Table 5: Summary of the quenched topological susceptibility (in the continuum) as determined in some recent studies. The results $\chi_{\text{top}}/\sigma^2=0.0355(33)$ [49] and $\chi_{\text{top}}/\sigma^2=0.0282(12)$ [43] have been converted to r_0^{-4} units by means of $\sigma^{1/2}r_0 = 1.193(10)$ [51]. In the second column the conversion to MeV has been done with our choice $r_0=0.49$ fm, while the third one contains the values given in the respective papers, but with our standard scale setting error throughout.

Comparing our final result to other recent⁸ determinations [49, 43, 39, 50, 42] (see table 5 for details) we see a downward trend over time. Our result in r_0^{-4} units is substantially more precise, since we work much closer to the continuum (our $\beta=6.0$ lattices are the coarsest ones used in the continuum extrapolation) and due to the large statistics.⁹ The conversion to MeV is, of course, limited by the basic scale setting ambiguity in the quenched theory. In any case our final result (5.1) supports the Witten Veneziano scenario for the origin of the η' mass.

Acknowledgments

We thank Ferenc Niedermayer for useful discussions. Computations were carried out on ALiCENext and a Pentium4 cluster at the University of Wuppertal. This work was supported by the Swiss NSF and under the Hungarian grant OTKA-AT049652.

A. Parametrization of r_0/a

In ref. [41] Necco and Sommer use a polynomial approximation for $\log(r_0/a)$. Specifically

$$\log(r_0/a) \Big|_{\text{NS}} = c_0 + c_1(\beta - 6) + c_2(\beta - 6)^2 + c_3(\beta - 6)^3 \tag{A.1}$$

with $c_0 = 1.6804, c_1 = 1.7331, c_2 = -0.7849, c_3 = 0.4428$ is suggested in order to generate matched lattices for $5.7 \leq \beta \leq 6.92$, i.e. combinations of $(\beta, L/a)$ with fixed L/r_0 . We have repeated the analysis with their data [41] and find essentially the same $c_{0..3}$, with $\chi^2/\text{d.o.f.} = 1.39$.

Here, we wish to explore a rational approximation of $\log(r_0/a)$ which has the merit of being consistent with perturbation theory. In the weak coupling regime $a \propto \exp(-1/[2\beta_0 g_0^2])$

⁸We like to draw the readers attention to two studies [47, 48] with a fermionic charge where no continuum limit has been taken. Note that the charges Q_I in [49] and Q_g in figure 5 of [39] are similar in spirit to our q_{ren} .

⁹There is a similar high-statistics study, based on an overlap charge [52]. They have no continuum limit yet for χ_{top} , but they see a deviation from zero in $\langle q^4 \rangle - 3\langle q^2 \rangle^2$ or the kurtosis in a fixed volume.




Figure 7: Old polynomial $\chi^2/6 = 1.39$ and new rational $\chi^2/6 = 0.91$ fit to the data of ref. [41]. Asymptotic freedom demands that the curve eventually becomes linear with slope $4\pi^2/33$, and this constraint is built into the new rational ansatz (A.2).

L/a	10	12	14	16	18	20
β	5.8996	6.0000	6.0926	6.1790	6.2601	6.3362
L/r_0	2.2357	2.2356	2.2357	2.2357	2.2356	2.2357
β	5.8980	6.0000	6.0938	6.1802	6.2602	6.3344
L/r_0	2.2394	2.2394	2.2394	2.2395	2.2393	2.2393

Table 6: Upper part: $(\beta, L/a)$ combinations which match $(6.0, 12)$ [via achieving $L/r_0 = 2.2356$ as accurately as possible], based on the interpolation formula (A.1) of [41]. Lower part: Same [via achieving $L/r_0 = 2.2394$ as accurately as possible], based on the interpolation formula (A.2).

with the universal coefficient $\beta_0 = 1/(4\pi)^2 \cdot [11N_c/3 - 2N_f/3]$. In SU(3) gluodynamics one has thus $a \propto \exp(-16\pi^2/132 \cdot \beta)$ or $\log(r_0/a) = 4\pi^2/33 \cdot \beta + \dots$ and we employ the ansatz

$$\log(r_0/a) \Big|_{\text{new}} = \frac{4\pi^2}{33} \beta \cdot \frac{1 + d_1/\beta + d_2/\beta^2}{1 + d_3/\beta + d_4/\beta^2} \quad (\text{A.2})$$

where this constraint is built in. Using again their data we find that the best fit is given through $d_1 = -8.2384, d_2 = 15.310, d_3 = -2.7395, d_4 = -11.526$ with $\chi^2/\text{d.o.f.} = 0.91$.

It is straightforward to find the β -values which match a simulation with $(\beta, L/a) = (6.0, 12)$. The results are given in table 6, both for (A.1) and for (A.2). It turns out that the matched β -values are more or less the same with either formula and the pertinent L/r_0 (and hence our estimates for the physical box-length) differ by 0.17% only. This should not come as a surprise, since the two curves in figure 7 are rather close. Still, since the new interpolation has a better theoretical foundation and the $\chi^2/\text{d.o.f.}$ is lower, we choose the new parameterization (A.2). Nonetheless, it is clear that the original parameterization (A.1)

remains a legitimate and good one in the interval in which it is given.

References

- [1] H. Leutwyler and A. Smilga, *Spectrum of Dirac operator and role of winding number in QCD*, *Phys. Rev. D* **46** (1992) 5607.
- [2] S. Durr, *Topological susceptibility in full QCD: lattice results versus the prediction from the QCD partition function with granularity*, *Nucl. Phys. B* **611** (2001) 281 [[hep-lat/0103011](#)].
- [3] TXL collaboration, G.S. Bali et al., *Quark mass effects on the topological susceptibility in QCD*, *Phys. Rev. D* **64** (2001) 054502 [[hep-lat/0102002](#)].
- [4] CP-PACS collaboration, A. Ali Khan et al., *Topological susceptibility in lattice QCD with two flavors of dynamical quarks*, *Phys. Rev. D* **64** (2001) 114501 [[hep-lat/0106010](#)].
- [5] UKQCD collaboration, A. Hart and M. Teper, *The topological susceptibility and $F(\pi)$ from lattice QCD*, *Phys. Lett. B* **523** (2001) 280 [[hep-lat/0108006](#)].
- [6] T.G. Kovacs, *The topological susceptibility with dynamical overlap fermions*, [hep-lat/0111021](#).
- [7] C. Bernard et al., *Topological susceptibility with the improved Asqtad action*, *Phys. Rev. D* **68** (2003) 114501 [[hep-lat/0308019](#)].
- [8] T.A. DeGrand and S. Schaefer, *Chiral properties of two-flavor QCD in small volume and at large lattice spacing*, *Phys. Rev. D* **72** (2005) 054503 [[hep-lat/0506021](#)].
- [9] G.I. Egri, Z. Fodor, S.D. Katz and K.K. Szabo, *Topology with dynamical overlap fermions*, *JHEP* **01** (2006) 049 [[hep-lat/0510117](#)].
- [10] A. Hasenfratz and R. Hoffmann, *Mixed action simulations on staggered background: interpretation and result for the 2-flavor QCD chiral condensate*, *Phys. Rev. D* **74** (2006) 114509 [[hep-lat/0609067](#)].
- [11] E. Witten, *Current algebra theorems for the U(1) 'Goldstone Boson'*, *Nucl. Phys. B* **156** (1979) 269.
- [12] G. Veneziano, *U(1) without instantons*, *Nucl. Phys. B* **159** (1979) 213.
- [13] P. Minkowski, *The eta-prime meson disobeys the Zweig rule up to infinite color*, *Phys. Lett. B* **237** (1990) 531.
- [14] P. Minkowski, *Note on the mass square of the η' meson*, *Phys. Lett. B* **423** (1998) 157 [[hep-ph/9711431](#)].
- [15] L. Girlanda, J. Stern and P. Talavera, *η' mass and chiral symmetry breaking at large- $N(C)$ and $N(f)$* , *Phys. Rev. Lett.* **86** (2001) 5858 [[hep-ph/0103221](#)].
- [16] R. Sommer, *A new way to set the energy scale in lattice gauge theories and its applications to the static force and α_s in SU(2) Yang-Mills theory*, *Nucl. Phys. B* **411** (1994) 839 [[hep-lat/9310022](#)].
- [17] M.F. Atiyah and I.M. Singer, *The index of elliptic operators. 1*, *Ann. Math.* **87** (1968) 484.
- [18] M. Lüscher, *Topology of lattice gauge fields*, *Commun. Math. Phys.* **85** (1982) 39.
- [19] A. Phillips and D. Stone, *Lattice gauge fields, principal bundles and the calculation of topological charge*, *Commun. Math. Phys.* **103** (1986) 599.

- [20] MILC collaboration, T.A. DeGrand, *A variant approach to the overlap action*, *Phys. Rev. D* **63** (2001) 034503 [[hep-lat/0007046](#)].
- [21] W. Bietenholz, *Optimised dirac operators on the lattice: construction, properties and applications*, [hep-lat/0611030](#).
- [22] H. Neuberger, *Exactly massless quarks on the lattice*, *Phys. Lett. B* **417** (1998) 141 [[hep-lat/9707022](#)].
- [23] P.H. Ginsparg and K.G. Wilson, *A remnant of chiral symmetry on the lattice*, *Phys. Rev. D* **25** (1982) 2649.
- [24] P. Hasenfratz, V. Laliena and F. Niedermayer, *The index theorem in QCD with a finite cut-off*, *Phys. Lett. B* **427** (1998) 125 [[hep-lat/9801021](#)].
- [25] M. Lüscher, *Exact chiral symmetry on the lattice and the Ginsparg-Wilson relation*, *Phys. Lett. B* **428** (1998) 342 [[hep-lat/9802011](#)].
- [26] M. Gockeler, A.S. Kronfeld, M.L. Laursen, G. Schierholz and U.J. Wiese, *Can the topological susceptibility be calculated from SU(N) lattice gauge theories?*, *Phys. Lett. B* **233** (1989) 192.
- [27] M. Campostrini, A. Di Giacomo, H. Panagopoulos and E. Vicari, *Topological charge, renormalization and cooling on the lattice*, *Nucl. Phys. B* **329** (1990) 683.
- [28] A. Di Giacomo and E. Vicari, *Renormalization and topological susceptibility on the lattice*, *Phys. Lett. B* **275** (1992) 429.
- [29] B. Alles, M. D'Elia, A. Di Giacomo and R. Kirchner, *A critical comparison of different definitions of topological charge on the lattice*, *Phys. Rev. D* **58** (1998) 114506 [[hep-lat/9711026](#)].
- [30] E. Seiler and I.O. Stamatescu, *Some remarks on the Witten-Veneziano formula for the η' mass* MPI-PAE/PTH10/87.
- [31] S. Chandrasekharan, *Lattice QCD with Ginsparg-Wilson fermions*, *Phys. Rev. D* **60** (1999) 074503 [[hep-lat/9805015](#)].
- [32] F. Niedermayer, *Exact chiral symmetry, topological charge and related topics*, *Nucl. Phys. B* **73** (Proc. Suppl.) (1999) 105 [[hep-lat/9810026](#)].
- [33] L. Giusti, G.C. Rossi and M. Testa, *Topological susceptibility in full QCD with Ginsparg-Wilson fermions*, *Phys. Lett. B* **587** (2004) 157 [[hep-lat/0402027](#)].
- [34] A. Hasenfratz and F. Knechtli, *Flavor symmetry and the static potential with hypercubic blocking*, *Phys. Rev. D* **64** (2001) 034504 [[hep-lat/0103029](#)].
- [35] S. Durr, *Gauge action improvement and smearing*, *Comput. Phys. Commun.* **172** (2005) 163 [[hep-lat/0409141](#)].
- [36] C. Christou, A. Di Giacomo, H. Panagopoulos and E. Vicari, *Improved lattice operators: the case of the topological charge density*, *Phys. Rev. D* **53** (1996) 2619 [[hep-lat/9510023](#)].
- [37] C.R. Gatttringer, I. Hip and C.B. Lang, *Topological charge and the spectrum of the fermion matrix in lattice QED(2)*, *Nucl. Phys. B* **508** (1997) 329 [[hep-lat/9707011](#)].
- [38] N. Cundy, M. Teper and U. Wenger, *Topology and chiral symmetry breaking in SU(N(c)) gauge theories*, *Phys. Rev. D* **66** (2002) 094505 [[hep-lat/0203030](#)].

- [39] L. Del Debbio and C. Pica, *Topological susceptibility from the overlap*, *JHEP* **02** (2004) 003 [[hep-lat/0309145](#)].
- [40] Online at <http://www.physics.utah.edu/~detar/milc>.
- [41] S. Necco and R. Sommer, *The $N(f) = 0$ heavy quark potential from short to intermediate distances*, *Nucl. Phys.* **B 622** (2002) 328 [[hep-lat/0108008](#)].
- [42] L. Del Debbio, L. Giusti and C. Pica, *Topological susceptibility in the SU(3) gauge theory*, *Phys. Rev. Lett.* **94** (2005) 032003 [[hep-th/0407052](#)].
- [43] L. Del Debbio, H. Panagopoulos and E. Vicari, *Theta dependence of SU(N) gauge theories*, *JHEP* **08** (2002) 044 [[hep-th/0204125](#)].
- [44] A.A. Khan et al., *Axial coupling constant of the nucleon for two flavours of dynamical quarks in finite and infinite volume*, *Phys. Rev.* **D 74** (2006) 094508 [[hep-lat/0603028](#)].
- [45] C. Aubin et al., *Light hadrons with improved staggered quarks: approaching the continuum limit*, *Phys. Rev.* **D 70** (2004) 094505 [[hep-lat/0402030](#)].
- [46] ALPHA collaboration, J. Garden, J. Heitger, R. Sommer and H. Wittig, *Precision computation of the strange quark's mass in quenched QCD*, *Nucl. Phys.* **B 571** (2000) 237 [[hep-lat/9906013](#)].
- [47] C. Gattringer, R. Hoffmann and S. Schaefer, *The topological susceptibility of SU(3) gauge theory near $T(c)$* , *Phys. Lett.* **B 535** (2002) 358 [[hep-lat/0203013](#)].
- [48] P. Hasenfratz, S. Hauswirth, T. Jorg, F. Niedermayer and K. Holland, *Testing the fixed-point QCD action and the construction of chiral currents*, *Nucl. Phys.* **B 643** (2002) 280 [[hep-lat/0205010](#)].
- [49] B. Lucini and M. Teper, *SU(n) gauge theories in four dimensions: exploring the approach to $N = \infty$* , *JHEP* **06** (2001) 050 [[hep-lat/0103027](#)].
- [50] L. Giusti, M. Lüscher, P. Weisz and H. Wittig, *Lattice QCD in the epsilon-regime and random matrix theory*, *JHEP* **11** (2003) 023 [[hep-lat/0309189](#)].
- [51] F. Niedermayer, P. Rufenacht and U. Wenger, *Fixed point gauge actions with fat links: scaling and glueballs*, *Nucl. Phys.* **B 597** (2001) 413 [[hep-lat/0007007](#)].
- [52] L. Giusti, B. Taglienti and S. Petrarca, *Large statistics study of QCD topological charge distribution*, *PoS(LAT2006)* 058.