

Giant gravitons on deformed pp-waves

Alex Hamilton

*Department of Physics, Columbia University, New York,
NY 10027 U.S.A.*

E-mail: hamilton@phys.columbia.edu

Jeff Murugan

*Department of Mathematics and Applied Mathematics, University of Cape Town,
Private Bag, Rondebosch 7700, South Africa*

E-mail: jeff@nassp.uct.ac.za

ABSTRACT: The recently constructed Lunin-Maldacena deformation of $AdS_5 \times S^5$ is known to support two inequivalent Penrose limits that lead to BPS pp-wave geometries. In this note, we construct new giant graviton solutions on these backgrounds. A detailed study of the spectra of small fluctuations about these solutions reveals a remarkably rich structure. In particular, the giants that we construct fall into two classes, one of which appears to remain stable in the Penrose limit independently of the strength of the deformation. The other class of giants, while more difficult to treat analytically, seems to exhibit a shape deformation not unlike the so-called “squashed giants” seen in the pp-wave with a constant NS B -field turned on. Some consideration is also given to the associated giant operators in the BMN limit of the dual $\mathcal{N} = 1$ SYM gauge theory.

KEYWORDS: AdS-CFT Correspondence, Penrose limit and pp-wave background, D-branes.

Contents

1. Introduction	1
2. Giant gravitons, pp-waves and orbits of S^5.	4
3. The Lunin-Maldacena geometry and its Penrose limits	8
4. Giants on the deformed pp-wave	12
4.1 $(J, 0, 0), (0, J, 0)$ and $(0, 0, J)$ orbits	12
4.2 (J, J, J) orbits	16
5. Small fluctuations and stability of giants	19
5.1 Undeformed giants	20
5.2 Deformed giants	22
6. Comments on the dual operators	24
7. Summary and outlook	27

1. Introduction

By now, the gauge theory/gravity correspondence needs little in the way of motivation. Indeed, since its concrete realization in Maldacena’s AdS/CFT conjecture [1] nearly a decade ago, this remarkable duality has revolutionized the ways in which we think both about quantum field theories and their gravity duals. From an understanding of the emergence of the entropy of a black hole from its constituent microstates [2], to the emergence of spacetime itself [3], the power of the correspondence is surpassed only by its stubborn resistance to direct proof. To date though, not only has the conjecture successfully withstood an enormous barrage of tests and checks, it has also emerged as one of the most promising approaches to understanding the strongly coupled behaviour of non-Abelian gauge theories. Indeed, the eventual goal of this program is nothing less than a weakly coupled gravitational dual to strongly coupled QCD, one that will be able to shed light on the non-perturbative structure of strong interactions (see, for example, [4] and references therein).

Fuelled largely by the discovery of D -brane degrees of freedom in string theory [5], development on the gravity side has been equally impressive. Among this class of extended objects, the giant gravitons of the $AdS_5 \times S^5$ solution of type IIB supergravity [6, 7] furnish a particularly rich background in which to study non-perturbative effects in the string theory [8]. As BPS objects blown up on an S^3 contained in either the AdS_5 or 5-sphere

parts of the geometry, computations associated to these giant gravitons are protected by powerful non-renormalization theorems and can be extrapolated from weak to strong coupling. Moreover, while giants carry all the same quantum numbers as their energetically degenerate point counterparts, as expanded states in the theory, their interactions are much softer. These interactions however, are difficult to calculate in the supergravity theory. Fortunately, the operators corresponding to giant gravitons have been identified and studied in some detail in the dual $\mathcal{N} = 4$ super Yang-Mills theory with gauge groups $U(N)$ [10] and $SU(N)$ [11]. In the $U(N)$ gauge theory at least, these giant operators are Schur polynomials of the form¹

$$\chi_R(X) = \frac{1}{n!} \sum_{\sigma \in S_n} \chi_R(\sigma) \text{tr}(\sigma X) ,$$

constructed from the Higgs fields of the SYM multiplet and characterised by Young diagrams of the representation R of $U(N)$. Consequently, the dynamics of the giant graviton is encoded in the Feynman diagrams of the gauge theory via the multi-point correlators of these Schur operators. The study of these operators and the realisation that these states afford a description in terms of free fermions of a single matrix model (see the first ref. of [10] and [12]) has led to a remarkable surge of activity focussed on extracting information about the supergravity background from the gauge theory (see [13] and references thereof) culminating in a complete classification of all the classical half-BPS solutions of the type IIB supergravity [14]. This said though, most of what is known about giant gravitons and their dual operators is known in the highly supersymmetric setting of the $AdS_5 \times S^5/\mathcal{N} = 4$ SYM duality. If we are to believe that the duality is an exact one eventually leading to a gravity dual of full 4-dimensional QCD then it is crucial that we understand and extend the gauge/gravity dictionary to less supersymmetric backgrounds.²

A significant step toward a systematic study of less supersymmetric systems was taken by Leigh and Strassler [15] in their construction of a three parameter family of $\mathcal{N} = 1$ SYM gauge theories obtained by an exactly marginal deformation of the $\mathcal{N} = 4$ superpotential. Then, by exploiting the fact that, at least for a one parameter subset, this Leigh-Strassler deformation may be rewritten as a Moyal-like deformation,³

$$\Phi_i * \Phi_j = e^{i\pi\gamma(Q_i^1 Q_j^2 - Q_i^2 Q_j^1)} \Phi_i \Phi_j ,$$

Lunin and Maldacena showed that the dual supergravity solution [30] could be constructed from the type IIB $AdS_5 \times S^5$ by a suitable single-parameter deformation of the 5-sphere. This realization has stimulated an enormous resurgence of interest in string theory on this deformed background including, but not limited to, extensions of the Lunin-Maldacena deformation to the eleven dimensional geometries of the Einstein-Sasaki form $AdS_5 \times Y^{p,q}$ [17],

¹For the special case of giants blown up on the $S^3 \subset S^5$, these Schur polynomials reduce to the more familiar subdeterminant operators identified in the second of refs [10].

²Ideally, we would like to be able to extend it further to non-conformal theories as well but we will adopt the time-honored tradition of learning to walk before trying to fly.

³For a recent extension of this star product to capture a more general form of the Leigh Strassler deformation see [16]

deformations of $\mathcal{N} = 1$ and $\mathcal{N} = 2$ theories [18] and a number of studies of semiclassical strings in this background [19]. A particularly insightful contribution to this set of ideas came in [20] where it was shown that the Lunin-Maldacena geometry can equivalently be reached by a sequence of T-dualities and shifts of the angular coordinates by the deformation parameter γ - so-called TsT transformations. If, on the other hand, the different angular directions are shifted by *different* parameter values, say γ_i , the resulting supergravity background is a completely non-supersymmetric deformation⁴ of $AdS_5 \times S^5$ whose gauge theory dual can be systematically constructed from an appropriate generalization of the Leigh-Strassler deformation [21]. Remarkably, this multi-parameter deformation is, in some ways, simpler than the single-parameter Lunin-Maldacena deformation. This is particularly true of the study of giant gravitons on these geometries. Generally very difficult to construct, it was rather astutely observed in [22] that many of the computations simplify significantly when the deformation in the direction of motion of the $D3$ -brane decouples ($\hat{\gamma}_1 = 0$). Unlike the undeformed geometries previously explored, these giant gravitons were found to no longer be energetically degenerate with the point graviton and consequently unstable, a conclusion supported in the spectrum of small fluctuations about the giant. Curiously though, even in the light of this instability, there seemed to be perfect quantitative agreement between the gauge theory and semiclassical open string dynamics.

In this article, we revisit the construction of giant graviton solutions in the single-parameter Lunin-Maldacena background. With the deformation coupling to the direction of motion of the D -brane, a direct construction of the giant graviton solution in the supergravity proves to be very difficult problem and, until recently, had remained an unsolved problem. There is, however, a limit - the Penrose limit - in which the analysis simplifies significantly [23]. This in itself is hardly surprising; after all the lightcone gauge string action is already known to drastically simplify in this background, enough that the perturbative string spectrum is exactly solvable [24]. The Penrose limit of a given geometry is obtained by focussing on a null geodesic in the background. The limiting geometry is the so-called pp-wave. For $AdS_5 \times S^5$, the Penrose limit comes from focussing on a great circle of the S^5 and then boosting to the infinite momentum frame along this geodesic. There are, of course, an infinite number of limits that can be taken, one for each great circle. However, since any great circle may be rotated into any other by the $SO(6)$ isometry of the 5-sphere, there is essentially only one resulting pp-wave. On the other hand, the Penrose limit of the deformed background is not unique. There are, in fact, two different limits that result in BPS backgrounds. One of these is the usual pp-wave background in magnetic coordinates and the other, a homogeneous pp-wave [31, 35, 36]. We would like to determine whether these backgrounds support giant graviton solutions.

Toward this end, the next section of this article is devoted to a summary of the construction of giant gravitons on the maximally supersymmetric type IIB pp-wave. We pay particular attention to the different null geodesics about which the Penrose limit can be taken and how they are related. Thereafter, in the interests of self-containment, we dis-

⁴The superconformal Lunin-Maldacena deformation then, is a special case in which all three tori are deformed by the same amount, $\hat{\gamma}_1 = \hat{\gamma}_2 = \hat{\gamma}_3 = \hat{\gamma}$

discuss the Lunin-Maldacena marginal deformation of $AdS_5 \times S^5$ and the two Penrose limits that result in BPS geometries. In section 4. we show that, indeed both of these deformed pp-waves support giant gravitons exhibiting a remarkably rich structure. In section 5, we initiate an exploration of some of this structure in the spectrum of small fluctuations about the giant graviton equilibrium configuration and conclude, finally, with some comments on the dual gauge theory operators and speculation on future directions.

Soon after this article appeared on the arXiv, we were made aware of [39] which has some overlap with our work and resolves the apparent puzzle in [22]. In the former, giant graviton configurations were investigated in the non-supersymmetric 3-parameter Frolov-Roiban-Tseytlin geometry and through a detailed study of the fluctuation spectra of giants blown up in both the sphere and AdS directions, it was shown that giant gravitons with angular momenta along one of the equatorial directions - what we call $(J, 0, 0)$, $(0, J, 0)$ and $(0, 0, J)$ configurations - are stable and independent of the deformation. This is in excellent agreement with our findings in section 4. and we return to this point there.

2. Giant gravitons, pp-waves and orbits of S^5 .

In the context of $AdS_5 \times S^5$ giant gravitons are, by now, well known to be a KK- mode blown up along a transverse S^3 contained in the 5-sphere⁵ part of the geometry. This blowing up is a consequence of the motion of the graviton along some geodesic - a great circle - of the S^5 through which a 5-form flux is threaded. On the other hand, the pp-wave background is obtained from the AdS geometry by focussing in on a particular null geodesic, ostensibly by asking what geometry a particle boosted to the infinite momentum frame along the geodesic “sees”. It makes sense then, to ask if the giants found on the $AdS_5 \times S^5$ background survive the Penrose limit. As far as we are aware, this question was first posed and answered in the affirmative in [25]. There, the authors found that giant gravitons in the maximally supersymmetric type IIB pp-wave divide into two types [26] depending on the geodesic along which the Penrose limit is taken. If the orbit of the graviton coincides with the geodesic along which the Penrose limit is taken the resulting giant graviton is a spherical $D3$ -brane whose worldvolume extends along x^+ and the spatial directions of an S^3 contained in the 5-sphere. These D -branes couple directly to the RR-field and, while they don't permit an exact worldsheet description, they do afford an alternate description in terms of multiple D -strings whose moduli are argued to be the fluctuations of the giant graviton [9]. If, however, the Penrose limit is taken along a geodesic different to the one that the $D3$ is moving along, the resulting configuration has Neumann boundary conditions in the x^+ and x^- directions as well as two spatial directions (which could be in either the 5-sphere or the AdS part of the geometry). A full light-cone gauge worldsheet analysis of these D -branes reveal that they preserve 16 supersymmetries but only if they are rotating in some transverse two-plane in the pp-wave background. As fascinating as these latter solutions are, we find the structure of the former (giant graviton) solutions on the deformed pp-wave rich enough on its own that, at least for the purposes

⁵Or, in the case of “dual giants”, the AdS_5

of this note, we will restrict our attention to them alone. A study of the second class of D -branes in the Lunin-Maldacena deformed background remains an open problem.

More detailed studies of these configurations, their supersymmetries, fluctuations and responses to certain non-vanishing B -fields may be found in refs. [28, 29]. In order to facilitate comparison with our later results, we now, very briefly but systematically review the construction of giant graviton solutions in the maximally supersymmetric type IIB pp-wave background. The construction works equally well when the Penrose limit is taken about $(J, 0, 0), (0, J, 0), (0, 0, J)$ or (J, J, J) geodesics. In particular, we show how the two sets of solutions are related by a light-cone-time-dependent rotation of coordinates (or, in the jargon of general relativists, a coordinate transformation between Brinkmann and Rosen coordinates). In the undeformed pp-wave background this transformation is more-or-less trivial, reflecting the fact that the Penrose limit is essentially unique - any orbit on the 5-sphere may be rotated into any other via the $SO(6)$ isometry group. This, we will find, is no longer the case for the pp-waves obtained from the deformed Lunin-Maldacena background.

The maximally supersymmetric type IIB pp-wave⁶

$$ds^2 = -2dx^+ dx^- - \left(\sum_{i=1}^4 (x^i)^2 + \sum_{a=5}^8 (x^a)^2 \right) (dx^+)^2 + \sum_{i=1}^4 (dx^i)^2 + \sum_{a=4}^8 (dx^a)^2, \quad (2.1)$$

obtains from the 10-dimensional $AdS_5 \times S^5$ geometry with metric (written in global coordinates)

$$ds^2 = R^2 \left(-\cosh^2 \rho dt^2 + d\rho^2 + \sinh^2 \rho d\Omega_3^2 + \cos^2 \alpha d\phi_1^2 + d\alpha^2 + \sin^2 \alpha d\tilde{\Omega}_3^2 \right),$$

by rescaling⁷

$$x^+ = \frac{1}{2}(t + \phi_1), \quad x^- = R^2(t - \phi_1), \quad \rho = \frac{r}{R}, \quad \alpha = \frac{y}{R}, \quad (2.2)$$

and sending $R \rightarrow \infty$. Additionally, the only components of the 5-form flux $F_{(5)}$ that survives the limit are those with a plus index,

$$F_{+1234} = F_{+5768} = \text{const.}, \quad (2.3)$$

with the constant fully determined once a normalisation of $F_{(5)}$ is specified. Physically, this is the geometry near the trajectory - the $(J, 0, 0)$ orbit - of a particle moving with large angular momentum along the ϕ_1 direction and sitting at $\rho = \alpha = 0$. When this particle is a $D3$ -brane, its action, in lightcone gauge and with a spherical ansatz for the brane worldvolume, can be written

$$S = -\frac{T_3 \lambda}{g_s} \int d\tau d\theta d\phi_2 d\phi_3 \left[r^3 \sqrt{2\lambda\nu + \lambda^2} - \lambda r^4 \right], \quad (2.4)$$

⁶In this and what follows, we have absorbed the usual mass parameter in the pp-wave into the definition of x^-

⁷Here, $r^2 = \sum_i (x^i)^2$ and $y^2 = \sum_a (x^a)^2$.

where λ and ν are defined through $X^+ = \lambda\tau$ and $X^- = \nu\tau$ respectively. The corresponding Hamiltonian, as a function of the radius of the $D3$ worldvolume, is minimized at $r = 0$, the point graviton, and $r = \sqrt{p^+ g_s / 2\pi^2 T_3}$, the giant graviton.

In the above discussion, the S^5 part of the geometry is parameterized by the five angles $(\alpha, \theta, \phi_1, \phi_2, \phi_3)$ in terms of which the 3–sphere metric is explicitly written as

$$d\tilde{\Omega}_3^2 = d\theta^2 + \cos^2 \theta d\phi_2^2 + \sin^2 \theta d\phi_3^2.$$

There are clearly three $U(1)$ isometries related to translations along each of ϕ_1 , ϕ_2 and ϕ_3 . A general orbit in this background is a linear superposition of these three isometries. A particle moving on the (J, J, J) orbit then, has equal angular momentum in each of these directions. To take a Penrose limit along a null geodesic associated to this orbit, we define the new angular variable $\psi = (\phi_1 + \phi_2 + \phi_3)/3$ with periodicity 2π . The geodesic we will focus on is given by

$$t = \psi, \quad \rho = 0, \quad \alpha = \alpha_0, \quad \theta = \frac{\pi}{4}, \quad \psi = \phi_1 = \phi_2 = \phi_3. \quad (2.5)$$

The geometry in the neighbourhood of this geodesic is recovered by defining lightcone coordinates as

$$x^+ = \frac{1}{2}(t + \psi), \quad x^- = R^2(t - \psi), \quad (2.6)$$

rescaling appropriate coordinates as

$$\begin{aligned} \rho &= \frac{r}{R}, & \alpha &= \alpha_0 + \frac{y^1}{R}, & \theta &= \frac{\pi}{4} + \sqrt{\frac{3}{2}} \frac{y^2}{R}, & \phi_1 &= \psi - \sqrt{2} \frac{y^3}{R}, \\ \phi_2 &= \psi + \frac{y^3 - \sqrt{3}y^4}{\sqrt{2}R}, & \phi_3 &= \psi + \frac{y^3 + \sqrt{3}y^4}{\sqrt{2}R}, \end{aligned} \quad (2.7)$$

and taking $R \rightarrow \infty$. The result is, of course, a pp-wave, this time with metric⁸

$$\begin{aligned} ds^2 &= -2dx^+ dx^- - \tilde{y}^2 (dx^+)^2 + 4(y^1 dy^3 + y^2 dy^4) dx^+ + d\tilde{y}^2 + d\tilde{r}^2 \\ &\quad + \tilde{y}^2 d\Omega_3^2 + \tilde{r}^2 d\tilde{\Omega}_3^2. \end{aligned} \quad (2.8)$$

The non-vanishing components of the 5–form in this background remains the same as in eq. (2.3). String theory on this background was previously studied in [31, 32] where the similarity between this metric and that of the motion of a particle in a magnetic field earned it the moniker of “magnetic pp-wave”. As expected, all 32 of the supersymmetries of $AdS_5 \times S^5$ survive the Penrose limit and, since the large J limit of the dual gauge theory is independent of the choice of $U(1)$ R-charge group, the gauge dual of this background is the same truncation of $\mathcal{N} = 4$ SYM as in [24].

Even though the two backgrounds in eqs. (2.1) and (2.8) *seem* very different, they are essentially the same, just written in different coordinates. This is not unexpected since

⁸Now, $\tilde{r}^2 = \sum_i (y^i)^2$ and $\tilde{y}^2 = \sum_a (y^a)^2$ and, incidentally, we have absorbed the “mass” parameter μ into the definition of the lightcone variables.

this magnetic pp-wave is just the geometry near one particular great circle (the (J, J, J) orbit) and any such great circle can be rotated into any other under the action of the $SO(6)$ isometry of the S^5 . This rotation may be made more explicit first by making the change of variables

$$\begin{aligned} x^+ &\rightarrow x^+, \\ x^- &\rightarrow x^- + y^1 y^3 + y^2 y^4, \end{aligned}$$

followed by the lightcone time-dependent coordinate rotation

$$\begin{bmatrix} z^1 \\ z^2 \\ z^3 \\ z^4 \end{bmatrix} = \begin{bmatrix} \cos x^+ & 0 & -\sin x^+ & 0 \\ 0 & \cos x^+ & 0 & -\sin x^+ \\ \sin x^+ & 0 & \cos x^+ & 0 \\ 0 & \sin x^+ & 0 & \cos x^+ \end{bmatrix} \begin{bmatrix} \tilde{y}^1 \\ \tilde{y}^2 \\ \tilde{y}^3 \\ \tilde{y}^4 \end{bmatrix}. \tag{2.9}$$

Implementing this change of variables in the magnetic pp-wave metric gives

$$ds^2 = -2dx^+ dx^- - (\tilde{y}^2 + z^2) (dx^+)^2 + d\tilde{y}^2 + \tilde{y}^2 d\Omega_3^2 + dz^2 + z^2 d\tilde{\Omega}_3^2;$$

precisely the $(J, 0, 0)$ metric of (2.1). The corresponding 5-form field is

$$F_{(5)} = \frac{4}{g_s} dx^+ \wedge \left(dz^1 \wedge dz^2 \wedge dz^3 \wedge dz^4 + dy^5 \wedge dy^6 \wedge dy^7 \wedge dy^8 \right).$$

In these coordinates, the construction of giant gravitons in this background is relatively straightforward. After all, having made the rotation to the z coordinates, it remains only to

- Parameterize the 3-dimensional worldvolume of the giant in static coordinates as

$$\begin{aligned} Z^1 &= z \cos \theta \cos \phi_2, \\ Z^2 &= z \sin \theta \cos \phi_3, \\ Z^3 &= z \cos \theta \sin \phi_2, \\ Z^4 &= z \sin \theta \sin \phi_3. \end{aligned}$$

- Substitute this ansatz into the worldvolume metric in lightcone gauge and,
- Compute the corresponding energy functional $E(z)$ together with it's turning points.

This simplicity does not, unfortunately, translate to the $\hat{\gamma}$ -deformed pp-wave. Nevertheless we can still extract some useful lessons from this exercise. In particular, inverting the coordinate transformation (2.9) gives the appropriate coordinatization of the giant worldvolume in magnetic coordinates as

$$\begin{aligned} \tilde{Y}^1 &= z \cos \theta \cos(\phi_2 - x^+), \\ \tilde{Y}^2 &= z \sin \theta \cos(\phi_3 - x^+), \\ \tilde{Y}^3 &= z \cos \theta \sin(\phi_2 - x^+), \\ \tilde{Y}^4 &= z \sin \theta \sin(\phi_3 - x^+). \end{aligned}$$

These coordinates are manifestly x^+ , lightcone time dependent. Physically, the giant on this pp-wave background also rotates with equal angular momentum in both the $(\tilde{y}^1, \tilde{y}^3)$ and $(\tilde{y}^2, \tilde{y}^4)$ planes. It is precisely this form of the magnetic coordinates that we will use of the construction of giant gravitons in the deformed pp-wave of [35, 36]. Finally, to close this section it is worth mentioning that in lightcone coordinates, the $D3$ -brane action is, not surprisingly, identical to (2.4).

3. The Lunin-Maldacena geometry and its Penrose limits

In principle, the Lunin-Maldacena construction [30] is identical to that used in generating the holographic duals of noncommutative field theories [33]. It hinges on the use of an $SL(2, \mathbb{R})$ transformation of the full $SL(3, \mathbb{R}) \times SL(2, \mathbb{R})$ duality group of type IIB supergravity compactified along a corresponding $U(1) \times U(1)$ two-torus. In practice, if the metric of this two-torus is denoted by g and the NS-NS 2-form by B then the Lunin-Maldacena deformation is implemented by making the replacement

$$\tau \equiv B + i\sqrt{g} \longrightarrow \tau_\gamma = \frac{\tau}{1 + \gamma\tau}. \tag{3.1}$$

Since this deformation is an exactly marginal one, the AdS factor remains unchanged. In order to apply this construction to the gravitational $AdS_5 \times S^5$ background,⁹

$$\begin{aligned} ds^2 &= R^2 \left(-\cosh^2 \rho dt^2 + d\rho^2 + \sinh^2 \rho d\Omega_3^2 + \sum_{i=1}^3 d\mu_i^2 + \mu_i^2 d\phi_i^2 \right), \\ F_{(5)} &= 4R^4 e^{-\phi_0} (\omega_{AdS_5} + \omega_{S^5}), \\ e^\phi &= e^{\phi_0}, \end{aligned} \tag{3.2}$$

we first define three new angles through

$$\phi_1 = \psi + \varphi_1 + \varphi_2, \quad \phi_2 = \psi - \varphi_1, \quad \phi_3 = \psi - \varphi_2, \tag{3.3}$$

then choose a $U(1) \times U(1)$ subgroup that acts by shifting φ_1 and φ_2 . The corresponding two-torus along which we compactify has τ parameter

$$\tau = iR\sqrt{\mu_1^2\mu_2^2 + \mu_1^2\mu_3^2 + \mu_2^2\mu_3^2},$$

Finally, applying the Lunin-Maldacena deformation (3.1) gives the type IIB supergravity

⁹Here, $(\mu_1, \mu_2, \mu_3) = (\cos \alpha, \sin \alpha \cos \theta, \sin \alpha \sin \theta)$ and the common radius of the AdS_5 and S^5 parts of the geometry, $R^4 = 4\pi e^{\phi_0} N l_s^4$. Note also that in the original undeformed geometry, the NS-NS 2-form $B = 0$. and the coordinates have been chosen to manifestly a $U(1)^3$ subgroup of the full $SO(6)$ isometry of the round 5-sphere.

multiplet,

$$\begin{aligned}
 ds^2 &= R^2 \left[-\cosh^2 \rho dt^2 + d\rho^2 + \sinh^2 \rho d\Omega_3^2 + d\alpha^2 + G \cos^2 \alpha d\phi_1^2 + \sin^2 \alpha (d\theta^2 \right. \\
 &\quad \left. + G \cos^2 \theta d\phi_2^2 + G \sin^2 \theta d\phi_3^2) + \hat{\gamma}^2 G \cos^2 \alpha \sin^4 \alpha \cos^2 \theta \sin^2 \theta (d\phi_1 + d\phi_2 + d\phi_3)^2 \right], \\
 B_{(2)} &= \hat{\gamma} R^2 G \left(\sin^2 \alpha \cos^2 \alpha \cos^2 \theta d\phi_1 \wedge d\phi_2 + \sin^2 \alpha \cos^2 \alpha \sin^2 \theta d\phi_3 \wedge d\phi_1 \right. \\
 &\quad \left. + \sin^4 \alpha \cos^2 \theta \sin^2 \theta d\phi_2 \wedge d\phi_3 \right), \\
 F_{(3)} &= -\frac{4\hat{\gamma}}{g_s} R^2 \cos^2 \alpha \sin^3 \alpha \cos \theta \sin \theta d\alpha \wedge d\theta \wedge (d\phi_1 + d\phi_2 + d\phi_3), \\
 F_{(5)} &= \frac{4}{g_5} R^4 \left(\cosh \rho \sinh^3 \rho dt \wedge d\rho \wedge d\Omega_3 + G \cos \alpha \sin^3 \alpha d\phi_1 \wedge d\alpha \wedge d\tilde{\Omega}_3 \right),
 \end{aligned}$$

where $G^{-1} = 1 + \hat{\gamma}^2 (\cos^2 \alpha \sin^2 \alpha + \sin^4 \alpha \cos^2 \theta \sin^2 \theta)$ and the rescaled $\hat{\gamma} = R^2 \gamma$. Since the deformation is continuous, this one-parameter family of backgrounds are topologically identical to that of the original round sphere. Clearly also, only the sphere part of the geometry suffers any deformation with the original $SO(6)$ isometry group breaking to a $U(1)^3$. This has important consequences for the Penrose limits of this background that we wish to consider. In the undeformed case, all null geodesics lying inside the 5-sphere can be rotated into each other by the action of the $SO(6)$ isometry group. Now, with only a $U(1)^3$ group remaining, this is no longer the case. Indeed there are two distinct classes of BPS geodesics whose Penrose limits result in non diffeomorphic pp-wave metrics. With τ as a worldline coordinate and setting $t = \phi_1 = \phi_2 = \phi_3 = \tau$, the maximum circles on the deformed S^5 with

$$(\mu_1^2, \mu_2^2, \mu_3^2) = (1, 0, 0), (0, 1, 0), (0, 0, 1),$$

furnish one set and the null geodesics with

$$(\mu_1^2, \mu_2^2, \mu_3^2) = (1/3, 1/3, 1/3),$$

the other. Since each of the $(J, 0, 0)$, $(0, J, 0)$ and $(0, 0, J)$ orbits may be rotated into each other with the residual isometry after the $SL(2, \mathbb{R})$ action on the 5-sphere, we need only focus on one of this set of orbits. To take the Penrose limit about the $(J, 0, 0)$ orbit say, on S_γ^5 , we set

$$\begin{aligned}
 \rho &= \frac{y}{R}, & \alpha &= \frac{r}{R}, \\
 t &= x^+ + \frac{x^-}{2R}, & \phi_1 &= x^+ - \frac{x^-}{2R},
 \end{aligned}$$

and then take $R \rightarrow \infty$. With a little algebra, it is easily seen that the resulting background

fields are

$$\begin{aligned}
 ds^2 &= -2dx^+ dx^- - (y^2 + (1 + \hat{\gamma}^2) r^2) (dx^+)^2 + dy^2 + y^2 d\Omega_3^2 + dr^2 + r^2 d\tilde{\Omega}_3^2, \\
 B_{(2)} &= \hat{\gamma} r^2 (\cos^2 \theta dx^+ \wedge d\phi_2 + \sin^2 \theta d\phi_3 \wedge dx^+), \\
 F_{(5)} &= \frac{4}{g_s} \left(y^3 dx^+ \wedge dy \wedge d\Omega + r^3 dx^+ \wedge dr \wedge d\tilde{\Omega}_3 \right) \\
 C_{(4)} &= -\frac{1}{g_s} \left(y^4 dx^+ \wedge d\Omega_3 + r^4 dx^+ \wedge d\tilde{\Omega}_3 \right) \\
 C_{(2)} &= 0, \\
 F_{(3)} &= dC_{(2)} = 0,
 \end{aligned} \tag{3.4}$$

with $d\tilde{\Omega}_3 = \sin \theta \cos \theta d\theta \wedge d\phi_1 \wedge d\phi_3$ the standard volume element on the round 3–sphere. Closed strings on this pp-wave were quantized, and their relation to the BMN limit of the dual $\mathcal{N} = 1$ field theory was studied initially in [31] and then in the current context in [30]. In a remarkable feat of reverse engineering, the authors of [31] found that, in lightcone gauge $x^+ = \tau$, the theory on the worldsheet is a massive one with a transverse oscillation spectrum in the deformed sphere directions of

$$\omega_n = \sqrt{1 + \left(\frac{n}{|p^+|} \pm \hat{\gamma} \right)^2}, \tag{3.5}$$

by working backwards from knowing the spectrum of anomalous dimensions of the corresponding gauge theory operators. This result was later confirmed by direct computation in [30].

An altogether different Penrose limit can be taken by focussing on states having charges near (J, J, J) . These live near the null geodesic $\tau = \psi$ with $\alpha_0 = \cos^{-1}(1/\sqrt{3})$ and $\theta_0 = \pi/4$. Setting

$$\begin{aligned}
 \theta &= \frac{\pi}{4} + \sqrt{\frac{2}{3}} \frac{x^1}{R}, & \alpha &= \alpha_0 - \frac{x^2}{R}, & \rho &= \frac{y}{R}, \\
 \varphi_1 &= \frac{\tilde{x}^3}{R} & \varphi_2 &= \frac{\tilde{x}^4}{R}, & t &= x^+ + \frac{x^-}{2R}, & \psi &= \frac{x^-}{2R} - x^+,
 \end{aligned}$$

redefining

$$x^3 = \sqrt{\frac{2}{3 + \hat{\gamma}^2}} \left(\tilde{x}^3 + \frac{1}{2} \tilde{x}^4 \right), \quad x^4 = \sqrt{\frac{3}{2(3 + \hat{\gamma}^2)}} \tilde{x}^4,$$

and taking the $R \rightarrow \infty$ limit gives the pp-wave metric

$$\begin{aligned}
 ds^2 &= -2dx^+ dx^- - \left(\sum_{a=5}^8 (x^a)^2 + \frac{4\hat{\gamma}^2}{3 + \hat{\gamma}^2} ((x^1)^2 + (x^2)^2) \right) (dx^+)^2 + \sum_{a=5}^8 (dx^a)^2 \\
 &+ \sum_{i=1}^4 (dx^i)^2 + \frac{4\sqrt{3}}{\sqrt{3 + \hat{\gamma}^2}} (x^1 dx^3 + x^2 dx^4) dx^+.
 \end{aligned} \tag{3.6}$$

In this same limit, the remaining fields in the IIB multiplet are

$$\begin{aligned}
 B_{(2)} &= \frac{\hat{\gamma}}{\sqrt{3}} dx^3 \wedge dx^4 + \frac{2\hat{\gamma}}{\sqrt{3 + \hat{\gamma}^2}} dx^+ \wedge (x^1 dx^4 - x^2 dx^3) , \\
 C_{(2)} &= \frac{2\hat{\gamma}}{\sqrt{3}g_s} dx^+ \wedge (x^2 dx^1 - x^1 dx^2) , \\
 F_{(5)} &= \frac{4}{g_s} dx^+ \wedge (dx^1 \wedge dx^2 \wedge dx^3 \wedge dx^4 + dx^5 \wedge dx^6 \wedge dx^7 \wedge dx^8) , \\
 e^{2\phi} &= \frac{1}{1 + \hat{\gamma}^2} e^{2\phi_0} .
 \end{aligned}
 \tag{3.7}$$

It is immediately evident from the non-vanishing RR 2-form and constant dilaton that this pp-wave differs rather nontrivially from the former pp-wave. Certainly, since the 3-brane couples to the RR 2-form, it might already be expected that the physics of giant gravitons is markedly more sophisticated on this background. Closed strings in this background, their supersymmetries and dual gauge theory operators were first studied in [35, 36]. There, it was noticed independently that a change of coordinates from x^- to $x^- + \sqrt{3/(3 + \hat{\gamma}^2)}(x^1 x^3 + x^2 x^4)$ brings the (J, J, J) pp-wave metric into the homogeneous plane wave form [34]

$$\begin{aligned}
 ds^2 &= -2dx^+ dx^- - \left[\sum_{a=5}^8 (x^a)^2 + \frac{4\hat{\gamma}^2}{3 + \hat{\gamma}^2} ((x^1)^2 + (x^2)^2) \right] (dx^+)^2 \\
 &\quad + \sum_{i=1}^8 (dx^i)^2 + 2\sqrt{\frac{3}{3 + \hat{\gamma}^2}} (x^1 dx^3 - x^3 dx^1 + x^2 dx^4 - x^4 dx^2) dx^+ ,
 \end{aligned}
 \tag{3.8}$$

after which it is a short, if somewhat technical, task to quantize the string sigma model and compute the full closed string spectrum which, in lightcone gauge and units of $2\pi\alpha' = 1$ reads

$$\omega_n = 1 \pm \sqrt{1 + 4n^2} .
 \tag{3.9}$$

Crucially, the closed string spectrum is completely independent of the deformation. Evidently the deformation of the geometry is exactly compensated for by the now non-vanishing B -field. Added support for this conclusion is given by a computation of the spectrum of anomalous dimensions of the corresponding operators in the dual $\mathcal{N} = 1$ gauge theory. The operators in question were proposed in [35] to be of the form

$$\mathcal{O}_{(p)}^{\hat{\gamma}} = \sum_{n,m} \text{tr} \left(\mathcal{O}_{n,m}^{\hat{\gamma}} \right) \delta_{m-n,p} ,
 \tag{3.10}$$

where $p = 0, 1, \dots, J$ and the $\mathcal{O}_{n,m}^{\hat{\gamma}}$ are built out of the complex Higgs's Φ^1, Φ^2 and Φ^3 in the gauge theory by taking one of the Φ 's to define a "background" lattice and inserting the other two as impurities hopped to the n - and m -th positions respectively. The matrix of anomalous dimensions of the $\mathcal{O}_{(p)}^{\hat{\gamma}}$ was then numerically checked to be identical, in the large J limit, to that of the corresponding undeformed operators [35]. This is in precise agreement with the string theory prediction.

One final point that deserves mentioning about this particular pp-wave geometry concerns its supersymmetry content. Our starting point, the type IIB $AdS_5 \times S^5$ background is maximally supersymmetric in 10–dimensions. It’s pp-wave limit is, famously, also maximally supersymmetric and preserves all 32 of the supersymmetries. The Lunin-Maldacena $\hat{\gamma}$ –deformation, of course, breaks a number of these. Exactly how many, can be computed by solving the Killing spinor equations for the backgrounds. Fortunately this rather technical exercise can be circumvented by using the fact that the $\hat{\gamma}$ –deformation and Penrose limits commute [30, 36]. In other words, starting from $AdS_5 \times S^5$, Penrose limits may be taken along $(J, 0, 0)$ and (J, J, J) orbits to get the respective pp-waves discussed in the previous section. An appropriate set of coordinates is then selected and compactified on a torus and the $SL(2, \mathbb{R})$ action applied along this torus. Finally, decompactifying the torus directions gives the $(J, 0, 0)$ and (J, J, J) deformed pp-waves respectively. Of the 32 Killing spinors associated to the original pp-wave backgrounds, only those that are independent of the torus coordinates will survive the $SL(2, \mathbb{R})$ transformation. For the $(J, 0, 0)$ deformed pp-wave, these include the standard 16 supersymmetries generically preserved by pp-waves plus an additional 8 so-called supernumerary supersymmetries giving a total of 24 altogether [31]. On the other hand, while the (J, J, J) deformed pp-wave preserves the same standard 16 Killing spinors, only 4 of the remaining 16 survive the $SL(2, \mathbb{R})$ transformation as supernumerary supersymmetries. This gives a total of 20 for this background.

4. Giants on the deformed pp-wave

Having developed the background on which we will work, we now proceed to the main focus of this work, namely the existence and stability of giant gravitons on these geometries. We begin with the deformed $(J, 0, 0)$ pp-wave of [30, 31] and then proceed to the more difficult case of the pp-wave constructed around the $J_{\phi_1} = J_{\phi_2} = J_{\phi_3}$ geodesic. The physics of the problem here consists of a $D3$ -brane wrapping the S^3 in the deformed 5-sphere and orbiting with fixed angular momentum along some great circle on S^5_γ . The Penrose limit in both cases is *along* this orbital direction - the ϕ_1 equatorial circle in the former case and the ψ orbit in the latter. Consequently, the worldvolume of the giant is of spherical topology (a $(+, 3, 0)$ brane in the notation of [26]). Had we boosted along a different direction, the brane worldvolume would have included both lightcone directions x^+ and x^- and two transverse spatial directions. We do not know whether these D –brane configurations are supported on this background and leave this for future consideration.

4.1 $(J, 0, 0)$, $(0, J, 0)$ and $(0, 0, J)$ orbits

On this class of geodesics, it will suffice to consider a $D3$ –brane moving along the ϕ_1 equatorial circle with fixed angular momentum and then take the Penrose limit along this same orbit. Consequently, we will take as an ansatz for the giant graviton solution,

$$\phi_1 = \omega t; \quad \sigma^1 = \theta; \quad \sigma^2 = \phi_2; \quad \sigma^3 = \phi_3. \tag{4.1}$$




Figure 1: The giant graviton lightcone Hamiltonian on the $(J, 0, 0)$ pp-wave exhibits two degenerate minima corresponding to the point and giant gravitons respectively.

with static gauge specified by

$$\begin{aligned}
 X^+ &= \lambda\tau; & X^- &= \mu\tau; \\
 X^1 &= r \cos \theta \cos \phi_2; & X^2 &= r \cos \theta \sin \phi_2; \\
 X^3 &= r \sin \theta \cos \phi_3; & X^4 &= r \sin \theta \sin \phi_3.
 \end{aligned}
 \tag{4.2}$$

The lightcone gauge constraint $X^+ = \tau$ is enforced by evaluating all expressions at $\lambda = 1$. Clearly, from the form of the deformed metric, any excitations in the AdS direction are unchanged by the deformation. Consequently, the background also supports AdS giants (i.e. those that are blown up along the S^3 of the AdS_5 part of the geometry). While these AdS giant solutions themselves are not affected by the deformation, their spectra of small fluctuations will be. However, we leave the study of these solutions for future work. For now, let's turn off any AdS fields;

$$X^a = 0 \quad a = 5, 6, 7, 8.$$

In this pp-wave limit, since the dilaton remains constant up to an overall factor of G and $G \rightarrow 1$, the $D3$ -brane action in this geometry is simply¹⁰

$$S = -\frac{T_3}{g_s} \int d^4\sigma \sqrt{-\det(P[g - B])} + T_3 \int P[C_{(4)}] \tag{4.3}$$

where, as usual, $P[X]$ denotes the pullback of the spacetime field X to the brane worldvolume and $g_s = \exp(\phi_0)$ is the 10-dimensional string coupling. In what follows, it will prove

¹⁰We thank M. Pirrone for pointing out that the supergravity conventions in [27] imply that the modified field strength $\mathcal{F} = F - B$. With the worldvolume gauge field turned off, this means that the NS- B -field enters the the Born-Infeld action as $\sqrt{-\det(P[g - B])}$ and the Chern-Simons term as $C_4 - C_2 \wedge B$.

more useful to rewrite the NS B -field in cartesian coordinates on the brane worldvolume. This is easily seen to be

$$B_{(2)} = \hat{\gamma} dX^+ \wedge (X^1 dX^2 - X^2 dX^1 + X^4 dX^3 - X^3 dX^4).$$

Focussing first on the Born-Infeld part of the action, we first define the pullback

$$D_{\mu\nu} = P[g - B]_{\mu\nu} = X_\mu^\alpha X_\nu^\beta g_{\alpha\beta} - X_\mu^\alpha X_\nu^\beta B_{\alpha\beta}, \quad (4.4)$$

where α, β, \dots are spacetime indices and $\mu\nu, \dots$ worldvolume ones. With a little tedious algebra, the components of $D_{\mu\nu}$ are readily computed as

$$D_{\mu\nu} = \begin{bmatrix} -2\lambda\mu - \lambda^2(1 + \hat{\gamma}^2)r^2 & 0 & -\hat{\gamma}\lambda r^2 \cos^2 \theta & \hat{\gamma}\lambda r^2 \sin^2 \theta \\ 0 & r^2 & 0 & 0 \\ \hat{\gamma}\lambda r^2 \cos^2 \theta & 0 & r^2 \cos^2 \theta & 0 \\ -\hat{\gamma}\lambda r^2 \sin^2 \theta & 0 & 0 & r^2 \sin^2 \theta \end{bmatrix}.$$

Notice that all off-diagonal contributions to $D_{\mu\nu}$ come from the deformation. Consequently, turning off $\hat{\gamma}$ manifestly reduces $D_{\mu\nu}$ to that of the undeformed pp-wave giant graviton. Computing the determinant of this matrix, we find that

$$\det D_{\mu\nu} = -r^6 \sin^2 \theta \cos^2 \theta (2\lambda\mu + \lambda^2 r^2). \quad (4.5)$$

On substituting this into the Born-Infeld action, the corresponding Lagrangian is read off as

$$L_{DBI} = -\frac{T_3 \tilde{\Omega}_3}{g_s} r^3 \sqrt{2\lambda\mu + \lambda^2 r^2}, \quad (4.6)$$

where $\tilde{\Omega}_3 = \int d\theta d\phi_2 d\phi_3 \sin \theta \cos \theta = 2\pi^2$ is the volume of the transverse 3-sphere that the giant expands into. That the volume element that arises in the DBI action is the standard one on the round sphere is a good consistency check of our spherical ansatz. We will return to this point a little later in the construction of giant gravitons moving along (J, J, J) orbits. Returning to our present construction, it remains to compute the Chern-Simons contribution to the full action. Since the only RR field that survives this Penrose limit is $C_{(4)}$, this is particularly easy

$$\begin{aligned} S_{CS} &= T_3 \int C_{(4)} = \frac{T_3}{g_s} \int r^4 \sin \theta \cos \theta dX^+ \wedge d\theta \wedge d\phi_2 \wedge d\phi_3 \\ &= \frac{T_3 \lambda}{g_s} \int d\tau \int d\theta d\phi_2 d\phi_3 r^4 \sin \theta \cos \theta. \end{aligned}$$

The corresponding Chern-Simons Lagrangian is

$$L_{CS} = M \lambda r^4, \quad (4.7)$$

where we have taken the liberty of defining $M \equiv T_3 \tilde{\Omega}_3 / g_s$. Putting this together then, gives the full 3-brane Lagrangian

$$L = -M \left[r^3 \sqrt{2\lambda\mu + \lambda^2 r^2} - \lambda r^4 \right]. \quad (4.8)$$

Notice that this Lagrangian is *independent of the deformation parameter*. This is not unlike what happens to closed strings quantized on the pp-wave of [35, 36]. Recall that there, the Lunin-Maldacena deformation had the effect of turning on an NS B -field as well as deforming the geometry and both these contributions exactly cancel so that the closed string spectrum is $\hat{\gamma}$ -independent. Of course, the deformation affects more than just the NS sector of the type IIB supergravity; it also turns on various form fields of the RR sector. Since strings couple only to the geometry and the B -field none of these form fields mattered in the study of the closed string spectrum. The 3-brane, on the other hand, also couples to the 2- and 4-form fields $C_{(2)}$ and $C_{(4)}$ respectively. However, while both $C_{(2)}$ and $C_{(4)}$ are affected by the deformation, it is only $C_{(4)}$ that survives the $(J, 0, 0)$ Penrose limit. It is hardly surprising then, that the Lagrangian (4.8) is independent of the deformation parameter. Importantly, this is in agreement with [39] where it was reported that giant gravitons with angular momentum along just one of the equatorial directions of the deformed 5-sphere not only survive the non-supersymmetric multi-parameter Frolov-Roiban-Tseytlin deformation (and by restriction also the single-parameter Lunin-Maldacena deformation) but that they are also independent of the deformation.

Returning to the construction of the giant graviton, the lightcone momentum of the 3-brane is

$$p^+ = -\frac{\delta L}{\delta \mu} = \frac{M\lambda r^3}{\sqrt{2\lambda\mu + \lambda^2 r^2}}, \tag{4.9}$$

with an associated lightcone Hamiltonian

$$p^- = H_{lc} = -\frac{\delta L}{\delta \lambda} = \frac{Mr^3(\mu + \lambda r^2)}{\sqrt{2\lambda\mu + \lambda^2 r^2}} - Mr^4.$$

This can be rewritten in terms of the lightcone momentum by solving the expression for the latter for μ and substituting into that for p^- . Consequently,

$$H_{lc} = \frac{M^2}{2p^+} r^6 + \frac{p^+}{2} r^2 - Mr^4. \tag{4.10}$$

As a function of the 3-brane radius, this Hamiltonian is extremized at

$$r = \begin{cases} 0 \\ \sqrt{\frac{p^+}{3M}} \\ \sqrt{\frac{p^+}{M}} \end{cases}. \tag{4.11}$$

As in the undeformed pp-wave, $r = 0$ is the point graviton while $R_G \equiv \sqrt{p^+/M}$ is the radius of the giant graviton. A plot of the lightcone Hamiltonian, H_{lc} , in units of $M = p^+ = 1$ is given in figure 1. With no way to energetically distinguish between the point and giant gravitons on this background, we expect that the giant solution should be stable. This expectation does, however, need to be verified in the spectrum of small fluctuations about the giant. We do so in the next section.

4.2 (J, J, J) orbits

The second of the two Penrose limits that result in BPS backgrounds - specifically that taken about the (J, J, J) orbit - is a little more difficult to treat. The IIB multiplet in this background is given by (3.6) and (3.7). The construction of giant gravitons in this background will be facilitated by choosing a slightly different gauge in which $x^- \rightarrow x^- + \sqrt{\frac{3}{3+\hat{\gamma}^2}}(x^1x^3 + x^2x^4)$. With this change of coordinates,

$$\begin{aligned}
 ds^2 &= -2dx^+dx^- - \left(\sum_{a=5}^8 (x^a)^2 + 4\frac{\hat{\gamma}^2}{3+\hat{\gamma}^2} \sum_{i=1}^2 (x^i)^2 \right) (dx^+)^2 + \sum_{a=5}^8 (dx^a)^2 \\
 &\quad + \sum_{i=1}^4 (dx^i)^2 + 2\sqrt{\frac{3}{3+\hat{\gamma}^2}} (x^1dx^3 - x^3dx^1 + x^2dx^4 - x^4dx^2), \\
 B_{(2)} &= \sqrt{\frac{\hat{\gamma}^2}{3}} dx^3 \wedge dx^4 + 2\sqrt{\frac{\hat{\gamma}^2}{3+\hat{\gamma}^2}} (x^1dx^4 - x^2dx^3), \\
 C_{(4)} - C_{(2)} \wedge B_{(2)} &= \frac{1}{3!} \frac{(3+\hat{\gamma}^2)}{3g_s} \epsilon_{ijkl} x^i dx^j \wedge dx^k \wedge dx^l, \\
 e^\phi &= \frac{3}{3+\hat{\gamma}^2} g_s.
 \end{aligned} \tag{4.12}$$

Note, in particular that the RR 2-form is no longer vanishing and so its coupling to the $D3$ -brane via the $C_{(2)} \wedge B_{(2)}$ Chern-Simons term is no longer trivial. Since the extended structure of the giant graviton is essentially supported by the RR flux, any change in the flux is likely to change the size and/or shape of the giant. Similar studies of pp-wave giant gravitons were carried out in [28, 29] where it was reported that when a constant transverse B -field is turned on in the background, giant gravitons respond by a deformation of their shape. The situation in this background is markedly different though. Here, in addition to a non-vanishing $C_{(2)}$, the NS B -field induced by the $\hat{\gamma}$ -deformation is certainly not constant. Proceeding with the construction then, our ansatz for the giant graviton on this background is

$$\psi = \omega t, \quad \sigma^1 = \theta, \quad \sigma^2 = \phi_2, \quad \sigma^3 = \phi_3. \tag{4.13}$$

Now, as promised, building on the lessons learnt from the undeformed pp-wave in magnetic coordinates, we parameterize the $D3$ -brane worldvolume by

$$\begin{aligned}
 X^+ &= \lambda\tau, & X^- &= \mu\tau; \\
 X^1 &= r \cos \theta \cos \left(\phi_2 - \sqrt{\frac{3}{3+\hat{\gamma}^2}} X^+ \right), & X^2 &= r \sin \theta \cos \left(\phi_3 - \sqrt{\frac{3}{3+\hat{\gamma}^2}} X^+ \right); \\
 X^3 &= r \cos \theta \sin \left(\phi_2 - \sqrt{\frac{3}{3+\hat{\gamma}^2}} X^+ \right), & X^4 &= r \sin \theta \sin \left(\phi_3 - \sqrt{\frac{3}{3+\hat{\gamma}^2}} X^+ \right).
 \end{aligned}$$

Note that, as in the undeformed pp-wave, this parameterization of the giant worldvolume is manifestly time-dependent. As in that case, we could perform a coordinate rotation to static Rosen coordinates but this comes at a cost of a significantly more cumbersome action

for the $D3$ -brane. Instead, we choose to work in the Brinkmann coordinates above, where at least the physics of the problem is slightly more transparent. As before, we denote the pullback of the combination of the metric and NS 2-form by $D_{\mu\nu} = P[g - B_{(2)}]_{\mu\nu}$. The time components of $D_{\mu\nu}$ are then easily computed as

$$\begin{aligned}
 D_{\tau\tau} &= -2\lambda\mu - \lambda^2 \frac{3}{3 + \hat{\gamma}^2} \left(r^2 + 4\frac{\hat{\gamma}}{3} (\cos^2 \theta \cos^2 \phi_2 + \sin^2 \theta \cos^2 \phi_3) \right), \\
 D_{\tau\theta} &= -D_{\theta\tau} = \lambda \sqrt{\frac{\hat{\gamma}^2}{3 + \hat{\gamma}^2}} r^2 (\cos^2 \theta \cos \phi_2 \sin \phi_3 + \sin^2 \theta \sin \phi_2 \cos \phi_3), \\
 D_{\tau\phi_2} &= -D_{\phi_2\tau} = -\lambda \sqrt{\frac{\hat{\gamma}^2}{3 + \hat{\gamma}^2}} r^2 \cos \theta \sin \theta \cos \phi_2 \cos \phi_3, \\
 D_{\tau\phi_3} &= -D_{\phi_3\tau} = \lambda \sqrt{\frac{\hat{\gamma}^2}{3 + \hat{\gamma}^2}} r^2 \cos \theta \sin \theta \cos \phi_2 \cos \phi_3,
 \end{aligned}$$

while its spatial components are

$$D_{rs} = \begin{bmatrix} r^2 & -\frac{\hat{\gamma}}{\sqrt{3}} r^2 \cos^2 \theta \cos \phi_2 \sin \phi_3 & -\frac{\hat{\gamma}}{\sqrt{3}} r^2 \sin^2 \theta \sin \phi_2 \cos \phi_3 \\ \frac{\hat{\gamma}}{\sqrt{3}} r^2 \cos^2 \theta \cos \phi_2 \sin \phi_3 & r^2 \cos^2 \theta & \frac{\hat{\gamma}}{\sqrt{3}} r^2 \cos \theta \sin \theta \cos \phi_2 \cos \phi_3 \\ \frac{\hat{\gamma}}{\sqrt{3}} r^2 \sin^2 \theta \sin \phi_2 \cos \phi_3 & -\frac{\hat{\gamma}}{\sqrt{3}} r^2 \cos \theta \sin \theta \cos \phi_2 \cos \phi_3 & r^2 \sin^2 \theta \end{bmatrix},$$

where $r, s = \theta, \phi_2, \phi_3$. For ease of notation, we have set $\tau = 0$ above. The time dependence of the solution can be restored by replacing ϕ_i with $\phi_i - \sqrt{\frac{3}{3 + \hat{\gamma}^2}} X^+$. Then, with $Y^i \equiv X^i / r$, i.e. the embedding coordinates of a unit 3-sphere and a little algebra, the determinant of the pullback can be expressed as

$$\begin{aligned}
 -\det D_{\mu\nu} &= r^6 \cos^2 \theta \sin^2 \theta \left[2\lambda\mu \left(1 + \frac{\hat{\gamma}^2}{3} ((Y^1)^2 + (Y^2)^2) \right) \right. \\
 &\quad \left. + \lambda^2 r^2 \frac{3}{3 + \hat{\gamma}^2} \left(1 + 2\frac{\hat{\gamma}^2}{3} ((Y^1)^2 + (Y^2)^2) \right)^2 + \lambda^2 \frac{\hat{\gamma}^2}{3 + \hat{\gamma}^2} r^2 (Y^1 Y^4 - Y^2 Y^3)^2 \right].
 \end{aligned}$$

The manifest dependence of this determinant on the Y^i - compared to, say the corresponding expression in the $(J, 0, 0)$ pp-wave - is strong evidence that the giant in this background no longer retains its round 3-sphere shape. Evidently, our ansatz for the giant is unable to capture the full shape deformation of the giant and requires some refinement. This is a nontrivial exercise best left for future work. Nevertheless, we can still make some progress - albeit limited - when the Lunin-Maldacena deformation is small. Toward this end, we will take $\hat{\gamma}$ small and expand all $\hat{\gamma}$ -dependent terms to lowest nontrivial order in $\hat{\gamma}$. Expanding to $\mathcal{O}(\hat{\gamma}^2)$ then, the $D3$ -brane Lagrangian

$$L = -T_3 e^{-\phi} \int d^3\sigma \sqrt{-\det D_{\mu\nu}} + T_3 \int P [C_{(4)} - C_{(2)} \wedge B_{(2)}],$$

becomes

$$\begin{aligned}
 L &= -\frac{T_3}{g_s} \left(1 + \frac{\hat{\gamma}^2}{3}\right) r^3 \sqrt{2\lambda\mu + \lambda^2 r^2} \int d\theta d\phi_2 d\phi_3 \cos\theta \sin\theta \\
 &\times \sqrt{1 + \frac{\hat{\gamma}^2}{3(2\lambda\mu + \lambda^2 r^2)} \left[(2\lambda\mu + 4\lambda^2 r^2)((Y^1)^2 + (Y^2)^2) + \lambda^2 r^2 (Y^1 Y^4 - Y^2 Y^3)^2 - \lambda^2 r^2 \right]} \\
 &+ 2\pi^2 \lambda \frac{T_3}{g_s} \left(1 + \frac{\hat{\gamma}^2}{3}\right) r^4 + \mathcal{O}(\hat{\gamma}^4),
 \end{aligned}$$

or, using $M = 2\pi^2 T_3 g_s^{-1}$ and the integrals

$$\int d\theta d\phi_2 d\phi_3 \cos\theta \sin\theta ((Y^1)^2 + (Y^2)^2) = \pi^2,$$

and

$$\begin{aligned}
 &\int d\theta d\phi_2 d\phi_3 \cos\theta \sin\theta (Y^1 Y^4 - Y^2 Y^3)^2 = \frac{\pi^2}{6}, \\
 L &= \lambda M r^4 - M r^3 \sqrt{2\lambda\mu + \lambda^2 r^2} + \frac{\hat{\gamma}^2}{3} \left[\lambda M r^4 - M r^3 \frac{\frac{5}{2}\lambda\mu + \frac{37}{24}\lambda^2 r^2}{\sqrt{2\lambda\mu + \lambda^2 r^2}} \right] + \mathcal{O}(\hat{\gamma}^4).
 \end{aligned}$$

The corresponding lightcone momentum

$$\begin{aligned}
 p^+ &= -p_- = -\frac{\delta L}{\delta \mu} \\
 &= \frac{\lambda M r^3}{\sqrt{2\lambda\mu + \lambda^2 r^2}} + \frac{\hat{\gamma}^2}{3} \frac{\lambda M r^3}{(2\lambda\mu + \lambda^2 r^2)^{3/2}} \left[\frac{5}{2}\lambda\mu + \frac{23}{24}\lambda^2 r^2 \right] + \mathcal{O}(\hat{\gamma}^4). \quad (4.14)
 \end{aligned}$$

To compute the lightcone Hamiltonian, we need to eliminate any explicit μ -dependence in the expression for p^- . In anticipation, solving (4.14) for μ to order $\hat{\gamma}^2$ we get,

$$\lambda\mu \sim \frac{1}{2} \left(\frac{\lambda M r^3}{p^+} \right)^2 - \frac{1}{2} \lambda^2 r^2 + \frac{\hat{\gamma}^2}{3} \left(\frac{5}{4} \left(\frac{\lambda M r^3}{p^+} \right)^2 - \frac{7}{24} \lambda^2 r^2 \right). \quad (4.15)$$

Finally, to lowest nontrivial order in $\hat{\gamma}$, the lightcone Hamiltonian

$$p^- \sim \frac{r^2}{2p^+} \left[M^2 r^4 \left(1 + \frac{5}{6}\hat{\gamma}^2\right) - 2M p^+ r^2 \left(1 + \frac{\hat{\gamma}^2}{3}\right) + (p^+)^2 \left(1 + \frac{7}{36}\hat{\gamma}^2\right) \right]. \quad (4.16)$$

As a rough check, note that this reduces to the correct undeformed Hamiltonian in the $\hat{\gamma} \rightarrow 0$ limit. In fact, writing it as

$$H_{lc} \sim \left[\frac{M^2}{2p^+} r^6 + \frac{p^+}{2} r^2 - M r^4 \right] + \hat{\gamma}^2 \left[\frac{5M^2 r^6}{12p^+} - \frac{M r^4}{3} + \frac{7(p^+)^2 r^2}{72} \right],$$

it is clear that, at least to $\mathcal{O}(\hat{\gamma}^2)$, the Hamiltonian is positive semi-definite, vanishing only at $r = 0$. We have checked numerically that this property persists even up to order $\hat{\gamma}^4$. Figure 2 is a plot of the lightcone Hamiltonian (4.16) as a function of r .




Figure 2: The deformation lifts the degenerate minima in the (J, J, J) pp-wave.

Notice now that the once degenerate vacuum at finite radius is lifted by the deformation. Consequently, the energy of the giant is greater than that of the point graviton on this background and the giant graviton is no longer a stable state in the theory. Indeed, beyond a certain critical value of the deformation parameter $\hat{\gamma} \sim 1$ it seems the second minimum vanishes and the giant is no longer a solution even. This is in good agreement with the results of [29] and, especially [22]. Here too, as the deformation is increased from zero, the $C_{(2)} \wedge B_{(2)}$ contribution to the Chern-Simons term grows faster than the four form flux term and eventually dominates. However it is important to bear in mind that our analysis is carried out perturbatively in $\hat{\gamma}$ and any conclusions drawn at large $\hat{\gamma}$ need to be done with care. So, although numerical results up to $\mathcal{O}(\hat{\gamma}^4)$ seem to support the reported behaviour of the $D3$ -brane lightcone Hamiltonian, there is probably not too much to be gained from estimating the critical value of $\hat{\gamma}$ at which the second minimum vanishes - at least until a more detailed study of the deformation of the giant on this pp-wave is completed.

5. Small fluctuations and stability of giants

The spectrum of small fluctuations about a giant graviton background encodes many interesting properties of the equilibrium configuration [9]. For instance, since zero energy modes in the vibration spectrum correspond to a family of solutions each having the same energy and angular momentum, the study of these modes is important in determining the ability of the giant ansatz in capturing all the classical BPS states. More relevantly for us, any perturbative instability in the giant configuration will appear in the fluctuation spectrum as a tachyonic mode. In this section we will systematically investigate the spectrum of small fluctuations about the equilibrium configurations of the pp-wave giant gravitons beginning first with the giant on the maximally supersymmetric type IIB pp-wave and

then proceeding to the $\hat{\gamma}$ -deformed $(J, 0, 0)$ background. While the vibration spectrum of giants on the undeformed pp-wave is not unknown [28, 29], we will rederive it here using the Lagrangian methods of [9] primarily by way of review and to establish our notation but also to facilitate comparison with the deformed giant spectrum. Essentially the Hamiltonian computation proceeds by fixing all the conserved quantities in the problem, diagonalising the Hamiltonian¹¹ and extracting the classical oscillation frequencies. The problem is that any mode for which the conserved quantities are not held fixed will necessarily be excluded from the spectrum. We will therefore proceed with the Lagrangian analysis of [9].

5.1 Undeformed giants

Our starting point here is the giant graviton configuration on the pp-wave (2.1) supported by the 5-form flux (2.3). This equilibrium configuration is specified by

$$\begin{aligned} X^+ &= \lambda\tau, & X^- &= \nu\tau, \\ X^1 &= r_0 \cos \theta_1 \cos \theta_2, & X^2 &= r_0 \sin \theta_1 \cos \theta_3, \\ X^3 &= r_0 \cos \theta_1 \sin \theta_2, & X^4 &= r_0 \sin \theta_1 \sin \theta_3. \end{aligned} \tag{5.1}$$

Correspondingly, the perturbed configuration has coordinates¹²

$$\begin{aligned} X^+ &= \tau, & X^- &= \nu\tau + \epsilon\delta x^-, \\ r &= r_0 + \epsilon\delta r, & X^a &= \epsilon\delta x^a, \end{aligned} \tag{5.2}$$

and $\sigma^i = \theta_i \in S^3$. This ansatz is then substituted into the $D3$ -brane action

$$S = -\frac{T_3}{g_s} \int d\tau d^3\sigma \sqrt{-\det(P[g])} + T_3 \int P[C_{(4)}],$$

and, as we are interested primarily in the fluctuation spectrum, expanded up to quadratic order in the perturbation parameter, ϵ . It is then quite straightforward to compute the induced metric on the brane worldvolume. With $D_{\mu\nu} \equiv P[g]_{\mu\nu}$, we find

$$\begin{aligned} D_{0i} &= D_{i0} \approx -2\epsilon\partial_i\delta x^- + \epsilon^2 [(\partial_\tau\delta r)(\partial_i\delta r) + (\partial_\tau\delta x^a)(\partial_i\delta x^a)], \\ D_{ij} &\approx r_0^2 g_{ij} + 2\epsilon r_0 \delta r g_{ij} + \epsilon^2 [\delta r^2 g_{ij} + (\partial_i\delta r)(\partial_j\delta r) + (\partial_i\delta x^a)(\partial_j\delta x^a)], \end{aligned}$$

and

$$\begin{aligned} D_{00} &\approx -(2\nu + \mu^2 r_0^2) \left[1 + \epsilon \left(\frac{2\partial_\tau\delta x^- + 2\mu^2 r_0 \delta r}{2\nu + \mu^2 r_0^2} \right) \right. \\ &\quad \left. + \epsilon^2 \left(\frac{\mu^2 \delta r^2 + \mu^2 (\delta x^a)^2 - (\partial_\tau\delta r)^2 - (\partial_\tau\delta x^a)^2}{2\nu + \mu^2 r_0^2} \right) \right]. \end{aligned}$$

A judicious application of Stokes' theorem on the $D3$ -brane worldvolume allows for the Chern-Simons contribution to the action to easily be written down as

$$S_{CS} = \frac{T_3}{g_s} \mu (r_0 + \epsilon\delta r)^4 \int d\tau d^3\sigma \sqrt{|g_{ij}|},$$

¹¹For small perturbations the Hamiltonian is a quadratic form in the coordinates and momenta representing the perturbation and may be diagonalised using standard methods for the diagonalisation of quadratic forms.

¹²Without loss of generality, λ can be set to unity.

Substituting this, together with the metric pullback terms above, into the worldvolume action, we can read off the term linear in ϵ as

$$\delta S = \epsilon \frac{T_3 \Omega_3}{g_s} \int d\tau \frac{r_0^2}{\sqrt{2\nu + \mu^2 r_0^2}} \left[\left(6\nu + 4\mu^2 r_0^2 - 4\mu r_0 \sqrt{2\nu + \mu^2 r_0^2} \right) \delta r + r_0 \frac{\partial \delta x^-}{\partial \tau} \right] \quad (5.3)$$

Since we vary with fixed boundary conditions (i.e. $\partial_\tau \delta x^-$ vanishes at the endpoints of τ) and the coefficient of $\partial_\tau \delta x^-$ is constant, this term gives no contribution to the first order variation of the action. In order for the radial perturbation to vanish, on the other hand, we must have $\nu = 0$. The other value of ν for which the δr coefficient vanishes, $\nu = -\frac{4}{9}\mu^2 r_0^2$, is a nonphysical solution and consequently discarded. Next we evaluate the perturbations second order in ϵ . With the choice of $\nu = 0$ above, these terms assemble as

$$\begin{aligned} \delta^2 S = & -\epsilon^2 \frac{T_3}{g_s} \mu r_0^3 \int d\tau d^3\sigma \sqrt{|g_{ij}|} \frac{\delta r}{2\mu^2 r_0^2} \left[\frac{\partial^2}{\partial \tau^2} - \mu^2 \frac{1}{\sqrt{|g_{kl}|}} \partial_i \left(\sqrt{|g_{kl}|} g^{ij} \partial_j \right) \right] \delta r \\ & - 2 \frac{\delta x^-}{2\mu^2 r_0^3} \frac{\partial \delta r}{\partial \tau} + 2 \frac{\delta r}{2\mu^2 r_0^3} \frac{\partial \delta x^-}{\partial \tau} \\ & + \frac{\delta x^-}{2\mu^4 r_0^4} \left[\frac{\partial^2}{\partial \tau^2} - \mu^2 \frac{1}{\sqrt{|g_{kl}|}} \partial_i \left(\sqrt{|g_{kl}|} g^{ij} \partial_j \right) \right] \delta x^- \\ & + \frac{\delta x^a}{2\mu^2 r_0^2} \left[\mu^2 + \frac{\partial^2}{\partial \tau^2} - \mu^2 \frac{1}{\sqrt{|g_{kl}|}} \partial_i \left(\sqrt{|g_{kl}|} g^{ij} \partial_j \right) \right] \delta x^a. \end{aligned} \quad (5.4)$$

In order to analyze the resulting vibration spectrum, the fluctuations are first decomposed into their respective Fourier components:

$$\delta x^\mu = \widetilde{\delta x^\mu} e^{-i\omega\tau} Y_l(\theta_1, \theta_2, \theta_3). \quad (5.5)$$

Here, the Y_l are 4-dimensional spherical harmonics satisfying

$$\frac{1}{\sqrt{|g_{mn}|}} \partial_i \left(\sqrt{|g_{mn}|} g^{ij} \partial_j \right) Y_l = -Q_l Y_l, \quad (5.6)$$

with the eigenvalues $Q_l = l(l+2)$. Substituting the Fourier expansion into (5.4) we see that the fluctuations in the X^a directions are separable. Their spectrum is easily determined to be

$$\omega_a^2 = \mu^2 (1 + Q_l). \quad (5.7)$$

The radial and X^- fluctuations are a bit more tricky. They are coupled, and satisfy the second order system

$$\begin{bmatrix} Q_l - \frac{\omega^2}{\mu^2} & \frac{-2i}{\mu^2 r_0} \omega \\ \frac{2i}{\mu^2 r_0} \omega & \frac{Q_l}{\mu^2 r_0^2} - \frac{\omega^2}{\mu^4 r_0^2} \end{bmatrix} \begin{bmatrix} \widetilde{\delta r} \\ \widetilde{\delta x^-} \end{bmatrix} = 0. \quad (5.8)$$

Solving this for ω leads to the spectrum

$$\omega_\pm^2 = \mu^2 \left(2 + Q_l \pm 2\sqrt{1 + Q_l} \right), \quad (5.9)$$

At this point some comments are in order. Most obvious in the above spectrum is its independence of the radius r_0 of the brane. This should be compared to the corresponding spectrum of sphere giants on $AdS_5 \times S^5$ where it was demonstrated in [9] that a suitable rescaling of the fluctuation fields completely scales out all r_0 dependence of the small fluctuation action. Secondly, note that as there are no complex frequencies, there are no tachyonic modes. Consequently, these giant gravitons on the pp-wave are stable. This is in perfect agreement with the results of [28, 25]. Moreover, there is a zero mode when $l = 0, Q_l = 0$ on the negative branch of (5.9). This corresponds to the statement that increasing the size of the giant (which relates to increasing the light-cone momentum) costs no energy, just as we expect.

5.2 Deformed giants

Moving on to the deformed case now, we focus on the giant gravitons on the $(J, 0, 0)$ pp-wave. As in the undeformed case above, we take a static, spherical ansatz for the giant, wrapping the deformed S^3 in the light cone gauge and allow fluctuations in the transverse directions. The ansatz for the perturbed giant is, as before

$$\begin{aligned} X^+ &= \tau, & X^- &= \nu\tau + \epsilon\delta x^-, \\ X^1 &= r \cos\theta \cos\phi_2, & X^2 &= r \cos\theta \sin\phi_2, \\ X^3 &= r \sin\theta \cos\phi_3, & X^4 &= r \sin\theta \sin\phi_3, \end{aligned} \tag{5.10}$$

where $r = r_0 + \epsilon\delta r$ and the AdS fluctuations are $X^a = \epsilon\delta x^a$. Expanded up to second order in ϵ the components of the pullback $D_{\mu\nu} = P[g - B_{(2)}]_{\mu\nu}$ appearing in the Born-Infeld contribution to the $D3$ -brane action,

$$S = -\frac{T_3}{g_s} \int d\tau d\Omega \sqrt{-\det D_{\mu\nu}} + T_3 \int P[C_{(4)} - C_{(2)} \wedge B_{(2)}] \tag{5.11}$$

are

$$\begin{aligned} D_{\tau\tau} &= -2\nu - (1 + \hat{\gamma}^2) r_0^2 + \epsilon [-2\partial_\tau\delta x^- - 2(1 + \hat{\gamma}^2) r_0\delta r] \\ &\quad + \epsilon^2 \left[-\sum_a (\delta x^a)^2 - (1 + \hat{\gamma}^2) \delta r^2 + \sum_{I=a+r} (\partial_\tau\delta x^I)^2 \right], \\ D_{\tau\theta} &= D_{\theta\tau} = -\epsilon\partial_\theta\delta x^- + \epsilon^2 \sum_{I=a+r} (\partial_\tau\delta x^I) (\partial_\theta\delta x^I), \\ D_{\tau\phi_2/\phi_2\tau} &= \pm\hat{\gamma}r_0^2 \cos^2\theta + \epsilon [-\partial_2\delta x^- \pm 2\hat{\gamma}r_0 \cos^2\theta\delta r] \\ &\quad + \epsilon^2 \left[\sum_{I=a+r} (\partial_\tau\delta x^I) (\partial_2\delta x^I) \pm \hat{\gamma} \cos^2\theta\delta r^2 \right], \\ D_{\tau\phi_3/\phi_3\tau} &= \mp\hat{\gamma}r_0^2 \sin^2\theta + \epsilon [-\partial_{\phi_3}\delta x^- \mp 2\hat{\gamma}r_0 \sin^2\theta\delta r] \\ &\quad + \epsilon^2 \left[\sum_{I=a+r} (\partial_\tau\delta x^I) (\partial_{\phi_3}\delta x^I) \mp \hat{\gamma} \sin^2\theta\delta r^2 \right], \\ D_{ij} &= r_0^2 g_{rs} + 2\epsilon r_0 g_{ij} \delta r + \epsilon^2 \left[g_{ij} \delta r^2 + \sum_{I=a+r} (\partial_i\delta x^I) (\partial_j\delta x^I) \right]. \end{aligned} \tag{5.12}$$

Here, $\sum_{I=a+r}$ is shorthand for a summation over both the AdS fluctuations, δx^a and radial perturbation δr and the undeformed spatial metric, g_{rs} is

$$g_{rs} = \begin{bmatrix} r^2 & & \\ & r^2 \cos^2 \theta & \\ & & r^2 \sin^2 \theta \end{bmatrix}.$$

Then, keeping terms second order and lower in ϵ , the determinant of $D_{\mu\nu}$ is

$$\begin{aligned} -\det D_{\mu\nu} = & r_0^6 \cos^2 \theta \sin^2 \theta (2\nu + r_0^2) + \epsilon (r_0^5 \cos^2 \theta \sin^2 \theta) \left[2r_0 \partial_\tau \delta x^- + (12\nu + 8r_0^2) \delta r \right] \\ & + \epsilon^2 (r_0^4 \cos^2 \theta \sin^2 \theta) \left[(30\nu + 28r_0^2) \delta r^2 + r_0^2 \sum_a (\delta x^a)^2 \right. \\ & + \sum_{I=a+r} \left((2\nu + r_0^2) \sum_s g^{ii} (\partial_i \delta x^I)^2 + \hat{\gamma}^2 r_0^2 (\partial_{\phi_2} \delta x^I - \partial_{\phi_3} \delta x^I)^2 - r_0^2 (\partial_\tau \delta x^I)^2 \right) \\ & \left. + 12 r_0 \delta r (\partial_\tau \delta x^-) + \sum_s g^{ii} (\partial_i \delta x^-)^2 \right]. \end{aligned} \quad (5.13)$$

Notice that the deformation has cancelled in the zeroth and first order fluctuations. This is just a manifestation, at the level of fluctuations, of the NS B -field exactly compensating for changes in the metric induced by the Lunin-Maldacena deformation. Interestingly though, this is not carried through to the higher order terms. At $\mathcal{O}(\epsilon^2)$ for example, the fluctuation spectrum exhibits a “magnetic field” like splitting due to the deformation. The Chern-Simons term on the other hand is trivially integrated over the sphere to give

$$S_{CS} = \frac{T_3}{g_s} \Omega_3 \int d\tau (r_0^4 + 4\epsilon r_0^3 \delta r + 6\epsilon^2 r_0^2 \delta r^2), \quad (5.14)$$

to second order in ϵ . Putting this all together, defining $m \equiv T_3/g_s$ and expanding the square root gives

$$\begin{aligned} S = & -m\Omega_3 r_0^3 \int d\tau \sqrt{2\nu + r_0^2} - r_0 \\ & - \epsilon \frac{m r_0^3}{\sqrt{2\nu + r_0^2}} \int d\Omega_3 d\tau \left[\partial_\tau \delta x^- + \frac{\delta r}{r_0} \left(6\nu + 4r_0^2 - 4r_0 \sqrt{2\nu + r_0^2} \right) \right] \\ & - \epsilon^2 \frac{m r_0}{2\sqrt{2\nu + r_0^2}} \int d\Omega_3 d\tau \left[\left(30\nu + 28r_0^2 - \frac{(6\nu + 4r_0^2)^2}{2\nu + r_0^2} - 12r_0 \sqrt{2\nu + r_0^2} \right) \delta r^2 \right. \\ & + \sum_{I=a+r} \left((2\nu + r_0^2) \sum_i g^{ii} (\partial_i \delta x^I)^2 + \hat{\gamma}^2 r_0^2 (\partial_{\phi_2} \delta x^I - \partial_{\phi_3} \delta x^I)^2 - r_0^2 (\partial_\tau \delta x^I)^2 \right) \\ & \left. + r_0^2 \sum_a (\delta x^a)^2 + 2 \left(\frac{6\nu + 2r_0^2}{2\nu + r_0^2} \right) r_0 \delta r (\partial_\tau \delta x^-) + \sum_i g^{ii} (\partial_i \delta x^-)^2 - \frac{r_0^2}{2\nu + r_0^2} (\partial_\tau \delta x^-)^2 \right]. \end{aligned}$$

The physical solution is found by minimizing this action. At first order in ϵ , the term $\partial_\tau \delta x^-$ vanishes since the endpoints in τ are held fixed. In addition, requiring that the remaining

term proportional to δr vanish leads to the constraint $\nu = 0$. There is, of course, a second solution, $\nu = -\frac{4}{9}r_0^2$ but this is a maximum of the action, so we disregard it.

Proceeding to the next order, to find the fluctuation spectrum, the fluctuations can be decomposed into eigenfunctions of the various differential operators in the second order action. Fluctuations with energy ω have time dependence $\sim e^{-i\omega\tau}$ and the spherical harmonics, $Y_l(\Omega_3)$ are eigenfunctions of

$$\frac{1}{\sqrt{|g_{kl}|}}\partial_i(\sqrt{|g_{kl}|}g^{ij}\partial_j)Y_l = -Q_l Y_l.$$

As observed above, the deformation introduces a splitting term which, after integrating by parts, is $\hat{\gamma}^2 r_0^2 (\partial_{\phi_2} - \partial_{\phi_3})^2$. This breaks some of the degeneracy of the spherical harmonics. Indeed, it is clear from its form that the eigenvalue of this operator is negative semi-definite. Consequently, the spherical harmonics are diagonalized as

$$\left(\frac{\partial}{\partial\phi_2} - \frac{\partial}{\partial\phi_3}\right)^2 Y_{l,\alpha} = -\alpha^2 Y_{l,\alpha}, \tag{5.15}$$

As in the undeformed case, vibrations of the giant in the AdS directions δx^a decouple from the rest of the fluctuations, and minimizing the second order action subject to the first order constraint $\nu = 0$ leads to the spectrum

$$\omega_a^2 = 1 + \hat{\gamma}^2 \alpha^2 + Q_l, \tag{5.16}$$

which is manifestly positive definite. The coupled radial and null fluctuations, δr and δx^- respectively, satisfy the linear system of equations

$$\begin{bmatrix} Q_l - \omega^2 + \hat{\gamma}^2 \alpha^2 & -2i\frac{\omega}{r_0} \\ 2i\frac{\omega}{r_0} & \frac{Q_l - \omega^2}{r_0^2} \end{bmatrix} \begin{bmatrix} \tilde{\delta r} \\ \tilde{\delta x^-} \end{bmatrix} = 0. \tag{5.17}$$

Solving this for ω yields the spectrum

$$\omega_{\pm}^2 = Q_l + \left(2 + \frac{1}{2}\hat{\gamma}^2 \alpha^2\right) \pm \sqrt{4Q_l + \left(2 + \frac{1}{2}\hat{\gamma}^2 \alpha^2\right)^2}. \tag{5.18}$$

Here, as in the case of the undeformed giant, the fluctuation spectrum is both manifestly independent of the size of the equilibrium configuration, r_0 and *dependent* on the deformation parameter. Since the latter enters into the spectrum as a positive contribution, it is easy to see that ω^2 is positive semi-definite and remains so irrespective of the deformation strength. Again, this lack of complex frequencies is a signal of the perturbative stability of these giants, exactly as was claimed in the previous section. Again, $\omega^2 = 0$ is one of the solutions of the $(\delta r, \delta x^-)$ system when $Q_l = 0$ reflecting the fact that the radius of the equilibrium configuration can be taken to have any value allowed by the pp-wave geometry.

6. Comments on the dual operators

In this section we make some brief comments on the gauge theory operators dual to pp-wave giants of both the deformed and undeformed kind and leave a more detailed study

of the dual giant operators and the open strings attached to them to future study [37]. Shortly after an observation by Balasubramanian *et.al.* [10] that the correct description of large single sphere-giant states in the dual $\mathcal{N} = 4$ conformal field theory are of the (sub)determinant form

$$\mathcal{O}_k = \frac{1}{k!} \epsilon_{i_1 \dots i_k a_1 \dots a_{N-k}}^{j_1 \dots j_k} \Phi_{j_1}^{i_1} \dots \Phi_{j_k}^{i_k}, \tag{6.1}$$

rather than single-trace operators of large R-charge, it was argued by Corley, Jevicki and Ramgoolam in [10] that, more generally, the operators in the $U(N)$ super Yang-Mills theory dual to giant gravitons are Schur polynomials in the Higgs fields of the SYM multiplet,

$$\chi_R(Z) = \frac{1}{n!} \sum_{\sigma \in S_n} \chi_R(\sigma) \text{tr}(\sigma Z). \tag{6.2}$$

Here, Z is one of three complex Higgs fields in the conformal field theory, $n \sim O(N)$ and

$$\text{tr}(\sigma Z) = \sum_{i_1 \dots i_n} Z_{i_{\sigma(1)}}^{i_1} Z_{i_{\sigma(2)}}^{i_2} \dots Z_{i_{\sigma(n)}}^{i_n}.$$


The Schur polynomials are labelled by n -box Young diagrams and the $\chi_R(\sigma)$ are characters of $\sigma \in S_n$ in the representation R . The extension to the $SU(N)$ gauge theory - and consequent disentanglement of bulk and boundary degrees of freedom in giant graviton dynamics - is non-trivial and may be found in [11]. It is not difficult to see that in the totally antisymmetric representation, the Schur polynomial reduces to the subdeterminant operator (6.1) - a single column Young diagram. Such column diagrams have a length bounded by N and so encode in a very natural way the momentum cutoff observed in the dynamics of sphere giants. Similarly, a giant graviton blown up along the AdS 3-sphere is naturally identified with the completely symmetric representation - a single row Young diagram whose length can grow without bound [10, 12]. Much of the power of the machinery developed in the first of [10] derives from the fact that the multipoint correlators of these Schur polynomial that encode much of the giant graviton dynamics can be computed *exactly*. The 2- and 3-point correlators, for example, are given by

$$\begin{aligned} \langle \chi_R(Z)(x_1) \chi_S(\bar{Z})(x_2) \rangle &= \delta_{RS} f_R \frac{1}{(x_1 - x_2)^{2n_R}}, \tag{6.3} \\ \langle \chi_{R_1}(Z)(x_1) \chi_{R_2}(Z)(x_2) \chi_{R_3}(\bar{Z})(x_3) \rangle &= g(R_1, R_2; R_3) f_{R_3} \frac{1}{\prod_{i=2}^3 (x_1 - x_i)^{2n_{R_i}}}, \end{aligned}$$

where, with the product running over the boxes of a Young diagram associated to the representation R , i labelling the rows and j the columns $f_R = \prod_{i,j} (N - i + j)$ and the factor $g(R_1, R_2; R_3)$ is a Littlewood-Richardson coefficient - the multiplicity with which the representations R_3 appears in the tensor product of representations R_1 and R_2 [38].

The Penrose limit of the gravity theory manifests in the gauge theory as the double scaling limit in which the $U(1)$ R-charge J and the number of colours N both go to infinity in such a way that the ratio J^2/N remains fixed [24]. This BMN limit of the giant operators in the undeformed $\mathcal{N} = 4$ gauge theory was first investigated in [25] and we

briefly summarize some of their results here. A sphere giant with large angular momentum J is dual to a Schur polynomial labelled by a single column young diagram with J rows.



With the technology developed in the second of [10] various extremal correlators obtained from considering the overlaps of combinations of Schur polynomials may be exactly computed. The normalized 3–point correlator, for example, is

$$\frac{\langle \chi_{R_{J_1}}(Z) \chi_{R_{J_2}}(Z) \chi_{R_J}(\bar{Z}) \rangle}{\|\chi_{R_{J_1}}(Z)\| \|\chi_{R_{J_2}}(Z)\| \|\chi_{R_J}(Z)\|} = \sqrt{\frac{(N - J_1)! (N - J_2)!}{(N - J)! N!}}. \tag{6.4}$$

Using Stirling’s approximation for $N!$ at large N and assuming that J^2/N is large, in the BMN limit

$$\frac{\langle \chi_{R_{J_1}}(Z) \chi_{R_{J_2}}(Z) \chi_{R_J}(\bar{Z}) \rangle}{\|\chi_{R_{J_1}}(Z)\| \|\chi_{R_{J_2}}(Z)\| \|\chi_{R_J}(Z)\|} \sim e^{-J_1 J_2 / 2N}. \tag{6.5}$$

Consequently, there is no mixing of multi-particle states and the single-particle giant graviton state dual to the Schur polynomial (6.2) is well defined. Similarly, the normalized overlap between a sphere giant with angular momentum J and a multi-particle Kaluza-Klein state

$$\frac{\langle (\chi_1(Z))^J \chi_J(\bar{Z}) \rangle}{\|(\chi_1(Z))^J\| \|\chi_J(Z)\|} = \sqrt{\frac{1}{N^J} \frac{N!}{J!(N - J)!}} \sim \sqrt{\frac{1}{J!}} e^{-J^2 / 2N}. \tag{6.6}$$

What about the giant gravitons in the deformed background studied above? The exactly marginal Leigh-Strassler deformation of the super Yang-Mills gauge theory [15], recall, can be realised as the Moyal-like deformation,

$$X * Y = e^{i\pi\hat{\gamma}(Q_X^1 Q_Y^2 - Q_Y^1 Q_X^2)} XY, \tag{6.7}$$

of the product of fields in the $\mathcal{N} = 4$ Lagrangian [30]. Here (Q_X^1, Q_Y^2) are the charges of the corresponding fields under the $U(1)_1 \times U(1)_2$ action. The Schur operator (6.2) dual to the giant graviton, however, is a holomorphic polynomial in Z so each factor in the product carries the same charge under $U(1)_1 \times U(1)_2$. Evidently then, the operator that we expect to be dual to a giant graviton on the deformed pp-wave obtained by taking a

Penrose limit about a $(J, 0, 0)$ geodesic, $\chi_R(Z)$, is blind to the $\hat{\gamma}$ -deformation. This is certainly in agreement with the supergravity analysis of section 4.1. but almost trivially so. To claim agreement between the two sides of the duality, we still need to reproduce the spectrum of fluctuations about the giant configuration (5.18) in the gauge theory. A much more formidable task, this is encoded in the dynamics of open strings attached to the giant operator¹³ (6.2). Open strings in the gauge theory are given by operators of the form $W_j^i = (M_1 M_2 \cdots M_n)_j^i$ where the M 's could be Higgs fields, their covariant derivatives or even fermions. Following [13], the prescription for attaching open strings to the giant graviton consists of removing some number of Z 's that make up the giant and replacing them with the same number of words W . For a maximal sphere giant with a single open string attached, for example, the resulting operator takes the form (up to a normalization)

$$\mathcal{O}_k = \epsilon_{i_1 \cdots i_N}^{j_1 \cdots j_N} Z_{j_1}^{i_1} \cdots Z_{j_{N-1}}^{i_{N-1}} \left(Y^k M Y^{J-k} \right)_{j_N}^{i_N}, \quad (6.8)$$

where the string of Y 's build up the worldvolume of the string excitation with angular momentum $J \sim O(\sqrt{N})$ and the impurities M are oscillator excitations of the string, either along the brane ($M = X$) or transverse to the giant worldvolume ($M = Z$). In any case, hopping of the impurity along the chain introduces γ -dependent phase factors into the correlators computed from the operators (6.8). It is the precise form of this γ -dependence that we need to match with the fluctuation spectrum about the pp-wave giant. For giants on the more general Frolov-Roiban-Tseytlin non-supersymmetric deformation of $AdS_5 \times S^5$, a detailed study of such operators was carried out in [22] where superb agreement between the gauge theory and supergravity was reported. Armed with some of the technology developed there and in [35], we return to these questions shortly [37].

7. Summary and outlook

In this article we have initiated a study of giant gravitons on the pp-wave geometries arising from the Lunin-Maldacena deformation of $AdS_5 \times S^5$. In particular we find two classes of giant graviton solutions, depending on whether the Penrose limit is taken about null geodesics carrying only one J_{ϕ_i} charge or those where all three J_{ϕ_i} 's are non-zero. In the former - of which the $(J, 0, 0)$ pp-wave is a prototype - the giant graviton is energetically degenerate with the Kaluza-Klein point-graviton irrespective of the strength of the deformation. Like their undeformed counterparts, these spherical $D3$ -branes exhibit a fluctuation spectrum that is independent of the size of the giant and absent of tachyonic modes indicating that they are stable objects in the supergravity theory. On the other hand, in section 4.2 we find that giants on the (J, J, J) pp-wave suffer a deformation - not unlike that of the squashed giants of [29] - due to non-vanishing contributions from the NS

¹³Fortunately, much of the difficulty in studying open strings attached to giant gravitons can be circumvented by following the remarkable proposal put forth in the series of articles [40]. In the first of these the authors systematically develop a powerful set of diagrammatic rules - based on the Young diagram labels of the Schur polynomials - for computing the two-point correlation functions of various giant operators and then use these to study the attachment of open strings to the giant worldvolume.

B -field and RR 2-form to the $D3$ -brane worldvolume action. Indeed, this deformation of the 3-sphere giant makes a full analytic treatment of the problem much more difficult than the $(J, 0, 0)$ case. Nevertheless, we are able to make some progress in the small $\hat{\gamma}$ limit. Up to $O(\hat{\gamma}^4)$, we find that the energy of the giant graviton is lifted with respect to the KK-graviton hinting that the giant graviton on this pp-wave wave background is energetically unstable. Indeed, for sufficiently large $\hat{\gamma}$ the second minimum of the lightcone Hamiltonian vanishes altogether. Finally, in section 6. we have concluded with some brief observations of the behaviour that we expect of the operators dual to these giant gravitons.

Arguably, we have only just scratched the surface in the study of these configurations and many questions remain. For us, the first among these is a more complete analysis of the (J, J, J) pp-wave giant. Clearly, this background deforms the 3-sphere worldvolume of the $D3$ -brane. The Lagrangian analysis carried out here has proven to be rather nontrivial. However, similar deformations have been observed in the slightly simpler context of giant gravitons in the maximally supersymmetric type IIB plane wave in the presence of a constant B -field in [29]. There, a fairly complete treatment of the problem was given using a Hamiltonian analysis and the emergence of a Nambu bracket structure in the worldvolume action. This then offers some hope that an analytic solution may be found here too. Secondly, this work began with the aim of determining the existence of giant graviton solutions on the Lunin-Maldacena background. To this end, we have made some progress in showing that at least the Penrose limit of this geometry supports these configurations. Following on from this work and the supergravity results of [22] two, more recent articles have since appeared that have succeeded in independently demonstrating the existence of giant gravitons in the non-supersymmetric Frolov-Roiban-Tseytlin (and by subsequent restriction to a single deformation parameter the Lunin-Maldacena) background. In [39] both sphere and AdS giants are constructed for giants with angular momentum along the $(J, 0, 0)$, $(0, J, 0)$ and $(0, 0, J)$ class of orbits. As is the case for the corresponding pp-wave studied in this article, these giant gravitons are not only blind to the deformation but their fluctuation spectra match nicely with ours. On the other hand, the authors of [41] begin from the giant graviton configurations in the undeformed $AdS_5 \times S^5$ and, by carefully following the boundary conditions defining the D3-brane in the supergravity through the TsT transformation that maps $AdS_5 \times S^5$ to $AdS_5 \times S^5_\gamma$, find that the D3-brane giant graviton not only survives the deformation but is described by a Hamiltonian (see eq. (4.11) in [41])

$$H = \frac{1}{R \cos \alpha} \sqrt{J_{\phi_1}^2 - 2J_{\phi_1} N \sin^4 \alpha + N^2 \sin^6 \alpha},$$

that is also manifestly independent of the deformation parameter. This is in excellent agreement with both our results and that of [39]. More intriguingly, it was also found that when the deformation parameter is rational, in addition to the D3-brane giants, the Lunin-Maldacena geometry also admits a class of D5-brane giants wrapping the torus on which the TsT transformation acts and with a non-vanishing world-volume gauge field strength inversely proportional to the deformation parameter.

Returning to the case at hand though, there are, of course, also giants that expand in the AdS directions of the pp-wave. Since the deformation is exactly marginal, these

are not affected and straightforward to construct. However, their spectrum of fluctuations *will!* Exactly how, remains to be seen in this case. Then, of course, there is the question of the operators dual to the deformed giants - some subset of the Schur polynomials that survive the γ -deformation - the open strings attached to them and the various correlators that need to be computed to match our supergravity results.

It seems clear then that the study of giant gravitons on these backgrounds is an extremely fruitful area and promises to shed much needed light on a complete understanding of the gauge theory/gravity correspondence.

Acknowledgments

J.M would like to dedicate this work to the memory of Thavarajoo Murugan. His presence will be greatly missed. Our gratitude is extended to Antal Jevicki, Dan Kabat, Horatiu Nastase, Marco Pirrone and Amanda Weltman for valuable discussions at various stages of this work. We are especially grateful to Robert de Mello Koch for his constant support and encouragement throughout this project. J.M. was supported by an NRF (South Africa) overseas postdoctoral research fellowship at Brown University where most of this work was carried out. A.H. was supported at Columbia University by DOE grant DEFG02-92ER40699. Finally, the faculty, staff and students of the High Energy Theory group at Brown are gratefully acknowledged for all their support.

References

- [1] J. Maldacena, *The large N limit of superconformal field theories and supergravity*, *Adv. Theor. Math. Phys.* **2** (1998) 231 [*Int. J. Theor. Phys.* **38** (1999) 1113] [[hep-th/9711200](#)].
- [2] O. Lunin and S.D. Mathur, *AdS/CFT duality and the black hole information paradox*, *Nucl. Phys.* **B 623** (2002) 342 [[hep-th/0109154](#)];
S.D. Mathur, *The fuzzball proposal for black holes: an elementary review*, *Fortschr. Phys.* **53** (2005) 793 [[hep-th/0502050](#)].
- [3] D. Berenstein, *Large- N BPS states and emergent quantum gravity*, *JHEP* **01** (2006) 125 [[hep-th/0507203](#)];
N. Seiberg, *Emergent spacetime*, [hep-th/0601234](#);
D. Berenstein and R. Cotta, *Aspects of emergent geometry in the AdS/CFT context*, *Phys. Rev.* **D 74** (2006) 026006 [[hep-th/0605220](#)].
- [4] H. Nastase, *The RHIC fireball as a dual black hole*, [hep-th/0501068](#);
S.S. Gubser, *Drag force in AdS/CFT*, *Phys. Rev.* **D 74** (2006) 126005 [[hep-th/0605182](#)];
J.J. Friess, S.S. Gubser and G. Michalogiorgakis, *Dissipation from a heavy quark moving through $N = 4$ super-Yang-Mills plasma*, *JHEP* **09** (2006) 072 [[hep-th/0605292](#)];
R.C. Brower, J. Polchinski, M.J. Strassler and C.-I. Tan, *The Pomeron and gauge/string duality*, [hep-th/0603115](#).
- [5] J. Polchinski, *Dirichlet-branes and Ramond-Ramond charges*, *Phys. Rev. Lett.* **75** (1995) 4724 [[hep-th/9510017](#)].
- [6] J. McGreevy, L. Susskind and N. Toumbas, *Invasion of the giant gravitons from anti-de Sitter space*, *JHEP* **06** (2000) 008 [[hep-th/0003075](#)].

- [7] M.T. Grisaru, R.C. Myers and O. Tafjord, *SUSY and Goliath*, *JHEP* **08** (2000) 040 [[hep-th/008016](#)];
A. Hashimoto, S. Hirano and N. Itzhaki, *Large branes in AdS and their field theory dual*, *JHEP* **08** (2000) 051 [[hep-th/0008016](#)].
- [8] S.R. Das, A. Jevicki and S.D. Mathur, *Giant gravitons, BPS bounds and noncommutativity*, *Phys. Rev. D* **63** (2001) 044001 [[hep-th/0008088](#)];
A. Mikhailov, *Giant gravitons from holomorphic surfaces*, *JHEP* **11** (2000) 027 [[hep-th/0010206](#)];
I. Bena and D.J. Smith, *Towards the solution to the giant graviton puzzle*, *Phys. Rev. D* **71** (2005) 025005 [[hep-th/0401173](#)];
T. Yoneya, *String theory and space-time uncertainty principle*, *Prog. Theor. Phys.* **103** (2000) 1081 [[hep-th/0004074](#)];
R.C. Myers and O. Tafjord, *Superstars and giant gravitons*, *JHEP* **11** (2001) 009 [[hep-th/0109127](#)];
B. Janssen and Y. Lozano, *A microscopical description of giant gravitons*, *Nucl. Phys. B* **658** (2003) 281 [[hep-th/0207199](#)].
- [9] S.R. Das, A. Jevicki and S.D. Mathur, *Vibration modes of giant gravitons*, *Phys. Rev. D* **63** (2001) 024013 [[hep-th/0009019](#)].
- [10] S. Corley, A. Jevicki and S. Ramgoolam, *Exact correlators of giant gravitons from dual $N = 4$ SYM theory*, *Adv. Theor. Math. Phys.* **5** (2002) 809 [[hep-th/0111222](#)];
V. Balasubramanian, M. Berkooz, A. Naqvi and M. Strassler, *Giant gravitons in conformal field theory*, *JHEP* **04** (2002) 034 [[hep-th/0107119](#)].
- [11] R. de Mello Koch and R. Gwyn, *Giant graviton correlators from dual $SU(N)$ super Yang-Mills theory*, *JHEP* **11** (2004) 081 [[hep-th/0410236](#)].
- [12] A. Jevicki, *Non-perturbative collective field theory*, *Nucl. Phys. B* **376** (1992) 75;
D. Berenstein, *A toy model for the AdS/CFT correspondence*, *JHEP* **07** (2004) 018 [[hep-th/0403110](#)].
- [13] D. Berenstein, *Shape and holography: studies of dual operators to giant gravitons*, *Nucl. Phys. B* **675** (2003) 179 [[hep-th/0306090](#)];
V. Balasubramanian, D. Berenstein, B. Feng and M-X. Huang, *D-branes in Yang-Mills theory and emergent gauge symmetry*, *JHEP* **03** (2005) 006 [[hep-th/0411205](#)].
- [14] H. Lin, O. Lunin and J. Maldacena, *Bubbling AdS space and 1/2 BPS geometries*, *JHEP* **10** (2004) 025 [[hep-th/0409174](#)].
- [15] R.G. Leigh and M.J. Strassler, *Exactly marginal operators and duality in four-dimensional $N = 1$ supersymmetric gauge theory*, *Nucl. Phys. B* **447** (1995) 95 [[hep-th/9503121](#)].
- [16] D. Bundzik, *Star product and the general Leigh-Strassler deformation*, *JHEP* **04** (2007) 035 [[hep-th/0608215](#)].
- [17] J.P. Gauntlett, S. Lee, T. Mateos and D. Waldram, *Marginal deformations of field theories with AdS_4 duals*, *JHEP* **08** (2005) 030 [[hep-th/0505207](#)];
C.-h. Ahn and J.F. Vazquez-Poritz, *Marginal deformations with $U(1)^3$ global symmetry*, *JHEP* **07** (2005) 032 [[hep-th/0505168](#)].
- [18] U. Gursoy and C. Núñez, *Dipole deformations of $N = 1$ sym and supergravity backgrounds with $U(1) \times U(1)$ global symmetry*, *Nucl. Phys. B* **725** (2005) 45 [[hep-th/0505100](#)];

- U. Gursoy, *Probing universality in the gravity duals of $N = 1$ SYM by gamma deformations*, *JHEP* **05** (2006) 014 [[hep-th/0602215](#)];
- N.P. Bobev, H. Dimov and R.C. Rashkov, *Semiclassical strings, dipole deformations of $N = 1$ SYM and decoupling of KK modes*, *JHEP* **02** (2006) 064 [[hep-th/0511216](#)].
- [19] S.A. Frolov, R. Roiban and A.A. Tseytlin, *Gauge - string duality for superconformal deformations of $N = 4$ super Yang-Mills theory*, *JHEP* **07** (2005) 045 [[hep-th/0503192](#)];
- N.P. Bobev, H. Dimov and R.C. Rashkov, *Semiclassical strings in Lunin-Maldacena background*, [hep-th/0506063](#);
- A.H. Prinsloo, *gamma(i) deformed lax pair for rotating strings in the fast motion limit*, *JHEP* **01** (2006) 050 [[hep-th/0510095](#)];
- H.-Y. Chen and S. Prem Kumar, *Precision test of AdS/CFT in Lunin-Maldacena background*, *JHEP* **03** (2006) 051 [[hep-th/0511164](#)];
- H.-Y. Chen and K. Okamura, *The anatomy of gauge/string duality in Lunin-Maldacena background*, *JHEP* **02** (2006) 054 [[hep-th/0601109](#)];
- S. Ryang, *Rotating strings with two unequal spins in Lunin-Maldacena background*, *JHEP* **11** (2005) 006 [[hep-th/0509195](#)].
- [20] S. Frolov, *Lax pair for strings in Lunin-Maldacena background*, *JHEP* **05** (2005) 069 [[hep-th/0503201](#)].
- [21] S.A. Frolov, R. Roiban and A.A. Tseytlin, *Gauge-string duality for (non)supersymmetric deformations of $N = 4$ super Yang-Mills theory*, *Nucl. Phys.* **B 731** (2005) 1 [[hep-th/0507021](#)].
- [22] R. de Mello Koch, N. Ives, J. Smolic and M. Smolic, *Unstable giants*, *Phys. Rev.* **D 73** (2006) 064007 [[hep-th/0509007](#)].
- [23] M. Blau, J. Figueroa-O'Farrill, C. Hull and G. Papadopoulos, *Penrose limits and maximal supersymmetry*, *Class. and Quant. Grav.* **19** (2002) L87 [[hep-th/0201081](#)].
- [24] D. Berenstein, J.M. Maldacena and H. Nastase, *Strings in flat space and pp waves from $N = 4$ super Yang-Mills*, *JHEP* **04** (2002) 013 [[hep-th/0202021](#)].
- [25] H. Takayanagi and T. Takayanagi, *Notes on giant gravitons on pp-waves*, *JHEP* **12** (2002) 018 [[hep-th/0209160](#)].
- [26] K. Skenderis and M. Taylor, *Branes in AdS and pp-wave spacetimes*, *JHEP* **06** (2002) 025 [[hep-th/0204054](#)]; *An overview of branes in the plane wave background*, *Class. Quant. Grav.* **20** (2003) S567 [[hep-th/0301221](#)].
- [27] J. Polchinski and M.J. Strassler, *The string dual of a confining four-dimensional gauge theory*, [hep-th/0003136](#).
- [28] D. Sadri and M.M. Sheikh-Jabbari, *Giant hedge-hogs: spikes on giant gravitons*, *Nucl. Phys.* **B 687** (2004) 161 [[hep-th/0312155](#)].
- [29] S. Prokushkin and M.M. Sheikh-Jabbari, *Squashed giants: bound states of giant gravitons*, *JHEP* **07** (2004) 077 [[hep-th/0406053](#)].
- [30] O. Lunin and J.M. Maldacena, *Deforming field theories with $U(1) \times U(1)$ global symmetry and their gravity duals*, *JHEP* **05** (2005) 033 [[hep-th/0502086](#)].
- [31] V. Niarchos and N. Prezas, *Bmn operators for $N = 1$ superconformal Yang-Mills theories and associated string backgrounds*, *JHEP* **06** (2003) 015 [[hep-th/0212111](#)].

- [32] L.A. Pando Zayas and J. Sonnenschein, *On Penrose limits and gauge theories*, *JHEP* **05** (2002) 010 [[hep-th/0202186](#)].
- [33] J.M. Maldacena and J.G. Russo, *Large- N limit of non-commutative gauge theories*, *JHEP* **09** (1999) 025 [[hep-th/9908134](#)].
- [34] G. Papadopoulos, J.G. Russo and A.A. Tseytlin, *Solvable model of strings in a time dependent plane wave background*, *Class. and Quant. Grav.* **20** (2003) 969 [[hep-th/0211289](#)]; M. Blau and M. O’Loughlin, *Homogeneous plane waves*, *Nucl. Phys.* **B 654** (2003) 135.
- [35] R. de Mello Koch, J. Murugan, J. Smolic and M. Smolic, *Deformed pp-waves from the Lunin-Maldacena background*, *JHEP* **08** (2005) 072 [[hep-th/0505227](#)].
- [36] T. Mateos, *Marginal deformation of $N = 4$ SYM and Penrose limits with continuum spectrum*, *JHEP* **08** (2005) 026 [[hep-th/0505243](#)].
- [37] A. Hamilton and J. Murugan, *A study of giant graviton operators in the Leigh-Strassler deformation of $\mathcal{N} = 4$ SYM*, in progress.
- [38] W. Fulton and J. Harris, *Representation theory*, Springer Verlag (1991).
- [39] M. Pirrone, *Giants on deformed backgrounds*, *JHEP* **12** (2006) 064 [[hep-th/0609173](#)].
- [40] R. de Mello Koch, J. Smolic and M. Smolic, *Giant gravitons - with strings attached (I)*, [hep-th/0701066](#);
R. de Mello Koch, J. Smolic and M. Smolic, *Giant gravitons - with strings attached. II*, [hep-th/0701067](#).
- [41] E. Imeroni and A. Naqvi, *Giants and loops in beta-deformed theories*, *JHEP* **03** (2007) 034 [[hep-th/0612032](#)].