

Landau-Lifshitz sigma-models, fermions and the AdS/CFT correspondence

Bogdan Stefański jr.

*Center for Theoretical Physics, Laboratory for Nuclear Science,
Massachusetts Institute of Technology,
Cambridge, MA 02139, U.S.A., and
Theoretical Physics Group, Blackett Laboratory, Imperial College,
London SW7 2BZ, U.K.
E-mail: b.stefanski@imperial.ac.uk*

ABSTRACT: We define Landau-Lifshitz sigma models on general coset space G/H , with H a maximal stability sub-group of G . These are non-relativistic models that have G -valued Nöther charges, local H invariance and are classically integrable. Using this definition, we construct the $PSU(2, 2|4)/PS(U(2|2)^2)$ Landau-Lifshitz sigma-model. This sigma model describes the thermodynamic limit of the spin-chain Hamiltonian obtained from the complete one-loop dilatation operator of the $N = 4$ super Yang-Mills (SYM) theory. In the second part of the paper, we identify a number of consistent truncations of the Type IIB Green-Schwarz action on $AdS_5 \times S^5$ whose field content consists of two real bosons and 4, 8 or 16 real fermions. We show that κ -symmetry acts trivially in these sub-sectors. In the context of the large spin limit of the AdS/CFT correspondence, we map the Lagrangians of these sub-sectors to corresponding truncations of the $PSU(2, 2|4)/PS(U(2|2)^2)$ Landau-Lifshitz sigma-model.

KEYWORDS: AdS-CFT Correspondence; Superstrings and Heterotic Strings.

Contents

1. Introduction	2
2. Landau-Lifshitz sigma models	4
2.1 The $\mathbf{U}(1 1)/\mathbf{U}(1)^2$ model	5
2.2 The $\mathrm{SU}(3)/S(\mathbf{U}(2) \times \mathbf{U}(1))$ model	6
2.3 The $\mathrm{SU}(2, 2 4)/S(\mathbf{U}(2 2) \times \mathbf{U}(2 2))$ model	7
2.3.1 Subsectors of the the $\mathrm{SU}(2, 2 4)/S(\mathbf{U}(2 2) \times \mathbf{U}(2 2))$ model	8
3. Green-Schwarz actions and fake κ-symmetry	9
3.1 Fermionic GS actions	10
3.2 The GS action on $\mathbf{U}(2 2)/\mathrm{SU}(2)^2$	11
3.3 The GS action on $\mathbf{U}(1 1)^2/\mathbf{U}(1)^2$	12
3.4 Fake κ -symmetry	14
4. Large charge limits of fermionic GS actions	16
4.1 Matching the $\mathbf{U}(1 1)^2$ sub-sectors	17
4.2 Large Charge Limit of fermionic GS actions	19
4.3 A gauge-theory inspired κ gauge	20
5. Conclusion	20
A. Some examples of Landau-Lifshitz sigma models	21
A.1 The $\mathrm{SU}(2 3)/S(\mathbf{U}(2 2) \times \mathbf{U}(1))$ model	21
A.2 The $\mathrm{SU}(4)/S(\mathbf{U}(2) \times \mathbf{U}(2))$ model	22
A.2.1 Subsectors of the $\mathrm{SU}(4)/S(\mathbf{U}(2) \times \mathbf{U}(2))$ model	23
A.3 The $\mathrm{SU}(2, 2)/S(\mathbf{U}(2) \times \mathbf{U}(2))$ model	24
A.3.1 Subsectors of the $\mathrm{SU}(2, 2)/S(\mathbf{U}(2) \times \mathbf{U}(2))$ model	26
A.4 The $\mathrm{SU}(2 2)/S(\mathbf{U}(1 1) \times \mathbf{U}(1 1))$ model	26
B. Quantising the action (3.18) in the $t + \alpha = \kappa\tau$ gauge	27
C. Comments on conformal invariance of the action (3.18)	30
D. Two dimensional spin connection	32
E. T-dual version of the action (3.18)	33

1. Introduction

The gauge/string correspondence [1] provides an amazing connection between quantum gauge and gravity theories. The correspondence is best understood in the case of the maximally supersymmetric dual pair of $\mathcal{N} = 4$ $SU(N)$ super-Yang-Mills (SYM) gauge theory and Type IIB string theory on $AdS_5 \times S^5$. Recent progress in understanding this duality has come from investigations of states in the dual theories with large charges [3–5]. In these large-charge limits (LCLs) it is possible to test the duality in sectors where quantities are not protected by supersymmetry. Typically, one compares the energy of some semi-classical string state with large charges (labelled schematically J) to the anomalous dimensions of the corresponding operator in the dual gauge theory, using $1/J$ as an expansion parameter which suppresses quantum corrections. A crucial ingredient, which made such comparisons possible, was the observation that computing anomalous dimensions in the $\mathcal{N} = 4$ SYM gauge theory is equivalent to finding the energy eigenvalues of certain integrable spin-chains [6] (following the earlier work on more generic gauge theories [7]). At the same time the classical Green-Schwarz (GS) action for the Type IIB string theory on $AdS_5 \times S^5$ was shown to be integrable [21]. The presence of integrable structures has led to an extensive use of Bethe ansatz-type techniques to investigate the gauge/string duality [8]. In particular, impressive results for matching the world-sheet S-matrix of the GS string sigma-model with the corresponding S-matrix of the spin-chain have been obtained [10].

The matching of anomalous dimensions of gauge theory operators with the energies of semi-classical string states was shown to work up to and including two loops in the 't Hooft coupling λ . At three loops it was shown that the string and gauge theory results differ. As has been noted many times in the literature, this result should not be interpreted as a falsification of the gauge/string correspondence conjecture. Indeed, while the (perturbative) gauge theory computations are done at small values in λ , they are compared to dual string theory energies which are computed at large values of λ and as such are not necessarily comparable. It has then been a fortunate coincidence that the one- and two-loop results do match.

This match was first established in a number of particular semi-classical string solutions and corresponding single-trace operators [5]. Later it was shown that, to leading order in the LCL, for some bosonic sub-sectors the string action reduced to a generalised Landau-Lifshitz (LL) sigma model, which also could be obtained as a thermodynamic limit of the corresponding spin-chain [13–16, 18] (see also [20]). In this way, by matching Lagrangians on both sides one can establish that energies of a wide class of string solutions do indeed match with the corresponding anomalous dimensions of gauge theory operators without having to compute these on a case-by-case basis.

A natural extension of this programme is to match, to leading order, the LCL of the full GS action of Type IIB string theory on $AdS_5 \times S^5$ to the thermodynamic limit of the spin-chain corresponding to the dilatation operator for the full $\mathcal{N} = 4$ SYM gauge theory; including fermions on both sides of the map is interesting given the different way in which they enter the respective actions. On the spin-chain side fermions are on equal footing to bosons [19, 17] - the LL equation, which describes the thermodynamic limit of

the system, relates to a super-coset manifold when fermions are included, as opposed to a coset manifold when there are no fermions. In particular, both fermions and bosons satisfy equations which are first order in τ and second order in σ . On the other hand, fermions in the GS action possess κ -symmetry [23–26] and their equations of motion are first order both in τ and σ . Previous progress on this question was able to match string and spin chain actions in a LCL up to quadratic level in fermions [20, 17, 19]. Roughly speaking, on the string side, κ -gauge fixed equations of motion for fermions typically come as $2n$ first order equations. From these one obtains n second-order equations for n by ‘integrating out’ half of the fermions. Taking a non-relativistic limit on the worldsheet one ends up with equations which are first order in τ and second order in σ which can be matched with the corresponding LL equations obtained from the spin chain side. Matching the terms quartic and higher in the fermions had so far not been achieved, though it is expected that this should be possible given the results of [9]. However, finding a suitable κ -gauge in which this matching could be done in a natural way remained an obstacle. Below we propose a κ gauge which appears to be natural from the point of view of the dual spin-chain and allows for a matching of higher order fermionic terms in the dual Lagrangians.

In this paper we first present a compact way of writing LL sigma models for quite general (super-)cosets G/H ; in particular we write down the full $PSU(2, 2|4)/PS(U(2|2)^2)$ LL sigma model which arises as the thermodynamic limit of the one-loop dilatation operator for the full $\mathcal{N} = 4$ SYM theory. This generalises earlier work by [11], and allows one to write down LL-type actions without having to go through the coherent-state [12] thermodynamic limit of the spin chain. We then identify a number of sub-sectors of the classical GS action¹ all of which have two real bosonic degrees of freedom and a larger number of fermionic degrees of freedom (specifically 4, 8 and 16 real fermionic d.o.f.s²). Finally, we define a LCL in which the GS actions for these fermionic sub-sectors reduce to corresponding LL actions. In this way we match the complete Lagrangians for these sub-sectors and not just the terms quadratic in fermions. Since the largest of these sectors contains the maximal number of fermions (sixteen) for a κ -fixed GS action the LCL matching to a LL model gives a clear indication of what the natural κ -gauge is from the point of view of the dual spin-chain.

The fermionic sub-sectors of the GS action that we find are quite interesting in themselves because on-shell κ -symmetry acts trivially on them - in particular the sub-sector containing 16 fermionic degrees of freedom contains the same number of fermions as the κ -fixed GS superstring on $AdS_5 \times S^5$. Since κ -symmetry acts trivially in this case one cannot use it to eliminate half of the fermions as one does in more conventional GS actions. Further, these fermionic sub-sectors naturally inherit the classical integrability of the full GS superstring on $AdS_5 \times S^5$ found in [21]. Integrating out the metric and the two bosonic

¹By a sub-sector we mean that the classical equations of motion for the full GS superstring on $AdS_5 \times S^5$ admit a truncation in which all other fields are set to zero in a manner which is consistent with their equations of motion. This is quite familiar in two cases: (i) when one sets all fermions in the GS action to zero and, (ii) when one further restricts the bosons to lie on some $AdS_p \times S^q$ sub-space ($1 \leq p, q \leq 5$).

²The 4 fermion model was previously postulated to be a sub-sector of the classical GS action in [29] and represents a starting point for our analysis.

degrees of freedom one then arrives at a new class of integrable differential equations for fermions only.

This paper is organised as follows. In section 2 we give a prescription for constructing a LL sigma model on a general coset G/H . We also present a number of explicit examples of LL sigma models most relevant to the gauge/string correspondence there and in appendix A. In section 3 we identify the fermionic sub-sectors of the GS superstring on $AdS_5 \times S^5$. In section 4 we define a LCL in which the GS action of the fermionic sub-sectors reduces, to leading order in J , to the LL sigma models for the corresponding gauge-theory fermionic sub-sectors. Since the GS action for the four fermion subsector is quadratic in the remaining appendices to this paper we present a more detailed discussion of it including a light-cone quantisation in appendix B, a discussion of its conformal invariance in appendix C and a T-dual form of the action in appendix E.

2. Landau-Lifshitz sigma models

In this section we construct the Lagrangian for a Landau-Lifshitz (LL) sigma model on a coset G/H .³ The Lagrangian will typically be first (second) order in the worldsheet time (space) coordinate, and so is non-relativistic on the worldsheet. We refer to such models as LL sigma models because in the case of $G/H = SU(2)/U(1)$ the equations of motion reduce to the usual LL equation

$$\partial_\tau n_i = \varepsilon_{ijk} n_j \partial_\sigma^2 n_k, \quad \text{where} \quad n_i n_i = 1. \quad (2.1)$$

The construction of LL Lagrangians is closely related to coherent states $|\omega, \Lambda\rangle$. Recall⁴ that to construct a coherent state $|\omega, \Lambda\rangle$ we need to specify a unitary irreducible representation Λ of G acting on a Hilbert space V_Λ and a vacuum state $|0\rangle$ on which H is a maximal stability sub-group, in other words for any $h \in H$

$$\Lambda(h) |0\rangle = e^{i\phi(h)} |0\rangle, \quad (2.2)$$

with $\phi(h) \in \mathbf{R}$. Given such a representation Λ and state $|0\rangle$ we define the operator Ω as

$$\Omega \equiv |0\rangle \langle 0|. \quad (2.3)$$

The LL sigma model Lagrangian on G/H is defined as

$$\mathcal{L}_{LL\ G/H} = \mathcal{L}_{LL\ G/H}^{WZ} + \mathcal{L}_{LL\ G/H}^{kin} \quad (2.4)$$

where

$$\mathcal{L}_{LL\ G/H}^{WZ} = -i \text{Tr} \left(\Omega g^\dagger \partial_\tau g \right), \quad (2.5)$$

$$\mathcal{L}_{LL\ G/H}^{kin} = \text{Tr} \left(g^\dagger D_\sigma g g^\dagger D_\sigma g \right). \quad (2.6)$$

³For earlier work on this see [11].

⁴For a detailed exposition of coherent states see [12]; a brief summary, using the same notation as in this paper, is also presented in appendix A of [18].

Above, $g^\dagger D_\sigma g \equiv g^\dagger \partial_\sigma g - g^\dagger \partial_\sigma g|_H$ is just the standard H -covariant current. It is then clear that $\mathcal{L}_{\text{LL } G/H}^{\text{kin}}$ is invariant under gauge transformations

$$g \rightarrow gh, \tag{2.7}$$

for any $h = h(\tau, \sigma) \in H$. We may also show that the same is true of $\mathcal{L}_{\text{LL } G/H}^{\text{WZ}}$. To see this note that the gauge variation of $\mathcal{L}_{\text{LL } G/H}^{\text{WZ}}$, using equation (2.2), is given by

$$\delta_H \mathcal{L}_{\text{LL } G/H}^{\text{WZ}} = e^{-i\phi(h)} \langle 0 | \partial_\tau h | 0 \rangle = e^{-i\phi(h)} \partial_\tau (\langle 0 | h | 0 \rangle) = i \partial_\tau \phi(h). \tag{2.8}$$

This in turn is a total derivative; and so the full action is invariant under local right H action. The Lagrangian also has a global G symmetry

$$g \rightarrow g_0 g, \tag{2.9}$$

for any $g_0 \in G$ with $\partial_\tau g_0 = \partial_\sigma g_0 = 0$, and the corresponding Nöther current is given by

$$(j_\tau, j_\sigma) = (g \Omega g^\dagger, 2i D_\sigma g g^\dagger). \tag{2.10}$$

In [18, 19] LL actions were written down in terms of Lie algebra matrices denoted typically by N . To make contact with the present notation we note that⁵

$$N \equiv g \Omega g^\dagger - \frac{1}{n} \mathbf{I}_n, \tag{2.11}$$

where the second term on the right hand side is included since N is traceless. Finally, let us note that these LL sigma models admit a Lax pair representation and as a result are integrable. This is most easily seen in terms of the matrix N for which the equations of motion are the LL matrix equation

$$\partial_\tau N = \frac{i}{2} [N, \partial_\sigma^2 N]. \tag{2.12}$$

This is equivalent to the zero-curvature condition on the following Lax pair

$$\mathcal{L} \longrightarrow \partial_\sigma - \frac{iN}{4\pi x}, \tag{2.13}$$

$$\mathcal{M} \longrightarrow \partial_\tau - \frac{iN}{4\pi^2 x^2} - \frac{[N, \partial_\sigma N]}{8\pi x}, \tag{2.14}$$

where $[\cdot, \cdot]$ is the (super)-commutator. In the remainder of this section we construct a number of explicit examples of LL sigma models. Further examples of interest in the gauge/string correspondence are relegated to appendix A. The reader who is not interested in the details of these examples should skip the remainder of this section.

2.1 The $\mathbf{U}(1|1)/\mathbf{U}(1)^2$ model

This is one of the simplest LL sigma models,⁶ in that the Lagrangian is quadratic

$$\mathcal{L}_{\text{LL } \mathbf{U}(1|1)/\mathbf{U}(1)^2} = i\bar{\psi} \partial_\tau \psi + \partial_\sigma \bar{\psi} \partial_\sigma \psi, \tag{2.15}$$

with ψ a complex Grassmann-odd field and $\bar{\psi}$ its complex conjugate. Notice that this result can be obtained using the explicit 2×2 supermatrix representation of $\mathbf{U}(1|1)$, with the vacuum state $|0\rangle$ being the super-vector $(0, 1)$.

⁵The following equation is due to Charles Young.

⁶There is also the equally simple bosonic $\mathbf{U}(1)$ LL sigma model.

2.2 The $SU(3)/S(U(2) \times U(1))$ model

Before proceeding to our main example - the $PSU(2,2|4)$ model - in this subsection we show how the above formal prescription applies to the well known $SU(3)$ Landau-Lifshitz model [18, 16]. Recall that the Lagrangian for this is

$$\mathcal{L}_{SU(3)/S(U(2) \times U(1))} = -iU^i \partial_\tau U_i - \frac{1}{2} |D_\sigma U_i|^2 + \Lambda(U_i U^i - 1), \quad (2.16)$$

where

$$D_\mu U_i \equiv \partial_\mu U_i - iC_\mu U_i, \quad C_\mu = -iU^i \partial_\mu U_i, \quad (2.17)$$

for $\mu = \tau, \sigma$ and $U^i \equiv U_i^*$. To show that we can obtain this from our general expression (2.4) we write elements of the group $SU(3)$ as 3×3 matrix g , split into a 3×2 matrix X and a vector Y

$$g = (X, Y), \quad (2.18)$$

and because g is in $SU(3)$ (i.e. $g^\dagger g = 1$) we have

$$X^\dagger X = \mathbf{1}_2, \quad Y^\dagger Y = \mathbf{1}_1, \quad X^\dagger Y = 0, \quad Y^\dagger X = 0, \quad (2.19)$$

$$XX^\dagger + YY^\dagger = \mathbf{1}_3. \quad (2.20)$$

The kinetic part of the Lagrangian (2.4) is then given by

$$\begin{aligned} \mathcal{L}_{\text{kin } SU(3)/S(U(2) \times U(1))} &= \frac{1}{4} \text{Tr} \left((g^{-1} D_1 g) (g^{-1} D_1 g) \right) \\ &= \frac{1}{4} \text{Tr} \left[\left(\begin{array}{cc} X^\dagger D_1 X & X^\dagger D_1 Y \\ Y^\dagger D_1 X & Y^\dagger D_1 Y \end{array} \right)^2 \right] = \frac{1}{4} \text{Tr} \left[\left(\begin{array}{cc} 0 & X^\dagger \partial_1 Y \\ Y^\dagger \partial_1 X & 0 \end{array} \right)^2 \right] \\ &= \frac{1}{2} \text{Tr} \left[X^\dagger \partial_1 Y Y^\dagger \partial_1 X \right] \\ &= -\frac{1}{2} \text{Tr} \left[\partial_1 X^\dagger Y Y^\dagger \partial_1 X \right] = -\frac{1}{2} \text{Tr} \left[\partial_1 Y^\dagger X X^\dagger \partial_1 Y \right] \\ &= \frac{1}{2} - \text{Tr} \left[\partial_1 X^\dagger (1 - X X^\dagger) \partial_1 X \right] = -\frac{1}{2} \partial_1 Y^i (\delta_i^j - Y_i Y^j) \partial_1 Y_j \\ &= -\frac{1}{2} \text{Tr} \left[\bar{D}_1 X^\dagger D_1 X \right] = -\frac{1}{2} \bar{D}_1 Y^i D_1 Y_i. \end{aligned} \quad (2.21)$$

The final expression is the same as the kinetic term of the usual $SU(3)$ Landau-Lifshitz Lagrangian (2.16) upon identifying Y_i with U_i (above $Y^i \equiv Y^\dagger$). Above, we have defined

$$D_1 Y_i = \partial_1 Y_i - Y_i Y^j \partial_1 Y_j, \quad \bar{D}_1 Y^i \equiv (D_1 Y_i)^\dagger \quad (2.22)$$

$$D_1 X = \partial_1 X - X X^\dagger \partial_1 X, \quad \bar{D}_1 X \equiv (D_1 X)^\dagger. \quad (2.23)$$

The WZ term of the Lagrangian is given by equation (2.5) and can be written as

$$\mathcal{L}_{\text{WZ } SU(3)/S(U(2) \times U(1))} = i \text{Tr} (X^\dagger \partial_0 X) = -i Y^i \partial_0 Y_i. \quad (2.24)$$

This follows from the fact that $g^{-1} \partial_0 g$ is traceless and so

$$\text{Tr} (X^\dagger \partial_0 X) = -Y^i \partial_0 Y_i. \quad (2.25)$$

Upon identifying Y_i with U_i , the WZ term above is the same as the usual SU(3) Landau-Lifshitz one (2.16). Notice that we have also given an alternate parametrisation of the SU(3) Landau-Lifshitz model in terms of X

$$\mathcal{L}_{\text{SU}(3)/S(\text{U}(2) \times \text{U}(1))} = i \text{Tr}(X^\dagger \partial_0 X) - \frac{1}{2} \text{Tr} \left[\bar{D}_1 X^\dagger D_1 X \right] + \Lambda(X^\dagger X - 1_2), \quad (2.26)$$

which has an explicit SU(2) gauge invariance.

Finally, out of X and Y we may define a matrix which takes values in the SU(3) Lie algebra

$$N^i_j = 3Y^i Y_j - \delta_j^i = -3X_{ja} X^{ai} + 2\delta_j^i, \quad (2.27)$$

where $a = 1, 2$. This matrix is however, not a general SU(3) matrix but rather satisfies the identity

$$N^2 = N + 2. \quad (2.28)$$

In terms of N the equations of motion take the form of the matrix Landau-Lifshitz equation

$$\partial_0 N = -\frac{i}{9} [N, \partial_1^2 N]. \quad (2.29)$$

These are equivalent to the consistency of the following linear problem

$$\mathcal{L}\psi = \left[\partial_\sigma - \frac{i}{4\pi x} N \right] \psi = 0, \quad (2.30)$$

$$\mathcal{M}\psi = \left[\partial_\tau - \frac{i}{4\pi^2 x^2} N - \frac{b}{4\pi x} [N, \partial_1 N] \right] \psi = 0. \quad (2.31)$$

2.3 The SU(2, 2|4)/S(U(2|2) × U(2|2)) model

In this sub-section we present an explicit Lagrangian for the complete PSU(2, 2|4) Landau-Lifshitz sigma model Lagrangian following the general discussion at the start of the present section. The action we are interested in is the Landau Lifshitz model as defined in equation (2.4) on the coset

$$\frac{PSU(2, 2|4)}{PS(\text{U}(2|2) \times \text{U}(2|2))}, \quad (2.32)$$

or on the coset

$$\frac{SU(2, 2|4)}{S(\text{U}(2|2) \times \text{U}(2|2))}, \quad (2.33)$$

both of which have 32 real components. The derivation is very similar to the SU(3) Lagrangian derived in the previous sub-section, and so we will simply state our results. A general group element g can be written as (X, Y) where now X (Y) is a 8×4 supermatrix, with the diagonal 4×4 blocks bosonic (fermionic) and the off-diagonal 4×4 blocks fermionic (bosonic). The Lagrangian is then given by

$$\mathcal{L}_{\text{LL } PSU(2, 2|4)/PS(\text{U}(2|2) \times \text{U}(2|2))} = i \text{STr}(X^\dagger \partial_0 X) - \frac{1}{2} \text{STr}(\bar{D}_1 X^\dagger D_1 X) + \Lambda(X^\dagger X - 1). \quad (2.34)$$

Note that there are 32 complex degrees of freedom in X , which the constraints reduce to 48 real degrees of freedom. The action also has a local U(2|2) gauge invariance, so in total the above Lagrangian has 32 degrees of freedom - the same as the coset.

In fact we may write X as

$$X = (\tilde{U}_a, \tilde{V}_a, U_a, V_a), \quad X^\dagger \equiv (\tilde{U}^a, \tilde{V}^a, U^a, V^a), \quad (2.35)$$

where $a = 1, \dots, 8$, and

$$\tilde{U}^a \tilde{U}_a = -1, \quad \tilde{V}^a \tilde{V}_a = -1, \quad \tilde{V}^a \tilde{U}_a = 0, \quad \tilde{U}^a \tilde{V}_a = 0, \quad (2.36)$$

$$U^a U_a = 1, \quad V^a V_a = 1, \quad V^a U_a = 0, \quad U^a V_a = 0, \quad (2.37)$$

$$U^a \tilde{U}_a = 0, \quad U^a \tilde{V}_a = 0, \quad V^a \tilde{U}_a = 0, \quad V^a \tilde{V}_a = 0, \quad (2.38)$$

$$\tilde{U}^a U_a = 0, \quad \tilde{U}^a V_a = 0, \quad \tilde{V}^a U_a = 0, \quad \tilde{V}^a V_a = 0. \quad (2.39)$$

Above we have defined

$$U^a = U_b^* C^{ba}, \quad V^a = V_b^* C^{ba}, \quad \tilde{U}^a = -\tilde{U}_b^* C^{ba}, \quad \tilde{V}^a = -\tilde{V}_b^* C^{ba}, \quad (2.40)$$

where $C^{ab} = \text{diag}(-1, -1, 1, 1, 1, 1, 1, 1)$.

The Lagrangian (2.41) written in terms of $\tilde{U}_a, \tilde{V}_a, U_a, V_a$ is

$$\begin{aligned} \mathcal{L}_{\text{LL}} \text{PSU}(2, 2|4)/\text{PS}(\text{U}(2|2)^2) &= -i\tilde{U}^a \partial_0 \tilde{U}_a - i\tilde{V}^a \partial_0 \tilde{V}_a - iU^a \partial_0 U_a - iV^a \partial_0 V_a \\ &\quad - \frac{1}{2} \left(\partial_1 \tilde{U}^a \partial_1 \tilde{U}_a + \partial_1 \tilde{V}^a \partial_1 \tilde{V}_a + \partial_1 U^a \partial_1 U_a + \partial_1 V^a \partial_1 V_a \right. \\ &\quad \left. - \tilde{U}^a \partial_1 \tilde{U}_a \tilde{U}^b \partial_1 \tilde{U}_b - \tilde{V}^a \partial_1 \tilde{V}_a \tilde{V}^b \partial_1 \tilde{V}_b \right. \\ &\quad \left. + V^a \partial_1 V_a V^b \partial_1 V_b + U^a \partial_1 U_a U^b \partial_1 U_b \right. \\ &\quad \left. + 2V^a \partial_1 U_a U^b \partial_1 V_b - 2\tilde{V}^a \partial_1 \tilde{U}_a \tilde{U}^b \partial_1 \tilde{V}_b + 2\tilde{U}^a \partial_1 U_a U^b \partial_1 \tilde{U}_b \right. \\ &\quad \left. + 2\tilde{U}^a \partial_1 V_a V^b \partial_1 \tilde{U}_b + 2\tilde{V}^a \partial_1 U_a U^b \partial_1 \tilde{V}_b + 2\tilde{V}^a \partial_1 V_a V^b \partial_1 \tilde{V}_b \right) \end{aligned} \quad (2.41)$$

One can check explicitly that this action has local $\text{U}(2|2)$ invariance

$$(\tilde{U}_a, \tilde{V}_a, U_a, V_a) \rightarrow (\tilde{U}_a, \tilde{V}_a, U_a, V_a) \mathbf{U}(\tau, \sigma), \quad (2.42)$$

for U a $\text{U}(2|2)$ matrix.

2.3.1 Subsectors of the the $\text{SU}(2, 2|4)/\text{S}(\text{U}(2|2) \times \text{U}(2|2))$ model

In the above Lagrangian we may set

$$\tilde{U}_a = (1, 0^7), \quad \tilde{V}_a = (0, 1, 0^6), \quad U_a = (0^2, U_3, \dots, U_8), \quad V_a = (0^2, V_3, \dots, V_8), \quad (2.43)$$

where

$$U^a U_a = 1, \quad V^a V_a = 1, \quad V^a U_a = 0, \quad U^a V_a = 0. \quad (2.44)$$

The resulting Lagrangian is that of the $\text{SU}(2|4)$ sector. If we further set

$$0 = U_3 = U_4 = V_3 = V_4, \quad (2.45)$$

we can recover the SO(6) Lagrangian ([18]). Details of this are presented in appendix B. We may further consistently set

$$0 = U_8 = V_3 = V_4 = V_5 = V_6 = V_7, \quad V_8 = 1, \quad (2.46)$$

in which case we obtain the SU(2|3) Lagrangian ([19]), with the identification $(U_3, U_4) \equiv (\psi_1, \psi_2)$.

We may instead set

$$U_a = (0^7, 1), \quad V_a = (0^6, 1, 0), \quad \tilde{U}_a = (U_1, \dots, U_6, 0^2), \quad \tilde{V}_a = (V_1, \dots, V_6, 0^2), \quad (2.47)$$

where

$$\tilde{U}^a \tilde{U}_a = -1, \quad \tilde{V}^a \tilde{V}_a = -1, \quad \tilde{V}^a \tilde{U}_a = 0, \quad \tilde{U}^a \tilde{V}_a = 0. \quad (2.48)$$

The resulting Lagrangian is that of the SU(2,2|2) sector. If we further set

$$0 = U_3 = U_4 = V_3 = V_4, \quad (2.49)$$

we recover the SO(2,4) Lagrangian, which is the Wick rotated version of the SO(6) Lagrangian ([18]). In appendix B we write out this Lagrangian explicitly.

A final interesting choice is to set

$$U_a = (0^7, 1), \quad V_a = (0, V_2, \dots, V_7, 0), \quad \tilde{U}_a = (0, U_2, \dots, U_7, 0), \quad \tilde{V}_a = (1, 0^7), \quad (2.50)$$

where

$$\tilde{U}^a \tilde{U}_a = -1, \quad V^a V_a = 1, \quad V^a \tilde{U}_a = 0, \quad \tilde{U}^a V_a = 0. \quad (2.51)$$

The resulting Lagrangian is that of the SU(1,2|3) sector. If we further set

$$0 = V_2 = V_7 = \tilde{U}_2 = \tilde{U}_7, \quad (2.52)$$

we get the SU(2|2) Lagrangian. In appendix B we write out this Lagrangian explicitly.

3. Green-Schwarz actions and fake κ -symmetry

in this section we construct GS sigma model actions whose field content are two real bosons and 4,8 or 16 real fermions. These models all come from consistent truncations of the equations of motion for the full Type IIB GS action on $AdS_5 \times S^5$.⁷ Just as any GS sigma model these fermionic actions have a κ -symmetry. However, we show that for these models κ -symmetry is trivial on-shell. As a result one cannot use it to reduce the fermionic degrees of freedom of these models by fixing a κ -gauge as one does in more conventional GS actions.

Let us briefly recall the construction of the GS action on a super-coset G/H . We require that: (i) H be bosonic and, (ii) G admit a \mathbb{Z}_4 automorphism that leaves H invariant, acts

⁷Other truncations including non-zero fermions were found in [22, 19] in the Metsaev-Tseytlin κ -gauge. The truncations considered in this section are found without fixing κ -symmetry.

by -1 on the remaining bosonic part of G/H , and by $\pm i$ on the fermionic part of G/H . The currents $j_\mu = g^\dagger \partial_\mu g$ can then be decomposed as

$$j_\mu = j_\mu^{(0)} + j_\mu^{(1)} + j_\mu^{(2)} + j_\mu^{(3)}, \quad (3.1)$$

where $j^{(k)}$ has eigenvalue i^k under the \mathbb{Z}_4 automorphism. In terms of these the GS action can be written as

$$\mathcal{L}_{\text{GS } G/H} = \int d^2\sigma \sqrt{-g} g^{\mu\nu} \text{Str}(j_\mu^{(2)} j_\nu^{(2)}) + \epsilon^{\mu\nu} \text{Str}(j_\mu^{(1)} j_\nu^{(3)}), \quad (3.2)$$

from which the equations of motion are

$$0 = \partial_\alpha (\sqrt{-g} g^{\alpha\beta} j_\beta^{(2)}) - \sqrt{-g} g^{\alpha\beta} [j_\alpha^{(0)}, j_\beta^{(2)}] + \frac{1}{2} \epsilon^{\alpha\beta} \left([j_\alpha^{(1)}, j_\beta^{(1)}] - [j_\alpha^{(3)}, j_\beta^{(3)}] \right), \quad (3.3)$$

$$0 = \left(\sqrt{-g} g^{\alpha\beta} + \epsilon^{\alpha\beta} \right) [j_\alpha^{(3)}, j_\beta^{(2)}], \quad (3.4)$$

$$0 = \left(\sqrt{-g} g^{\alpha\beta} - \epsilon^{\alpha\beta} \right) [j_\alpha^{(1)}, j_\beta^{(2)}]. \quad (3.5)$$

3.1 Fermionic GS actions

Having briefly reviewed the general construction of GS actions on G/H super-cosets, we now turn to the main focus of this section which is identifying GS actions with a large number of fermionic degrees of freedom, which are consistent truncations of the full $AdS_5 \times S^5$ GS action. To do this consider the following sequence of super-cosets

$$\frac{\mathbf{U}(1|1) \times \mathbf{U}(1|1)}{\mathbf{U}(1) \times \mathbf{U}(1)} \subset \frac{\mathbf{U}(2|2)}{\text{SU}(2) \times \text{SU}(2)} \subset \frac{PS(\mathbf{U}(1, 1|2) \times \mathbf{U}(2|2))}{\text{SU}(1, 1) \times \text{SU}(2)^3} \subset \frac{PSU(2, 2|4)}{\text{SO}(1, 4) \times \text{SO}(5)}. \quad (3.6)$$

The \subset symbols are valid both for the numerators and denominators and hence for the cosets as written above. Notice that the right-most of these cosets is just the usual Type IIB on $AdS_5 \times S^5$ super-coset. Further, it is easy to convince oneself that each of the cosets above admits a \mathbb{Z}_4 automorphism which is compatible with the \mathbb{Z}_4 automorphism of the Type IIB on $AdS_5 \times S^5$ super-coset. The \mathbb{Z}_4 automorphisms may be used to write down GS actions for each of these cosets. The fact that the cosets embed into each other as shown above in a manner compatible with the \mathbb{Z}_4 automorphism implies that their GS actions can be thought of as coming from a consistent truncation of the GS action of any coset to the right of it in the above sequence. In particular this reasoning shows that the GS actions for $\mathbf{U}(1|1)^2/\mathbf{U}(1)^2$, $\mathbf{U}(2|2)/\text{SU}(2)^2$ and $\mathbf{U}(1, 1|2) \times \mathbf{U}(2|2)/(\text{SU}(1, 1) \times \text{SU}(2)^3)$ can all be thought of as coming from consistent truncations of the Type IIB GS action on $AdS_5 \times S^5$.

Counting the number of bosonic and fermionic components of the three cosets $\mathbf{U}(1|1)^2/\mathbf{U}(1)^2$, $\mathbf{U}(2|2)/\text{SU}(2)^2$ and $\mathbf{U}(1, 1|2) \times \mathbf{U}(2|2)/(\text{SU}(1, 1) \times \text{SU}(2)^3)$ we see immediately that they each have 2 real bosonic components and, respectively, 4, 8 and 16 real fermionic components - which is why we refer to these actions as fermionic GS actions. We might expect that some of the fermionic degrees of freedom could be eliminated from the GS actions by fixing κ -symmetry. In fact, it turns out that for these models κ -symmetry acts trivially on-shell and so cannot be used to eliminate some of the fermionic degrees of

freedom. Indeed, the GS actions on the above-mentioned cosets do have 4,8 and 16 real fermionic degrees of freedom, respectively.

In the remainder of this sub-section we write down explicitly the GS actions for $\mathbf{U}(2|2)/\mathrm{SU}(2)^2$ and $\mathbf{U}(1|1)^2/\mathbf{U}(1)^2$ and discuss their κ and gauge transformations; the GS action for $\mathbf{U}(1,1|2) \times \mathbf{U}(2|2)/\mathrm{SU}(1,1) \times \mathrm{SU}(2)^3$ may also be written down in an analogous fashion but since we will not need its explicit form later we refrain from writing it out in full.

3.2 The GS action on $\mathbf{U}(2|2)/\mathrm{SU}(2)^2$

The GS action on action on $\mathbf{U}(2|2)/\mathrm{SU}(2)^2$ can be written down in terms of the parametrisation of the $\mathbf{U}(2|2)$ supergroup-valued matrix written as

$$g = (X, Y; \tilde{X}, \tilde{Y}), \tag{3.7}$$

where X, Y (\tilde{X}, \tilde{Y}) are four-component super-vectors with the first (last) two entries Grassmann even and the last (first) two entries Grassmann odd. Since the matrix g is unitary we must have

$$\begin{aligned} 1 &= X^\dagger X = Y^\dagger Y = \tilde{X}^\dagger \tilde{X} = \tilde{Y}^\dagger \tilde{Y}, \\ 0 &= X^\dagger Y = Y^\dagger X = X^\dagger \tilde{X} = \tilde{X}^\dagger X = X^\dagger \tilde{Y} = \tilde{Y}^\dagger X \\ &= Y^\dagger \tilde{X} = \tilde{X}^\dagger Y = Y^\dagger \tilde{Y} = \tilde{Y}^\dagger Y = \tilde{X}^\dagger \tilde{Y} = \tilde{Y}^\dagger \tilde{X}, \\ 1_{(2|2)} &= XX^\dagger + YY^\dagger + \tilde{X}\tilde{X}^\dagger + \tilde{Y}\tilde{Y}^\dagger, \end{aligned} \tag{3.8}$$

where the matrix $1_{(2|2)}$ is just the 4×4 identity matrix. The \mathbb{Z}_4 automorphism is given by

$$\Omega : M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \longrightarrow \begin{pmatrix} \sigma^2 & 0 \\ 0 & \sigma^2 \end{pmatrix} \begin{pmatrix} -A^T & C^T \\ -B^T & -D^T \end{pmatrix} \begin{pmatrix} \sigma^2 & 0 \\ 0 & \sigma^2 \end{pmatrix}, \tag{3.9}$$

which acts on the current as

$$\Omega(j_\mu) = \begin{pmatrix} -Y^\dagger \partial_\mu Y & X^\dagger \partial_\mu Y & -\tilde{Y}^\dagger \partial_\mu Y & \tilde{X}^\dagger \partial_\mu Y \\ Y^\dagger \partial_\mu X & -X^\dagger \partial_\mu X & \tilde{Y}^\dagger \partial_\mu X & -\tilde{X}^\dagger \partial_\mu X \\ Y^\dagger \partial_\mu \tilde{Y} & -X^\dagger \partial_\mu \tilde{Y} & -\tilde{Y}^\dagger \partial_\mu \tilde{Y} & \tilde{X}^\dagger \partial_\mu \tilde{Y} \\ -Y^\dagger \partial_\mu \tilde{X} & X^\dagger \partial_\mu \tilde{X} & \tilde{Y}^\dagger \partial_\mu \tilde{X} & -\tilde{X}^\dagger \partial_\mu \tilde{X} \end{pmatrix}. \tag{3.10}$$

The Green-Schwarz action then is

$$\begin{aligned} \mathcal{L}_{\mathrm{GS} \mathbf{U}(2|2)/(\mathrm{SU}(2) \times \mathrm{SU}(2))} &= \frac{1}{2} \int d^2\sigma \sqrt{g} g^{\mu\nu} \left((X^\dagger \partial_\mu X + Y^\dagger \partial_\mu Y)(X^\dagger \partial_\nu X + Y^\dagger \partial_\nu Y) \right. \\ &\quad \left. - (\tilde{X}^\dagger \partial_\mu \tilde{X} + \tilde{Y}^\dagger \partial_\mu \tilde{Y})(\tilde{X}^\dagger \partial_\nu \tilde{X} + \tilde{Y}^\dagger \partial_\nu \tilde{Y}) \right) \\ &\quad + 2i\epsilon^{\mu\nu} \left(X^\dagger \partial_\mu \tilde{X} Y^\dagger \partial_\nu \tilde{Y} + \tilde{Y}^\dagger \partial_\mu Y \tilde{X}^\dagger \partial_\nu X \right. \\ &\quad \left. - X^\dagger \partial_\mu \tilde{Y} Y^\dagger \partial_\nu \tilde{X} - \tilde{X}^\dagger \partial_\mu Y \tilde{Y}^\dagger \partial_\nu X \right). \end{aligned} \tag{3.11}$$

One can easily check that this action has a local $SU(2) \times SU(2)$ invariance which acts on the doublets (X, Y) and (\tilde{X}, \tilde{Y}) . The action also has κ -symmetry which acts on the fields as⁸

$$\begin{aligned}\delta_\kappa X &= -\tilde{X}(\bar{\epsilon}_1 + \tilde{\epsilon}_1) - \tilde{Y}(\bar{\epsilon}_2 + \tilde{\epsilon}_2) \\ \delta_\kappa Y &= i\tilde{X}(\epsilon_2 - \tilde{\epsilon}_2) - i\tilde{Y}(\epsilon_1 - \tilde{\epsilon}_1) \\ \delta_\kappa \tilde{X} &= X(\epsilon_1 + \tilde{\epsilon}_1) + iY(\bar{\epsilon}_2 - \tilde{\epsilon}_2) \\ \delta_\kappa \tilde{Y} &= X(\epsilon_2 + \tilde{\epsilon}_2) - iY(\bar{\epsilon}_1 - \tilde{\epsilon}_1),\end{aligned}\tag{3.12}$$

where

$$\begin{aligned}\epsilon_i &= \Pi_+^{\alpha\beta} (X^\dagger \partial_\alpha X + Y^\dagger \partial_\alpha Y + \tilde{X}^\dagger \partial_\alpha \tilde{X} + \tilde{Y}^\dagger \partial_\alpha \tilde{Y}) \kappa_{i,\beta} \\ \tilde{\epsilon}_i &= \Pi_-^{\alpha\beta} (X^\dagger \partial_\alpha X + Y^\dagger \partial_\alpha Y + \tilde{X}^\dagger \partial_\alpha \tilde{X} + \tilde{Y}^\dagger \partial_\alpha \tilde{Y}) \tilde{\kappa}_{i,\beta},\end{aligned}\tag{3.13}$$

for $i = 1, 2$ with $\kappa_{i,\beta}$ and $\tilde{\kappa}_{i,\beta}$ local Grassmann-odd parameters. The world-sheet metric also varies as

$$\begin{aligned}\delta_\kappa(\sqrt{-g}g^{\alpha\beta}) &= \Pi_+^{\alpha\gamma} \left(\kappa_{1,+}^\beta (\tilde{X}^\dagger \partial_\gamma X - iY^\dagger \partial_\gamma \tilde{Y}) + \kappa_{2,+}^\beta (\tilde{Y}^\dagger \partial_\gamma X + iY^\dagger \partial_\gamma \tilde{X}) + \text{c.c.} \right) + \alpha \leftrightarrow \beta \\ &+ \Pi_-^{\alpha\gamma} \left(\tilde{\kappa}_{1,-}^\beta (\tilde{X}^\dagger \partial_\gamma X + iY^\dagger \partial_\gamma \tilde{Y}) + \tilde{\kappa}_{2,-}^\beta (\tilde{Y}^\dagger \partial_\gamma X - iY^\dagger \partial_\gamma \tilde{X}) + \text{c.c.} \right) + \alpha \leftrightarrow \beta.\end{aligned}\tag{3.14}$$

Notice that the above variation is consistent with the symmetries and the unimodularity of $\sqrt{-g}g^{\alpha\beta}$ as long as

$$\kappa_i^\alpha = \Pi_+^{\alpha\beta} \kappa_{i,\beta}, \quad \tilde{\kappa}_i^\alpha = \Pi_-^{\alpha\beta} \tilde{\kappa}_{i,\beta}.\tag{3.15}$$

In the above formulas we have decomposed two-component vectors v_α as

$$v_\pm^\alpha \equiv \Pi_\pm^{\alpha\beta} v_\beta \equiv \frac{1}{2} \left(\sqrt{-g}g^{\alpha\beta} \pm \epsilon^{\alpha\beta} \right) v_\beta.\tag{3.16}$$

3.3 The GS action on $U(1|1)^2/U(1)^2$

To obtain the GS action on $U(1|1)^2/U(1)^2$ we may simply set

$$0 = X_3 = Y_4 = \tilde{X}_1 = \tilde{Y}_2.\tag{3.17}$$

in the action (3.11). This is because now the group element g given in equation (3.7) belongs to $U(1|1)^2 \subset U(2|2)$; this truncation is also consistent with the \mathbb{Z}_4 automorphism (3.9). As was argued at the start of this sub-section these facts imply that setting the above components to zero is a consistent truncation of the equations of motion for the

⁸The κ -action below has the nice feature of acting as a local fermionic group action by multiplication from the right. Such a representation was originally suggested in [27] and was developed more fully for the $AdS_5 \times S^5$ GS action in [28]; the formulas below are a simple extension of this latter construction to the coset at hand.

action (3.11). The GS action for the truncated theory then is

$$\begin{aligned} \mathcal{L}_{\text{GS } \mathbf{U}(1|1)^2/\mathbf{U}(1)^2} = \frac{1}{2} \int d^2\sigma \sqrt{g} g^{\mu\nu} & \left((X^\dagger \partial_\mu X + Y^\dagger \partial_\mu Y)(X^\dagger \partial_\nu X + Y^\dagger \partial_\nu Y) \right. \\ & \left. - (\tilde{X}^\dagger \partial_\mu \tilde{X} + \tilde{Y}^\dagger \partial_\mu \tilde{Y})(\tilde{X}^\dagger \partial_\nu \tilde{X} + \tilde{Y}^\dagger \partial_\nu \tilde{Y}) \right) \\ & - 2i\epsilon^{\mu\nu} \left(X^\dagger \partial_\mu \tilde{Y} Y^\dagger \partial_\nu \tilde{X} + \tilde{X}^\dagger \partial_\mu Y \tilde{Y}^\dagger \partial_\nu X \right). \end{aligned} \quad (3.18)$$

It has two $\mathbf{U}(1)$ gauge invariances

$$X \rightarrow e^{i\theta_1} X, \quad Y \rightarrow e^{i\theta_1} Y, \quad (3.19)$$

$$\tilde{X} \rightarrow e^{i\theta_2} \tilde{X}, \quad \tilde{Y} \rightarrow e^{i\theta_2} \tilde{Y}, \quad (3.20)$$

as well as κ -symmetry which is simply the restriction of equations (3.12) and (3.14).

If we parametrise the group element $g = (X, Y, \tilde{X}, \tilde{Y}) \in \mathbf{U}(1|1)^2$ by

$$X = \left(e^{it/2} \left(1 + \frac{1}{2} \psi^2 \right), 0, 0, -e^{-i\alpha/2} \bar{\psi} \right), \quad Y = \left(0, e^{it/2} \left(1 + \frac{1}{2} \eta^2 \right), -e^{-i\alpha/2} \bar{\eta}, 0 \right), \quad (3.21)$$

$$\tilde{X} = \left(0, e^{it/2} \eta, e^{-i\alpha/2} \left(1 - \frac{1}{2} \eta^2 \right), 0 \right), \quad \tilde{Y} = \left(e^{it/2} \psi, 0, 0, e^{-i\alpha/2} \left(1 - \frac{1}{2} \psi^2 \right) \right), \quad (3.22)$$

where $\psi^2 \equiv \bar{\psi}\psi$ and $\eta^2 \equiv \bar{\eta}\eta$, the action (3.18) becomes

$$\begin{aligned} \mathcal{L}_{\text{GS } \mathbf{U}(1|1)^2/\mathbf{U}(1)^2} = \int d^2\sigma \sqrt{g} g^{\mu\nu} & \left(-\partial_\mu \phi_+ \partial_\nu \phi_- + i\partial_\mu \phi_+ \eta_i \overleftrightarrow{\partial}_\nu \eta^i - \partial_\mu \phi_+ \partial_\nu \phi_+ \eta^i \eta_i \right) \\ & - e^{\mu\nu} \partial_\mu \phi_+ (\eta_1 \overleftrightarrow{\partial}_\nu \eta_2 - \eta^1 \overleftrightarrow{\partial}_\nu \eta^2), \end{aligned} \quad (3.23)$$

This action was postulated in [29] to be a consistent truncation of the full Type IIB GS action on $AdS_5 \times S^5$, by checking the absence of certain cubic terms in the latter action, using an explicit non-unitary representation for $PSU(2,2|4)$. Here we have shown that on group-theoretic grounds this action is indeed such a consistent truncation, and have obtained its form using a unitary representation of the group.

On the local coordinates defined above κ -symmetry acts as

$$\delta\eta_i = \epsilon_i, \quad \delta t = -\delta\alpha = i(\eta^i \epsilon_i + \eta_i \epsilon^i). \quad (3.24)$$

In particular notice that $\delta\phi_+ = 0$. The parameters ϵ_i are not however free, instead they are given by

$$\epsilon_j = \frac{i}{2} \left(\eta_i \overleftrightarrow{\partial}_\alpha \eta^i + i\partial_\alpha \phi_- + i\eta^i \eta_i \partial_\alpha \phi_+ \right) \kappa_j^\alpha. \quad (3.25)$$

Above κ_i^α are complex-valued Grassmann functions of the world-sheet; their complex conjugates are denoted by $\kappa^{\alpha i}$. We will also require that the metric vary under κ -symmetry

as

$$\begin{aligned}
 \delta(\sqrt{-g}g^{\alpha\beta}) &= -\frac{i}{2} \left[\kappa^{(\alpha} P_+^{\beta)\gamma} (-\eta_1 \partial_\gamma \phi_+ - i\eta^2 \partial_\gamma \phi_+ + 2i\partial_\gamma \eta_1 + 2\partial_\gamma \eta^2) \right. \\
 &\quad + \bar{\kappa}^{(\alpha} P_+^{\beta)\gamma} (-\eta^1 \partial_\gamma \phi_+ + i\eta_2 \partial_\gamma \phi_+ - 2i\partial_\gamma \eta^1 + 2\partial_\gamma \eta_2) \\
 &\quad + \tilde{\kappa}^{(\alpha} P_-^{\beta)\gamma} (\eta_1 \partial_\gamma \phi_+ - i\eta^2 \partial_\gamma \phi_+ - 2i\partial_\gamma \eta_1 + 2\partial_\gamma \eta^2) \\
 &\quad \left. + \bar{\tilde{\kappa}}^{(\alpha} P_-^{\beta)\gamma} (\eta^1 \partial_\gamma \phi_+ + i\eta_2 \partial_\gamma \phi_+ + 2i\partial_\gamma \eta^1 + 2\partial_\gamma \eta_2) \right] \\
 &= -\frac{i}{2} \left[\sqrt{-g}g^{\alpha\gamma} \left(\kappa^{\beta i} \partial_\gamma \eta_i + \kappa_i^\beta \partial_\gamma \eta^i + \frac{i}{2} \partial_\gamma \phi_+ (\kappa^{\beta i} \eta_i - \kappa_i^\beta \eta^i) \right) \right. \\
 &\quad + i\epsilon^{\alpha\gamma} \left(\kappa_1^\beta \partial_\gamma \eta_2 + \kappa_2^\beta \partial_\gamma \eta_1 - \kappa^{\beta 1} \partial_\gamma \eta^2 - \kappa^{\beta 2} \partial_\gamma \eta^1 \right) \\
 &\quad \left. - \frac{1}{2} \epsilon^{\alpha\gamma} \partial_\gamma \phi_+ \left(\kappa_1^\beta \eta_2 + \kappa_2^\beta \eta_1 + \kappa^{\beta 1} \eta^2 + \kappa^{\beta 2} \eta^1 \right) \right]. \quad (3.26)
 \end{aligned}$$

where $a^{(\alpha} b^{\beta)} = a^\alpha b^\beta + a^\beta b^\alpha$ and

$$\kappa_1^\alpha = \frac{i}{2} (\bar{\tilde{\kappa}}^\alpha - \bar{\kappa}^\alpha), \quad \kappa_2^\alpha = \frac{1}{2} (\tilde{\kappa}^\alpha + \kappa^\alpha), \quad (3.27)$$

with the complex conjugates defined as $\kappa^\dagger \equiv \bar{\kappa}$ and $\tilde{\kappa}^\dagger \equiv \bar{\tilde{\kappa}}$. The above variation of the metric is symmetric and since $\sqrt{-g}g^{\alpha\beta}$ has unit determinant (is uni-modular) we require that

$$\kappa^\alpha = P_+^{\alpha\beta} \kappa_\beta, \quad \tilde{\kappa}^\alpha = P_-^{\alpha\beta} \tilde{\kappa}_\beta. \quad (3.28)$$

Using the above formulas one can check that the action (3.23) is indeed invariant under this symmetry. However, as we show below this local symmetry is trivial on-shell.

3.4 Fake κ -symmetry

In this sub-section we show that κ -symmetry acts trivially on-shell on the fermionic GS actions studied in this section. To see this most easily we will first consider the particle limit (in other words we remove all σ dependence of fields) for the action $\mathcal{L}_{\text{GS } \text{U}(1|1)^2/\text{U}(1)^2}$. This gives

$$\mathcal{L}_{\text{particle}} = - \int d\tau e^{-1} \dot{\phi}_+ \left(\dot{\phi}_- + \dot{\phi}_+ \eta^i \eta_i - i\eta_i \dot{\eta}^i - i\eta^i \dot{\eta}_i \right) = - \int d\tau e^{-1} \dot{\phi}_+ a, \quad (3.29)$$

where for convenience we have defined⁹

$$a = \left(\dot{\phi}_- + \dot{\phi}_+ \eta^i \eta_i - i\eta_i \dot{\eta}^i - i\eta^i \dot{\eta}_i \right). \quad (3.30)$$

Setting $e = \text{constant}$, we may solve the the ϕ_+ , ϕ_- and η_i equations of motion to get

$$\phi_+ = 2\kappa\tau, \quad \phi_- = \lambda\tau, \quad \eta_i = e^{-i\kappa\tau} \eta_{0i}, \quad (3.31)$$

⁹As an aside note that the fermion index i can now run over any number and is not restricted to $i = 1, 2$ as is the case for the super-string. This is quite typical of κ -invariant particle actions.

where κ , λ (respectively, η_{0i}) are complex constant Grassmann-even (odd) numbers.¹⁰ Finally, we turn to the equation for the einbein e which reduces to

$$\kappa\lambda = 0. \tag{3.32}$$

or in other words forces us to set either κ or λ to zero. As a result the theory consists of two sectors, one with $\kappa = 0$ and the other with $\lambda = 0$. The former sector is trivial and uninteresting as all fields apart from ϕ_- are constant and the energy is zero. The physically more relevant sector has $\lambda = 0$ and $\kappa \neq 0$.

Let us now turn to the κ invariance of the action (3.29). It is easy to see that this action is invariant under

$$\begin{aligned} \delta\phi_+ &= 0, & \delta\eta_i &= a\kappa_i, & \delta\phi_- &= ia(\eta^i\kappa_i + \eta_i\kappa^i), \\ \delta(e^{-1}) &= 2i(\dot{\eta}^i\kappa_i + \dot{\eta}_i\kappa^i) + \dot{\phi}_+(\eta^i\kappa_i + \eta_i\kappa^i), \end{aligned} \tag{3.33}$$

where κ_i are arbitrary Grassmann-odd functions of τ . Since we are free to pick the parameters κ_i one might think that we could simply gauge away the fermionic degrees of freedom using this symmetry; had the κ variations been of the form

$$\delta\eta_i = \kappa_i,$$

we would have been able to gauge away the fermions. In fact this is not the case: the κ variation of the fermions instead reads

$$\delta\eta_i = a\kappa_i, \tag{3.34}$$

From the equation for the einbein e we see that in fact $a = 0$ (in the physically important sector for which $\kappa \neq 0$ as discussed above) and so on-shell the above κ symmetry acts trivially on all fields except the einbein itself. But any κ variation of the einbein e can be compensated for by a diffeomorphism. We conclude that while the actions (3.29) and (3.18) formally have a κ -symmetry, this has a trivial action on-shell and so cannot be used to eliminate any fermions.

The argument in the above paragraph relies on the fact that on fermions κ -symmetry was acting as $\delta\eta_i = a\kappa_i$ and on-shell $a = 0$. Returning to the fermionic GS superstring actions discussed in this section we see from equation (3.13) that here too κ -symmetry acts as $\delta\eta_i = a_{\text{string}}\kappa_i$, where now

$$a_{\text{string}} = (X^\dagger\partial_\alpha X + Y^\dagger\partial_\alpha Y + \tilde{X}^\dagger\partial_\alpha\tilde{X} + \tilde{Y}^\dagger\partial_\alpha\tilde{Y}). \tag{3.35}$$

It is easy to check that because of the Virasoro constraints a_{string} is also zero on-shell. We conclude that the κ -symmetry of the action (3.23) is trivial on-shell and so cannot be used to eliminate any fermions.

¹⁰In the above solution we have, without loss of generality, set the constant parts of ϕ_+ and ϕ_- to zero.

4. Large charge limits of fermionic GS actions

Given a \mathbb{Z}_4 automorphism on some coset G/H we may construct a Green-Schwarz Lagrangian for it (3.2). On general grounds the large charge limit of this Lagrangian should be a generalised Landau Lifshitz sigma model. Further, since we expect the global charges of the two actions to map onto one another, this LL sigma model should be constructed on a coset G/\tilde{H} . In this section we will attempt to identify \tilde{H} .

One step in this direction is to count the number of degrees of freedom that the GS action has and compare it with that of the LL model. For example in the case of the Type IIB superstring on $AdS_5 \times S^5$ there are 10 real bosonic degrees of freedom, and there are $32/2 = 16$ fermionic degrees of freedom (where the factor of $1/2$ comes from κ symmetry). In the large charge limit two of the bosonic degrees of freedom are eliminated; the remaining eight are 'doubled' since the LL Lagrangian should be thought of as a Lagrangian on phase space. The 16 fermions are described by coupled first order equations. When taking the LCL we integrate out half of the fermions, in order to arrive at second order equations [19], leaving us with 8 real fermionic degrees freedom; as in the case of the bosons this should also be 'doubled', leaving us with 16 fermionic degrees of freedom. At this point we may simply guess what \tilde{H} is in the case of $G = PSU(2, 2|4)$, since the only coset of the form G/\tilde{H} with 16 bosonic and fermionic degrees of freedom each is

$$\tilde{H} = PS(\mathbf{U}(1, 1|2) \times \mathbf{U}(2|2)), \tag{4.1}$$

though of course in this case \tilde{H} is well known from gauge theory.

Let us pursue this counting argument further and consider the GS action on

$$\frac{\mathbf{U}(1|1)^2}{\mathbf{U}(1)^2}. \tag{4.2}$$

This is a sub-sector of the classical GS string action on $AdS_5 \times S^5$. It has 2 real bosonic degrees of freedom and 4 real fermionic degrees of freedom. As was shown in section 3.4, κ -symmetry in this case is trivial on-shell, and so, following the counting argument in the previous paragraph,¹¹ we expect the LL sigma model corresponding to the LCL of this GS action to have 4 real fermionic degrees of freedom and no bosonic degrees of freedom. The only such coset is

$$\frac{\mathbf{U}(1|1)^2}{\mathbf{U}(1)^4}, \tag{4.3}$$

in other words $\tilde{H} = \mathbf{U}(1)^4$.

¹¹For the bosons we subtract two real degrees of freedom in the LCL and double the remaining ones. In the present case this gives $2 \times (2 - 2) = 0$ d.o.f. For the fermions, the number of d.o.f. in the LL sigma model should be the same as that of the GS string once κ -symmetry is fixed. This is because, once κ -symmetry is fixed, we halve the number of d.o.f. since the GS action gives first order differential equations, and the LL action gives second order differential equations; we then double it because the LL action is an action on phase space. In the present case, since κ -symmetry is trivial on-shell we end up with $2 \times 4/2 = 4$ fermionic d.o.f.

Similarly, we may consider the bigger sub-sector of the full classical superstring on $AdS_5 \times S^5$

$$\frac{\mathbf{U}(2|2)}{\mathbf{SU}(2)^2}, \tag{4.4}$$

for which κ -symmetry is also trivial on-shell. This sub-sector has 2 bosonic and 8 fermionic d.o.f. As a result we expect the LL sigma-model to have no bosonic d.o.f. and 8 fermionic d.o.f. Again this is enough for us to identify

$$\frac{\mathbf{U}(2|2)}{\mathbf{U}(2)^2}, \tag{4.5}$$

as the coset on which the LL sigma model is constructed. Finally, the largest classical sub-sector of the GS string action on $AdS_5 \times S^5$ for which κ -symmetry is trivial is the GS action on

$$\frac{PS(\mathbf{U}(1,1|2) \times \mathbf{U}(2|2))}{\mathbf{SU}(1,1) \times \mathbf{SU}(2)^3}. \tag{4.6}$$

By our counting argument the corresponding LCL coset should have 16 fermionic and no bosonic d.o.f. As a result, the LL sigma model which corresponds to the LCL limit of the GS action on $(\mathbf{U}(1,1|2) \times \mathbf{U}(2|2))/\mathbf{SU}(1,1) \times \mathbf{SU}(2)^3$ is constructed over the coset

$$\frac{PS(\mathbf{U}(1,1|2) \times \mathbf{U}(2|2))}{\mathbf{U}(1,1) \times \mathbf{U}(2)^3}. \tag{4.7}$$

While this counting argument shows how to identify \tilde{H} , it is not very clear how the LCL should be taken in practice and in particular how starting from a GS action one arrives at a LL action. The rest of this section will address these issues in the three cases of $G = \mathbf{U}(1|1)^2$, $\mathbf{U}(2|2)$ and $\mathbf{U}(1,1|2) \times \mathbf{U}(2|2)$. We will restrict our discussion to the leading order term in the LCL and leave the matching of sub-leading terms to a future publication.

4.1 Matching the $\mathbf{U}(1|1)^2$ sub-sectors

In this subsection we will argue that the large charge limit of the Lagrangian given in equations (3.18) and (3.23) which describes the Green-Schwarz string on the coset

$$\frac{\mathbf{U}(1|1)^2}{\mathbf{U}(1)^2}, \tag{4.8}$$

is given by the Landau-Lifshitz Lagrangian on the coset¹²

$$\frac{\mathbf{U}(1|1)^2}{\mathbf{U}(1)^4}. \tag{4.9}$$

We will first arrive at this result in a very pedestrian way. Since general solutions to both the LL and GS cosets can be given explicitly in full generality we will write them down using unconstrained coordinates. On the GS side,

$$\phi_+ = \kappa\tau, \tag{4.10}$$

¹²This is somewhat different to the comparison between gauge and string theory done in ([29]) where it was argued that on the gauge theory side the coset should be $\mathbf{U}(1|1)/\mathbf{U}(1)^2$.

the general solution takes the form

$$\eta_1 = \sum_{n=-\infty}^{\infty} e^{in\sigma} (e^{i\omega_n\tau} \psi_n^+ + e^{-i\omega_n\tau} \psi_n^-), \quad (4.11)$$

where ψ_n^\pm are constant Grassmann-odd numbers, and

$$\omega_n = \sqrt{n^2 + \kappa^2/4}. \quad (4.12)$$

η_2 is completely determined via the equation of motion

$$\partial_\sigma \eta_2 = i\partial_\tau \eta^1 - \frac{\kappa}{2} \eta^1. \quad (4.13)$$

In the LCL we take $\kappa \rightarrow \infty$ in which case we have

$$\begin{aligned} \eta_1 &\sim \sum_{n=-\infty}^{\infty} e^{in\sigma} \left(e^{i(\kappa/2+n^2/\kappa)\tau} \psi_n^+ + e^{-i(\kappa/2+n^2/\kappa)\tau} \psi_n^- \right) \\ &= e^{i\kappa\tau/2} \left[\psi_0^+ + \sum_{n=1}^{\infty} e^{in^2\tau/\kappa} (\psi_n^+ e^{in\sigma} + \psi_{-n}^+ e^{-in\sigma}) \right] \\ &\quad + e^{-i\kappa\tau/2} \left[\psi_0^- + \sum_{n=1}^{\infty} e^{-in^2\tau/\kappa} (\psi_n^- e^{in\sigma} + \psi_{-n}^- e^{-in\sigma}) \right] \\ &\equiv e^{i\kappa\tau/2} \psi_{1\text{ LL}} + e^{-i\kappa\tau/2} \bar{\psi}_{2\text{ LL}}, \end{aligned} \quad (4.14)$$

where $\psi_{1\text{ LL}}$ and $\bar{\psi}_{2\text{ LL}}$ are the 2 complex fermionic d.o.f. for the LL sigma model on (see equation (2.15))

$$\frac{\mathbf{U}(1|1)^2}{\mathbf{U}(1)^4}. \quad (4.15)$$

In particular, after rescaling $\tau \rightarrow \kappa\tau$, they satisfy the equations of motion

$$0 = (\partial_\sigma^2 - i\partial_\tau) \psi_{1,2\text{ LL}}. \quad (4.16)$$

In this way we match, to leading order in the LCL, the classical string Lagrangian with the corresponding coherent state continuum limit of the gauge theory dilatation operator in the $\mathbf{U}(1|1)^2$ sub-sector.

Notice that physical string solutions have to satisfy the level-matching condition

$$\int_0^{2\pi} \partial_1 \phi_- = 2\pi m, \quad \text{for } m \in \mathbb{Z}. \quad (4.17)$$

The winding parameter m does not, however, enter the LCL Lagrangian. Rather, it gives a constraint on its solutions. This matches the spin-chain side where m enters as a constraint on the Bethe roots, but does not enter the algebraic Bethe equations or the LL sigma-model action. This feature is very similar to the $\text{SL}(2)$ sector discussed in [30].

4.2 Large Charge Limit of fermionic GS actions

In this section we re-phrase the above discussion in terms of the embedding coordinates $X, Y \dots$, and the currents $j_\mu^{(k)}$. This allows for a straightforward generalisation from the $\mathbf{U}(1|1)^2$ sub-sector to the $\mathbf{U}(2|2)$ and $\mathbf{U}(1, 1|2) \times \mathbf{U}(2|2)$ sub-sectors. We present the explicit discussion only for the case of $\mathbf{U}(2|2)$, but the other case follows almost trivially.

The first thing to note is that the equation of motion for one of the two bosonic fields, ϕ_+ , is particularly simple in the GS models presently considered. This can be obtained as the super-trace of equation (3.3). As a result we may set

$$X^\dagger \partial_\mu X + Y^\dagger \partial_\mu Y - \tilde{X}^\dagger \partial_\mu \tilde{X} - \tilde{Y}^\dagger \partial_\mu \tilde{Y} = i\kappa \delta_{\mu,0}. \quad (4.18)$$

Using this, in conformal gauge the equation of motion for the off-diagonal component of the worldsheet metric implies that

$$X^\dagger \partial_\sigma X + Y^\dagger \partial_\sigma Y + \tilde{X}^\dagger \partial_\sigma \tilde{X} + \tilde{Y}^\dagger \partial_\sigma \tilde{Y} = 0, \quad (4.19)$$

while the fermionic equations of motion (3.4), (3.5) reduce to¹³

$$0 = \kappa(j_\tau^{(3)} - j_\sigma^{(3)}) + \dots, \quad 0 = \kappa(j_\tau^{(1)} + j_\sigma^{(1)}) + \dots. \quad (4.20)$$

As a result of these relations the WZ term does not contribute to the bosonic equation of motion (3.3).¹⁴ This fact allows us to check explicitly that the bosonic equations of motion, together with the ansatz (4.18), are consistent with the equations of motion for the metric $g_{\mu\nu}$ in conformal gauge. In fact these Virasoro constraints then imply that

$$D_\mu t = \delta_{\mu,0} \frac{\kappa}{2}, \quad \tilde{D}_\mu \alpha = -\delta_{\mu,0} \frac{\kappa}{2}. \quad (4.21)$$

As in the discussion around equation (4.17) above, the level matching condition that follows from the Virasoro constraints does not enter the LCL action.

Using equations (4.18), (4.19) and (4.20) together with a rescaling $\tau \rightarrow \kappa\tau$ we may re-write the GS Lagrangian in conformal gauge as follows

$$\begin{aligned} \mathcal{L}_{\text{GS } \mathbf{U}(2|2)/\text{SU}(2)^2} &= \eta^{\mu\nu} \text{Str}(j_\mu^{(2)} j_\nu^{(2)}) + \epsilon^{\mu\nu} \text{Str}(j_\mu^{(1)} j_\nu^{(3)}) \\ &= \eta^{\mu\nu} \left(X^\dagger \partial_\mu X + Y^\dagger \partial_\mu Y - \tilde{X}^\dagger \partial_\mu \tilde{X} - \tilde{Y}^\dagger \partial_\mu \tilde{Y} \right) \\ &\quad \times \left(X^\dagger \partial_\mu X + Y^\dagger \partial_\mu Y + \tilde{X}^\dagger \partial_\mu \tilde{X} + \tilde{Y}^\dagger \partial_\mu \tilde{Y} \right) \\ &\quad - 2 \text{Str} \left(j_\sigma^{(1)} j_\sigma^{(3)} \right) \\ &= i \left(X^\dagger \partial_\tau X + Y^\dagger \partial_\tau Y + \tilde{X}^\dagger \partial_\tau \tilde{X} + \tilde{Y}^\dagger \partial_\tau \tilde{Y} \right) - \text{STr} \left((j_\sigma^{(1)} + j_\sigma^{(3)}) (j_\sigma^{(1)} + j_\sigma^{(3)}) \right) \\ &= \mathcal{L}_{\text{LL } \mathbf{U}(2|2)/\mathbf{U}(2)^2} \end{aligned} \quad (4.22)$$

¹³In terms of $X, Y, \tilde{X}, \tilde{Y}$ this implies that we have relations of the form

$$X^\dagger \partial_\tau \tilde{Y} = i\tilde{X}^\dagger \partial_\sigma Y, \quad \tilde{X}^\dagger \partial_\tau Y = -iX^\dagger \partial_\sigma \tilde{Y}, \quad \text{etc.}$$

¹⁴This is easy to see since the WZ term's contribution to these equations is proportional to $\left[j_\tau^{(1)}, j_\sigma^{(1)} \right] - \left[j_\tau^{(3)}, j_\sigma^{(3)} \right]$. However, since $j_\tau^{(1)} = -j_\sigma^{(1)}$ and $j_\tau^{(3)} = j_\sigma^{(3)}$ each of these commutators vanishes separately.

The right-hand side of the above equation is nothing but the LL sigma model Lagrangian defined on G/\tilde{H} , where \tilde{H} is fixed under the \mathbb{Z}_2 automorphism which is the square of the \mathbb{Z}_4 automorphism used in the construction of the GS action. We have thus shown that to leading order in the LCL the fermionic GS actions constructed in section 3 above reduce to LL sigma model actions in the manner anticipated by the general argument presented at the start of the present section. It would be interesting to consider sub-leading corrections to this LCL for example in a manner similar to [15].

4.3 A gauge-theory inspired κ gauge

The GS sigma model on $AdS_5 \times S^5$ has κ -symmetry. This, as well as other symmetries of the string action, such as world-sheet diffeomorphisms, are not manifest in the corresponding spin-chain simply because this latter system keeps track only of the physical degrees of freedom. One of the challenges of defining a LCL is to identify suitable gauges for these stringy symmetries in which the physical degrees of freedom are written in the most natural coordinates for the spin-chain: while all gauges should be in principle equivalent it may be much more difficult to define a LCL between the two theories if we pick an unnatural gauge. In the previous sub-section we have defined an LCL which matches all 16 fermionic degrees of freedom from the GS action to the corresponding LL model in a very natural way. This strongly suggests what κ -gauge should be used in the full $AdS_5 \times S^5$ string action when comparing to gauge theory. Specifically it should be the gauge which keeps non-zero the 16 fermions of the coset $PS(U(1,1|2) \times U(2|2))/(SU(1,1) \times SU(2)^3)$. In fact this is the gauge used recently in [31] and the above argument can be interpreted as one motivation for their κ -gauge choice.

5. Conclusion

Over the last few years our understanding of the gauge/string correspondence has been greatly improved by using the Large Charge Limit. In this approach, the reciprocal of a large charge is employed as an expansion parameter that gives a handle on the two sides of the duality. One may compare the anomalous dimensions of gauge invariant operators with the energies of corresponding string states - this is done most efficiently by comparing the string Lagrangian in a suitable LCL with the thermodynamic limit of the spin-chain Lagrangian used to compute gauge theory anomalous dimensions. In the first part of this paper we presented a construction of a general Landau-Lifshitz sigma model on a coset G/H . On general grounds one expects that the energies of states in the LL sigma model correspond, in the thermodynamic limit, to the anomalous dimensions of the one-loop gauge theory dilatation operator. We applied this construction to give an explicit Lagrangian for the thermodynamic limit of the complete one-loop dilatation operator of $\mathcal{N} = 4$ SYM.

Using this construction we matched, for the first time, fermionic parts of the string and spin-chain Lagrangians beyond the quadratic level. In order to match the LL Lagrangian with the LCL of the GS string action, we identified a suitable κ -gauge for the GS action in which the comparison is most natural. While all κ -gauges should in principle be equivalent the presently chosen gauge does not require any complicated field redefinitions when

matching the fermions in the string and spin-chain Lagrangians, making such a comparison possible.

The above-mentioned choice of κ -gauge was identified in the following way. The GS action on $AdS_5 \times S^5$ has, at the level of classical equations of motion, a number of consistent truncations; these include the well-known truncations to various purely bosonic $AdS_p \times S^q$ subsectors as well as $AdS_p \times S^p$ super-string sub-sectors for $p = 2, 3$. In the present papers we identified a number of new sub-sectors, whose field content is that of two real bosons (which one can think of as $AdS_1 \times S^1$) and 4, 8, and 16 real fermions. What is more, we have shown that on these sub-sectors κ -symmetry acts trivially on-shell, as a result one is indeed left with 4, 8, and 16 physical fermionic degrees of freedom in these sub-sectors, and the issue of fixing a κ -gauge does not arise. We then showed how to match in the LCL these truncated GS Lagrangians, including higher-order fermionic terms, with the corresponding LL Lagrangian constructed earlier in the paper.

The dual theories have 16 real physical fermionic degrees of freedom which we have matched with one another in a straightforward way. This allows us to identify the κ -gauge for the GS action on $AdS_5 \times S^5$ most suited to the dual gauge-theory description. Namely, it is the gauge which keeps those 16 fermions which form the largest of the fermionic sub-sectors we have identified. We intend to extend this comparison to the complete one-loop Lagrangians, including higher-order mixed bosonic and fermionic terms in the near future.

Acknowledgments

I am grateful to Arkady Tseytlin for many stimulating discussions throughout this project and to Chris Hull for a number of detailed conversations on κ -symmetry. I would also like to thank Charles Young for discussions and Gleb Arutyunov for providing a copy of his notes [28]. This work is funded by EPSRC, MCOIF and is supported in part by funds provided by the U.S. Department of Energy (D.O.E.) under cooperative research agreement DEFG02-05ER41360

A. Some examples of Landau-Lifshitz sigma models

In this appendix we collect some expressions for a number of relevant Landau-Lifshitz sigma models.

A.1 The $SU(2|3)/S(U(2|2) \times U(1))$ model

The $SU(2|3)$ sub-sector sigma model Lagrangian is [18]

$$\mathcal{L}_{LL \text{ } SU(2|3)} = -iU^i \partial_\tau U_i - i\psi^\alpha \partial_\tau \psi_\alpha - \frac{1}{2} |D_\sigma U_i|^2 - \frac{1}{2} \bar{D}_\sigma \psi^a D_\sigma \psi_a + \Lambda(U_i U^i + \psi_a \psi^a - 1), \quad (\text{A.1})$$

where

$$D_\mu \equiv \partial_\mu - iC_\mu, \quad \bar{D}_\mu \equiv \partial_\mu + iC_\mu, \quad C_\mu = -iU^i \partial_\mu U_i - i\psi^a \partial_\mu \psi_a, \quad (\text{A.2})$$

and $\psi^a = \psi_a^*$ and $a = 1, 2$.

A.2 The $SU(4)/S(U(2) \times U(2))$ model

The $SO(6) \sim SU(4)$ sub-sector sigma model Lagrangian is [18]

$$\begin{aligned} \mathcal{L}_{\text{LL } SU(4)} &= \mathcal{L}_{SU(4) \text{ WZ}} - \frac{1}{8} \text{Tr}(\partial_1 m)^2 - \frac{1}{32} \text{Tr}(m \partial_1 m)^2 + \Lambda(m - m^3) \\ &= -iV^i \partial_\tau V_i - \frac{1}{2} |D_\sigma V_i|^2 + \Lambda_1(V^i V_i - 1) + \Lambda_2(V_i V_i - 1) + \Lambda_2^*(V^i V^i - 1) \end{aligned} \quad (\text{A.3})$$

where m_{ij} is a 6×6 matrix, related to V_i by

$$m_{ij} = V_i V^j - V_j V^i, \quad (\text{A.4})$$

and

$$D_\mu \equiv \partial_\mu - iC_\mu, \quad C_\mu = -iV^i \partial_\mu V_i. \quad (\text{A.5})$$

Let us define

$$M^A_B = \frac{1}{2} m_{ij} \rho^{ijA}_B, \quad m_{ij} = \frac{1}{4} \text{tr}(M \rho^{ij}), \quad (\text{A.6})$$

where ρ are the usual $SU(4)$ ρ -matrices. Notice that

$$\text{Tr} M = 0, \quad M^\dagger = M, \quad M^2 = M. \quad (\text{A.7})$$

and so we can write it as

$$M = 2X X^\dagger - 1 = -2Y Y^\dagger + 1, \quad (\text{A.8})$$

where now X and Y are 4×2 matrices which satisfy

$$X^\dagger X = \mathbf{1}_2, \quad Y^\dagger Y = \mathbf{1}_2, \quad X^\dagger Y = 0, \quad Y^\dagger X = 0, \quad (\text{A.9})$$

$$X X^\dagger + Y Y^\dagger = \mathbf{1}_4. \quad (\text{A.10})$$

Further we can write the 4×2 matrix X as two four-component vectors u_A and v_A

$$X = (u_A, v_A), \quad (\text{A.11})$$

in terms of which M^A_B can be written as

$$M^A_B = 2u^A u_B + 2v^A v_B - \delta_B^A, \quad (\text{A.12})$$

with

$$u^A u_A = 1, \quad v^A v_A = 1, \quad u^A v_A = 0. \quad (\text{A.13})$$

We can relate u_A and v_A to V_i by

$$V_i = \frac{1}{\sqrt{2}} u^A \rho_{AB}^i v^B, \quad V^i = \frac{1}{\sqrt{2}} v_A \rho^{iAB} u_B. \quad (\text{A.14})$$

It is an easy check to see that these are consistent with

$$V_i V^i = 1, \quad V_i V_i = 0, \quad M^A_B = V_i V^j \rho^{ijA}_B. \quad (\text{A.15})$$

In terms of these, the Lagrangian is

$$\begin{aligned} \mathcal{L}_{\text{LL SU}(4)} &= -iu^A \partial_0 u_A - iv^A \partial_0 v_A - \frac{1}{2} \left(\partial_1 u^A \partial_1 u_A + \partial_1 v^A \partial_1 v_A \right. \\ &\quad \left. + u^A \partial_1 u_A u^B \partial_1 u_B + v^A \partial_1 v_A v^B \partial_1 v_B + 2u^A \partial_1 v_A v^B \partial_1 u_B \right) \\ &= -i \text{Tr}(X^\dagger \partial_0 X) - \frac{1}{2} \text{Tr}(\bar{D}_1 X^\dagger D_1 X), \end{aligned} \quad (\text{A.16})$$

As before

$$X = (u_A, v_A), \quad X^\dagger \equiv \begin{pmatrix} u^A \\ v^A \end{pmatrix}, \quad (\text{A.17})$$

$$D_\mu X = \partial_\mu X - X X^\dagger \partial_\mu X. \quad (\text{A.18})$$

The action (A.16) has a local U(2) invariance

$$X \rightarrow XU(\tau, \sigma), \quad (\text{A.19})$$

for $\mathbf{U}(\tau, \sigma)$ a general U(2) matrix

$$U^\dagger(\tau, \sigma)U(\tau, \sigma) = U(\tau, \sigma)U^\dagger(\tau, \sigma) = 1_2. \quad (\text{A.20})$$

In terms of the u_A and v_A the action (A.16) is invariant with respect to the following local transformations

$$\begin{aligned} (u_A, v_A) &\rightarrow (\cos \theta(\tau, \sigma) u_A + \sin \theta(\tau, \sigma) v_A, -\sin \theta(\tau, \sigma) u_A + \cos \theta(\tau, \sigma) v_A), \\ (u_A, v_A) &\rightarrow (e^{i\phi_1(\tau, \sigma)} u_A, e^{i\phi_1(\tau, \sigma)} v_A), \\ (u_A, v_A) &\rightarrow (e^{i\phi_2(\tau, \sigma)} u_A, e^{-i\phi_2(\tau, \sigma)} v_A), \\ (u_A, v_A) &\rightarrow (e^{i\phi_3(\tau, \sigma)} v_A, -e^{i\phi_3(\tau, \sigma)} u_A), \end{aligned} \quad (\text{A.21})$$

A.2.1 Subsectors of the SU(4)/S(U(2) × U(2)) model

When written in terms of the V_i , the Lagrangian $\mathcal{L}_{\text{SU}(4)}$ can be reduced to the SU(3) sub-sector by requiring

$$V^{2a} = -iV^{2a-1} \equiv \frac{1}{\sqrt{2}} U^a, \quad a = 1, 2, 3, \quad (\text{A.22})$$

which can further be restricted to the SU(2) subsector for $V^5 = 0 = V^6$. In terms of the u_A and v_A this restriction is easily enforced by setting for example

$$u_A = (U_1, U_2, U_3, 0), \quad v_A = (0, 0, 0, 1). \quad (\text{A.23})$$

Since $U_a U^a = 1$, this choice satisfies the constraints (A.13). Restricting to the SU(2) sector is achieved by setting $u_3 = U_3 = 0$. Upon inserting these ansatze, the Lagrangian (A.16) reduces to the Lagrangian (2.16).

Another interesting sub-sector is obtained by setting

$$u_A = (U_1, U_2, 0, 0), \quad v_A = (0, 0, V_3, V_4), \quad (\text{A.24})$$

together with the conditions

$$U^1 U_1 + U^2 U_2 = 1, \quad V^1 V_1 + V^2 V_2 = 1. \quad (\text{A.25})$$

This results in $SU(2) \times SU(2)$ subsector consisting of two decoupled $SU(2)$ Landau Lifshitz Lagrangians.

A.3 The $SU(2, 2)/S(U(2) \times U(2))$ model

For later convenience we present here the $SO(2, 4)/S(O(2) \times O(4)) \sim SU(2, 2)/S(U(2) \times U(2))$ Landau-Lifshitz Lagrangian

$$\begin{aligned} \mathcal{L}_{\text{LL } SU(2,2)} &= -i\tilde{V}^i \partial_0 \tilde{V}_i - \frac{1}{2} |D_\sigma \tilde{V}_i|^2 \\ &= -i\tilde{u}^A \partial_0 \tilde{u}_A - i\tilde{v}^A \partial_0 \tilde{v}_A - \frac{1}{2} \left(\partial_1 \tilde{u}^A \partial_1 \tilde{u}_A + \partial_1 \tilde{v}^A \partial_1 \tilde{v}_A \right. \\ &\quad \left. - \tilde{u}^A \partial_1 \tilde{u}_A \tilde{u}^B \partial_1 \tilde{u}_B - \tilde{v}^A \partial_1 \tilde{v}_A \tilde{v}^B \partial_1 \tilde{v}_B - 2\tilde{u}^A \partial_1 \tilde{v}_A \tilde{v}^B \partial_1 \tilde{u}_B \right) \\ &= i\text{Tr}(\tilde{X}^\dagger \partial_0 \tilde{X}) + \frac{1}{2} \text{Tr}(\tilde{D}_1 \tilde{X}^\dagger D_1 \tilde{X}), \end{aligned} \quad (\text{A.26})$$

where

$$\tilde{V}^i \equiv \tilde{V}_j^* \eta^{ji}, \quad \text{where} \quad \eta_{ij} = \text{diag}(-1, -1, 1, 1, 1, 1), \quad (\text{A.27})$$

and

$$\tilde{u}^A \equiv \tilde{u}_B^* C^{BA}, \quad \tilde{v}^A \equiv \tilde{v}_B^* C^{BA}, \quad \text{where} \quad C_{AB} = (1, 1, -1, -1). \quad (\text{A.28})$$

The 4×2 matrix \tilde{X} has two columns

$$\tilde{X} = (\tilde{u}_A, \tilde{v}_A), \quad (\text{A.29})$$

and the covariant derivatives are

$$D_\mu \tilde{V}_i = \partial_\mu \tilde{V}_i + \tilde{V}^j \partial_\mu \tilde{V}_j \tilde{V}_i, \quad (\text{A.30})$$

$$D_\mu \tilde{X} = \partial_\mu \tilde{X} - \tilde{X} \tilde{X}^\dagger \partial_\mu \tilde{X}. \quad (\text{A.31})$$

We define

$$\tilde{X}^\dagger \equiv - \begin{pmatrix} \tilde{u}^A \\ \tilde{v}^A \end{pmatrix}. \quad (\text{A.32})$$

This is done for convenience, so that the form of the action in terms of X is independent of the signature. The fields in the Lagrangian (A.26) now satisfy the constraints

$$\tilde{X}^\dagger \tilde{X} = 1_2, \quad (\text{A.33})$$

$$\tilde{V}^i \tilde{V}_i = -1, \quad \tilde{V}_i \tilde{V}_i = 0, \quad (\text{A.34})$$

$$\tilde{u}^A \tilde{u}_A = -1, \quad \tilde{v}^A \tilde{v}_A = -1, \quad \tilde{u}^A \tilde{v}_A = 0. \quad (\text{A.35})$$

The action (A.26) has a local non-compact $U(2)$ invariance

$$\tilde{X} \rightarrow \tilde{X} U(\tau, \sigma), \quad (\text{A.36})$$

for $U(\tau, \sigma)$ a general $U(2)$ matrix

$$U^\dagger(\tau, \sigma)U(\tau, \sigma) = U(\tau, \sigma)U^\dagger(\tau, \sigma) = 1_2. \quad (\text{A.37})$$

In terms of the \tilde{u}_A and \tilde{v}_A the action (A.26) is invariant with respect to the following local transformations

$$\begin{aligned} (\tilde{u}_A, \tilde{v}_A) &\rightarrow (\cos \theta(\tau, \sigma) \tilde{u}_A + \sin \theta(\tau, \sigma) \tilde{v}_A, -\sin \theta(\tau, \sigma) \tilde{u}_A + \cos \theta(\tau, \sigma) \tilde{v}_A), \\ (\tilde{u}_A, \tilde{v}_A) &\rightarrow (e^{i\phi_1(\tau, \sigma)} \tilde{u}_A, e^{i\phi_1(\tau, \sigma)} \tilde{v}_A), \\ (\tilde{u}_A, \tilde{v}_A) &\rightarrow (e^{i\phi_2(\tau, \sigma)} \tilde{u}_A, e^{-i\phi_2(\tau, \sigma)} \tilde{v}_A), \\ (\tilde{u}_A, \tilde{v}_A) &\rightarrow (e^{i\phi_3(\tau, \sigma)} \tilde{v}_A, -e^{i\phi_3(\tau, \sigma)} \tilde{u}_A), \end{aligned} \quad (\text{A.38})$$

To relate the \tilde{V}_i coordinates to the \tilde{u}_A, \tilde{v}_A coordinates recall that the $SU(4)$ ρ matrices could be combined into 8×8 γ matrices of $SO(6)$ as follows

$$\gamma^i = \begin{pmatrix} 0 & \rho_{AB}^i \\ \rho^{iAB} & 0 \end{pmatrix}, \quad i = 1, \dots, 6, \quad (\text{A.39})$$

with the γ^i satisfying the $SO(6)$ anti-commutation relations

$$\{\gamma^i, \gamma^j\} = 2\delta^{ij}. \quad (\text{A.40})$$

The $SO(2, 4)$ γ -matrix algebra is instead

$$\{\tilde{\gamma}^i, \tilde{\gamma}^j\} = -2\eta^{ij}. \quad (\text{A.41})$$

Given a set of $SO(6)$ γ matrices we can define

$$\tilde{\gamma}^i = \begin{cases} \gamma^i, & i = 1, 2, \\ i\gamma^i, & i = 3, \dots, 6, \end{cases} \quad (\text{A.42})$$

which satisfy (A.41). Similarly we will define

$$\tilde{\rho}_{AB}^i = \begin{cases} \rho_{AB}^i, & i = 1, 2, \\ i\rho_{AB}^i, & i = 3, \dots, 6, \end{cases} \quad \text{and} \quad \tilde{\rho}^{iAB} = \begin{cases} \rho^{iAB}, & i = 1, 2, \\ i\rho^{iAB}, & i = 3, \dots, 6, \end{cases} \quad (\text{A.43})$$

which now satisfy

$$\tilde{\rho}_{AB}^i \tilde{\rho}^{jBC} + \tilde{\rho}_{AB}^j \tilde{\rho}^{iBC} = -2\delta_A^C \eta^{ij}, \quad (\text{A.44})$$

as well as

$$\eta_{ij} \tilde{\rho}_{AB}^i \tilde{\rho}^{jCD} = 2(\delta_A^C \delta_B^D - \delta_A^D \delta_B^C). \quad (\text{A.45})$$

Note also that for $SU(4)$ ρ matrices we had

$$(\rho_{AB}^i)^* = -\rho^{iAB}, \quad (\text{A.46})$$

while for the $SU(2, 2)$ $\tilde{\rho}$ matrices we have

$$(\tilde{\rho}_{AB}^i)^* = \eta^{ij} \rho^{jAB}. \quad (\text{A.47})$$

The relationship between the \tilde{v}_A, \tilde{u}_A and the \tilde{V}_i is

$$\tilde{V}_i = \frac{1}{\sqrt{2}} \tilde{u}^A \tilde{\rho}_{AB}^i \tilde{v}^B, \quad \tilde{V}^i = \frac{1}{\sqrt{2}} \tilde{v}_A \tilde{\rho}^{iAB} \tilde{u}_B. \quad (\text{A.48})$$

This can be used to derive the equality between the first and second lines in equation (A.26).

A.3.1 Subsectors of the $SU(2,2)/S(U(2) \times U(2))$ model

When written in terms of the V_i , the Lagrangian $\mathcal{L}_{\text{LL } SU(2,2)}$ can be reduced to the $SU(1,2)$ sub-sector by requiring

$$\tilde{V}^{2a} = -i\tilde{V}^{2a-1} \equiv \frac{1}{\sqrt{2}}\tilde{U}^a, \quad a = 1, 2, 3, \quad (\text{A.49})$$

which can further be restricted to the $SU(2)$ subsector for $\tilde{V}^5 = 0 = \tilde{V}^6$. In terms of the \tilde{u}_A and \tilde{v}_A this restriction is easily enforced by setting for example

$$\tilde{u}_A = (\tilde{U}_1, \tilde{U}_2, \tilde{U}_3, 0), \quad \tilde{v}_A = (0, 0, 0, 1). \quad (\text{A.50})$$

We require

$$\sum_{a=1}^3 \eta^{ab} \tilde{U}_a^* \tilde{U}_b = -1, \quad (\text{A.51})$$

so as to satisfy the constraints (A.35). Restricting to the $SU(1,1)$ sector is achieved by setting $\tilde{u}_3 = \tilde{U}_3 = 0$. Upon inserting these ansatze, the Lagrangian (A.26) reduces to the standard $SU(1,2)$ Landau-Lifshitz Lagrangian [18]

$$\mathcal{L}_{\text{SU}(1,2)} = -i\tilde{U}^a \partial_0 \tilde{U}_a - \frac{1}{2} |D_\sigma \tilde{U}_a|^2 + \Lambda(\tilde{U}^a \tilde{U}_a + 1), \quad (\text{A.52})$$

with $a = 1, 2, 3$ and $\tilde{U}^a \equiv \eta^{ab} \tilde{U}_b^*$.

A.4 The $SU(2|2)/S(U(1|1) \times U(1|1))$ model

Lets construct the LL model on $SU(2|2)/S(U(1|1) \times U(1|1))$. Starting from equation (2.4), with Tr now replaced by STr we may define

$$g = (X, Y), \quad (\text{A.53})$$

with X and Y super-matrices which satisfy

$$X^\dagger X = 1_2, \quad Y^\dagger Y = 1_2, \quad XX^\dagger + YY^\dagger = \begin{pmatrix} 1_2 & 0 \\ 0 & 1_2 \end{pmatrix}. \quad (\text{A.54})$$

The LL Lagrangian for this model is then

$$\begin{aligned} \mathcal{L}_{\text{LL } SU(2|2)} &= \frac{i}{2} \text{STr} \left[\begin{pmatrix} 1_n & 0 \\ 0 & 1_m \end{pmatrix} g^{-1} \partial_0 g \right] - \frac{1}{4} \text{STr} ((g^{-1} D_1 g)(g^{-1} D_1 g)) \\ &= i \text{STr}(X^\dagger \partial_0 X) - \frac{1}{2} \text{STr} [\bar{D}_1 X^\dagger D_1 X], \end{aligned} \quad (\text{A.55})$$

where

$$D_1 X \equiv \partial_1 X - X X^\dagger \partial_1 X. \quad (\text{A.56})$$

The bosonic base of $SU(2|2)$ is $SU(2) \times SU(2)$, where in the case of interest to us we write

$$X = (\tilde{u}_A, v_A), \quad A = 1 \dots, 4, \quad (\text{A.57})$$

with

$$\tilde{u}^A \tilde{u}_A = -1, \quad v^A v_A = 1, \quad (\text{A.58})$$

and

$$\tilde{u}^A \equiv \tilde{u}_B^* C^{BA}, \quad v^A \equiv v_B^* C^{BA}, \quad X^\dagger \equiv - \begin{pmatrix} \tilde{u}^A \\ v^A \end{pmatrix}, \quad (\text{A.59})$$

where $C^{BA} \equiv \text{diag}(-1, -1, 1, 1)$. Note that the first (last) two components of u_A (v_A) are bosonic and the last (first) two components of u_A (v_A) are fermionic.

$$\begin{aligned} \mathcal{L}_{\text{LL SU}(2|2)} = & -i\tilde{u}^A \partial_0 \tilde{u}_A - iv^A \partial_0 v_A - \frac{1}{2} \left(\partial_1 \tilde{u}^A \partial_1 \tilde{u}_A + \partial_1 v^A \partial_1 v_A \right. \\ & \left. - \tilde{u}^A \partial_1 \tilde{u}_A \tilde{u}^B \partial_1 \tilde{u}_B + v^A \partial_1 v_A v^B \partial_1 v_B + 2\tilde{u}^A \partial_1 v_A v^B \partial_1 \tilde{u}_B \right). \end{aligned} \quad (\text{A.60})$$

The action (A.60) has a local non-compact $U(1|1)$ invariance

$$\tilde{X} \rightarrow \tilde{X} U(\tau, \sigma), \quad (\text{A.61})$$

for $U(\tau, \sigma)$ a general $U(1|1)$ matrix

$$U^\dagger(\tau, \sigma) U(\tau, \sigma) = U(\tau, \sigma) U^\dagger(\tau, \sigma) = 1_2. \quad (\text{A.62})$$

In terms of the \tilde{u}_A and v_A the action (A.60) is invariant with respect to the following local transformations

$$\begin{aligned} (\tilde{u}_A, v_A) & \rightarrow (\tilde{u}_A + v_A \theta_1(\tau, \sigma), v_A + \tilde{u}_A \theta_1(\tau, \sigma)), \\ (\tilde{u}_A, v_A) & \rightarrow (\tilde{u}_A - iv_A \theta_2(\tau, \sigma), v_A + i\tilde{u}_A \theta_2(\tau, \sigma)), \\ (\tilde{u}_A, v_A) & \rightarrow (e^{i\phi_1(\tau, \sigma)} \tilde{u}_A, e^{i\phi_1(\tau, \sigma)} v_A), \\ (\tilde{u}_A, v_A) & \rightarrow (e^{i\phi_2(\tau, \sigma)} \tilde{u}_A, e^{-i\phi_2(\tau, \sigma)} v_A), \end{aligned} \quad (\text{A.63})$$

where ψ_1, ψ_2 (θ_1, θ_2) are real Grassmann-even (-odd) valued function.

B. Quantising the action (3.18) in the $t + \alpha = \kappa\tau$ gauge

Given the simple form of the action (3.18), (3.23) we present a brief light-cone quantisation of it here. The main point is that, as expected, the Hamiltonian has a non-zero normal ordering constant (B.18).

Since the equation of motion for ϕ_+ is

$$0 = \partial_\mu (\sqrt{g} g^{\mu\nu} \partial_\nu \phi_+), \quad (\text{B.1})$$

we may impose conformal gauge ($g_{\mu\nu} = \eta_{\mu\nu}$) and set

$$\phi_+ = 2\kappa\tau. \quad (\text{B.2})$$

The fermionic equations of motion then reduce to

$$0 = (i\partial_0 + \kappa)\eta^i + \partial_1 \eta_j, \quad (\text{B.3})$$

where $i \neq j$. The fermionic fields have the following periodicity conditions

$$\eta_1(\tau, 2\pi) = e^{i\alpha} \eta_1(\tau, 0), \quad \eta_2(\tau, 2\pi) = e^{-i\alpha} \eta_2(\tau, 0), \quad (\text{B.4})$$

The fermionic equations of motion then are solved by

$$\eta_1 = \sum_{n=-\infty}^{\infty} \theta_n e^{i(n\sigma + \omega_n \tau)} + \tilde{\theta}_n e^{i(n\sigma - \omega_n \tau)}, \quad (\text{B.5})$$

$$\eta_2 = \sum_{n=-\infty}^{\infty} \xi_n e^{-i(n\sigma + \omega_n \tau)} + \tilde{\xi}_n e^{-i(n\sigma - \omega_n \tau)}, \quad (\text{B.6})$$

where

$$\omega_n = \sqrt{n^2 + \frac{\kappa^2}{4}}, \quad (\text{B.7})$$

and for $n \neq 0$

$$\bar{\theta} = \frac{in}{\omega_n + \kappa} \xi_n, \quad \bar{\tilde{\theta}} = \frac{-in}{\omega_n - \kappa} \tilde{\xi}_n, \quad (\text{B.8})$$

while for $n = 0$

$$0 = \theta_0 = \tilde{\xi}_0. \quad (\text{B.9})$$

The Virasoro constraints can be used to find ϕ_- in terms of the other fields

$$0 = \partial_0 \phi_- + \frac{i}{2} \eta_i \overleftrightarrow{\partial}_0 \eta^i - \frac{\kappa}{2} \eta^i \eta_i, \quad (\text{B.10})$$

$$0 = \partial_1 \phi_- + \frac{i}{2} \eta_i \overleftrightarrow{\partial}_1 \eta^i. \quad (\text{B.11})$$

The Nöther current for time translations $t \rightarrow t + \epsilon$ is

$$j_\mu^t = -\partial_\mu \phi_+ - \partial_\mu \phi_- + i \eta_i \overleftrightarrow{\partial}_\mu \eta^i - 2 \partial_\mu \phi_+ \eta^i \eta_i - \epsilon_\mu{}^\nu \left(\eta_1 \overleftrightarrow{\partial}_\nu \eta_2 - \eta^1 \overleftrightarrow{\partial}_\nu \eta^2 \right). \quad (\text{B.12})$$

We can use the equations of motion to write the Hamiltonian of the system as

$$H_\kappa \equiv -\frac{1}{2\pi} \int d\sigma j_0^t = 2\kappa + \frac{i}{2\pi} \int d\sigma \eta_i \overleftrightarrow{\partial}_0 \eta^i. \quad (\text{B.13})$$

The canonical momentum conjugate to η_i is $4i\kappa\eta^i$ and so upon quantisation we must have

$$\{\eta^i(\tau, \sigma), \eta_j(\tau, \sigma')\} = -\frac{1}{4\kappa} \delta_j^i \delta(\sigma - \sigma'). \quad (\text{B.14})$$

As a consequence the mode oscillators have the following non-zero anti-commutators

$$\{\bar{\xi}_n, \xi_m\} = -\delta_{nm} \frac{\omega_n + \kappa}{16\pi\kappa\omega_n}, \quad \{\bar{\tilde{\xi}}_n, \tilde{\xi}_m\} = -\delta_{nm} \frac{\omega_n - \kappa}{16\pi\kappa\omega_n}, \quad (\text{B.15})$$

together with

$$\{\bar{\xi}_0, \xi_0\} = -\frac{1}{8\pi\kappa}, \quad \{\bar{\tilde{\theta}}_0, \tilde{\theta}_0\} = -\frac{1}{8\pi\kappa}, \quad (\text{B.16})$$

with all other anti-commutators equal to zero. With the convention that $\xi_n, \bar{\xi}_n, \xi_0$ and $\bar{\theta}_0$ are the annihilation operators the normal ordered expression for H_κ in the quantum theory is

$$H_\kappa = 2\kappa(\bar{\xi}_0\xi_0 + \bar{\theta}_0\tilde{\theta}_0) + 4 \sum_{n \neq 0} \omega_n^2 \left(\frac{\bar{\xi}_n\xi_n}{\omega_n + \kappa} + \frac{\tilde{\xi}_n\bar{\xi}_n}{\omega_n - \kappa} \right) + a_\kappa. \quad (\text{B.17})$$

The normal ordering constant a_κ is

$$a_\kappa = \frac{1}{4\pi\kappa} \sum_{n=-\infty}^{\infty} \omega_n. \quad (\text{B.18})$$

The remaining non-trivial bosonic Nöther current for the rotations

$$\eta_1 \rightarrow e^{i\epsilon}\eta_1, \quad \eta_2 \rightarrow e^{-i\epsilon}\eta_2, \quad (\text{B.19})$$

is

$$j_\mu^c = \partial_\mu\phi_+(\eta_1\eta^1 - \eta_2\eta^2) + 2i\eta_\mu^\nu\partial_\nu\phi_+(\eta_2\eta_1 + \eta^2\eta^1). \quad (\text{B.20})$$

The corresponding normal-ordered conserved current is

$$\begin{aligned} J &= -\frac{1}{2\pi} \int d\sigma j_0^c = \frac{\kappa}{\pi} \int d\sigma(\eta_2\eta^2 - \eta_1\eta^1) \\ &= 2\kappa\bar{\theta}_0\tilde{\theta}_0 - 2\kappa\bar{\xi}_0\xi_0 + \sum_{n \neq 0} \omega_n \left(\frac{\tilde{\xi}_n\bar{\xi}_n}{\omega_n - \kappa} - \frac{\bar{\xi}_n\xi_n}{\omega_n + \kappa} \right). \end{aligned} \quad (\text{B.21})$$

In this case the normal ordering constant is zero. Since ϕ_- is periodic in σ we require that

$$0 = \int_0^{2\pi} d\sigma \partial_1\phi_- = i \int_0^{2\pi} d\sigma \eta_i \overleftrightarrow{\partial}_1 \eta^i, \quad (\text{B.22})$$

In the quantum theory this is equivalent to the level matching requirement

$$0 = \sum_{n \neq 0} n\omega_n \left(\frac{\tilde{\xi}_n\bar{\xi}_n}{\omega_n - \kappa} - \frac{\bar{\xi}_n\xi_n}{\omega_n + \kappa} \right) | \text{physical} \rangle. \quad (\text{B.23})$$

Finally, we may compute the four non-zero supercharges

$$Q_1 \equiv i \int_0^{2\pi} \frac{d\sigma}{2\pi} Q_{04}^1 = \kappa e^{-i\kappa\tau} \frac{d\sigma}{2\pi} \eta^1 = \kappa\bar{\theta}_0, \quad (\text{B.24})$$

$$Q_2 \equiv i \int_0^{2\pi} \frac{d\sigma}{2\pi} Q_{03}^2 = \kappa e^{-i\kappa\tau} \frac{d\sigma}{2\pi} \eta^2 = \kappa\bar{\xi}_0, \quad (\text{B.25})$$

$$\bar{Q}_1 \equiv i \int_0^{2\pi} \frac{d\sigma}{2\pi} Q_{01}^4 = \kappa e^{i\kappa\tau} \frac{d\sigma}{2\pi} \eta_1 = \kappa\tilde{\theta}_0, \quad (\text{B.26})$$

$$\bar{Q}_2 \equiv i \int_0^{2\pi} \frac{d\sigma}{2\pi} Q_{02}^3 = \kappa e^{i\kappa\tau} \frac{d\sigma}{2\pi} \eta_2 = \kappa\xi_0, \quad (\text{B.27})$$

The above (super-)charges form a $\mathbf{U}(1|1)^2$ algebra and in particular we find

$$[H_\kappa, J] = 0, \quad \{Q_i, \bar{Q}_j\} = -\frac{\kappa}{8\pi} \delta_{ij}. \quad (\text{B.28})$$

C. Comments on conformal invariance of the action (3.18)

In this appendix we entertain the possibility of using the action (3.18), (3.23) as a Polyakov string action. As a warm-up let us integrate out ϕ_- in the action (3.23). We arrive at an effective action for the fermions in which we may set

$$\phi_+ = 2\kappa\tau. \quad (\text{C.1})$$

Explicitly we then have

$$\mathcal{L}_{\text{eff}} = (-\kappa) \int d^2\sigma i\eta_i \overleftrightarrow{\partial}_0 \eta^i - 2\kappa\eta^i \eta_i + \eta_1 \overleftrightarrow{\partial}_1 \eta_2 - \eta^1 \overleftrightarrow{\partial}_1 \eta^2. \quad (\text{C.2})$$

We may represent the worldsheet gamma matrices as

$$\rho^0 = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \rho^1 = \begin{pmatrix} 0 & -i \\ -i & 0 \end{pmatrix}, \quad (\text{C.3})$$

and define a world-sheet Dirac spinor as

$$\psi_\alpha = \begin{pmatrix} \eta_1 \\ \eta^2 \end{pmatrix}, \quad \alpha = 1, 2. \quad (\text{C.4})$$

The conjugate spinor is then

$$\bar{\psi}_\alpha = (\psi^\dagger \rho^0)_\alpha = (-\eta^1, \eta_2)_\alpha, \quad (\text{C.5})$$

and the effective action may be written as

$$\mathcal{L}_{\text{eff}} = (-\kappa) \int d^2\sigma i\delta_a^\mu \bar{\psi} \rho^a \overleftrightarrow{\partial}_\mu \psi + 2\kappa \bar{\psi} \psi, \quad (\text{C.6})$$

where

$$\bar{\psi} \rho^a \overleftrightarrow{\partial}_\mu \psi \equiv \bar{\psi} \rho^a \partial_\mu \psi - \partial_\mu \bar{\psi} \gamma^a \psi. \quad (\text{C.7})$$

This is simply the Lagrangian for a worldsheet Dirac fermion of mass 2κ . Since such fermions are not conformal, we get the first indication that the Lagrangian (3.18) is also not conformal.

With the above definitions for ρ^a , ψ_α and $\bar{\psi}_\alpha$ we can re-write the action (3.18) as

$$\mathcal{L} = \int d^2\sigma \sqrt{-g} \left(g^{\mu\nu} \partial_\mu \phi_+ \partial_\nu \phi_- + i e_a^\mu \bar{\psi} \rho^a \overleftrightarrow{\partial}_\mu \psi + 2m \bar{\psi} \psi \right), \quad (\text{C.8})$$

where

$$e_a^\mu = \left(g^{\mu\nu} \partial_\nu \phi_+, \frac{1}{\sqrt{-g}} \epsilon^{\mu\nu} \partial_\nu \phi_+ \right). \quad (\text{C.9})$$

We have written the above expression in the form of an inverse zweibein; we will see shortly that this is indeed justified. The corresponding zweibein is

$$e_\mu^a = (\partial_\mu \phi_+, \epsilon_{\mu\nu} \partial^\nu \phi_+), \quad (\text{C.10})$$

and the metric is

$$G_{\mu\nu} \equiv e_\mu^a e_\nu^b \eta_{ab} = -\frac{1}{m} g_{\mu\nu}. \quad (\text{C.11})$$

Above

$$m \equiv g^{\mu\nu} \partial_\mu \phi_+ \partial_\nu \phi_+ = \sqrt{-G^{\mu\nu} \partial_\mu \phi_+ \partial_\nu \phi_+}, \quad (\text{C.12})$$

is the norm of ϕ_+ , which needs to be non-zero. For completeness note that the determinant of the metric and the zwei-bein are

$$G \equiv \det G_{\mu\nu} = \frac{g}{m^2}, \quad e \equiv \det(e_\mu^a) = -\frac{\sqrt{-g}}{m}. \quad (\text{C.13})$$

Rescaling fermions in the action (C.8) by

$$\psi \rightarrow m^{-1/2} \psi, \quad (\text{C.14})$$

gives

$$\mathcal{L} = \int d^2\sigma \sqrt{-g} \left(g^{\mu\nu} \partial_\mu \phi_+ \partial_\nu \phi_- + im^{-1} e_a^\mu \bar{\psi} \rho^a \overleftrightarrow{\partial}_\mu \psi + 2\bar{\psi} \psi \right). \quad (\text{C.15})$$

Integrating by parts this can be written as

$$\begin{aligned} \mathcal{L} &= \int d^2\sigma \sqrt{-g} \left(g^{\mu\nu} \partial_\mu \phi_+ \partial_\nu \phi_- + 2im^{-1} e_a^\mu \bar{\psi} \rho^a \partial_\mu \psi + \frac{\partial_\mu (e_a^\mu \sqrt{-g} m^{-1})}{\sqrt{-g}} \bar{\psi} \rho^a \psi + 2\bar{\psi} \psi \right) \\ &= \int d^2\sigma \sqrt{-g} \left(g^{\mu\nu} \partial_\mu \phi_+ \partial_\nu \phi_- + 2im^{-1} e_a^\mu \bar{\psi} \rho^a \partial_\mu \psi + m^{-1} \frac{\partial_\mu (e_a^\mu \sqrt{G})}{\sqrt{G}} \bar{\psi} \rho^a \psi + 2\bar{\psi} \psi \right) \\ &= \int d^2\sigma \sqrt{-g} \left(g^{\mu\nu} \partial_\mu \phi_+ \partial_\nu \phi_- + 2im^{-1} e_a^\mu \bar{\psi} \rho^a \partial_\mu \psi + m^{-1} \omega_a^{01} \bar{\psi} \rho^a \psi + 2\bar{\psi} \psi \right) \\ &= \int d^2\sigma \sqrt{-G} \left(-G^{\mu\nu} \partial_\mu \phi_+ \partial_\nu \phi_- + 2\bar{\psi} (i\rho^\mu D_\mu + m) \psi \right). \end{aligned} \quad (\text{C.16})$$

The final form of the action is that of a world-sheet Dirac fermion of mass m together with the fields ϕ_\pm moving in a curved metric $G_{\mu\nu}$. Above we have used the fact that in two dimensions for any zwei-bein \hat{e}_μ^a and corresponding metric $\hat{g}_{\mu\nu}$, the spin connection $\hat{\omega}_\mu^{ab}$ can be written as

$$\hat{\omega}_\mu^{ab} = -\epsilon^{ab} \frac{1}{\sqrt{\hat{g}}} \hat{e}_\mu^c \epsilon_c^d \partial_\nu \left(\hat{e}_d^\nu \sqrt{\hat{g}} \right), \quad (\text{C.17})$$

where ϵ^{ab} (ϵ_c^d) is the flat Minkowski space ϵ -tensor with non-zero components $\epsilon^{01} = -\epsilon^{10} = 1$ ($\epsilon_0^1 = \epsilon_1^0 = -1$). This formula can be derived from the expressions presented in appendix D.

We may now want to define a string theory path integral for this Lagrangian. To do so we consider the Polyakov path integral for the Lagrangian (C.16). Since the path integral integrates over metrics $g_{\mu\nu}$, and the Lagrangian is a function of the metric $G_{\mu\nu} = -m^{-1} g_{\mu\nu}$ we first rescale

$$g_{\mu\nu} \rightarrow -m^{1/2} g_{\mu\nu}, \quad (\text{C.18})$$

in order to eliminate the metric $G_{\mu\nu}$. We arrive at a Polyakov-type path-integral with action

$$\mathcal{L} \rightarrow \int d^2\sigma \sqrt{-g} \left(g^{\mu\nu} \partial_\mu \phi_+ \partial_\nu \phi_- + 2\bar{\psi} (i\rho^\mu D_\mu + \sqrt{m}) \psi \right). \quad (\text{C.19})$$

This Lagrangian is conformally invariant. One way to see this is to generalise the argument presented in [32] which considered sigma-models on plane-wave backgrounds. Let us integrate out the fermions to obtain an effective Lagrangian for ϕ_{\pm} ¹⁵

$$\begin{aligned}
 \mathcal{L}_{\text{eff}} &\sim \eta^{\mu\nu} \partial_{\mu} \phi_{+} \partial_{\mu} \phi_{-} + \frac{i}{2} \log \det \left[-\frac{\delta^2 \mathcal{L}}{\delta \eta \delta \eta} \right] \\
 &= \eta^{\mu\nu} \partial_{\mu} \phi_{+} \partial_{\mu} \phi_{-} + \frac{i}{2} \log \det \left[\partial_{\mu_1} \phi_{+} \Pi_{+}^{\mu_1 \mu_2} \partial_{\mu_2} \partial_{\nu_1} \phi_{+} \Pi_{-}^{\nu_1 \nu_2} \partial_{\nu_2} + \frac{1}{4} m^2 \right] \\
 &= \eta^{\mu\nu} \partial_{\mu} \phi_{+} \partial_{\mu} \phi_{-} + \frac{i}{2} \det \left[\partial^2 + \frac{1}{4} m^2 \right] + \frac{i}{2} \log(m^2) \\
 &\sim \eta^{\mu\nu} \partial_{\mu} \phi_{+} \partial_{\mu} \phi_{-} + \eta^{\mu\nu} \partial_{\mu} \phi_{+} \partial_{\mu} \phi_{+} \ln \Lambda,
 \end{aligned}
 \tag{C.20}$$

where Λ is the cut-off. We can re-absorb this divergent piece by re-defining ϕ_{-}

$$\phi_{-} \rightarrow \phi_{-} - \phi_{+} \ln \Lambda.
 \tag{C.21}$$

This shows that the Lagrangian (C.19) is conformal. As it stands however, this Lagrangian is not Weyl invariant and, just as in [32], we need to turn on a dilaton

$$\Phi = \phi_{+}^2.
 \tag{C.22}$$

D. Two dimensional spin connection

Let us consider a general Lorentzian two dimensional metric $g_{\mu\nu}$ which we will parametrise for convenience as

$$g_{\mu\nu} = \begin{pmatrix} a^2 & b \\ b & d^2 \end{pmatrix},
 \tag{D.1}$$

where a , b and d are complex functions of τ and σ the coordinates on the manifold. The zweibein from which this follows is given by

$$e_{\mu}^1 = (a \sinh \rho, -d \sinh \rho), \quad e_{\mu}^2 = (a \cosh \rho, d \cosh \rho),
 \tag{D.2}$$

where

$$\cosh \frac{1}{2} \rho = \frac{b}{ad}.
 \tag{D.3}$$

The Christoffel symbols

$$\Gamma_{\nu\lambda}^{\mu} = \frac{1}{2} g^{\mu\kappa} (g_{\kappa\nu, \lambda} + g_{\kappa\lambda, \nu} - g_{\nu\lambda, \kappa})
 \tag{D.4}$$

are given by

$$\Gamma_{11}^1 = g^{-1} (aba_{,1} + ad^2 a_{,0} - bb_{,0}),
 \tag{D.5}$$

$$\Gamma_{12}^1 = \Gamma_{21}^1 = g^{-1} (d^2 aa_{,1} - bdd_{,0}),
 \tag{D.6}$$

$$\Gamma_{22}^1 = g^{-1} (d^2 b_{,1} - bdd_{,1} - d^3 d_{,0}),
 \tag{D.7}$$

$$\Gamma_{11}^2 = g^{-1} (-a^3 a_{,1} - aba_{,0} + a^2 b_{,0}),
 \tag{D.8}$$

$$\Gamma_{12}^2 = \Gamma_{21}^2 = g^{-1} (-aba_{,1} + a^2 dd_{,0}),
 \tag{D.9}$$

$$\Gamma_{22}^2 = g^{-1} (a^2 dd_{,1} - bb_{,1} + bdd_{,0}).
 \tag{D.10}$$

¹⁵I am grateful to A. Tseytlin for a number of discussions and explanations of these issues.

where $g = \det g_{\mu\nu}$. It is easy to check that these satisfy the defining equation

$$g_{\mu\nu,\lambda} - g_{\kappa\nu}\Gamma_{\mu\lambda}^{\kappa} - g_{\kappa\mu}\Gamma_{\nu\lambda}^{\kappa} = 0. \quad (\text{D.11})$$

The spin connection ω_{μ}^{mn} can be determined from the following equation

$$D_{\mu}e_{\nu}^m = \partial_{\mu}e_{\nu}^m + \omega_{\mu}^m{}_n e_{\nu}^n - \Gamma_{\mu\nu}^{\kappa} e_{\kappa}^m = 0. \quad (\text{D.12})$$

Since ω_{μ}^{mn} is anti-symmetric in (m, n) the non-zero components are given by

$$\omega_0^{01} = -\omega_0^{10} = \frac{-2a^2 da_{,1} - bda_{,0} + adb_{,0} + abd_{,0}}{2ad\sqrt{-g}}, \quad (\text{D.13})$$

$$\omega_1^{01} = -\omega_1^{10} = \frac{-bda_{,1} - adb_{,1} + abd_{,1} + 2ad^2 d_{,0}}{2ad\sqrt{-g}}. \quad (\text{D.14})$$

E. T-dual version of the action (3.18)

Performing T-duality for the action (3.18) along α leads to a very simple form for an equivalent action. In this appendix we briefly present these results. To T-dualise along α we replace $\partial_{\mu}\alpha$ by A_{α} and adding the Lagrange multiplier term $\epsilon^{\mu\nu} A_{\mu}\partial_{\nu}\tilde{\alpha}$. The A_{μ} are then integrated out and we obtain the action

$$\begin{aligned} \mathcal{L}_{\kappa}^d = & \int d^2\sigma \frac{\sqrt{g}g^{\mu\nu}}{1 - \eta^i\eta_i} \left(-(\partial_{\mu}t - \frac{i}{2}\eta_i \overleftrightarrow{\partial}_{\mu}\eta^i)(\partial_{\nu}t - \frac{i}{2}\eta_i \overleftrightarrow{\partial}_{\nu}\eta^i) \right. \\ & \left. + (\partial_{\mu}\tilde{\alpha} - (\eta_1 \overleftrightarrow{\partial}_{\mu}\eta_2 - \eta^1 \overleftrightarrow{\partial}_{\mu}\eta^2))(\partial_{\nu}\tilde{\alpha} - (\eta_1 \overleftrightarrow{\partial}_{\nu}\eta_2 - \eta^1 \overleftrightarrow{\partial}_{\nu}\eta^2)) \right) \\ & + \frac{2\epsilon^{\mu\nu}}{1 - \eta^i\eta_i} \left((\partial_{\mu}t - \frac{i}{2}\eta_i \overleftrightarrow{\partial}_{\mu}\eta^i)(\partial_{\nu}\tilde{\alpha} - (\eta_1 \overleftrightarrow{\partial}_{\nu}\eta_2 - \eta^1 \overleftrightarrow{\partial}_{\nu}\eta^2)) \right) \end{aligned} \quad (\text{E.1})$$

where we have used the fact that up to total derivatives

$$\int d^2\sigma \frac{\epsilon^{\mu\nu}}{1 - \eta^i\eta_i} \partial_{\mu}t \partial_{\nu}\tilde{\alpha} \eta^i \eta_i = \int d^2\sigma \frac{\epsilon^{\mu\nu}}{1 - \eta^i\eta_i} \partial_{\mu}t \partial_{\nu}\tilde{\alpha}. \quad (\text{E.2})$$

At the level of classical equations of motion we may integrate out the metric to get a Nambu-Goto type action

$$\mathcal{L}_{\kappa}^{dNG} = \int d^2\sigma \frac{\epsilon^{\mu\nu}}{1 - \eta^i\eta_i} \left((\partial_{\mu}t - i\eta_i \overleftrightarrow{\partial}_{\mu}\eta^i)(\partial_{\nu}\tilde{\alpha} - (\eta_1 \overleftrightarrow{\partial}_{\nu}\eta_2 - \eta^1 \overleftrightarrow{\partial}_{\nu}\eta^2)) \right), \quad (\text{E.3})$$

where we have rescaled $t \rightarrow t/2$ and multiplied the whole action by a factor of 2. The Nambu-Goto form of the action is particularly simple due to the 'two-dimensional' target space form of the action (E.1). The equations of motion for $\tilde{\alpha}$ and t imply that

$$\frac{1}{1 - \eta^i\eta_i} (\partial_{\mu}t - \frac{i}{2}\eta_i \overleftrightarrow{\partial}_{\mu}\eta^i) = \partial_{\mu}\chi_1, \quad (\text{E.4})$$

$$\frac{1}{1 - \eta^i\eta_i} (\partial_{\nu}\tilde{\alpha} - (\eta_1 \overleftrightarrow{\partial}_{\nu}\eta_2 - \eta^1 \overleftrightarrow{\partial}_{\nu}\eta^2)) = \partial_{\nu}\chi_2, \quad (\text{E.5})$$

where χ_i are arbitrary Grassmann-even functions of τ and σ . The fermion equations of motion can then be written in form notation as

$$0 = \eta_i d\chi_{1\wedge} d\chi_2 - id\eta_i \wedge d\chi_2 + id\chi_{1\wedge} d\eta^j, \quad (\text{E.6})$$

where $i \neq j$.

Let us combine the (1 + 1 dimensional) spacetime coordinates into a two-vector

$$x^i = (t, \tilde{\alpha})^i, \quad i = 1, 2, \quad (\text{E.7})$$

and represent the spacetime gamma matrices as

$$\gamma^0 = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \gamma^1 = \begin{pmatrix} 0 & -i \\ -i & 0 \end{pmatrix}, \quad (\text{E.8})$$

a spacetime Dirac spinor as

$$\Psi_\alpha = \begin{pmatrix} \eta_1 \\ \eta^2 \end{pmatrix}, \quad \alpha = 1, 2. \quad (\text{E.9})$$

The conjugate spinor is then

$$\bar{\Psi}_\alpha = (\Psi^\dagger \gamma^0)_\alpha = (-\eta^1, \eta_2)_\alpha. \quad (\text{E.10})$$

With these definitions the Nambu-Goto action can be written as

$$\mathcal{L}_\kappa^{dNG} = \frac{1}{2} \int d^2\sigma \frac{\epsilon_{ij} \epsilon^{\mu\nu}}{1 + \Psi^2} \Pi_\mu^i \Pi_\nu^j, \quad (\text{E.11})$$

where we define

$$\Pi_\mu^i \equiv (\partial_\mu x^i - i\bar{\Psi}\gamma^i \overleftrightarrow{\partial}_\mu \Psi), \quad \Psi^2 = \bar{\Psi}\Psi, \quad (\text{E.12})$$

and

$$\bar{\Psi}\gamma^i \overleftrightarrow{\partial}_\mu \Psi \equiv \bar{\Psi}\gamma^i \partial_\mu \Psi - \partial_\mu \bar{\Psi}\gamma^i \Psi. \quad (\text{E.13})$$

References

- [1] J.M. Maldacena, *The large- N limit of superconformal field theories and supergravity*, *Adv. Theor. Math. Phys.* **2** (1998) 231 [*Int. J. Theor. Phys.* **38** (1999) 1113] [[hep-th/9711200](#)]; S.S. Gubser, I.R. Klebanov and A.M. Polyakov, *Gauge theory correlators from non-critical string theory*, *Phys. Lett.* **B 428** (1998) 105 [[hep-th/9802109](#)].
- [2] E. Witten, *Anti-de Sitter space and holography*, *Adv. Theor. Math. Phys.* **2** (1998) 253 [[hep-th/9802150](#)].
- [3] D. Berenstein, J.M. Maldacena and H. Nastase, *Strings in flat space and pp waves from $N = 4$ super Yang-Mills*, *JHEP* **04** (2002) 013 [[hep-th/0202021](#)].
- [4] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, *A semi-classical limit of the gauge/string correspondence*, *Nucl. Phys.* **B 636** (2002) 99 [[hep-th/0204051](#)].

- [5] S. Frolov and A.A. Tseytlin, *Semiclassical quantization of rotating superstring in $AdS_5 \times S^5$* , *JHEP* **06** (2002) 007 [[hep-th/0204226](#)]; *Multi-spin string solutions in $AdS_5 \times S^5$* , *Nucl. Phys.* **B 668** (2003) 77 [[hep-th/0304255](#)]; *Quantizing three-spin string solution in $AdS_5 \times S^5$* , *JHEP* **07** (2003) 016 [[hep-th/0306130](#)]; *Rotating string solutions: AdS/CFT duality in non-supersymmetric sectors*, *Phys. Lett.* **B 570** (2003) 96 [[hep-th/0306143](#)]; G. Arutyunov, S. Frolov, J. Russo and A.A. Tseytlin, *Spinning strings in $AdS_5 \times S^5$ and integrable systems*, *Nucl. Phys.* **B 671** (2003) 3 [[hep-th/0307191](#)]; G. Arutyunov, J. Russo and A.A. Tseytlin, *Spinning strings in $AdS_5 \times S^5$: new integrable system relations*, *Phys. Rev.* **D 69** (2004) 086009 [[hep-th/0311004](#)]; N. Beisert, S. Frolov, M. Staudacher and A.A. Tseytlin, *Precision spectroscopy of AdS/CFT*, *JHEP* **10** (2003) 037 [[hep-th/0308117](#)].
- [6] J.A. Minahan and K. Zarembo, *The Bethe-ansatz for $N = 4$ super Yang-Mills*, *JHEP* **03** (2003) 013 [[hep-th/0212208](#)]; N. Beisert, C. Kristjansen and M. Staudacher, *The dilatation operator of $N = 4$ super Yang-Mills theory*, *Nucl. Phys.* **B 664** (2003) 131 [[hep-th/0303060](#)]; N. Beisert, J.A. Minahan, M. Staudacher and K. Zarembo, *Stringing spins and spinning strings*, *JHEP* **09** (2003) 010 [[hep-th/0306139](#)]; N. Beisert, *The complete one-loop dilatation operator of $N = 4$ super Yang-Mills theory*, *Nucl. Phys.* **B 676** (2004) 3 [[hep-th/0307015](#)]; N. Beisert and M. Staudacher, *The $N = 4$ SYM integrable super spin chain*, *Nucl. Phys.* **B 670** (2003) 439 [[hep-th/0307042](#)]; N. Beisert, *The $SU(2|3)$ dynamic spin chain*, *Nucl. Phys.* **B 682** (2004) 487 [[hep-th/0310252](#)].
- [7] A.V. Belitsky, A.S. Gorsky and G.P. Korchemsky, *Gauge/string duality for QCD conformal operators*, *Nucl. Phys.* **B 667** (2003) 3 [[hep-th/0304028](#)]; A.V. Belitsky, S.E. Derkachov, G.P. Korchemsky and A.N. Manashov, *Superconformal operators in $N = 4$ super-Yang-Mills theory*, *Phys. Rev.* **D 70** (2004) 045021 [[hep-th/0311104](#)]; *Quantum integrability in (super) Yang-Mills theory on the light-cone*, *Phys. Lett.* **B 594** (2004) 385 [[hep-th/0403085](#)].
- [8] J.A. Minahan and K. Zarembo, *The Bethe-ansatz for $N = 4$ super Yang-Mills*, *JHEP* **03** (2003) 013 [[hep-th/0212208](#)]; V.A. Kazakov, A. Marshakov, J.A. Minahan and K. Zarembo, *Classical / quantum integrability in AdS/CFT*, *JHEP* **05** (2004) 024 [[hep-th/0402207](#)]; N. Beisert, V. Dippel and M. Staudacher, *A novel long range spin chain and planar $N = 4$ super Yang-Mills*, *JHEP* **07** (2004) 075 [[hep-th/0405001](#)]; G. Arutyunov, S. Frolov and M. Staudacher, *Bethe ansatz for quantum strings*, *JHEP* **10** (2004) 016 [[hep-th/0406256](#)]; V.A. Kazakov and K. Zarembo, *Classical/quantum integrability in non-compact sector of AdS/CFT*, *JHEP* **10** (2004) 060 [[hep-th/0410105](#)]; N. Beisert, V.A. Kazakov and K. Sakai, *Algebraic curve for the $SO(6)$ sector of AdS/CFT*, *Commun. Math. Phys.* **263** (2006) 611 [[hep-th/0410253](#)]; S. Schafer-Nameki, *The algebraic curve of 1-loop planar $N = 4$ SYM*, *Nucl. Phys.* **B 714** (2005) 3 [[hep-th/0412254](#)]; N. Beisert and M. Staudacher, *Long-range PSU(2, 2|4) Bethe ansätze for gauge theory and strings*, *Nucl. Phys.* **B 727** (2005) 1 [[hep-th/0504190](#)]; S. Frolov, J. Plefka and M. Zamaklar, *The $AdS_5 \times S^5$ superstring in light-cone gauge and its Bethe equations*, *J. Phys.* **A 39** (2006) 13037 [[hep-th/0603008](#)].

- [9] N. Beisert, V.A. Kazakov, K. Sakai and K. Zarembo, *The algebraic curve of classical superstrings on $AdS_5 \times S^5$* , *Commun. Math. Phys.* **263** (2006) 659 [[hep-th/0502226](#)]; *Complete spectrum of long operators in $N = 4$ SYM at one loop*, *JHEP* **07** (2005) 030 [[hep-th/0503200](#)].
- [10] M. Staudacher, *The factorized S-matrix of CFT/ads*, *JHEP* **05** (2005) 054 [[hep-th/0412188](#)]; N. Beisert, *An $SU(1|1)$ -invariant S-matrix with dynamic representations*, *Bulg. J. Phys.* **33S1** (2006) 371 [[hep-th/0511013](#)]; *The $SU(2|2)$ dynamic S-matrix*, [hep-th/0511082](#); R.A. Janik, *The $AdS_5 \times S^5$ superstring worldsheet S-matrix and crossing symmetry*, *Phys. Rev. D* **73** (2006) 086006 [[hep-th/0603038](#)]; G. Arutyunov and S. Frolov, *On $AdS_5 \times S^5$ string S-matrix*, *Phys. Lett. B* **639** (2006) 378 [[hep-th/0604043](#)]; J. Plefka, F. Spill and A. Torrielli, *On the Hopf algebra structure of the AdS/CFT S-matrix*, *Phys. Rev. D* **74** (2006) 066008 [[hep-th/0608038](#)]; N. Beisert, R. Hernandez and E. Lopez, *A crossing-symmetric phase for $AdS_5 \times S^5$ strings*, *JHEP* **11** (2006) 070 [[hep-th/0609044](#)]; N. Beisert, B. Eden and M. Staudacher, *Transcendentality and crossing*, *J. Stat. Mech.* (2007) P01021 [[hep-th/0610251](#)]; T. Klose, T. McLoughlin, R. Roiban and K. Zarembo, *Worldsheet scattering in $AdS_5 \times S^5$* , *JHEP* **03** (2007) 094 [[hep-th/0611169](#)]; N. Dorey, D.M. Hofman and J. Maldacena, *On the singularities of the magnon S-matrix*, [hep-th/0703104](#).
- [11] S. Randjbar-Daemi, A. Salam and J.A. Strathdee, *Generalized spin systems and sigma models*, *Phys. Rev. B* **48** (1993) 3190 [[hep-th/9210145](#)].
- [12] A. Perelomov, *Generalized coherent states and their applications*, Springer, Berlin, Germany (1986); W.M. Zhang, D.H. Feng and R. Gilmore, *Coherent states: theory and some applications*, *Rev. Mod. Phys.* **62** (1990) 867.
- [13] M. Kruczenski, *Spin chains and string theory*, *Phys. Rev. Lett.* **93** (2004) 161602 [[hep-th/0311203](#)].
- [14] M. Kruczenski, A.V. Ryzhov and A.A. Tseytlin, *Large spin limit of $AdS_5 \times S^5$ string theory and low energy expansion of ferromagnetic spin chains*, *Nucl. Phys. B* **692** (2004) 3 [[hep-th/0403120](#)].
- [15] M. Kruczenski and A.A. Tseytlin, *Semiclassical relativistic strings in S^5 and long coherent operators in $N = 4$ SYM theory*, *JHEP* **09** (2004) 038 [[hep-th/0406189](#)].
- [16] R. Hernandez and E. Lopez, *The $SU(3)$ spin chain sigma model and string theory*, *JHEP* **04** (2004) 052 [[hep-th/0403139](#)].
- [17] R. Hernandez and E. Lopez, *Spin chain sigma models with fermions*, *JHEP* **11** (2004) 079 [[hep-th/0410022](#)].
- [18] B. Stefański and A.A. Tseytlin, *Large spin limits of AdS/CFT and generalized Landau-Lifshitz equations*, *JHEP* **05** (2004) 042 [[hep-th/0404133](#)].
- [19] B. Stefański and A.A. Tseytlin, *Super spin chain coherent state actions and $AdS_5 \times S^5$ superstring*, *Nucl. Phys. B* **718** (2005) 83 [[hep-th/0503185](#)].

- [20] A. Mikhailov, *Speeding strings*, *JHEP* **12** (2003) 058 [[hep-th/0311019](#)]; *Slow evolution of nearly-degenerate extremal surfaces*, *J. Geom. Phys.* **54** (2005) 228 [[hep-th/0402067](#)]; *Supersymmetric null-surfaces*, *JHEP* **09** (2004) 068 [[hep-th/0404173](#)]; *Notes on fast moving strings*, *Commun. Math. Phys.* **264** (2006) 683 [[hep-th/0409040](#)].
- [21] I. Bena, J. Polchinski and R. Roiban, *Hidden symmetries of the $AdS_5 \times S^5$ superstring*, *Phys. Rev. D* **69** (2004) 046002 [[hep-th/0305116](#)].
- [22] L.F. Alday, G. Arutyunov and A.A. Tseytlin, *On integrability of classical superstrings in $AdS_5 \times S^5$* , *JHEP* **07** (2005) 002 [[hep-th/0502240](#)].
- [23] W. Siegel, *Hidden local supersymmetry in the supersymmetric particle action*, *Phys. Lett. B* **128** (1983) 397.
- [24] M.B. Green and J.H. Schwarz, *Covariant description of superstrings*, *Phys. Lett. B* **136** (1984) 367; *Properties of the covariant formulation of superstring theories*, *Nucl. Phys. B* **243** (1984) 285.
- [25] M. Henneaux and L. Mezincescu, *A sigma model interpretation of Green-Schwarz covariant superstring action*, *Phys. Lett. B* **152** (1985) 340.
- [26] R.R. Metsaev and A.A. Tseytlin, *Type IIB superstring action in $AdS_5 \times S^5$ background*, *Nucl. Phys. B* **533** (1998) 109 [[hep-th/9805028](#)].
- [27] I.N. McArthur, *Kappa-symmetry of Green-Schwarz actions in coset superspaces*, *Nucl. Phys. B* **573** (2000) 811 [[hep-th/9908045](#)].
- [28] G. Arutyunov, unpublished notes.
- [29] L.F. Alday, G. Arutyunov and S. Frolov, *New integrable system of 2dim fermions from strings on $AdS_5 \times S^5$* , *JHEP* **01** (2006) 078 [[hep-th/0508140](#)].
- [30] I.Y. Park, A. Tirziu and A.A. Tseytlin, *Spinning strings in $AdS_5 \times S^5$: one-loop correction to energy in $SL(2)$ sector*, *JHEP* **03** (2005) 013 [[hep-th/0501203](#)].
- [31] S. Frolov, J. Plefka and M. Zamaklar, *The $AdS_5 \times S^5$ superstring in light-cone gauge and its Bethe equations*, *J. Phys. A* **39** (2006) 13037 [[hep-th/0603008](#)];
G. Arutyunov, S. Frolov, J. Plefka and M. Zamaklar, *The off-shell symmetry algebra of the light-cone $AdS_5 \times S^5$ superstring*, *J. Phys. A* **40** (2007) 3583 [[hep-th/0609157](#)].
- [32] G. Papadopoulos, J.G. Russo and A.A. Tseytlin, *Solvable model of strings in a time-dependent plane-wave background*, *Class. and Quant. Grav.* **20** (2003) 969 [[hep-th/0211289](#)].