

# On thermodynamics of $\mathcal{N} = 6$ superconformal Chern-Simons theories at strong coupling

---

**Ahmad Ghodsi, Mohammad R. Garousi and Mehran Khosravi**

*Department of Physics, Ferdowsi University, P.O. Box 1436, Mashhad, Iran, and  
Institute for Studies in Theoretical Physics and Mathematics (IPM),*

*P.O. Box 19395-5531, Tehran, Iran*

*E-mail: [ahmad@mail.ipm.ir](mailto:ahmad@mail.ipm.ir), [garousi@mail.ipm.ir](mailto:garousi@mail.ipm.ir), [me\\_kh30@stu-mail.um.ac.ir](mailto:me_kh30@stu-mail.um.ac.ir)*

ABSTRACT: Recently it has been conjectured that  $\mathcal{N} = 6$ ,  $U(N)_k \times U(N)_{-k}$  Chern-Simons theory is dual to M-theory on  $AdS_4 \times S^7/Z_k$ . By studying one-loop correction to the M-theory effective action, we calculate the correction to the entropy of thermal field theory at strong coupling. For large  $k$  level, we have also found the  $\alpha'$  correction to the entropy from the string correction of the type IIA effective action. The structure of these two corrections at strong 't Hooft coupling are different.

KEYWORDS: M-Theory, AdS-CFT Correspondence.

---

## Contents

<b>1.</b>	<b>Introduction</b>	<b>1</b>
<b>2.</b>	<b>Review of M-theory</b>	<b>2</b>
<b>3.</b>	<b>M2-branes in flat space</b>	<b>3</b>
3.1	Entropy from the Wald formula	6
3.2	Entropy from free energy	7
<b>4.</b>	<b>M2 branes in orbifold space</b>	<b>8</b>
4.1	Large $k$ limit	10
<b>5.</b>	<b>M5-branes in flat space</b>	<b>11</b>

---

## 1. Introduction

Following the idea that the Chern-Simons gauge theory may be used to describe the coincident M2-branes [1], Bagger and Lambert [2] as well as Gustavsson [3] have constructed three dimensional  $\mathcal{N} = 8$  superconformal SO(4) Chern-Simons gauge theory based on 3-algebra. It is believed that the BLG theory at level one describes two M2-branes on  $R^8/Z_2$  orbifold [4].

The signature of the metric on 3-algebra in the BLG model is positive definite. This assumption has been relaxed in [5] to study  $N$  coincident M2-branes with  $N > 2$ . The so called BF membrane theory with arbitrary semi-simple Lie group has been proposed in [5]. This theory has ghost fields, however, the unitarity returns by gauging the global shift symmetry for the Bosonic and Fermionic ghost fields [6]. This gauged BF membrane theory is argued to be equivalent to the three dimensional  $\mathcal{N} = 8$  SYM theory [7].

Another approach to study multiple coincident M2-branes is proposed by Aharony-Bergman-Jafferis-Maldacena (ABJM) [8]. They consider a particular brane configuration which preserves  $\mathcal{N} = 3$  supersymmetry. At low energy, by integrating out the massive modes of the brane configuration one finds  $U(N)_k \times U(N)_{-k}$  Chern-Simons conformal field theory which preserves  $\mathcal{N} = 6$  supersymmetry. This theory is renormalizable and is consistent even at high energies. By lifting the brane configuration to M-theory they have shown that the gauge theory is equivalent to the low energy theory of  $N$  coincident M2-branes in orbifold  $R^8/Z_k$ . Using the AdS-CFT correspondence, then they have proposed the duality between  $\mathcal{N} = 6$  Chern-Simons theory at level  $k$  and M-theory on  $AdS_4 \times S^7/Z_k$ . For further study of the theory of multiple coincident M2-branes see [11].

In this paper, we are interested in studying the entropy of the above gauge theory at finite temperature. At weak coupling,  $\lambda \ll 1$ , the entropy is given by the free field theory [8]

$$S = 3V_2 N^2 T^2 \left( \frac{7\zeta(3)}{\pi} + O(\lambda) \right). \quad (1.1)$$

At strong coupling and at small  $k$  level, the entropy is given by the area of the Schwarzschild  $AdS_4 \times S^7/Z_k$  [8]

$$S = V_2 T^2 \frac{2^{\frac{7}{2}}}{3^3} \pi^2 N^2 \frac{1}{\sqrt{\lambda}}. \quad (1.2)$$

The correction to the above entropy in which we are interested, is coming from one loop corrections to the Schwarzschild  $AdS_4 \times S^7/Z_k$  solution. At strong coupling and at large  $k$ , one expects similar behavior for the entropy from the type IIA supergravity. We will also find (1.2) from type IIA supergravity and its  $\alpha'$  correction. In the presence of higher derivative corrections, the entropy can be calculated using the Wald formula [9] or the free energy method [10].

An outline of the paper is as follows. In section two we briefly review the p-brane solutions of M-theory at tree-level. In third section we compute the corrections to the Schwarzschild  $AdS_4 \times S^7$  in the presence of higher derivative corrections to the eleven dimensional supergravity action. In subsection (3.1) we calculate the entropy using the Wald formula and in subsection (3.2) we find the same result for entropy using the free energy method. In section four, we find the entropy of the  $U(N)_k \times U(N)_{-k}$  Chern-Simons conformal field theory at strong coupling and at small  $k$ , which is given by the the entropy of Schwarzschild  $AdS_4 \times S^7/Z_k$  at one-loop. In subsection (4.1), we calculate the entropy of the thermal field theory at strong coupling and at large level  $k$  from studying the type IIA supergravity, and find its  $\alpha'$  correction. Finally in the last section, we calculate the entropy of Schwarzschild  $AdS_7 \times S^4$  at one-loop which is the entropy of the world-volume theory of  $N$  coincident M5-branes at strong coupling.

## 2. Review of M-theory

The eleven dimensional supergravity action and its one loop corrections is given by

$$S = \frac{1}{2\kappa_{11}^2} \int d^{11}x \sqrt{-g} \left( R - \frac{1}{2n!} F_{(n)}^2 + \gamma W \right), \quad (2.1)$$

where  $\gamma = 4\pi^2 \kappa_{11}^{\frac{4}{3}}/3$  and  $W$  in terms of the Weyl tensors is

$$W = C^{hmnk} C_{pmnq} C_h{}^{rsp} C^q{}_{rsk} + \frac{1}{2} C^{hkmm} C_{pqmn} C_h{}^{rsp} C^q{}_{rsk}. \quad (2.2)$$

One set of solutions to the tree-level part of the above action is the non-extremal p-brane solutions in  $D = 11$  dimensional space-time (see e.g. [13])

$$ds^2 = H(r)^{-\frac{d-2}{D-2}} \left( -f(r) dt^2 + \sum_{i=1}^p (dx^i)^2 \right) + H(r)^{\frac{p+1}{D-2}} \left( f(r)^{-1} dr^2 + r^2 d\Omega_{d-1}^2 \right), \quad (2.3)$$

$$F_{ti_1 \dots i_p r} = \epsilon_{ti_1 \dots i_p r} H(r)^{-2} \frac{Q}{r^{d-1}},$$

where  $D = (p + 1) + d$  and  $d$  is the number of dimensions transverse to the p-branes. The functions  $H$  and  $f$  are  $H(r) = 1 + (\frac{L}{r})^{d-2}$  and  $f(r) = 1 - (\frac{r_0}{r})^{d-2}$ . In the above equation the relation between  $L$  and  $Q$  is

$$L^{2(d-2)} + L^{d-2}r_0^{d-2} = \frac{Q^2}{(d-2)^2}. \tag{2.4}$$

For  $r_0 = 0$  we obtain the extremal solution, depending only on a single parameter,  $Q$ , related to the charge density of the BPS p-brane. For  $r_0 \neq 0$  a horizon develops at  $r = r_0$ . To obtain the near horizon geometry, we take the limit  $r \ll L$ . In this limit the relation (2.4) simplifies to  $L^{d-2} = Q/(d-2)$ , and the non-extremal solution becomes

$$ds^2 = \left(\frac{r}{L}\right)^{\frac{2(d-2)}{p+1}} \left[ - \left(1 - \left(\frac{r_0}{r}\right)^{d-2}\right) dt^2 + \sum_{i=1}^p (dx^i)^2 \right] + \frac{\left(\frac{L}{r}\right)^2}{\left(1 - \left(\frac{r_0}{r}\right)^{d-2}\right)} dr^2 + L^2(d\Omega_{d-1})^2, \\ F_{ti_1 \dots i_p r} = (d-2)\epsilon_{ti_1 \dots i_p r} \frac{r^{d-3}}{h^{d-2}}, \tag{2.5}$$

which is the product space of  $S^{d-1}$  with the Schwarzschild black hole in  $AdS_{D-d+1}$ . According to the AdS/CFT correspondence, there is a finite temperature field theory dual to the M-theory on Schwarzschild black hole in  $AdS_{D-d+1} \times S^{d-1}$ .

### 3. M2-branes in flat space

The field theory of large  $N$  M2-branes in flat space is dual to M-theory on  $AdS_4 \times S^7$  and its finite temperature is dual to M-theory on Schwarzschild  $AdS_4 \times S^7$ . At strong coupling, the field theory is dual to eleven dimensional supergravity on Schwarzschild  $AdS_4 \times S^7$ . The entropy and the temperature of field theory are given by the corresponding quantities in Schwarzschild  $AdS_4 \times S^7$ ,

$$S = V_2 T^2 \frac{2^{\frac{7}{2}}}{3^3} \pi^2 N^{\frac{3}{2}}, \tag{3.1}$$

which is the area of the horizon of Schwarzschild  $AdS_4 \times S^7$ . The background and subsequently the entropy and temperature of the field theory are modified by the one-loop corrections. By including the higher derivative terms one needs to consider a general ansatz for the metric and field strength, and then finds the deformations of the near horizon geometry of M2-branes. We start with the following ansatz for the metric

$$ds^2 = S^{2n}(r)H^2(r) \left( K^2(r)d\tau^2 + P^2(r)dr^2 + \sum_{i=1}^2 dx_i^2 \right) + L^2 S^2(r) d\Omega_7^2, \tag{3.2}$$

where  $H(r) = \frac{r}{L}$ . At tree-level, the above functions are

$$K(r) = \left(1 - \frac{r_0^3}{r^3}\right)^{\frac{1}{2}}, \quad P(r) = \frac{L^2}{2r^2} \left(1 - \frac{r_0^3}{r^3}\right)^{-\frac{1}{2}}, \quad S(r) = 1, \tag{3.3}$$

where we have made the space-time Euclidean and changed the variable  $r$  in (2.5) to  $r^2/L$ . The electric field strength at tree-level is also given by  $F_{\tau x_1 x_2 r} = 3\frac{r^2}{L^3}$ . Considering the fact that the charge of membranes is independent of the loop corrections, we restrict the field strength of M2-branes to the following form

$$F_{\tau x_1 x_2 r} = \frac{6r^4}{L^5} K(r)P(r)S^{4n-7}(r). \quad (3.4)$$

Using the above ansatz, one writes the action (2.1) as an integration over  $r$

$$I = -\frac{\beta V_2 V_7}{2\kappa_{11}^2} \int dr \left( \mathcal{L} + \gamma \mathcal{W} \right), \quad (3.5)$$

where

$$\mathcal{L} = S^{4n+7} r^4 L^3 K P \left( R - \frac{1}{2} \frac{1}{4!} F_{(4)}^2 \right), \quad \mathcal{W} = S^{4n+7} r^4 L^3 K P W. \quad (3.6)$$

The Euler-Lagrange equations are

$$\frac{\partial \mathcal{L}}{\partial \Phi} - \frac{d}{dr} \frac{\partial \mathcal{L}}{\partial \Phi'} + \frac{d^2}{dr^2} \frac{\partial \mathcal{L}}{\partial \Phi''} = -\gamma \left( \frac{\partial \mathcal{W}}{\partial \Phi} - \frac{d}{dr} \frac{\partial \mathcal{W}}{\partial \Phi'} + \frac{d^2}{dr^2} \frac{\partial \mathcal{W}}{\partial \Phi''} \right) \equiv -\gamma \omega_{\Phi}, \quad (3.7)$$

where  $(\Phi = \{K, P, S\})$ . The left hand side can be computed by inserting the ansatz (3.2) and (3.4) into the supergravity action. The right hand side of the Euler-Lagrange equation belong to the next order of perturbations so we just need to compute it by inserting the tree-level solutions. The value of  $W$  is

$$W = \frac{1152}{L^8} \left( \frac{r_0^{12}}{r^{12}} - \frac{7}{50} \frac{r_0^6}{r^6} + \frac{1533}{4000} \right). \quad (3.8)$$

In the above relation the first term is coming from  $AdS_4$  and the other terms result from the fact that we have considered the eleven dimensional metric in which the radii of  $AdS_4$  and  $S^7$  are not equal. These terms makes the entropy to be different from the one that has been found in [12] in which only the  $AdS_4$  part has been considered.<sup>1</sup>

The right hand sides for the Euler-Lagrange equations (3.7) are given by

$$\begin{aligned} \omega_K &= -\frac{6}{25} \frac{1}{L^3 r^{\frac{17}{2}} (r^3 - r_0^3)^{\frac{1}{2}}} \left( 1533r^{12} + 3136r_0^6 r^6 + 129472r^3 r_0^9 - 160800r_0^{12} \right), \\ \omega_P &= -\frac{12}{25} \frac{(r^3 - r_0^3)^{\frac{1}{2}}}{L^5 r^{\frac{19}{2}}} \left( 1533r^{12} - 3136r_0^6 r^6 + 1344r^3 r_0^9 - 26400r_0^{12} \right), \\ \omega_S &= -\frac{36792}{25} \frac{1}{L^3 r^{10}} \left( \left( n - \frac{29}{20} \right) r^{12} + \frac{224}{219} \left( n - \frac{47}{56} \right) r_0^6 r^6 \right. \\ &\quad \left. - \frac{192}{73} (n-1) r^3 r_0^9 + \frac{800}{511} \left( n - \frac{7}{4} \right) r_0^{12} \right). \end{aligned} \quad (3.9)$$

---

<sup>1</sup>Notice that in D3-brane case the radii of  $AdS_5$  and  $S^5$  are equal so that the value of  $W$  for 5 dimension and 10 dimension are the same.

In order to solve the Euler-Lagrange equations we use the perturbative approach. We consider the solutions to be just some corrections to the tree-level solutions

$$\begin{aligned}
 K(r) &= \left(1 - \frac{r_0^3}{r^3}\right)^{\frac{1}{2}} (1 + \gamma X(r)) , \\
 P(r) &= \frac{L^2}{2r^2} \left(1 - \frac{r_0^3}{r^3}\right)^{-\frac{1}{2}} (1 + \gamma Y(r)) , \\
 S(r) &= (1 + \gamma Z(r)) .
 \end{aligned} \tag{3.10}$$

Inserting them into the left hand side of the Euler-Lagrange equations one finds the following differential equations for perturbed functions:

$$\begin{aligned}
 0 &= 100L^6 r^{10} \left(n + \frac{7}{2}\right) \left[ r(r^3 - r_0^3) Z''(r) + 3 \left(r^3 - \frac{r_0^3}{2}\right) Z'(r) - 3r^2 Z(r) \right] - 300L^6 r^{12} Y(r) \\
 &\quad - 100L^6 r^{10} (r^3 - r_0^3) Y'(r) + 4599r^{12} + 9408r^6 r_0^6 + 388416r^3 r_0^9 - 482400r_0^{12} , \\
 0 &= 300L^6 r^{10} \left(n + \frac{7}{2}\right) \left[ \left(r^3 - \frac{r_0^3}{2}\right) Z'(r) - r^2 Z(r) \right] - 300L^6 r^{12} Y(r) \\
 &\quad + 100L^6 r^{10} (r^3 - r_0^3) X'(r) + 4599r^{12} - 9408r^6 r_0^6 + 4032r^3 r_0^9 - 79200r_0^{12} , \\
 0 &= 1500(n^2 + 7n + 7)L^6 r^{10} \left[ r(r^3 - r_0^3) Z''(r) + 4 \left(r^3 - \frac{r_0^3}{4}\right) Z'(r) - 4 \frac{n^2 + 7n - \frac{91}{8}}{n^2 + 7n + 7} r^2 Z(r) \right] \\
 &\quad + 500 \left(n + \frac{7}{2}\right) L^6 r^{10} \left[ r(r^3 - r_0^3) X''(r) + 5 \left(r^3 - \frac{r_0^3}{10}\right) X'(r) \right. \\
 &\quad \quad \quad \left. - 3 \left(r^3 - \frac{r_0^3}{2}\right) Y'(r) - 12r^2 Y(r) \right] \\
 &\quad + (91980n - 133371)r^{12} + 94080 \left(n - \frac{47}{56}\right) r^6 r_0^6 \\
 &\quad - 241920(n - 1)r^9 r_0^3 + 144000 \left(n - \frac{7}{4}\right) r_0^{12} .
 \end{aligned} \tag{3.11}$$

There is an interesting solution to the above equations. If one considers

$$n = -\frac{7}{2}, \tag{3.12}$$

the above equations simplify drastically, i.e. the  $Z(r)$  dependence in the first two equations will be canceled. Moreover, for  $n = -\frac{7}{2}$  the area of the horizon in terms of  $r_0$  is independent of  $S(r)$ . We shall show that this simplifies the Wald formula when calculating the entropy.

To find the solution for above differential equations one should use the boundary condition at the horizon. Solving the first equation gives rise to the following exact value for  $Y(r)$

$$Y(r) = \frac{1}{L^6} \left( \frac{1533}{100} + \frac{3568}{25} \frac{r_0^3}{r^3} + \frac{2784}{25} \frac{r_0^6}{r^6} - 536 \frac{r_0^9}{r^9} \right), \tag{3.13}$$

where the constant of differential equation has been fixed by imposing the fact that the value of  $Y(r)$  at horizon,  $r = r_0$ , must be finite. Inserting the value of  $Y(r)$  into the second equation, one will find the following solution for  $X(r)$ :

$$X(r) = \frac{1}{L^6} \left( c - \frac{3568}{25} \frac{r_0^3}{r^3} - \frac{3568}{25} \frac{r_0^6}{r^6} + 88 \frac{r_0^9}{r^9} \right), \quad (3.14)$$

where  $c$  is a constant which can not be fixed by the boundary conditions. Instead, up to the first order of  $\gamma$ , it can be set to zero by time scaling. Finally replacing all the above values in the third equation, one finds the following second order differential equation for  $Z(r)$

$$0 = -7875L^6 r^{11} (r^3 - r_0^3) Z''(r) - 31500L^6 r^{10} \left( r^3 - \frac{1}{4} r_0^3 \right) Z'(r) + 141750L^6 r^{12} Z(r) - 455301r^{12} - 408240r^6 r_0^6 + 1088640r^3 r_0^9 - 756000r_0^{12}. \quad (3.15)$$

Unlike  $X(r)$  and  $Y(r)$  which have a simple series solution, here one can not find such a behavior for  $Z(r)$ . One may try to solve this differential equation with the well defined boundary conditions at the horizon and at infinity. The solution is needed for studying the Kaluza-Klein modes. However as we will see we do not need to know the exact form of  $Z(r)$  for calculating the thermodynamical quantities such as temperature and entropy in which we are interested in this paper.

### 3.1 Entropy from the Wald formula

One way of calculating the entropy in a higher derivative theory of gravity is the Wald formula [9]. It is given by

$$S = \frac{4\pi}{2\kappa_{11}^2} \int dx^H \sqrt{g^H} \frac{\partial}{\partial R_{\mu\nu\rho\lambda}} (L + \gamma W) g_{\mu\rho}^\perp g_{\nu\lambda}^\perp, \quad (3.16)$$

where the superscript  $H$  refers to the horizon and  $g^\perp$  is the orthogonal metric to the horizon. In our model the orthogonal directions are  $\tau$  and  $r$ , i.e.  $g_{rr}^\perp = g_{rr}$  and  $g_{\tau\tau}^\perp = g_{\tau\tau}$ . As we have mentioned before for  $n = -\frac{7}{2}$  the horizon area in terms of  $r_0$  does not modify by the higher derivative corrections, so the first term in above formula which is the area of horizon at leading order does not modify either. So we need only the tree-level solution to calculate the first term. On the other hand, the second term is proportional to  $\gamma$ , so to the first order of  $\gamma$  one has to replace the tree-level solution into the second term too. Therefor the first term is

$$S_1 = \frac{4\pi}{2\kappa_{11}^2} V_2 V_7 L^5 r_0^2, \quad (3.17)$$

and the second term is

$$S_2 = \frac{4\pi}{2\kappa_{11}^2} V_2 V_7 L^5 r_0^2 \left( \frac{34616}{250} \frac{\gamma}{L^6} \right). \quad (3.18)$$

The final result for entropy will be

$$S = \frac{4\pi}{2\kappa_{11}^2} V_2 V_7 L^5 r_0^2 \left( 1 + \frac{34616}{250} \frac{\gamma}{L^6} \right). \quad (3.19)$$

To write the above entropy in terms of the temperature, we recall that the temperature of the black hole is given by the surface gravity at the horizon

$$\hat{\kappa} = 2\pi T = \sqrt{g^{rr}} \frac{d}{dr} \sqrt{g_{\tau\tau}}|_{r=r_0} = \frac{1}{rP} \frac{d}{dr}(rK) + \frac{nK}{PS} \frac{d}{dr} S|_{r=r_0}. \quad (3.20)$$

As we have anticipated before the temperature is independent of  $Z(r)$  because the second term is zero. Note that at horizon  $\frac{dS}{dr} = \gamma \frac{dZ}{dr}$  is finite (3.15) and  $K/P = 0$ . However the temperature does depend on corrections of  $K$  and  $P$ . So the entropy in terms of temperature depends on the loop correction of  $K$  and  $P$  and is independent of  $Z$ . The temperature is

$$T = \frac{3}{2\pi} \frac{r_0}{L^2} \left( 1 + \frac{\gamma}{L^6} \frac{1383}{20} \right), \quad (3.21)$$

and the entropy in terms of temperature is

$$S = \frac{V_2 V_7}{2\kappa_{11}^2} \left( \frac{16}{9} \pi^3 L^9 T^2 \right) \left( 1 + \frac{41}{250} \frac{\gamma}{L^6} \right), \quad (3.22)$$

Using the relations  $L^9 = N^{\frac{3}{2}} \kappa_{11}^2 \frac{\sqrt{2}}{\pi^5}$ , where  $N$  is the total number of membranes and  $V_7 = \frac{\pi^4}{3}$  one finds

$$S = V_2 T^2 \left\{ \frac{2^{\frac{7}{2}}}{3^3} \pi^2 N^{\frac{3}{2}} + \frac{41}{250} \left( \frac{2\pi}{3} \right)^5 2^{\frac{1}{6}} 3\pi^{\frac{7}{3}} N^{\frac{1}{2}} \right\}. \quad (3.23)$$

Note that the coefficient of the last term,  $\frac{41}{250}$ , is different from the value found in [12]. This is resulted from the fact that we have considered eleven dimensional metric in which  $W$  is given by (3.8) instead of four dimensional metric considered in [12] in which  $W$  is given by the first term of (3.8). To double check our results we calculate the entropy from the free energy in the next section.

### 3.2 Entropy from free energy

Another way of finding the entropy of a higher derivative theory is to use the free energy. Following [10], one can identify the free energy of the theory with the Euclidean gravitational action,  $I$ , times the temperature,  $T$ , i.e.

$$I = \beta F, \quad (3.24)$$

where  $\beta = \frac{1}{T}$ . The calculation of the Euclidean action is divergent at large distances,  $r_{\max}$ , and requires a subtraction. The integral must be regulated by subtracting off its zero temperature limit, i.e.

$$F = \lim_{r_{\max} \rightarrow \infty} \frac{I - I_0}{\beta}, \quad (3.25)$$

where  $I_0$  is the zero temperature limit in which the periodicity of the Euclidean time is defined by  $\beta'$ . One must adjust  $\beta'$  so that the geometry at  $r = r_{\max}$  is the same in the two cases, i.e. the black hole and its zero temperature. This can be done by equating the circumference of the Euclidean time in two cases

$$\int_0^{\beta'} \sqrt{g_{\tau\tau}^{(\text{zero temperature})}}|_{r_{\max}} d\tau = \int_0^{\beta} \sqrt{g_{\tau\tau}^{(\text{black hole})}}|_{r_{\max}} d\tau. \quad (3.26)$$



Having found  $F$ , the entropy in terms of the free energy is then given by  $S = -\frac{\partial F}{\partial T}$ .

Let us now try to find the entropy of membranes with this approach. To simplify the calculations we use the observation made in the previous section that the entropy and temperature are independent of  $Z(r)$ . So we set  $Z(r)$  equal to zero in this approach. The Euclidean action and its regulator are

$$\begin{aligned}
 I &= -\frac{\beta V_2 V_7}{2\kappa_{11}^2} \int_{r_0}^{r_{\max}} dr (\mathcal{L} + \gamma \mathcal{W}) \\
 &= 4 \frac{\beta V_2 V_7}{2\kappa_{11}^2} L^3 (r_{\max}^3 - r_0^3) \left[ 1 - \frac{\gamma}{L^6} \left( \frac{32193}{500} + \frac{6968}{25} \frac{r_0^3}{r_{\max}^3} + \frac{4784}{25} \frac{r_0^6}{r_{\max}^6} + 48 \frac{r_0^9}{r_{\max}^9} \right) \right], \\
 I_0 &= -\frac{\beta' V_2 V_7}{2\kappa_{11}^2} \int_0^{r_{\max}} dr (\mathcal{L} + \gamma \mathcal{W}) \\
 &= 4 \frac{\beta' V_2 V_7}{2\kappa_{11}^2} L^3 r_{\max}^3 \left[ 1 - \frac{\gamma}{L^6} \left( \frac{32193}{500} \right) \right], \tag{3.27}
 \end{aligned}$$

and the relation between  $\beta$  and  $\beta'$  is given by  $\beta' = \beta K(r)|_{r_{\max}}$ . The free energy then becomes

$$F = \lim_{r_{\max} \rightarrow \infty} \frac{I - I_0}{\beta} = -\frac{V_2 V_7}{2\kappa_{11}^2} 2r_0^3 L^3 \left( 1 + \frac{103807}{500} \frac{\gamma}{L^6} \right), \tag{3.28}$$

In terms of temperature it becomes

$$F = -\frac{V_2 V_7}{2\kappa_{11}^2} \left( \frac{16}{27} \pi^3 L^9 T^3 \right) \left( 1 + \frac{41}{250} \frac{\gamma}{L^6} \right). \tag{3.29}$$

Finally, the entropy is given by

$$S = 3 \frac{V_2 V_7}{2\kappa_{11}^2} \left( \frac{16}{27} \pi^3 L^9 T^2 \right) \left( 1 + \frac{41}{250} \frac{\gamma}{L^6} \right). \tag{3.30}$$

This is exactly the entropy that we have found in (3.22) using the Wald formula.

#### 4. M2 branes in orbifold space

It has been conjectured that  $\mathcal{N} = 6$  superconformal Chern-Simons matter theories at level  $k$  with gauge group  $U_k(N) \times U_{-k}(N)$  is dual to M-theory on  $AdS_4 \times S^7/Z_k$  [8]. At finite temperature and at strong 't Hooft coupling  $\lambda = N/k$ , the entropy of field theory is given by the entropy of Schwarzschild black hole in  $AdS_4 \times S^7/Z_k$ , (1.2). In this section we are going to find the correction to this entropy from higher derivative corrections in the gravity side. In the presence of higher derivative terms, the Schwarzschild  $AdS_4 \times S^7/Z_k$  background is changing. We consider the same ansatz for the  $AdS_4$  part as before. So we choose the following ansatz for the eleven dimensional metric

$$ds^2 = S^{-7}(r) H^2(r) \left( K^2(r) d\tau^2 + P^2(r) dr^2 + \sum_{i=1}^2 dx_i^2 \right) + L^2 S^2(r) ds_{S^7/Z_k}^2, \tag{4.1}$$

where the orbifold  $S^7/Z_k$  can be written as follows (see e.g. [14] or [15])

$$ds_{S^7/Z_k}^2 = ds_{CP^3}^2 + (d\alpha + \omega)^2, \quad (4.2)$$

where

$$\begin{aligned} \omega &= \frac{1}{2} \left( \cos^2 \xi - \sin^2 \xi \right) d\psi + \frac{1}{2} \cos^2 \xi \cos \theta_1 d\phi_1 + \frac{1}{2} \sin^2 \xi \cos \theta_2 d\phi_2, \\ ds_{CP^3}^2 &= d\xi^2 + \cos^2 \xi \sin^2 \xi \left( d\psi + \frac{\cos \theta_1}{2} d\phi_1 - \frac{\cos \theta_2}{2} d\phi_2 \right)^2 \\ &\quad + \frac{1}{4} \cos^2 \xi \left( d\theta_1^2 + \sin^2 \theta_1 d\phi_1^2 \right) + \frac{1}{4} \sin^2 \xi \left( d\theta_2^2 + \sin^2 \theta_2 d\phi_2^2 \right). \end{aligned} \quad (4.3)$$

and the variables run the values  $0 \leq \xi < \frac{\pi}{2}$ ,  $0 \leq \chi_i < 4\pi$ ,  $0 \leq \phi_i \leq 2\pi$  and  $0 \leq \theta_i < \pi$ . The  $Z_k$  orbifold in  $\alpha$  direction gives a periodicity as  $\alpha \sim \alpha + \frac{2\pi}{k}$ . The metric of  $S^7/Z_k$  in terms of the above variables simplifies to

$$\begin{aligned} ds_{S^7/Z_k}^2 &= d\alpha^2 + d\xi^2 + \frac{1}{4} (d\psi^2 + \cos^2 \xi d\phi_1^2 + \sin^2 \xi d\phi_2^2 + \cos^2 \xi d\theta_1^2 + \sin^2 \xi d\theta_2^2) \\ &\quad + \frac{1}{2} (\cos^2 \xi - \sin^2 \xi) d\alpha d\psi + \frac{1}{2} \cos^2 \xi \cos \theta_1 d\alpha d\phi_1 + \frac{1}{2} \sin^2 \xi \cos \theta_2 d\alpha d\phi_2 \\ &\quad + \frac{1}{4} \cos^2 \xi \cos \theta_1 d\psi d\phi_1 - \frac{1}{4} \sin^2 \xi \cos \theta_2 d\psi d\phi_2. \end{aligned} \quad (4.4)$$

Note that the volume of  $S^7/Z_k$  in terms of the volume of  $CP^3$  is  $\text{Vol}(S^7/Z_k) = \frac{2\pi}{k} \text{Vol}(CP^3)$ .

By looking at the metric one finds that the scalar curvatures behave as  $\mathcal{R}_{\text{AdS}} \sim \mathcal{R}_{S^7/Z_k} \sim 1/L^2$ . Using the fact that 1-loop correction behaves as  $W \sim \mathcal{R}^4$ , one observes that the ratio of one loop correction to supergravity is of order  $W/\mathcal{R} \sim 1/L^6 \sim 1/(Nk) \sim \lambda/N^2$ . To find the numeric factor, we note that with the above ansatz the differential equations in (3.12) and the temperature do not change. However the free energy (3.29) changes by a factor of  $\frac{1}{k}$  due to integration over  $\alpha$ , i.e.

$$F = -\frac{1}{k} \frac{V_2 V_7}{2\kappa_{11}^2} \left( \frac{16}{27} \pi^3 L^9 T^3 \right) \left( 1 + \frac{41}{250} \frac{\gamma}{L^6} \right). \quad (4.5)$$

To write the free energy in terms of the number of M2-branes, we note that the total number of M2-branes is given by

$$N = \frac{1}{2\kappa_{11}^2} \int_{S^7/Z_k} *F_{(4)} = \frac{1}{k} \frac{1}{2\kappa_{11}^2} \int_{S^7} *F_{(4)}, \quad (4.6)$$

this gives  $L^9 = (kN)^{\frac{3}{2}} \kappa_{11}^2 \frac{\sqrt{2}}{\pi^5}$ . So the free energy becomes

$$F = -\frac{1}{k} V_2 T^3 \left\{ \frac{2^{\frac{7}{2}}}{3^4} \pi^2 (kN)^{\frac{3}{2}} + \frac{41}{250} \left( \frac{2\pi}{3} \right)^5 2^{\frac{1}{6}} \pi^{\frac{7}{3}} (kN)^{\frac{1}{2}} \right\}. \quad (4.7)$$

In terms of the 't Hooft coupling it becomes

$$F = -V_2 T^3 \left\{ \frac{2^{\frac{7}{2}}}{3^4} \pi^2 \frac{N^2}{\sqrt{\lambda}} + \frac{41}{250} \left( \frac{2\pi}{3} \right)^5 2^{\frac{1}{6}} \pi^{\frac{7}{3}} \sqrt{\lambda} \right\}. \quad (4.8)$$

The first term is the same entropy as in (1.2), and the second term is the one loop correction in the gravity side. Note that this expression is valid for strong 't Hooft coupling and for large  $N$ , i.e. the second term is small with comparing to the first term.

### 4.1 Large $k$ limit

The radius of  $\alpha$  circle is large whenever  $k^5 \ll N$ , so the M-theory description is valid for small  $k$ . However for larger value of  $k$  the circle becomes small, the curvature becomes large, and so all higher derivative terms of M-theory are relevant to our calculation of entropy. Hence, in this case, one should reduce the M-theory to a weakly coupled IIA string theory. So consider the reduction of the M-theory background to the string frame of IIA as

$$ds_{D=11}^2 = e^{-2\phi/3} ds_{\text{IIA}}^2 + e^{4\phi/3} (R_{11} d\tilde{\alpha} + A)^2, \quad (4.9)$$

where  $R_{11} = g_s^{2/3} l_p$  is the radius of the eleventh direction with angle  $\tilde{\alpha}$ . Comparing the above metric with (4.1), one finds  $\tilde{\alpha} = k\alpha$ , the string coupling to be  $g_s^2 e^{2\phi} = g_s^2 L^3 / (R_{11}^3 k^3) = 2^{5/2} \pi \sqrt{N/k^5}$ , and the metric in IIA string theory to be  $(kR_{11}/L) ds_{\text{IIA}}^2 = ds_{AdS_4}^2 + L^2 ds_{CP^3}^2$ . Reduction also gives a two form magnetic field strength as

$$F_{(2)} = dA = kd\omega. \quad (4.10)$$

Additionally, we have a four form electric field strength as before which is  $F_{\tau x_1 x_2 r} = \frac{3r^2}{L^3}$ .

The type IIA prescription is valid if  $k^5 \gg N$ . By looking at the metric one finds that the scalar curvatures behave as  $\mathcal{R}_{AdS} \sim \mathcal{R}_{CP^3} \sim kR_{11}/L^3 \sim \sqrt{k/N}$ . The first  $\alpha'$  correction to the type IIA effective action can be written in terms of  $W$ . Using the fact that  $W$  behaves as  $W \sim \mathcal{R}^4$ , one observes that the ratio of  $\alpha'$ -correction to supergravity is of order  $W/\mathcal{R} \sim (k/N)^{3/2}$ . On the other hand, the ratio of one loop string correction to supergravity is of order  $\mathcal{R}^4/(e^{-2\phi}\mathcal{R}) \sim 1/Nk$ . Since loop correction of M-theory and type IIA both behave as  $1/Nk$ , one expects that the numerical value of string loop correction to be the same as the M-theory loop correction (4.8). So we concentrate on  $\alpha'$  correction. If  $N \sim k \gg 1$ , then string loop correction is suppressed and  $\alpha'$ -correction is important. Considering the  $\alpha'$  corrections to the effective action, we are going to find their effects on the background solution and subsequently on the entropy. So consider the Lagrangian of type II theory in the presence of first  $\alpha'$  correction which is given by

$$S = -\frac{1}{16\pi G_{10}} \int d^{10}x \sqrt{g} \left\{ e^{-2\phi} (R + 4(\partial\phi)^2) - \frac{1}{2} \frac{1}{4!} F_{(4)}^2 - \frac{1}{2} \frac{1}{2!} F_{(2)}^2 + \gamma e^{-2\phi} W \right\}, \quad (4.11)$$

where  $\gamma = \frac{1}{8}\zeta(3)(\alpha')^3$ . We choose the following ansatz for the metric

$$ds_{\text{IIA}}^2 = \left( \frac{L}{kR_{11}} \right) \left[ S^{-6}(r) \left( \frac{r}{L} \right)^2 \left( K^2(r) d\tau^2 + P^2(r) dr^2 + \sum_{i=1}^2 dx_i^2 \right) + L^2 S^2(r) ds_{CP^3}^2 \right], \quad (4.12)$$

and for the electric four form

$$F_{\tau x_1 x_2 r} = \frac{6r^4}{L^5} K(r) P(r) S^{-18}(r). \quad (4.13)$$

Again using the perturbative method (3.10) we are able to compute the corrections to the supergravity solution, i.e.,

$$\begin{aligned}
 K(r) &= \left(1 - \frac{r_0^3}{r^3}\right)^{\frac{1}{2}} \left[1 + \frac{\gamma}{L^6} \left(\frac{kR_{11}}{L}\right)^3 \left(-136\frac{r_0^3}{r^3} - 136\frac{r_0^6}{r^6} + 88\frac{r_0^9}{r^9}\right)\right], \\
 P(r) &= \frac{L^2}{2r^2} \left(1 - \frac{r_0^3}{r^3}\right)^{-\frac{1}{2}} \left[1 + \frac{\gamma}{L^6} \left(\frac{kR_{11}}{L}\right)^3 \left(4 + 136\frac{r_0^3}{r^3} + 136\frac{r_0^6}{r^6} - 536\frac{r_0^9}{r^9}\right)\right], \quad (4.14)
 \end{aligned}$$

and the differential equation for  $Z(r)$  is given by

$$0 = -3r^{11}L^6(r^3 - r_0^3)Z''(r) - 3L^6r^{10}(r^3 - r_0^3)Z'(r) - 420r^{12}L^6Z(r) + 816r^{12} + 2592r_0^{12}. \quad (4.15)$$

The correction of the dilaton field does not enter into the entropy in which we are interested.

We are going to calculate the entropy from the Wald formula (3.16). In terms of  $r_0$  it is

$$S = \frac{4\pi V_2 V_{CP^3} L^5 r_0^2}{2kR_{11}\kappa_{10}^2} \left(1 + 112\frac{\gamma}{L^6} \left(\frac{kR_{11}}{L}\right)^3\right), \quad (4.16)$$

where the first term is the area of the horizon and the second term is coming from the  $W$  terms. The temperature in terms of  $r_0$  is

$$T = \frac{3r_0}{2\pi L^2} \left(1 + 76\frac{\gamma}{L^6} \left(\frac{kR_{11}}{L}\right)^3\right), \quad (4.17)$$

and the entropy in terms of temperature becomes

$$S = \frac{8\pi^3 L^9 V_2 V_{CP^3} T^2}{9kR_{11}\kappa_{10}^2} \left(1 - 40\frac{\gamma}{L^6} \left(\frac{kR_{11}}{L}\right)^3\right). \quad (4.18)$$

Using the relations  $\frac{2\pi R_{11}}{\kappa_{11}^2} = \frac{1}{\kappa_{10}^2}$ ,  $2\pi \text{Vol}(CP^3) = \text{Vol}(S^7)$ ,  $l_{11} = g_2^{1/3} l_s$  and  $2\kappa_{11}^2 = (2\pi l_{11})^9 / 2\pi$  one finds

$$S = \frac{2^{\frac{7}{2}} \pi^2 N^2}{3^3 \sqrt{\lambda}} V_2 T^2 \left\{1 - \frac{5\zeta(3)}{\pi^3 2^{\frac{17}{2}} \lambda^{3/2}}\right\}. \quad (4.19)$$

Here also the result is valid for the strong 't Hooft coupling and for large  $N$ . The first term is the same as the corresponding term in the M-theory. The second term is resulted from the first  $\alpha'$  correction to the type IIA supergravity action. The structure of this term is very different from the one loop correction in the M-theory side, but the behavior of correction as inverse of the 't Hooft coupling is very similar to the familiar  $AdS_5 \times S^5$  case in [12].

## 5. M5-branes in flat space

The thermal field theory of  $N$  M5 branes in flat space is dual to M-theory on Schwarzschild  $AdS_7 \times S^4$ . The entropy of the field theory at the strong coupling is given by the entropy of the tree-level Schwarzschild AdS solution,

$$S = V_5 T^5 \frac{2^7}{3^6} \pi^3 N^3. \quad (5.1)$$

In this section we would like to find the one loop corrections to this entropy. The near horizon geometry of  $N$  M5-branes at the supergravity level is  $AdS_7$  Schwarzschild times  $S^4$ . By including the higher derivative terms we need to consider a general ansatz and again find deformations of this near horizon geometry. Using the observation in M2-brane case that there is an ansatz in which the area of horizon in terms of the parameter  $r_0$  is independent of the higher derivative corrections, we use the following ansatz for M5-branes

$$ds^2 = S^{-\frac{8}{5}}(r)H^2(r)\left(K^2(r)d\tau^2 + P^2(r)dr^2 + \sum_{i=1}^5 dx_i^2\right) + L^2S^2(r)d\Omega_4^2, \quad (5.2)$$

where  $H(r) = \frac{r}{L}$ . At tree-level the above functions are

$$K(r) = \left(1 - \frac{r_0^6}{r^6}\right)^{\frac{1}{2}}, \quad P(r) = \frac{2L^2}{r^2} \left(1 - \frac{r_0^6}{r^6}\right)^{-\frac{1}{2}}, \quad S(r) = 1, \quad (5.3)$$

where we have changed the variable  $r$  in (2.5) to  $\sqrt{rL}$  and made the space-time Euclidean. The electric field strength at tree-level is also given by  $F_{\tau x_1 \dots x_5 r} = 6\frac{r^5}{L^6}$ . Considering the fact that the charge of M5-branes is independent of the loop corrections, we restrict the field strength of M5-branes to the following form

$$F_{\tau x_1 \dots x_5 r} = \frac{3r^7}{L^8}K(r)P(r)S^{\frac{5}{3}}. \quad (5.4)$$

The Euclidean action then simplifies to

$$I = -\frac{\beta V_5 V_4}{2\kappa_{11}^2} \int dr \left( \mathcal{L} + \gamma \mathcal{W} \right), \quad (5.5)$$

where

$$\mathcal{L} = S^{-\frac{8}{5}}r^7L^{-3}KP\left(R - \frac{1}{2}\frac{1}{7!}F_{(7)}^2\right), \quad \mathcal{W} = S^{-\frac{8}{5}}r^7L^{-3}KPW. \quad (5.6)$$

Here also the right hand side of the Euler-Lagrange equation belong to the next order of perturbations so we just need to compute it by inserting the tree-level solutions. The value of  $W$  is

$$W = \frac{9}{8000L^8} \left( 1533 + 7000\frac{r_0^{12}}{r^{12}} - 11000\frac{r_0^{18}}{r^{18}} + 45625\frac{r_0^{24}}{r^{24}} \right). \quad (5.7)$$

Note that only the last term is the value of  $W$  for seven dimensional Schwarzschild  $AdS_7$  considered in [12]. The derivatives of  $\mathcal{W}$  in (3.7) which appear as source terms for the equations of motion are

$$\begin{aligned} \omega_K &= \frac{3}{4000} \frac{1}{L^9 r^{16} (r^6 - r_0^6)^{\frac{1}{2}}} \left( 6298875r_0^{24} - 5507000r_0^{18}r^6 + 121400r_0^{12}r^{12} + 6351r^{24} \right), \\ \omega_P &= \frac{3}{8000} \frac{(r^6 - r_0^6)^{\frac{1}{2}}}{L^{11}r^{20}} \left( 1030875r_0^{24} - 181000r_0^{18}r^6 + 63400r_0^{12}r^{12} + 6351r^{24} \right), \\ \omega_S &= -\frac{27}{2500} \frac{1}{L^9 r^{19}} \left( 4599r^{24} - 44000r^{12}r_0^{12} + 234000r^6r_0^{18} - 243625r_0^{24} \right). \end{aligned} \quad (5.8)$$

Using the perturbative method to solve the Euler-Lagrange equations, i.e.

$$\begin{aligned}
 K(r) &= \left(1 - \frac{r_0^6}{r^6}\right)^{\frac{1}{2}} (1 + \gamma X(r)) , \\
 P(r) &= \frac{2L^2}{r^2} \left(1 - \frac{r_0^6}{r^6}\right)^{-\frac{1}{2}} (1 + \gamma Y(r)) , \\
 S(r) &= (1 + \gamma Z(r)) ,
 \end{aligned} \tag{5.9}$$

one finds the following differential equations

$$\begin{aligned}
 0 &= -20000r^{19}(r^6 - r_0^6)L^6Y'(r) - 120000r^{24}L^6Y(r) \\
 &\quad - 19053r^{24} - 364200r^{12}r_0^{12} + 16521000r^6r_0^{18} - 18896625r_0^{24} , \\
 0 &= 20000r^{19}L^6(r^6 - r_0^6)^2X'(r) - 120000r^{24}L^6(r^6 - r_0^6)Y(r) \\
 &\quad - 19053r^{30} + 19053r^{24}r_0^6 + 733200r^{12}r_0^{18} - 3635625r^6r_0^{24} \\
 &\quad - 190200r^{18}r_0^{12} + 3092625r_0^{30} , \\
 0 &= -28800r^{19}L^6\left(r(r^6 - r_0^6)Z''(r) + (7r^6 - r_0^6)Z'(r) - 72r^5Z(r)\right) \\
 &\quad + \frac{993384}{5}r^{24} - 10524600r_0^{24} + 10108800r^6r_0^{18} - 1900800r^{12}r_0^{12} .
 \end{aligned} \tag{5.10}$$

The solution to the first two equations are

$$\begin{aligned}
 X(r) &= -\frac{1}{L^6} \left( c + \frac{21299}{1600} \frac{r_0^6}{r^6} + \frac{4999}{320} \frac{r_0^{12}}{r^{12}} - \frac{2749}{320} \frac{r_0^{18}}{r^{18}} \right) , \\
 Y(r) &= -\frac{1}{L^6} \left( \frac{6351}{40000} - \frac{21299}{1600} \frac{r_0^6}{r^6} - \frac{5231}{320} \frac{r_0^{12}}{r^{12}} + \frac{16797}{320} \frac{r_0^{18}}{r^{18}} \right) ,
 \end{aligned} \tag{5.11}$$

where  $c$  is a constant which can be set to zero by time scaling, up to first order of  $\gamma$ . Here also the differential equation for  $Z(r)$  does not have such a simple solution. However we do not need to know this solution for finding the thermodynamical quantities  $T$  and  $S$  in which we are interested. So we ignore  $Z(r)$  from now on in this section.

The temperature of the black hole can be found by surface gravity at the horizon

$$\hat{\kappa} = 2\pi T = \sqrt{g^{rr}} \frac{d}{dr} \sqrt{g_{\tau\tau}}|_H = \frac{3}{2} \frac{r_0}{L^2} \left( 1 + \frac{\gamma}{L^6} \frac{105901}{40000} \right) . \tag{5.12}$$

Using the ansatz (5.2), one observes that the horizon area in terms of  $r_0$  does not modify by the higher derivative corrections so the first term in the Wald formula (3.16), which is the area of horizon at leading order does not modify either. So we need only the tree-level solution to calculate the first term in the Wald formula (3.16)

$$S_1 = \frac{4\pi}{2\kappa_{11}^2} V_5 V_4 \frac{r_0^5}{L} . \tag{5.13}$$

On the other hand the second term is proportional to  $\gamma$ , so to first order of  $\gamma$  one has to replace the tree-level solution into the second term too, i.e.

$$S_2 = \frac{4\pi}{2\kappa_{11}^2} V_5 V_4 \frac{r_0^5}{L} \left( \frac{12597}{500} \frac{\gamma}{L^6} \right). \quad (5.14)$$

So the final result for the entropy in terms of the temperature is

$$S = \frac{V_5 V_4}{2\kappa_{11}^2} \left( \frac{2^{12}}{3^5} \pi^6 L^9 T^5 \right) \left( 1 + \frac{95651}{8000} \frac{\gamma}{L^6} \right). \quad (5.15)$$

Inserting  $L^9 = N^3 \kappa_{11}^2 2^{-7} \pi^{-5}$  where  $N$  is the total number of M5-branes and  $V_4 = 8 \frac{\pi^2}{3}$ , one finds

$$S = V_5 T^5 \left( \frac{2^7}{3^6} \pi^3 N^3 + \frac{95651}{8000} \left( \frac{2\pi}{3} \right)^8 3 \left( \frac{\pi}{2} \right)^{\frac{1}{3}} 2^6 N \right). \quad (5.16)$$

Using the same steps as before for obtaining the free energy, one finds the following result for the free energy

$$F = -\frac{V_5 V_4}{2\kappa_{11}^2} \frac{1}{2} \frac{r_0^6}{L^3} \left( 1 + \frac{1113661}{40000} \frac{\gamma}{L^6} \right) = -\frac{V_5 V_4}{2\kappa_{11}^2} \left( \frac{2048}{729} \pi^6 L^9 T^6 \right) \left( 1 + \frac{95651}{8000} \frac{\gamma}{L^6} \right), \quad (5.17)$$

which gives exactly the same entropy as in (5.15)

## References

- [1] J.H. Schwarz, *Superconformal Chern-Simons theories*, *JHEP* **11** (2004) 078 [[hep-th/0411077](#)];  
A. Basu and J.A. Harvey, *The M2-M5 brane system and a generalized Nahm's equation*, *Nucl. Phys. B* **713** (2005) 136 [[hep-th/0412310](#)].
- [2] J. Bagger and N. Lambert, *Comments on multiple M2-branes*, *JHEP* **02** (2008) 105 [[arXiv:0712.3738](#)]; *Gauge symmetry and supersymmetry of multiple M2-branes*, *Phys. Rev. D* **77** (2008) 065008 [[arXiv:0711.0955](#)]; *Modeling multiple M2's*, *Phys. Rev. D* **75** (2007) 045020 [[hep-th/0611108](#)].
- [3] A. Gustavsson, *Algebraic structures on parallel M2-branes*, [arXiv:0709.1260](#); *Selfdual strings and loop space Nahm equations*, *JHEP* **04** (2008) 083 [[arXiv:0802.3456](#)].
- [4] N. Lambert and D. Tong, *Membranes on an orbifold*, [arXiv:0804.1114](#);  
J. Distler, S. Mukhi, C. Papageorgakis and M. Van Raamsdonk, *M2-branes on M-folds*, *JHEP* **05** (2008) 038 [[arXiv:0804.1256](#)].
- [5] J. Gomis, G. Milanesi and J.G. Russo, *Bagger-Lambert theory for general Lie algebras*, *JHEP* **06** (2008) 075 [[arXiv:0805.1012](#)];  
S. Benvenuti, D. Rodriguez-Gomez, E. Tonni and H. Verlinde, *N = 8 superconformal gauge theories and M2 branes*, [arXiv:0805.1087](#);  
P.-M. Ho, Y. Imamura and Y. Matsuo, *M2 to D2 revisited*, *JHEP* **07** (2008) 003 [[arXiv:0805.1202](#)].
- [6] M.A. Bandres, A.E. Lipstein and J.H. Schwarz, *Ghost-free superconformal action for multiple M2-branes*, *JHEP* **07** (2008) 117 [[arXiv:0806.0054](#)];  
J. Gomis, D. Rodriguez-Gomez, M. Van Raamsdonk and H. Verlinde, *Supersymmetric Yang-Mills theory from lorentzian three-algebras*, [arXiv:0806.0738](#).

- [7] B. Ezhuthachan, S. Mukhi and C. Papageorgakis, *D2 to D2*, *JHEP* **07** (2008) 041 [arXiv:0806.1639].
- [8] O. Aharony, O. Bergman, D.L. Jafferis and J. Maldacena, *N=6 superconformal Chern-Simons-matter theories, M2-branes and their gravity duals*, arXiv:0806.1218.
- [9] R.M. Wald, *Black hole entropy is the Nöther charge*, *Phys. Rev. D* **48** (1993) 3427 [gr-qc/9307038];  
T. Jacobson and R.C. Myers, *Black hole entropy and higher curvature interactions*, *Phys. Rev. Lett.* **70** (1993) 3684 [hep-th/9305016];  
T. Jacobson, G. Kang and R.C. Myers, *Black hole entropy in higher curvature gravity*, gr-qc/9502009.
- [10] S.W. Hawking and D.N. Page, *Thermodynamics of black holes in Anti-de Sitter space*, *Commun. Math. Phys.* **87** (1983) 577;  
E. Witten, *Anti-de Sitter space, thermal phase transition and confinement in gauge theories*, *Adv. Theor. Math. Phys.* **2** (1998) 505 [hep-th/9803131].
- [11] S. Mukhi and C. Papageorgakis, *M2 to D2*, *JHEP* **05** (2008) 085 [arXiv:0803.3218];  
D.S. Berman, L.C. Tadowski and D.C. Thompson, *Aspects of multiple membranes*, *Nucl. Phys. B* **802** (2008) 106 [arXiv:0803.3611];  
M. Van Raamsdonk, *Comments on the Bagger-Lambert theory and multiple M2-branes*, *JHEP* **05** (2008) 105 [arXiv:0803.3803];  
A. Morozov, *On the problem of multiple M2 branes*, *JHEP* **05** (2008) 076 [arXiv:0804.0913];  
U. Gran, B.E.W. Nilsson and C. Petersson, *On relating multiple M2 and D2-branes*, arXiv:0804.1784;  
P.-M. Ho, R.-C. Hou and Y. Matsuo, *Lie 3-algebra and multiple M2-branes*, *JHEP* **06** (2008) 020 [arXiv:0804.2110];  
J. Gomis, A.J. Salim and F. Passerini, *Matrix theory of type IIB plane wave from membranes*, *JHEP* **08** (2008) 002 [arXiv:0804.2186];  
E.A. Bergshoeff, M. de Roo and O. Hohm, *Multiple M2-branes and the embedding tensor*, *Class. and Quant. Grav.* **25** (2008) 142001 [arXiv:0804.2201];  
K. Hosomichi, K.-M. Lee and S. Lee, *Mass-deformed Bagger-Lambert theory and its BPS objects*, arXiv:0804.2519;  
G. Papadopoulos, *M2-branes, 3-Lie Algebras and Plücker relations*, *JHEP* **05** (2008) 054 [arXiv:0804.2662];  
J.P. Gauntlett and J.B. Gutowski, *Constraining maximally supersymmetric membrane actions*, arXiv:0804.3078;  
G. Papadopoulos, *On the structure of k-Lie algebras*, *Class. and Quant. Grav.* **25** (2008) 142002 [arXiv:0804.3567];  
P.-M. Ho and Y. Matsuo, *M5 from M2*, *JHEP* **06** (2008) 105 [arXiv:0804.3629];  
A. Morozov, *From simplified BLG action to the first-quantized M-theory*, arXiv:0805.1703;  
Y. Honma, S. Iso, Y. Sumitomo and S. Zhang, *Janus field theories from multiple M2 branes*, *Phys. Rev. D* **78** (2008) 025027 [arXiv:0805.1895];  
H. Fujii, S. Terashima and M. Yamazaki, *A new N = 4 membrane action via orbifold*, arXiv:0805.1997;  
P.-M. Ho, Y. Imamura, Y. Matsuo and S. Shiba, *M5-brane in three-form flux and multiple M2-branes*, *JHEP* **08** (2008) 014 [arXiv:0805.2898];  
C. Krishnan and C. Maccaferri, *Membranes on calibrations*, *JHEP* **07** (2008) 005 [arXiv:0805.3125];  
Y. Song, *Mass deformation of the multiple M2 branes theory*, arXiv:0805.3193;



- I. Jeon, J. Kim, N. Kim, S.-W. Kim and J.-H. Park, *Classification of the BPS states in Bagger-Lambert theory*, *JHEP* **07** (2008) 056 [[arXiv:0805.3236](#)];
- M. Li and T. Wang, *M2-branes coupled to antisymmetric fluxes*, *JHEP* **07** (2008) 093 [[arXiv:0805.3427](#)];
- K. Hosomichi, K.-M. Lee, S. Lee, S. Lee and J. Park,  *$N = 4$  superconformal Chern-Simons theories with hyper and twisted hyper multiplets*, *JHEP* **07** (2008) 091 [[arXiv:0805.3662](#)];
- S. Banerjee and A. Sen, *Interpreting the M2-brane action*, [arXiv:0805.3930](#);
- H. Lin, *Kac-Moody extensions of 3-algebras and M2-branes*, *JHEP* **07** (2008) 136 [[arXiv:0805.4003](#)];
- P. De Medeiros, J.M. Figueroa-O'Farrill and E. Mendez-Escobar, *Lorentzian Lie 3-algebras and their Bagger-Lambert moduli space*, *JHEP* **07** (2008) 111 [[arXiv:0805.4363](#)];
- A. Gustavsson, *One-loop corrections to Bagger-Lambert theory*, [arXiv:0805.4443](#);
- J.-H. Park and C. Sochichiu, *Single M5 to multiple M2: taking off the square root of Nambu-Goto action*, [arXiv:0806.0335](#);
- F. Passerini, *M2-brane superalgebra from Bagger-Lambert theory*, [arXiv:0806.0363](#);
- C. Ahn, *Holographic supergravity dual to three dimensional  $N = 2$  gauge theory*, [arXiv:0806.1420](#);
- M. Benna, I. Klebanov, T. Klose and M. Smedback, *Superconformal Chern-Simons theories and  $AdS_4/CFT_3$  correspondence*, [arXiv:0806.1519](#);
- S. Cecotti and A. Sen, *Coulomb branch of the lorentzian three algebra theory*, [arXiv:0806.1990](#);
- A. Mauri and A.C. Petkou, *An  $N = 1$  superfield action for M2 branes*, [arXiv:0806.2270](#);
- E.A. Bergshoeff, M. de Roo, O. Hohm and D. Roest, *Multiple membranes from gauged supergravity*, [arXiv:0806.2584](#);
- P. de Medeiros, J.M. Figueroa-O'Farrill and E. Mendez-Escobar, *Metric Lie 3-algebras in Bagger-Lambert theory*, [arXiv:0806.3242](#);
- M. Blau and M. O'Loughlin, *Multiple M2-Branes and Plane Waves*, [arXiv:0806.3253](#);
- Y. Honma, S. Iso, Y. Sumitomo and S. Zhang, *Scaling limit of  $N = 6$  superconformal Chern-Simons theories and lorentzian Bagger-Lambert theories*, [arXiv:0806.3498](#);
- T.L. Curtright, D.B. Fairlie and C.K. Zachos, *Ternary Virasoro-Witt algebra*, [arXiv:0806.3515](#);
- C. Sochichiu, *On Nambu-Lie 3-algebra representations*, [arXiv:0806.3520](#);
- Y. Imamura and K. Kimura, *Coulomb branch of generalized ABJM models*, [arXiv:0806.3727](#);
- J.A. Minahan and K. Zarembo, *The Bethe ansatz for superconformal Chern-Simons*, [arXiv:0806.3951](#);
- T.A. Larsson, *Virasoro 3-algebra from scalar densities*, [arXiv:0806.4039](#);
- K. Furuuchi, S.-Y.D. Shih and T. Takimi, *M-theory superalgebra from multiple membranes*, [arXiv:0806.4044](#);
- A. Armoni and A. Naqvi, *A non-supersymmetric large- $N$  3D CFT and its gravity dual*, [arXiv:0806.4068](#);
- A. Agarwal, *Mass deformations of super Yang-Mills theories in  $D = 2 + 1$  and super-membranes: a note*, [arXiv:0806.4292](#);
- D. Gaiotto, S. Giombi and X. Yin, *Spin chains in  $N = 6$  superconformal Chern-Simons-matter theory*, [arXiv:0806.4589](#);
- I.A. Bandos and P.K. Townsend, *Light-cone M5 and multiple M2-branes*, [arXiv:0806.4777](#);
- C. Ahn, *Towards holographic gravity dual of  $N = 1$  superconformal Chern-Simons gauge theory*, *JHEP* **07** (2008) 101 [[arXiv:0806.4807](#)];

- J. Bedford and D. Berman, *A note on quantum aspects of multiple membranes*, arXiv:0806.4900;
- G. Arutyunov and S. Frolov, *Superstrings on  $AdS_4 \times CP^3$  as a coset  $\sigma$ -model*, arXiv:0806.4940;
- B.J. Stefanski, B., *Green-Schwarz action for Type IIA strings on  $AdS_4 \times CP^3$* , arXiv:0806.4948;
- G. Grignani, T. Harmark and M. Orselli, *The  $SU(2) \times SU(2)$  sector in the string dual of  $N = 6$  superconformal Chern-Simons theory*, arXiv:0806.4959;
- K. Hosomichi, K.-M. Lee, S. Lee, S. Lee and J. Park,  *$N = 5, 6$  superconformal Chern-Simons Theories and  $M2$ -branes on orbifolds*, arXiv:0806.4977;
- P. Fré and P.A. Grassi, *Pure spinor formalism for  $Osp(N|4)$  backgrounds*, arXiv:0807.0044;
- K. Okuyama, *Linear  $\sigma$ -models for the  $R^8/Z_k$  orbifold*, arXiv:0807.0047;
- J. Bagger and N. Lambert, *Three-algebras and  $N = 6$  Chern-Simons gauge theories*, arXiv:0807.0163;
- S. Terashima, *On  $M5$ -branes in  $N = 6$  membrane action*, arXiv:0807.0197;
- G. Grignani, T. Harmark, M. Orselli and G.W. Semenoff, *Finite size giant magnons in the string dual of  $N = 6$  superconformal Chern-Simons theory*, arXiv:0807.0205;
- S. Chakraborty, A. Kumar and S. Jain,  *$w_\infty$  3-algebra*, arXiv:0807.0284;
- S. Terashima and F. Yagi, *Orbifolding the membrane action*, arXiv:0807.0368;
- N. Gromov and P. Vieira, *The  $AdS_4/CFT_3$  algebraic curve*, arXiv:0807.0437;
- C. Ahn and P. Bozhilov, *Finite-size effects of membranes on  $AdS_4 \times S_7$* , arXiv:0807.0566;
- N. Gromov and P. Vieira, *The all loop  $AdS_4/CFT_3$  Bethe ansatz*, arXiv:0807.0777;
- B. Chen and J.-B. Wu, *Semi-classical strings in  $AdS_4 \times CP^3$* , arXiv:0807.0802;
- S. Cherkis and C. Sämann, *Multiple  $M2$ -branes and generalized 3-Lie algebras*, arXiv:0807.0808;
- C.-S. Chu, P.-M. Ho, Y. Matsuo and S. Shiba, *Truncated Nambu-Poisson bracket and entropy formula for multiple membranes*, arXiv:0807.0812;
- M.A. Bandres, A.E. Lipstein and J.H. Schwarz, *Studies of the ABJM theory in a formulation with manifest  $SU(4)$   $R$ -symmetry*, arXiv:0807.0880;
- Y. Zhou, *Interpreting  $N$   $M2$  branes*, arXiv:0807.0890;
- J. Gomis, D. Rodriguez-Gomez, M. Van Raamsdonk and H. Verlinde, *A massive study of  $M2$ -brane proposals*, arXiv:0807.1074;
- T. Li, Y. Liu and D. Xie, *Multiple  $D2$ -brane action from  $M2$ -branes*, arXiv:0807.1183.
- [12] S.S. Gubser, I.R. Klebanov and A.A. Tseytlin, *Coupling constant dependence in the thermodynamics of  $N = 4$  supersymmetric Yang-Mills theory*, Nucl. Phys. **B 534** (1998) 202 [hep-th/9805156].
- [13] J.L. Petersen, *Introduction to the Maldacena conjecture on AdS/CFT*, Int. J. Mod. Phys. **A 14** (1999) 3597 [hep-th/9902131].
- [14] M. Cvetič, H. Lü and C.N. Pope, *Consistent warped-space Kaluza-Klein reductions, half-maximal gauged supergravities and  $CP(n)$  constructions*, Nucl. Phys. **B 597** (2001) 172 [hep-th/0007109].
- [15] T. Nishioka and T. Takayanagi, *On type IIA Penrose limit and  $N = 6$  Chern-Simons theories*, JHEP **08** (2008) 001 [arXiv:0806.3391].