

The rational dressing for abelian twisted loop Toda systems

Kh. S. Nirov

*Institute for Nuclear Research of the Russian Academy of Sciences,
60th October Anniversary Prospect 7a, 117312 Moscow, Russia
E-mail: knirov@physik.uni-wuppertal.de*

A.V. Razumov

*Institute for High Energy Physics,
142281 Protvino, Moscow Region, Russia
E-mail: alexander.razumov@ihep.ru*

ABSTRACT: We consider abelian twisted loop Toda equations associated with the complex general linear groups. The Dodd-Bullough-Mikhailov equation is a simplest particular case of the equations under consideration. We construct new soliton solutions of these Toda equations using the method of rational dressing.

KEYWORDS: Integrable Field Theories, Differential and Algebraic Geometry.

Contents

1. Introduction	1
2. Loop Toda equations	2
3. Abelian Toda equations associated with loop groups of complex general linear groups	7
3.1 First type: untwisted loop Toda equations	7
3.2 Second type: twisted loop Toda equations, odd-dimensional case	7
3.3 Third type: twisted loop Toda equations, even-dimensional case	9
4. Rational dressing	10
5. Soliton solutions	21
5.1 Odd-dimensional case	21
5.2 Even-dimensional case	23
6. Conclusion	26

1. Introduction

The comprehensive investigation of completely integrable systems has at least two reasons. First, such systems serve as a testing area for developing various methods to solve nonlinear partial differential equations. And second, they possess an interesting class of solutions, called solitons, which have properties attractive from the point of view of possible physical applications.

The two-dimensional loop Toda equations provide an illustrative and very rich example of completely integrable nonlinear equations, see, for example, the monographs [1, 2]. Different methods applicable to loop Toda equations for constructing their soliton-like solutions were analysed in the paper [3]. Namely, multi-soliton solutions of abelian *untwisted* loop Toda equations associated with the general linear groups were explicitly constructed by means of the Hirota's [4–9] and the rational dressing [10, 11] methods, and a direct relationship between these approaches was established.¹

In this paper we continue the investigation of abelian loop Toda equations associated with the complex general linear groups started in the paper [3]. Here we consider abelian *twisted* loop Toda equations. It is interesting that the famous Dodd-Bullough-Mikhailov

¹It is worth to note that sometimes it is helpful to employ a combination of such complementary methods, see, for example, [12, 13]

equation is a simplest particular case of such twisted loop Toda equations. We develop the rational dressing method in application to these classes of nonlinear equations and construct for them new soliton solutions.

Here we start with the classification of the abelian loop Toda equations, associated with the complex general linear groups, reducing them from the general classification of the loop Toda systems performed in [14, 15]. In this way we find direct relationships between different loop Toda equations and define proper places for the twisted loop Toda systems to be subsequently solved. For the latter purpose we use the rational dressing method that proved to be very efficient and favourable already in the untwisted case [3]. Indeed, the method of rational dressing, consistently developed for the twisted loop Toda equations under consideration, allows us to construct more general classes of solutions where the Hirota solitons form a substantially smaller subset. Besides, it is an essential advantage of the rational dressing, being actually a version of the inverse scattering method, that it does not rely on guessing at an appropriate change of the dependent field variables as in the Hirota's approach.

2. Loop Toda equations

In this section, mainly following the monographs [1, 2] and our papers [14, 15], we recall basic notions and introduce notations to be used below. We start our consideration with a Lie group \mathcal{G} whose Lie algebra \mathfrak{G} is endowed with a \mathbb{Z} -gradation,

$$\mathfrak{G} = \bigoplus_{k \in \mathbb{Z}} \mathfrak{G}_k, \quad [\mathfrak{G}_k, \mathfrak{G}_l] \subset \mathfrak{G}_{k+l},$$

and denote by L such a positive integer that the grading subspaces \mathfrak{G}_k , where $0 < |k| < L$, are trivial. We denote by \mathcal{G}_0 the closed Lie subgroup of \mathcal{G} corresponding to the zero-grade Lie subalgebra \mathfrak{G}_0 . Then, the Toda equation associated with \mathcal{G} is an equation for a mapping Ξ of the Euclidean plane \mathbb{R}^2 to \mathcal{G}_0 , explicitly of the form

$$\partial_+(\Xi^{-1}\partial_-\Xi) = [\mathcal{F}_-, \Xi^{-1}\mathcal{F}_+\Xi], \tag{2.1}$$

where \mathcal{F}_- and \mathcal{F}_+ are some fixed mappings of \mathbb{R}^2 to \mathfrak{G}_{-L} and \mathfrak{G}_{+L} , respectively, satisfying the relations

$$\partial_+\mathcal{F}_- = 0, \quad \partial_-\mathcal{F}_+ = 0. \tag{2.2}$$

Here we use the customary notation $\partial_- = \partial/\partial z^-$, $\partial_+ = \partial/\partial z^+$ for the partial derivatives over the standard coordinates on \mathbb{R}^2 . Certainly, to obtain a nontrivial Toda equation we have to assume that the subspaces \mathfrak{G}_{-L} and \mathfrak{G}_{+L} are nontrivial.

When the Lie group \mathcal{G}_0 is abelian, the corresponding Toda equation is said to be *abelian*, otherwise we deal with a *non-abelian Toda equation*.

We see that a Toda equation is specified by a choice of a \mathbb{Z} -gradation of the Lie algebra \mathfrak{G} of \mathcal{G} and mappings \mathcal{F}_- , \mathcal{F}_+ satisfying the conditions (2.2). Therefore, to classify the Toda equations associated with a Lie group \mathcal{G} we should classify \mathbb{Z} -gradations of the Lie algebra \mathfrak{G} of \mathcal{G} up to isomorphisms.

It is essential for our purposes that the Toda equation (2.1) together with the relations (2.2) are equivalent to the zero-curvature condition for a flat connection on the trivial fiber bundle $\mathbb{R}^2 \times \mathcal{G} \rightarrow \mathbb{R}^2$. Indeed, writing the zero-curvature condition as the equation

$$\partial_- \mathcal{O}_+ - \partial_+ \mathcal{O}_- + [\mathcal{O}_-, \mathcal{O}_+] = 0 \tag{2.3}$$

for the \mathfrak{G} -valued components of the flat connection under consideration, imposing the grading conditions

$$\mathcal{O}_- = \mathcal{O}_{-0} + \mathcal{O}_{-L}, \quad \mathcal{O}_+ = \mathcal{O}_{+0} + \mathcal{O}_{+L},$$

and destroying the residual gauge invariance by the condition

$$\mathcal{O}_{+0} = 0,$$

we bring the connection components to the form

$$\mathcal{O}_- = \Xi^{-1} \partial_- \Xi + \mathcal{F}_-, \quad \mathcal{O}_+ = \Xi^{-1} \mathcal{F}_+ \Xi, \tag{2.4}$$

and then derive the equation (2.1) and the relations (2.2) directly from the zero-curvature condition (2.3), as well as vice versa [16, 2, 17].

It follows from the equality (2.3) that there is a mapping Φ of \mathbb{R}^2 to \mathcal{G} such that

$$\Phi^{-1} \partial_- \Phi = \mathcal{O}_-, \quad \Phi^{-1} \partial_+ \Phi = \mathcal{O}_+. \tag{2.5}$$

We say in this situation that the connection with the components \mathcal{O}_- and \mathcal{O}_+ is generated by the mapping Φ .

We consider the case where \mathcal{G} is a twisted loop group of a complex classical Lie group G which is defined as follows. Let a be an automorphism of G satisfying the relation $a^M = \text{id}_G$ for some positive integer M .² The twisted loop group $\mathcal{L}_{a,M}(G)$ is formed by the mappings χ of the unit circle S^1 to G satisfying the equality

$$\chi(\epsilon_M \bar{p}) = a(\chi(\bar{p})),$$

where $\epsilon_M = e^{2\pi i/M}$ is the M th principal root of unity. We think the circle S^1 as consisting of complex numbers of modulus one. The group law in $\mathcal{L}_{a,M}(G)$ is defined pointwise. The Lie algebra of $\mathcal{L}_{a,M}(G)$ is the twisted loop Lie algebra $\mathcal{L}_{A,M}(\mathfrak{g})$, where \mathfrak{g} is the Lie algebra of G and A is the automorphism of \mathfrak{g} corresponding to the automorphism a of G . The Lie algebra $\mathcal{L}_{A,M}(\mathfrak{g})$ is formed by the mappings ξ of S^1 to \mathfrak{g} satisfying the equality

$$\xi(\epsilon_M \bar{p}) = A(\xi(\bar{p}))$$

with the Lie algebra operation defined pointwise. Note that the relation $A^M = \text{id}_{\mathfrak{g}}$ is satisfied automatically.

In the paper [18] we classified a wide class of the so-called integrable \mathbb{Z} -gradations with finite-dimensional grading subspaces of the twisted loop Lie algebras of the finite-dimensional complex simple Lie algebras. Namely, we showed that any such \mathbb{Z} -gradation of

²Here M is not necessarily the order of the automorphism a , but can be its arbitrary multiple.

a loop Lie algebra $\mathcal{L}_{A,M}(\mathfrak{g})$ is conjugated by an isomorphism to the standard \mathbb{Z} -gradation of another loop Lie algebra $\mathcal{L}_{A',M'}(\mathfrak{g})$, where the automorphisms A and A' differ by an inner automorphism of \mathfrak{g} .

Recall that for the *standard \mathbb{Z} -gradation* of the Lie algebra $\mathcal{L}_{A,M}(\mathfrak{g})$ the grading subspaces are

$$\mathcal{L}_{A,M}(\mathfrak{g})_k = \{\xi = \lambda^k x \in \mathcal{L}_{A,M}(\mathfrak{g}) \mid x \in \mathfrak{g}, A(x) = \epsilon_M^k x\},$$

where by λ we denote the restriction of the standard coordinate on \mathbb{C} to S^1 .

It is well known that twisted loop Lie algebras defined by automorphisms which differ by an inner automorphism are isomorphic, and really different twisted loop Lie algebras can be labeled by the elements of the corresponding outer automorphism group. In particular, if A is an inner automorphism, the loop Lie algebra $\mathcal{L}_{A,M}(\mathfrak{g})$ is isomorphic to an untwisted loop Lie algebra $\mathcal{L}(\mathfrak{g}) = \mathcal{L}_{\text{id}_{\mathfrak{g}},1}(\mathfrak{g})$. Therefore, in this case a Toda equation associated with $\mathcal{L}_{a,M}(G)$ and specified by some choice of a \mathbb{Z} -gradation of $\mathcal{L}_{A,M}(\mathfrak{g})$ is equivalent to a Toda equation associated with $\mathcal{L}(G) = \mathcal{L}_{\text{id}_G,1}(G)$ and specified by the corresponding choice of a \mathbb{Z} -gradation of $\mathcal{L}(\mathfrak{g})$.

Thus, to describe Toda equations associated with the loop groups $\mathcal{L}_{a,M}(G)$, where a is an inner automorphism of G , it suffices to describe the Toda equations associated with the untwisted loop groups $\mathcal{L}(G)$. However, due to simplicity of the standard \mathbb{Z} -gradation, to study Toda equations it is more convenient, instead of using one untwisted loop group $\mathcal{L}(G)$ and different \mathbb{Z} -gradations of $\mathcal{L}(\mathfrak{g})$, to use different twisted loop groups $\mathcal{L}_{a,M}(G)$ and the standard \mathbb{Z} -gradation of $\mathcal{L}_{A,M}(\mathfrak{g})$. Similarly, to describe the Toda equations associated with the loop groups $\mathcal{L}_{a,M}(G)$, where a is an outer automorphism of G satisfying the relation $a^M = \text{id}_G$, it suffices to use only the standard \mathbb{Z} -gradation of the loop Lie algebras $\mathcal{L}_{A,M}(\mathfrak{g})$. Having all this in mind and slightly abusing terminology, we say that when a is an outer automorphism of G then a Toda equation associated with $\mathcal{L}_{a,M}(G)$ is a *twisted loop Toda equation associated with G* , and when a is an inner automorphism of G then a Toda equation associated with $\mathcal{L}_{a,M}(G)$ is an *untwisted loop Toda equation associated with G* .³

The group $\mathcal{L}_{a,M}(G)$ and its Lie algebra $\mathcal{L}_{A,M}(\mathfrak{g})$ are infinite-dimensional manifolds. Nevertheless, using the so-called exponential law [19, 20], it is possible to write the zero-curvature representation of the Toda equations associated with $\mathcal{L}_{a,M}(G)$ in terms of finite-dimensional manifolds. The essence of this useful law can be expressed by the canonical identification $C^\infty(\mathcal{M}, C^\infty(\mathcal{N}, \mathcal{P})) = C^\infty(\mathcal{M} \times \mathcal{N}, \mathcal{P})$, where \mathcal{M} , \mathcal{N} and \mathcal{P} are finite-dimensional manifolds, among which \mathcal{N} is compact.

The connection components \mathcal{O}_- and \mathcal{O}_+ entering the equality (2.3) are mappings of \mathbb{R}^2 to the loop Lie algebra $\mathcal{L}_{A,M}(\mathfrak{g})$. We denote the mappings of $\mathbb{R}^2 \times S^1$ to \mathfrak{g} , corresponding to \mathcal{O}_- and \mathcal{O}_+ in accordance with the exponential law, by ω_- and ω_+ , and call them also the connection components. The mapping Φ generating the connection under consideration is a mapping of \mathbb{R}^2 to $\mathcal{L}_{a,M}(G)$. Denoting the respective mapping of $\mathbb{R}^2 \times S^1$ to G by φ we can write

$$\varphi^{-1} \partial_- \varphi = \omega_-, \quad \varphi^{-1} \partial_+ \varphi = \omega_+, \tag{2.6}$$

³It is common to omit the word ‘untwisted’.

which is equivalent to the expressions (2.5). Having in mind that the mapping φ uniquely determines the connection generating mapping Φ , we say that the mapping φ also generates the connection under consideration. We introduce, according to the exponential law, the smooth mapping γ of $\mathbb{R}^2 \times S^1$ to G respective to Ξ , and smooth mappings of $\mathbb{R}^2 \times S^1$ to \mathfrak{g} respective to \mathcal{F}_- and \mathcal{F}_+ .

Now, explicitly describing the grading subspaces of the standard \mathbb{Z} -gradation of the loop Lie algebra $\mathcal{L}_{A,M}(\mathfrak{g})$, we find that the subalgebra $\mathcal{L}_{A,M}(\mathfrak{g})_0$ is isomorphic to the subalgebra $\mathfrak{g}_{[0]_M}$ of \mathfrak{g} , and the Lie group $\mathcal{L}_{a,M}(G)_0$ is isomorphic to the connected Lie subgroup G_0 of G corresponding to the Lie algebra $\mathfrak{g}_{[0]_M}$. Here $\mathfrak{g}_{[k]_M}$ mean the grading subspaces of the \mathbb{Z}_M -gradation of \mathfrak{g} induced by the automorphism A , and $[k]_M$ denotes the element of the ring \mathbb{Z}_M corresponding to the integer k . For the connection components ω_- and ω_+ we can write the expressions

$$\omega_- = \gamma^{-1} \partial_- \gamma + \lambda^{-L} c_-, \quad \omega_+ = \lambda^L \gamma^{-1} c_+ \gamma, \quad (2.7)$$

which are equivalent to the equalities (2.4). Here c_- and c_+ are mappings of \mathbb{R}^2 to $\mathfrak{g}_{-[L]_M}$ and $\mathfrak{g}_{+[L]_M}$ respectively. Hence, the Toda equation (2.1) can be written as

$$\partial_+(\gamma^{-1} \partial_- \gamma) = [c_-, \gamma^{-1} c_+ \gamma], \quad (2.8)$$

and the conditions (2.2) imply that

$$\partial_+ c_- = 0, \quad \partial_- c_+ = 0. \quad (2.9)$$

We call an equation of the form (2.8) also a loop Toda equation.

Our classification of loop Toda equations is based on a convenient block-matrix representation of the grading subspaces [14, 15] we have implemented. Each element x of the complex classical Lie algebra \mathfrak{g} under consideration is treated as a $p \times p$ block matrix $(x_{\alpha\beta})$, where $x_{\alpha\beta}$ is an $n_\alpha \times n_\beta$ matrix. The sum of the positive integers n_α is the size n of the matrices representing the elements of \mathfrak{g} . For the case of Toda equations associated with the loop groups $\mathcal{L}_{a,M}(\mathrm{GL}_n(\mathbb{C}))$, where a is an inner automorphism of $\mathrm{GL}_n(\mathbb{C})$, the integers n_α are arbitrary. For the other cases they should satisfy some restrictions dictated by the structure of the Lie algebra \mathfrak{g} .

The mapping γ has the block-diagonal form

$$\gamma = \begin{pmatrix} \Gamma_1 & & & \\ & \Gamma_2 & & \\ & & \ddots & \\ & & & \Gamma_p \end{pmatrix}. \quad (2.10)$$

For each $\alpha = 1, \dots, p$ the mapping Γ_α is a mapping of \mathbb{R}^2 to the Lie group $\mathrm{GL}_{n_\alpha}(\mathbb{C})$. For the case of Toda equations associated with the loop groups $\mathcal{L}_{a,M}(\mathrm{GL}_n(\mathbb{C}))$, where a is an inner automorphism of $\mathrm{GL}_n(\mathbb{C})$, the mappings Γ_α are arbitrary. For the other cases they satisfy some additional restrictions.

The mapping c_+ has the following block-matrix structure:

$$c_+ = \begin{pmatrix} 0 & C_{+1} & & & \\ & 0 & \ddots & & \\ & & \ddots & \ddots & \\ & & & 0 & C_{+(p-1)} \\ C_{+0} & & & & 0 \end{pmatrix}, \quad (2.11)$$

where for each $\alpha = 1, \dots, p - 1$ the mapping $C_{+\alpha}$ is a mapping of \mathbb{R}^2 to the space of $n_\alpha \times n_{\alpha+1}$ complex matrices, and C_{+0} is a mapping of \mathbb{R}^2 to the space of $n_p \times n_1$ complex matrices. The mapping c_- has a similar block-matrix structure:

$$c_- = \begin{pmatrix} 0 & & & & C_{-0} \\ C_{-1} & 0 & & & \\ & \ddots & \ddots & & \\ & & \ddots & 0 & \\ & & & C_{-(p-1)} & 0 \end{pmatrix}, \quad (2.12)$$

where for each $\alpha = 1, \dots, p - 1$ the mapping $C_{-\alpha}$ is a mapping of \mathbb{R}^2 to the space of $n_{\alpha+1} \times n_\alpha$ complex matrices, and C_{-0} is a mapping of \mathbb{R}^2 to the space of $n_1 \times n_p$ complex matrices. The conditions (2.9) imply

$$\partial_+ C_{-\alpha} = 0, \quad \partial_- C_{+\alpha} = 0, \quad \alpha = 0, 1, \dots, p - 1. \quad (2.13)$$

For the case of Toda equations associated with the loop groups $\mathcal{L}_{a,M}(\mathrm{GL}_n(\mathbb{C}))$, where a is an inner automorphism of $\mathrm{GL}_n(\mathbb{C})$, the mappings $C_{\pm\alpha}$ are arbitrary. For the other cases they should satisfy some additional restrictions.

The Toda equation (2.8) is equivalent to the following system of equations for the mappings Γ_α :

$$\begin{aligned} \partial_+ (\Gamma_1^{-1} \partial_- \Gamma_1) &= -\Gamma_1^{-1} C_{+1} \Gamma_2 C_{-1} + C_{-0} \Gamma_p^{-1} C_{+0} \Gamma_1, \\ \partial_+ (\Gamma_2^{-1} \partial_- \Gamma_2) &= -\Gamma_2^{-1} C_{+2} \Gamma_3 C_{-2} + C_{-1} \Gamma_1^{-1} C_{+1} \Gamma_2, \\ &\vdots \\ \partial_+ (\Gamma_{p-1}^{-1} \partial_- \Gamma_{p-1}) &= -\Gamma_{p-1}^{-1} C_{+(p-1)} \Gamma_p C_{-(p-1)} + C_{-(p-2)} \Gamma_{p-2}^{-1} C_{+(p-2)} \Gamma_{p-1}, \\ \partial_+ (\Gamma_p^{-1} \partial_- \Gamma_p) &= -\Gamma_p^{-1} C_{+0} \Gamma_1 C_{-0} + C_{-(p-1)} \Gamma_{p-1}^{-1} C_{+(p-1)} \Gamma_p. \end{aligned} \quad (2.14)$$

It appears that in the case under consideration without any loss of generality we can assume that the positive integer L , entering the construction of Toda equations, is equal to 1. Note also that if any of the mappings $C_{+\alpha}$ or $C_{-\alpha}$ is a zero mapping, then the equations (2.14) are equivalent to a Toda equation associated with a finite-dimensional group or to a set of two such equations.

3. Abelian Toda equations associated with loop groups of complex general linear groups

It can be shown that there are three types of abelian loop Toda equations associated with the groups $GL_n(\mathbb{C})$, see, for example, the paper [3].

3.1 First type: untwisted loop Toda equations

The abelian Toda equations of the first type arise when the automorphism A is defined by the equality

$$A(x) = hxh^{-1}, \quad x \in \mathfrak{gl}_n(\mathbb{C}),$$

where h is a diagonal matrix with the diagonal matrix elements

$$h_{kk} = \epsilon_n^{n-k+1}, \quad k = 1, \dots, n. \tag{3.1}$$

The corresponding automorphism a of $GL_n(\mathbb{C})$ is defined by the equality

$$a(g) = hgh^{-1}, \quad g \in GL_n(\mathbb{C}),$$

where again h is a diagonal matrix determined by the relation (3.1). Here the integer M is equal to n , and A is an inner automorphism which generates a \mathbb{Z}_n -gradation of $\mathfrak{gl}_n(\mathbb{C})$. The block-matrix structure related to this gradation is the matrix structure itself. In other words, all blocks are of size one by one. The mappings Γ_α are mappings of \mathbb{R}^2 to the Lie group $GL_1(\mathbb{C})$ which is isomorphic to the Lie group $\mathbb{C}^\times = \mathbb{C} \setminus \{0\}$. The mappings $C_{\pm\alpha}$ are just complex functions on \mathbb{R}^2 . The Toda equations under consideration have the form (2.14) with $p = n$.

We have shown in the paper [3] that if the functions $C_{-\alpha}$ and $C_{+\alpha}$ have no zeros then the Toda equations (2.14) are equivalent to the same equations, but where $C_{-\alpha} = C_-$ and $C_{+\alpha} = C_+$ for some functions C_- and C_+ which have no zeros. If these functions are real, then with the help of an appropriate change of the coordinates z^- and z^+ we can come to the Toda equations with $C_{\pm\alpha}$ equal to a nonzero constant m . This system of equations gives the Toda equations associated with untwisted loop groups of general linear groups. In the paper [3] we investigated the soliton solutions of the above Toda equations obtained by two different approaches, the Hirota's and rational dressing methods, and established explicit relationships between these methods.

3.2 Second type: twisted loop Toda equations, odd-dimensional case

The abelian Toda equations of the other two types arise when we use outer automorphisms of $\mathfrak{gl}_n(\mathbb{C})$. For the equations of the second type n is odd, and for the equations of the third type n is even.

Consider first the case of an odd $n = 2s - 1$, $s \geq 2$. In this case an abelian Toda equation arises when the automorphism A is defined by the equality

$$A(x) = -h(B^{-1t_x}B)h^{-1}, \tag{3.2}$$

where ${}^t x$ means the transpose of x , h is a diagonal matrix with the diagonal matrix elements

$$h_{kk} = \epsilon_{2n}^{n-k+1} = \epsilon_{4s-2}^{2s-k},$$

and B is an $n \times n$ matrix of the form

$$B = \left(\begin{array}{c|cccc} 1 & & & & \\ \hline & & & & 1 \\ & & & & \dots \\ & & & 1 & \dots \\ & & \dots & -1 & \\ & \dots & & & \\ -1 & & & & \end{array} \right).$$

The corresponding group automorphism a is defined as

$$a(g) = h(B^{-1}t_g^{-1}B)h^{-1}. \tag{3.3}$$

The order M of the automorphism A is $2N = 4s - 2$ and the integer p is $2s - 1$. The mapping γ is a diagonal matrix of the form (2.10), where the mappings Γ_α are mappings of \mathbb{R}^2 to \mathbb{C}^\times , subject to the constraints

$$\Gamma_1 = 1, \quad \Gamma_{2s-\alpha+1} = \Gamma_\alpha^{-1}, \quad \alpha = 2, \dots, s.$$

The mappings $C_{\pm\alpha}$ in the relations (2.11) and (2.12) are complex functions satisfying the equality

$$C_{\pm 0} = C_{\pm 1}, \tag{3.4}$$

and for $s > 2$ the equalities

$$C_{\pm(2s-\alpha)} = -C_{\pm\alpha}, \quad \alpha = 2, \dots, s - 1. \tag{3.5}$$

Let us choose the mappings Γ_α , $\alpha = 2, \dots, s$, as a complete set of mappings parameterizing the mapping γ . Taking into account the equalities (3.4) and (3.5) we come to a set of $s - 1$ independent equations equivalent to the Toda equation under consideration. As well as in the untwisted case, under appropriate conditions the Toda equations under consideration are equivalent to the same equations, but where

$$C_{\pm 0} = C_{\pm 1} = C_{\pm s} = m, \tag{3.6}$$

and

$$C_{\pm\alpha} = -C_{\pm(2s-\alpha)} = m, \quad \alpha = 2, \dots, s - 1. \tag{3.7}$$

Explicitly, we have the equations

$$\begin{aligned} \partial_+(\Gamma_2^{-1}\partial_-\Gamma_2) &= -m^2(\Gamma_2^{-1}\Gamma_3 - \Gamma_2), \\ \partial_+(\Gamma_3^{-1}\partial_-\Gamma_3) &= -m^2(\Gamma_3^{-1}\Gamma_4 - \Gamma_2^{-1}\Gamma_3), \\ &\vdots \\ \partial_+(\Gamma_{s-1}^{-1}\partial_-\Gamma_{s-1}) &= -m^2(\Gamma_{s-1}^{-1}\Gamma_s - \Gamma_{s-2}^{-1}\Gamma_{s-1}), \\ \partial_+(\Gamma_s^{-1}\partial_-\Gamma_s) &= -m^2(\Gamma_s^{-2} - \Gamma_{s-1}^{-1}\Gamma_s), \end{aligned} \tag{3.8}$$

The mappings C_{-1}, C_{+0} are complex 1×2 matrix-valued functions, the mappings C_{-0}, C_{+1} are complex 2×1 matrix-valued functions. Here we have

$$C_{-0} = J_2^{-1t} C_{-1}, \quad C_{+0} = {}^t C_{+1} J_2. \quad (3.9)$$

The mappings $C_{\pm\alpha}, \alpha = 2, \dots, p-1 = 2s-2$, are just complex functions, satisfying for $s > 2$ the equalities

$$C_{\pm(2s-\alpha)} = -C_{\pm\alpha}, \quad \alpha = 2, \dots, s-1. \quad (3.10)$$

The mappings $(\Gamma_1)_{11}$ and $\Gamma_\alpha, \alpha = 2, \dots, s$, form a complete set of mappings parameterizing the mapping γ . Taking into account the equalities (3.9) and (3.10) we come to a set of s independent equations equivalent to the Toda equation under consideration. As well as for the first two types, under appropriate conditions these equations can be reduced to equations with

$$C_{-\alpha} = m, \quad C_{+\alpha} = m, \quad \alpha = 2, \dots, s, \quad (3.11)$$

and

$$(C_{-1})_{11} = (C_{-1})_{12} = m/\sqrt{2}, \quad (C_{+1})_{11} = (C_{+1})_{21} = m/\sqrt{2}, \quad (3.12)$$

where m is a nonzero constant. Thus, we come to the equations

$$\begin{aligned} \partial_+(\Gamma_1^{-1}\partial_-\Gamma_1) &= -\frac{m^2}{2}(\Gamma_1^{-1} - \Gamma_1)\Gamma_2, \\ \partial_+(\Gamma_2^{-1}\partial_-\Gamma_2) &= -m^2\Gamma_2^{-1}\Gamma_3 + \frac{m^2}{2}(\Gamma_1^{-1} + \Gamma_1)\Gamma_2, \\ \partial_+(\Gamma_3^{-1}\partial_-\Gamma_3) &= -m^2(\Gamma_3^{-1}\Gamma_4 - \Gamma_2^{-1}\Gamma_3), \\ &\vdots \\ \partial_+(\Gamma_{s-1}^{-1}\partial_-\Gamma_{s-1}) &= -m^2(\Gamma_{s-1}^{-1}\Gamma_s - \Gamma_{s-2}^{-1}\Gamma_{s-1}), \\ \partial_+(\Gamma_s^{-1}\partial_-\Gamma_s) &= -m^2(\Gamma_s^{-2} - \Gamma_{s-1}^{-1}\Gamma_s), \end{aligned} \quad (3.13)$$

where, with a slight abuse of notation, we have denoted $(\Gamma_1)_{11}$ by Γ_1 .

We also note that all three systems of Toda equations described above can be represented in standard forms with explicit indication of the Cartan matrices of the corresponding affine Lie algebras of the types $A_{n-1}^{(1)}, A_{2s-2}^{(2)}$ and $A_{2s-1}^{(2)}$, respectively, see, for example, the paper [3].

4. Rational dressing

In this section we apply the method of rational dressing to construct solutions of the abelian Toda systems associated with the loop groups of the complex general linear groups. Here we solve the abelian Toda equations of the second and third types which have the forms (3.8) and (3.13) respectively. In fact, some preliminary relations of the rational dressing formalism can be introduced on a common basis in application to the both types of abelian Toda systems.

Because in the cases under consideration the matrices c_- and c_+ are commuting, it is obvious that

$$\gamma = I_n, \tag{4.1}$$

where I_n is the $n \times n$ unit matrix, is a solution to the Toda equation (2.8). Denote a mapping of $\mathbb{R}^2 \times S^1$ to $GL_n(\mathbb{C})$, which generates the corresponding connection, by φ . Using the equalities (2.6) and (2.7) and remembering that in our case $L = 1$, we write

$$\varphi^{-1}\partial_-\varphi = \lambda^{-1}c_-, \quad \varphi^{-1}\partial_+\varphi = \lambda c_+, \tag{4.2}$$

where the matrices c_+ and c_- having generally the forms (2.11) and (2.12), are specified by the relations (3.6), (3.7) for the Toda equations of the second type, and by the relations (3.11), (3.12) for the Toda equations of the third type.

To construct some other solutions to the Toda equations we will look for a mapping ψ , such that the mapping

$$\varphi' = \varphi\psi \tag{4.3}$$

would generate a connection satisfying the grading condition

$$\omega_- = \omega_{-0} + \omega_{-1}, \quad \omega_+ = \omega_{+0} + \omega_{+1} \tag{4.4}$$

and the gauge-fixing constraint

$$\omega_{+0} = 0. \tag{4.5}$$

For any $\bar{m} \in \mathbb{R}^2$ the mapping $\tilde{\psi}_m$ defined by the equality $\tilde{\psi}_m(\bar{p}) = \psi(\bar{m}, \bar{p})$, $\bar{p} \in S^1$, is a smooth mapping of S^1 to $GL_n(\mathbb{C})$. We treat S^1 as a subset of the complex plane which, in turn, will be treated as a subset of the Riemann sphere. Assume that it is possible to extend analytically each mapping $\tilde{\psi}_m$ to all of the Riemann sphere. As the result we obtain a mapping of the direct product of \mathbb{R}^2 and the Riemann sphere to $GL_n(\mathbb{C})$, which we also denote by ψ . Suppose that for any $\bar{m} \in \mathbb{R}^2$ the analytic extension of $\tilde{\psi}_m$ results in a rational mapping regular at the points 0 and ∞ , hence the name *rational dressing*. Below, for each point \bar{p} of the Riemann sphere we denote by ψ_p the mapping of \mathbb{R}^2 to $GL_n(\mathbb{C})$ defined by the equality $\psi_p(\bar{m}) = \psi(\bar{m}, \bar{p})$.

Since we deal with the Toda equations described in sections 3.2 and 3.3, for any $\bar{m} \in \mathbb{R}^2$ and $\bar{p} \in S^1$ we should have

$$\psi(\bar{m}, \epsilon_{2N}\bar{p}) = hB^{-1}t\psi^{-1}(\bar{m}, \bar{p})Bh^{-1}, \tag{4.6}$$

where h is a block-diagonal matrix described by the relation

$$h_{\alpha\beta} = \epsilon_{2N}^{N-\alpha+1} I_{n_\alpha} \delta_{\alpha\beta}, \quad \alpha, \beta = 1, \dots, p,$$

with $n_1 = 1$ for the Toda equations of the second type, and $n_1 = 2$ for the Toda equations of the third type, while for all other indices $\alpha = 2, \dots, p$ we always have $n_\alpha = 1$. Note that

$h_{11} = -I_{n_1}$. Here we also use the notation

$$B = \left(\begin{array}{c|cccc} J_{n_1} & & & & \\ \hline & & & & 1 \\ & & & & \cdot \\ & & & 1 & \cdot \\ & & -1 & & \cdot \\ & \cdot & & & \\ -1 & \cdot & & & \end{array} \right)$$

common for the both cases. The equality (4.6) means that for any $\bar{m} \in \mathbb{R}^2$ two rational mappings coincide on S^1 , therefore, they must coincide on the entire Riemann sphere.

We define a linear mapping \hat{a} acting on a mapping χ of the direct product of \mathbb{R}^2 and the Riemann sphere to the algebra $\text{Mat}_n(\mathbb{C})$ of $n \times n$ complex matrices as⁴

$$\hat{a}\chi(\bar{m}, \bar{p}) = hB^{-1t}\chi^{-1}(\bar{m}, \epsilon_{2N}^{-1}\bar{p})Bh^{-1}.$$

The relation (4.6) is equivalent to the equality $\hat{a}\psi = \psi$. To construct rational mappings satisfying this relation we will use the following procedure. First, we construct a family of mappings ψ satisfying the relation $\hat{a}^2\psi = \psi$, and then select from it the mappings satisfying the equality $\hat{a}\psi = \psi$.

It is easy to see that the mapping

$$\psi = \sum_{k=1}^N \hat{a}^{2k}\chi \tag{4.7}$$

satisfies the relation $\hat{a}^2\psi = \psi$. It is worth to note that $\hat{a}^{2N}\chi = \chi$. We start with a rational mapping χ regular at the points 0 and ∞ and having poles at r different nonzero points $\mu_i, i = 1, \dots, r$. More specifically, we consider a mapping χ of the form

$$\chi = \left(I_n + N \sum_{i=1}^r \frac{\lambda}{\lambda - \mu_i} P_i \right) \chi_0,$$

where P_i are some smooth mappings of \mathbb{R}^2 to the algebra $\text{Mat}_n(\mathbb{C})$ and χ_0 is a mapping of \mathbb{R}^2 to the Lie subgroup of $\text{GL}_n(\mathbb{C})$ formed by the elements $g \in \text{GL}_n(\mathbb{C})$ satisfying the equality

$$h^2gh^{-2} = g. \tag{4.8}$$

With account of the equality

$$\hat{a}^2\chi(\bar{m}, \bar{p}) = h^2\chi(\bar{m}, \epsilon_N^{-1}\bar{p})h^{-2}$$

the averaging procedure (4.7) leads to the mapping

$$\psi = \left(I_n + \sum_{i=1}^r \sum_{k=1}^N \frac{\lambda}{\lambda - \epsilon_{2N}^{2k}\mu_i} h^{2k} P_i h^{-2k} \right) \psi_0, \tag{4.9}$$

⁴Note that below χ is a mapping to the Lie group $\text{GL}_n(\mathbb{C})$, although to justify the relation (4.7) it is convenient to think $\text{GL}_n(\mathbb{C})$ as a subset of $\text{Mat}_n(\mathbb{C})$.

where $\psi_0 = N\chi_0$. We assume that $\mu_i^{2N} \neq \mu_j^{2N}$ for all $i \neq j$.

Denote by ψ^{-1} the mapping of $\mathbb{R}^2 \times S^1$ to $\text{GL}_n(\mathbb{C})$ defined by the relation

$$\psi^{-1}(\bar{m}, \bar{p}) = (\psi(\bar{m}, \bar{p}))^{-1}.$$

Suppose that for any fixed $\bar{m} \in \mathbb{R}^2$ the mapping $\tilde{\psi}_m^{-1}$ of S^1 to $\text{GL}_n(\mathbb{C})$, defined by the equality $\tilde{\psi}_m^{-1}(\bar{p}) = \psi^{-1}(\bar{m}, \bar{p})$, can be extended analytically to a mapping of the Riemann sphere to $\text{GL}_n(\mathbb{C})$, which we also denote by ψ^{-1} , and as the result we obtain a rational mapping of the same structure as the mapping ψ ,

$$\psi^{-1} = \psi_0^{-1} \left(I_n + \sum_{i=1}^r \sum_{k=1}^N \frac{\lambda}{\lambda - \epsilon_{2N}^{2k} \nu_i} h^{2k} Q_i h^{-2k} \right), \quad (4.10)$$

with the pole positions satisfying the conditions $\nu_i \neq 0$, $\nu_i^{2N} \neq \nu_j^{2N}$ for all $i \neq j$, and additionally $\nu_i^N \neq \mu_j^N$ for any i and j .⁵

The mappings ψ and ψ^{-1} given by the equalities (4.9) and (4.10), respectively, satisfy the relations $\hat{a}^2 \psi = \psi$ and $\hat{a}^2 \psi^{-1} = \psi^{-1}$. To satisfy the relations $\hat{a} \psi = \psi$ and $\hat{a} \psi^{-1} = \psi^{-1}$ we have to assume that the pole positions of the mappings ψ and ψ^{-1} are necessarily connected as

$$\nu_i = \mu_i / \epsilon_{2N}, \quad i = 1, \dots, r,$$

and the matrices P_i and Q_i are related as

$$Q_i = h^{-1} B^{-1} {}^t P_i B h, \quad i = 1, \dots, r, \quad (4.11)$$

By definition, the equality

$$\psi^{-1} \psi = I_n$$

is valid at all points of the direct product of \mathbb{R}^2 and S^1 . Since $\psi^{-1} \psi$ is a rational mapping, the above equality is valid at all points of the direct product of \mathbb{R}^2 and the Riemann sphere. Hence, the residues of $\psi^{-1} \psi$ at the points $\nu_i = \mu_i / \epsilon_{2N}$ and μ_i should be equal to zero. Explicitly we have

$$h^{-1} B^{-1} {}^t P_i B h \left(I_n + \sum_{j=1}^r \sum_{k=1}^N \frac{\mu_i / \epsilon_{2N}}{\mu_i / \epsilon_{2N} - \epsilon_{2N}^{2k} \mu_j} h^{2k} P_j h^{-2k} \right) = 0, \quad (4.12)$$

$$\left(I_n + \sum_{j=1}^r \sum_{k=1}^N \frac{\mu_i}{\mu_i - \epsilon_{2N}^{2k-1} \mu_j} h^{2k-1} B^{-1} {}^t P_j B h^{-2k+1} \right) P_i = 0. \quad (4.13)$$

We will discuss later how to satisfy these relations, and now let us consider what connection is generated by the mapping φ' defined by (4.3) with the mapping ψ possessing the properties described above.

⁵Actually, as it will be clear, for the extended mappings ψ and ψ^{-1} we have $\psi^{-1} \psi = I_n$. This justifies the notation used.

Using the equality (4.3) and the relations (4.2), we obtain for the components of the connection generated by φ' the expressions

$$\omega_- = \psi^{-1} \partial_- \psi + \lambda^{-1} \psi^{-1} c_- \psi, \quad (4.14)$$

$$\omega_+ = \psi^{-1} \partial_+ \psi + \lambda \psi^{-1} c_+ \psi. \quad (4.15)$$

We see that the component ω_- is a rational mapping which has simple poles at the points μ_i , $\nu_i = \mu_i/\epsilon_{2N}$ and zero.⁶ Similarly, the component ω_+ is a rational mapping which has simple poles at the points μ_i , $\nu_i = \mu_i/\epsilon_{2N}$ and infinity. We are looking for a connection which satisfies the grading condition (4.4) and the gauge-fixing condition (4.5). The grading condition in our case is the requirement that for each point of \mathbb{R}^2 the component ω_- is rational and has the only simple pole at zero, while the component ω_+ is rational and has the only simple pole at infinity. Hence, we demand that the residues of ω_- and ω_+ at the points μ_i and $\nu_i = \mu_i/\epsilon_{2N}$ should vanish.

The residues of ω_- and ω_+ at the points $\nu_i = \mu_i/\epsilon_{2N}$ are equal to zero if and only if

$$(\partial_- Q_i - \epsilon_{2N} \mu_i^{-1} Q_i c_-) \left(I_n + \sum_{j=1}^r \sum_{k=1}^N \frac{\mu_i/\epsilon_{2N}}{\mu_i/\epsilon_{2N} - \epsilon_{2N}^{2k} \mu_j} h^{2k} P_j h^{-2k} \right) = 0, \quad (4.16)$$

$$(\partial_+ Q_i - \epsilon_{2N}^{-1} \mu_i Q_i c_+) \left(I_n + \sum_{j=1}^r \sum_{k=1}^N \frac{\mu_i/\epsilon_{2N}}{\mu_i/\epsilon_{2N} - \epsilon_{2N}^{2k} \mu_j} h^{2k} P_j h^{-2k} \right) = 0, \quad (4.17)$$

respectively, with the equality (4.11) to be taken into account. Similarly, the requirement of vanishing of the residues at the points μ_i gives the relations

$$\left(I_n + \sum_{j=1}^r \sum_{k=1}^N \frac{\mu_i}{\mu_i - \epsilon_{2N}^{2k-1} \mu_j} h^{2k-1} B^{-1t} P_j B h^{-2k+1} \right) (\partial_- P_i + \mu_i^{-1} c_- P_i) = 0, \quad (4.18)$$

$$\left(I_n + \sum_{j=1}^r \sum_{k=1}^N \frac{\mu_i}{\mu_i - \epsilon_{2N}^{2k-1} \mu_j} h^{2k-1} B^{-1t} P_j B h^{-2k+1} \right) (\partial_+ P_i + \mu_i c_+ P_i) = 0. \quad (4.19)$$

To obtain the relations (4.16)–(4.19) we made use of the equalities (4.12), (4.13).

Suppose that we have succeeded in satisfying the relations (4.12), (4.13) and (4.16)–(4.19). In such a case from the equalities (4.14) and (4.15) it follows that the connection under consideration satisfies the grading condition.

It follows from the equality (4.15) that

$$\omega_+(\bar{m}, 0) = \psi_0^{-1}(\bar{m}) \partial_+ \psi_0(\bar{m}).$$

Taking into account that $\omega_{+0}(\bar{m}) = \omega_+(\bar{m}, 0)$, we conclude that the gauge-fixing constraint $\omega_{+0} = 0$ is equivalent to the relation

$$\partial_+ \psi_0 = 0. \quad (4.20)$$

⁶Here and below discussing the holomorphic properties of mappings and functions we assume that the point of the space \mathbb{R}^2 is arbitrary but fixed.

Assuming that this relation is satisfied, we come to a connection satisfying both the grading condition and the gauge-fixing condition.

Recall that if a flat connection ω satisfies the grading and gauge-fixing conditions, then there exist a mapping γ from \mathbb{R}^2 to G and mappings c_- and c_+ of \mathbb{R}^2 to \mathfrak{g}_{-1} and \mathfrak{g}_{+1} , respectively, such that the representation (2.7) for the components ω_- and ω_+ is valid. In general, the mappings c_- and c_+ parameterizing the connection components may be different from the mappings c_- and c_+ which determine the mapping φ . Let us denote the mappings corresponding to the connection under consideration by γ' , c'_- and c'_+ . Thus, we have

$$\psi^{-1}\partial_-\psi + \lambda^{-1}\psi^{-1}c_-\psi = \gamma'^{-1}\partial_-\gamma' + \lambda^{-1}c'_-, \tag{4.21}$$

$$\psi^{-1}\partial_+\psi + \lambda\psi^{-1}c_+\psi = \lambda\gamma'^{-1}c'_+\gamma'. \tag{4.22}$$

Note that ψ_∞ is a mapping of \mathbb{R}^2 to the Lie subgroup of $GL_n(\mathbb{C})$ defined by the relation (4.8). We recall that this subgroup coincides with G_0 and denote ψ_∞ by γ . From the relation (4.21) we obtain the equality

$$\gamma'^{-1}\partial_-\gamma' = \gamma^{-1}\partial_-\gamma.$$

The same relation (4.21) gives

$$\psi_0^{-1}c_-\psi_0 = c'_-.$$

Impose the condition $\psi_0 = I_n$, which is consistent with the condition (4.20). Here we have

$$c'_- = c_-.$$

Finally, from the equality (4.22) we obtain

$$\gamma'^{-1}c'_+\gamma' = \gamma^{-1}c_+\gamma.$$

We see that if we impose the condition $\psi_0 = I_n$, then the components of the connection under consideration have the form (2.7) where $\gamma = \psi_\infty$.

Thus, to find solutions to the Toda equations under consideration, we can use the following procedure. We fix $2r$ complex numbers μ_i and ν_i and find matrix-valued functions P_i and Q_i satisfying the relations (4.12), (4.13) and (4.16)–(4.19). With the help of the relations (4.9), (4.10), assuming that

$$\psi_0 = I_n,$$

we construct the mappings ψ and ψ^{-1} . Then, the mapping

$$\gamma = \psi_\infty \tag{4.23}$$

satisfies the Toda equation (2.8).

Let us return to the relations (4.12), (4.13). It is easy to see that they are equivalent, and so, we will use the relation (4.13) for further calculations. We can show that, if we suppose that the matrix P_i has the maximum rank, then we get the trivial solution of the

Toda equation given by (4.1). Hence, we will assume that P_i is not of maximum rank. The simplest case here is given by matrices of rank one which can be represented as

$$P_i = u_i {}^t w_i,$$

where u and w are n -dimensional column vectors. This representation allows writing the relations (4.13) as

$$u_i + \sum_{j=1}^r \sum_{k=1}^N \frac{\mu_i}{\mu_i - \epsilon_{2N}^{2k-1} \mu_j} h^{2k-1} B^{-1} w_j ({}^t u_j B h^{-2k+1} u_i) = 0. \quad (4.24)$$

Using the identity

$$\sum_{k=1}^N \frac{z \epsilon_{2N}^{-2kj}}{z - \epsilon_{2N}^{2k}} = N \frac{z^{N-|j|_N}}{z^N - 1},$$

where $|j|_N$ is the residue of division of j by N , we can rewrite the equality (4.24) in terms of the components of u_i as follows:

$${}^t u_{i,1} J_{n_1} + N \sum_{j=1}^r (R_1)_{ij} {}^t w_{j,1} = 0,$$

where $u_{i,1}$ and $w_{i,1}$ gather first n_1 components of the corresponding n -dimensional column vectors, so these are in fact n_1 -dimensional column vectors,⁷

$$u_{i,N+2-k} - N \sum_{j=1}^r (R_k)_{ij} w_{j,k} = 0, \quad k = 2, \dots, s,$$

and

$$u_{i,N+2-k} + N \sum_{j=1}^r (R_k)_{ij} w_{j,k} = 0, \quad k = s+1, \dots, p = N.$$

Here the $r \times r$ matrices R_1 and R_k are defined as

$$(R_1)_{ij} = \frac{1}{\mu_i^N + \mu_j^N} \left(\mu_i^N ({}^t u_{i,1} J_{n_1} u_{j,1}) - \sum_{\ell=2}^s \mu_i^{N-|\ell-1|_N} \mu_j^{|\ell-1|_N} (u_{i,N+2-\ell} u_{j,\ell}) + \sum_{\ell=s+1}^N \mu_i^{N-|\ell-1|_N} \mu_j^{|\ell-1|_N} (u_{i,N+2-\ell} u_{j,\ell}) \right), \quad (4.25)$$

$$(R_k)_{ij} = \frac{1}{\mu_i^N + \mu_j^N} \left(-\mu_i^{N-|1-k|_N} \mu_j^{|1-k|_N} ({}^t u_{i,1} J_{n_1} u_{j,1}) + \sum_{\ell=2}^{k-1} \mu_i^{N-|\ell-k|_N} \mu_j^{|\ell-k|_N} (u_{i,N+2-\ell} u_{j,\ell}) - \sum_{\ell=k}^s \mu_i^{N-|\ell-k|_N} \mu_j^{|\ell-k|_N} (u_{i,N+2-\ell} u_{j,\ell}) + \sum_{\ell=s+1}^N \mu_i^{N-|\ell-k|_N} \mu_j^{|\ell-k|_N} (u_{i,N+2-\ell} u_{j,\ell}) \right) \quad (4.26)$$

⁷We remember that either $n_1 = 1$ or $n_1 = 2$.

for $k = 2, \dots, s$, and

$$\begin{aligned}
 (R_k)_{ij} = & \frac{1}{\mu_i^N + \mu_j^N} \left(-\mu_i^{N-|1-k|N} \mu_j^{|1-k|N} ({}^t u_{i,1} J_{n_1} u_{j,1}) \right. \\
 & + \sum_{\ell=2}^s \mu_i^{N-|\ell-k|N} \mu_j^{|\ell-k|N} (u_{i,N+2-\ell} u_{j,\ell}) - \sum_{\ell=s+1}^{k-1} \mu_i^{N-|\ell-k|N} \mu_j^{|\ell-k|N} (u_{i,N+2-\ell} u_{j,\ell}) \\
 & \left. + \sum_{\ell=k}^N \mu_i^{N-|\ell-k|N} \mu_j^{|\ell-k|N} (u_{i,N+2-\ell} u_{j,\ell}) \right) \quad (4.27)
 \end{aligned}$$

for $k = s + 1, \dots, N$. Recall that for all cases considered here $N = p = 2s - 1$.

We use the equations (4.25), (4.26) and (4.27) to express the vectors w_i via the vectors u_i ,

$${}^t w_{i,1} = -\frac{1}{N} \sum_{j=1}^r (R_1^{-1})_{ij} {}^t u_{j,1} J_{n_1}, \quad w_{i,k} = \frac{1}{N} \sum_{j=1}^r (R_k^{-1})_{ij} u_{j,N+2-k}$$

for $k = 2, \dots, s$, and

$$w_{i,k} = -\frac{1}{N} \sum_{j=1}^r (R_k^{-1})_{ij} u_{j,N+2-k}$$

for $k = s + 1, \dots, N = p$. As the result, having expressed the matrices P_i and Q_i in terms of the components of the vectors u_i , we find a solution of the relations (4.12) and (4.13).

Further, it follows from the equality (4.24) that, to fulfill also the relations (4.16)–(4.19), it is sufficient to satisfy the equations

$$\partial_- u_i = -\mu_i^{-1} c_- u_i, \quad \partial_+ u_i = -\mu_i c_+ u_i.$$

The general solution to these equations is given formally by the expression

$$u_i(z^-, z^+) = \exp(-\mu_i^{-1} c_- z^- - \mu_i c_+ z^+) u_i^0, \quad (4.28)$$

where $u_i^0 = u_i(0, 0)$. We will make explicit this formal solution when later constructing soliton solutions.

Thus, we see that it is possible to satisfy the relations (4.12), (4.13) and (4.16)–(4.19). This gives us solutions of the Toda equation (2.8), and so, to the equations (3.8) and (3.13) by specifying the above formal expression of u_i for the two corresponding cases. Let us show that they can be written in a simple determinant form.

Using the equalities (4.23) and (4.9), we get

$$\gamma = \psi_\infty = I_n + \sum_{i=1}^r \sum_{k=1}^N h^{2k} P_i h^{-2k}.$$

For the matrix elements of γ this gives the expression

$$\gamma_{k\ell} = \delta_{k\ell} \left(1 + N \sum_{i=1}^r (P_i)_{kk} \right) = \delta_{k\ell} \Gamma_k.$$

Hence, we have

$$\Gamma_1 = I_{n_1} - \sum_{i,j=1}^r u_{i,1} (R_1^{-1})_{ij} {}^t u_{j,1} J_{n_1}, \quad \Gamma_\alpha = 1 + \sum_{i,j=1}^r u_{i,\alpha} (R_\alpha^{-1})_{ij} u_{j,2s+1-\alpha},$$

where $\alpha = 2, \dots, s$, and

$$\Gamma_\alpha = 1 - \sum_{i,j=1}^r u_{i,\alpha} (R_\alpha^{-1})_{ij} u_{j,2s+1-\alpha}, \quad \alpha = s+1, \dots, 2s-1.$$

We assume for convenience that the functions $u_{i,\alpha}$ are defined for arbitrary integral values of α so that

$$u_{i,2s-1+\alpha} = u_{i,\alpha}.$$

By definition the matrices R_α are periodic in the index α . It appears that it is more appropriate to use quasi-periodic quantities \tilde{u}_α and \tilde{R}_α defined as

$$\tilde{u}_\alpha = M^\alpha u_\alpha, \quad \tilde{R}_1 = M R_1 M^{2s}, \quad \tilde{R}_\alpha = M^{2s+1-\alpha} R_\alpha M^\alpha,$$

where $\alpha = 2, \dots, 2s-1$; here M is a diagonal $r \times r$ matrix given by

$$M_{ij} = \mu_i \delta_{ij}.$$

For these quantities we have quasi-periodicity conditions

$$\tilde{u}_{2s-1+\alpha} = M^{2s-1} \tilde{u}_\alpha, \quad \tilde{R}_{2s} = M^{2s-1} \tilde{R}_1 M^{2s-1}, \quad \tilde{R}_{2s-1+\alpha} = M^{-2s+1} \tilde{R}_\alpha M^{2s-1}.$$

The expression of the matrix elements of the matrices \tilde{R}_α through the quasi-periodic quantities $\tilde{u}_{i\alpha}$ has a remarkably simple form. We have for $\alpha = 1$

$$(\tilde{R}_1)_{ij} = \frac{1}{\mu_i^{2s-1} + \mu_j^{2s-1}} \left(\mu_i^{2s-1} ({}^t \tilde{u}_{i,1} J_{n_1} \tilde{u}_{j,1}) \mu_j^{2s-1} - \mu_j^{2s-1} \sum_{\beta=2}^s \tilde{u}_{i,2s+1-\beta} \tilde{u}_{j,\beta} + \mu_j^{2s-1} \sum_{\beta=s+1}^{2s-1} \tilde{u}_{i,2s+1-\beta} \tilde{u}_{j,\beta} \right).$$

Further, we have for $\alpha = 2, \dots, s$

$$(\tilde{R}_\alpha)_{ij} = \frac{1}{\mu_i^{2s-1} + \mu_j^{2s-1}} \left(-\mu_i^{2s-1} ({}^t \tilde{u}_{i,1} J_{n_1} \tilde{u}_{j,1}) \mu_j^{2s-1} + \mu_j^{2s-1} \sum_{\beta=2}^{\alpha-1} \tilde{u}_{i,2s+1-\beta} \tilde{u}_{j,\beta} - \mu_i^{2s-1} \sum_{\beta=\alpha}^s \tilde{u}_{i,2s+1-\beta} \tilde{u}_{j,\beta} + \mu_i^{2s-1} \sum_{\beta=s+1}^{2s-1} \tilde{u}_{i,2s+1-\beta} \tilde{u}_{j,\beta} \right),$$

and for $\alpha = s + 1, \dots, 2s - 1$

$$(\tilde{R}_\alpha)_{ij} = \frac{1}{\mu_i^{2s-1} + \mu_j^{2s-1}} \left(-\mu_i^{2s-1} ({}^t\tilde{u}_{i,1} J_{n_1} \tilde{u}_{j,1}) \mu_j^{2s-1} + \mu_j^{2s-1} \sum_{\beta=2}^s \tilde{u}_{i,2s+1-\beta} \tilde{u}_{j,\beta} - \mu_j^{2s-1} \sum_{\beta=s+1}^{\alpha-1} \tilde{u}_{i,2s+1-\beta} \tilde{u}_{j,\beta} + \mu_i^{2s-1} \sum_{\beta=\alpha}^{2s-1} \tilde{u}_{i,2s+1-\beta} \tilde{u}_{j,\beta} \right).$$

Here we used the identity $|-k|_N = N - 1 - |k - 1|_N$. The quasi-periodic functions have the following useful properties:

$$(\tilde{R}_{\alpha+1})_{ij} = (\tilde{R}_\alpha)_{ij} + \tilde{u}_{i,2s+1-\alpha} \tilde{u}_{j,\alpha}, \quad \alpha = 2, \dots, s, \quad (4.29)$$

$$(\tilde{R}_{\alpha+1})_{ij} = (\tilde{R}_\alpha)_{ij} - \tilde{u}_{i,2s+1-\alpha} \tilde{u}_{j,\alpha}, \quad \alpha = s + 1, \dots, 2s - 1, \quad (4.30)$$

and

$$(\tilde{R}_1)_{ij} = -(\tilde{R}_2)_{ji}, \quad (\tilde{R}_\alpha)_{ij} = (\tilde{R}_{2s+2-\alpha})_{ji}, \quad \alpha = 2, \dots, 2s - 1. \quad (4.31)$$

In terms of the quasi-periodic quantities, for the $n_1 \times n_1$ matrix-valued function Γ_1 and for the functions Γ_α we have

$$\Gamma_1 = I_{n_1} - \sum_{i,j=1}^r \mu_i^{2s-1} \tilde{u}_{i,1} (\tilde{R}_1^{-1})_{ij} {}^t\tilde{u}_{j,1} J_{n_1}, \quad \Gamma_\alpha = 1 + \sum_{i,j=1}^r \tilde{u}_{i,\alpha} (\tilde{R}_\alpha^{-1})_{ij} \tilde{u}_{j,2s+1-\alpha},$$

for $\alpha = 2, \dots, s$, and

$$\Gamma_\alpha = 1 - \sum_{i,j=1}^r \tilde{u}_{i,\alpha} (\tilde{R}_\alpha^{-1})_{ij} \tilde{u}_{j,2s+1-\alpha},$$

for $\alpha = s + 1, \dots, 2s - 1$. The expressions for the functions Γ_α for $\alpha > 1$ can be written as

$$\Gamma_\alpha = 1 + {}^t\tilde{u}_\alpha \tilde{R}_\alpha^{-1} \tilde{u}_{2s+1-\alpha}, \quad \alpha = 2, \dots, s,$$

and

$$\Gamma_\alpha = 1 - {}^t\tilde{u}_\alpha \tilde{R}_\alpha^{-1} \tilde{u}_{2s+1-\alpha}, \quad \alpha = s + 1, \dots, 2s - 1.$$

Here \tilde{R}_α is an $r \times r$ matrix and \tilde{u}_α is an r -dimensional column vector. We remember that in the cases under consideration we should have

$$J_{n_1}^{-1} {}^t\Gamma_1 J_{n_1} = \Gamma_1^{-1}, \quad \Gamma_{2s+1-\alpha} = \Gamma_\alpha^{-1}, \quad \alpha = 2, \dots, 2s - 1. \quad (4.32)$$

To verify these relations we use that

$$\gamma^{-1} = \psi_\infty^{-1} = I_n + \sum_{i=1}^r \sum_{k=1}^{2s-1} h^{2k} (h^{-1} B^{-1} {}^tP_i B h) h^{-2k},$$

therefore we find the following expression of Γ_1^{-1} in terms of quasi-periodic quantities,

$$\Gamma_1^{-1} = I_{n_1} - \sum_{i,j=1}^r \tilde{u}_{i,1} \mu_j^{2s-1} (\tilde{R}_1^{-1})_{ji} {}^t\tilde{u}_{j,1} J_{n_1}.$$

Comparing now this expression with what we have for Γ_1 above, we conclude that the first relation of equations (4.32) is satisfied.

The expressions just given above allow writing a remarkable determinant representation for the functions Γ_α . It can be shown that

$$\Gamma_\alpha = \frac{\det(\tilde{R}_\alpha + \tilde{u}_{2s+1-\alpha} {}^t \tilde{u}_\alpha)}{\det \tilde{R}_\alpha}, \quad \alpha = 2, \dots, s,$$

and

$$\Gamma_\alpha = \frac{\det(\tilde{R}_\alpha - \tilde{u}_{2s+1-\alpha} {}^t \tilde{u}_\alpha)}{\det \tilde{R}_\alpha}, \quad \alpha = s+1, \dots, 2s-1.$$

Using the properties (4.29) and (4.30) we can see

$$\Gamma_\alpha = \frac{\det \tilde{R}_{\alpha+1}}{\det \tilde{R}_\alpha}, \quad \alpha = 2, \dots, s, s+1, \dots, 2s-1.$$

For these functions we can also easily demonstrate that

$$\Gamma_{2s+1-\alpha} = \frac{\det \tilde{R}_{2s+2-\alpha}}{\det \tilde{R}_{2s+1-\alpha}} = \frac{\det({}^t \tilde{R}_\alpha)}{\det({}^t \tilde{R}_{\alpha+1})} = \frac{\det \tilde{R}_\alpha}{\det \tilde{R}_{\alpha+1}} = \Gamma_\alpha^{-1},$$

using for this purpose the relations (4.30). Hence all equations (4.32) are fulfilled.

We remember also that for the case $n_1 = 1$ corresponding to the second type of abelian twisted loop Toda equations considered here ($n = p = 2s - 1$), we have $I_1 = J_1 = 1$, and so, we can write for the function Γ_1 the expression

$$\Gamma_1 = 1 - {}^t \tilde{u}_1 M^{2s-1} \tilde{R}_1^{-1} \tilde{u}_1,$$

where \tilde{u}_1 is also an r -dimensional column vector. It can be shown that

$$\Gamma_1 = \frac{\det(\tilde{R}_1 - \tilde{u}_1 {}^t u_1 M^{2s-1})}{\det \tilde{R}_1},$$

We obtain from the expressions of \tilde{R}_1 and \tilde{R}_2 directly that

$$(\tilde{R}_1)_{ij} \mu_j^{-N} = {}^t \tilde{u}_{i,1} J_{n_1} \tilde{u}_{j,1} + \mu_i^{-N} (\tilde{R}_2)_{ij},$$

and so, for $n_1 = 1$ we can write

$$M^{-2s+1} \tilde{R}_2 + \tilde{u}_1 {}^t \tilde{u}_1 = \tilde{R}_1 M^{-2s+1}.$$

Using this relation in the above expression for Γ_1 as the ratio of determinants, we easily derive

$$\Gamma_1 = \frac{\det \tilde{R}_2}{\det \tilde{R}_1}. \tag{4.33}$$

But we have from the equality (4.31) that

$$\tilde{R}_1 = -{}^t \tilde{R}_2,$$

and so, for $n_1 = 1$ the expression (4.33) gives

$$\Gamma_1 = (-1)^r.$$

5. Soliton solutions

5.1 Odd-dimensional case

Here we consider the case of $n = p = N = 2s - 1$. It means also that we have $n_1 = 1$. The eigenvectors of the matrices ${}^t c_-$, ${}^t c_+$, c_- and c_+ are n -dimensional column vectors Ψ_ρ , $\rho = 1, \dots, 2s - 1$, satisfying the relations

$${}^t c_- \Psi_\rho = m \epsilon_{2N}^{s+2\rho} \Psi_\rho, \quad {}^t c_+ \Psi_\rho = m \epsilon_{2N}^{-s-2\rho} \Psi_\rho, \quad c_- \Psi_\rho = m \epsilon_{2N}^{-s-2\rho} \Psi_\rho, \quad c_+ \Psi_\rho = m \epsilon_{2N}^{s+2\rho} \Psi_\rho,$$

where the $2s - 1$ components of Ψ_ρ are defined as

$$\begin{aligned} (\Psi_\rho)_\alpha &= \epsilon_{2N}^{\alpha(s+2\rho)}, & \alpha &= 1, \dots, s, \\ (\Psi_\rho)_\alpha &= (-1)^{\alpha-s-1} \epsilon_{2N}^{\alpha(s+2\rho)}, & \alpha &= s+1, \dots, 2s-1. \end{aligned}$$

Consequently, we can give a concrete expression to the formal solution (4.28) as

$$\begin{aligned} u_{i,\alpha} &= \sum_{\rho=1}^{2s-1} c_{i\rho} \epsilon_{2N}^{\alpha(s+2\rho)} e^{-Z_\rho(\mu_i)}, & \alpha &= 1, \dots, s \\ u_{i,\alpha} &= \sum_{\rho=1}^{2s-1} c_{i\rho} (-1)^{\alpha-s-1} \epsilon_{2N}^{\alpha(s+2\rho)} e^{-Z_\rho(\mu_i)}, & \alpha &= s+1, \dots, 2s-1, \end{aligned}$$

where $c_{i\rho}$ are arbitrary constants and we have introduced the notation

$$Z_\rho(\mu_i) = m(\epsilon_{2N}^{-s-2\rho} \mu_i^{-1} z^- + \epsilon_{2N}^{s+2\rho} \mu_i z^+).$$

Then, after some calculation using, in particular, properties of ϵ_{2N} , we write for the matrix elements of \tilde{R}_α for $\alpha \geq 2$:

$$(\tilde{R}_\alpha)_{ij} = (-1)^\alpha \mu_i^{2s+1-\alpha} \mu_j^\alpha \sum_{\rho,\sigma=1}^{2s-1} c_{i\rho} c_{j\sigma} \frac{\epsilon_{2N}^{4\rho+1-2(\rho-\sigma)\alpha}}{1 + \mu_j \mu_i^{-1} \epsilon_{2N}^{-2(\rho-\sigma)}} e^{-Z_\rho(\mu_i) - Z_\sigma(\mu_j)}. \quad (5.1)$$

It is clear that to obtain nontrivial solutions to the Toda equations we should require that at least two coefficients $c_{i\rho}$ for any $i = 1, \dots, r$ are different from zero. In this, we construct solutions depending on only r combinations of independent variables z^- and z^+ . We denote such nonzero constants by C_{J_i} and C_{K_i} . The expression for the matrix elements (5.1) takes then the form

$$(\tilde{R}_\alpha)_{ij} = (-1)^\alpha \mu_i^{2s+1-\alpha} \epsilon_{2N}^{4J_i+1-2\alpha J_i} C_{J_i} e^{-Z_{J_i}(\mu_i)} (\tilde{R}'_\alpha)_{ij} \mu_j^\alpha C_{J_j} \epsilon_{2N}^{2\alpha J_j} e^{-Z_{J_j}(\mu_j)},$$

where

$$\begin{aligned} (\tilde{R}'_\alpha)_{ij} &= \frac{1}{1 + \mu_j \mu_i^{-1} \epsilon_{2N}^{2(J_j-J_i)}} + \frac{C_{K_j}}{C_{J_j}} \frac{\epsilon_{2N}^{2(K_j-J_j)\alpha}}{1 + \mu_j \mu_i^{-1} \epsilon_{2N}^{2(K_j-J_j)}} e^{Z_{J_j}(\mu_j) - Z_{K_j}(\mu_j)} \\ &+ \frac{C_{K_i}}{C_{J_i}} \frac{\epsilon_{2N}^{4(K_i-J_i)-2(K_i-J_i)\alpha}}{1 + \mu_j \mu_i^{-1} \epsilon_{2N}^{2(J_j-K_i)}} e^{Z_{J_i}(\mu_i) - Z_{K_i}(\mu_i)} \\ &+ \frac{C_{K_i} C_{K_j}}{C_{J_i} C_{J_j}} \frac{\epsilon_{2N}^{4(K_i-J_i)-2(K_i-J_i+K_j-J_j)\alpha}}{1 + \mu_j \mu_i^{-1} \epsilon_{2N}^{2(K_j-K_i)}} e^{Z_{J_i}(\mu_i) - Z_{K_i}(\mu_i) + Z_{J_j}(\mu_j) - Z_{K_j}(\mu_j)}. \end{aligned} \quad (5.2)$$

It is easy to show that

$$\Gamma_\alpha = \frac{\det \tilde{R}_{\alpha+1}}{\det \tilde{R}_\alpha} = (-1)^r \frac{\det \tilde{R}'_{\alpha+1}}{\det \tilde{R}'_\alpha}.$$

Recalling also that $\Gamma_1 = (-1)^r$, we see that we can take \tilde{R}'_α instead of \tilde{R}_α to construct solutions of the Toda equations using for that the above determinant representation.

Defining a new set of parameters

$$\begin{aligned} \rho_i &= J_i - K_i, & \theta_{\rho_i} &= \frac{\pi \rho_i}{2s - 1}, & \kappa_{\rho_i} &= -i(\epsilon_{2N}^{\rho_i} - \epsilon_{2N}^{-\rho_i}) = 2 \sin \theta_{\rho_i}, \\ \exp \delta_i &= \frac{C_{K_i}}{C_{J_i}}, & \zeta_i &= i \epsilon_{2N}^{s+J_i+K_i} \mu_i, & f_i &= \epsilon_{2N}^{\rho_i} \zeta_i, & \tilde{f}_i &= \epsilon_{2N}^{-\rho_i} \zeta_i, \end{aligned} \quad (5.3)$$

and introducing the notation

$$D_{ij}(f, g) = \frac{f_i}{f_i + g_j},$$

we can rewrite the expression for \tilde{R}'_α as

$$\begin{aligned} (\tilde{R}'_\alpha)_{ij} &= D_{ij}(f, f) + \epsilon_{2N}^{2\rho_i(\alpha-1)} e^{Z_i(\zeta) + \delta_i - 2i\theta_{\rho_i}} D_{ij}(\tilde{f}, f) + D_{ij}(f, \tilde{f}) e^{Z_j(\zeta) + \delta_j - 2i\theta_{\rho_j}} \epsilon_{2N}^{-2\rho_j(\alpha-1)} \\ &\quad + \epsilon_{2N}^{2\rho_i(\alpha-1)} e^{Z_i(\zeta) + \delta_i - 2i\theta_{\rho_i}} D_{ij}(\tilde{f}, \tilde{f}) e^{Z_j(\zeta) + \delta_j - 2i\theta_{\rho_j}} \epsilon_{2N}^{-2\rho_j(\alpha-1)}, \end{aligned} \quad (5.4)$$

where now the dependence on independent variables is given through

$$Z_i(\zeta) = m \kappa_{\rho_i} (\zeta_i^{-1} z^- + \zeta_i z^+).$$

In fact, it appears that it is appropriate to use the matrices $T_\alpha = D^{-1}(f, f) \tilde{R}'_\alpha$ and write the solutions under construction as

$$\Gamma_\alpha = \frac{\det T_{\alpha+1}}{\det T_\alpha}.$$

The problem of constructing multi-soliton solutions for the Toda equations (3.8) is thus reduced to calculating the determinant of the $r \times r$ matrix T_α .

To obtain a one-soliton solution we set $r = 1$. In this case T_α are ordinary functions, and we easily find

$$T_{\alpha+1} = 1 + 2 \frac{\cos(2\alpha - 1)\theta_\rho}{\cos \theta_\rho} e^{Z(\zeta) + \delta - 2i\theta_\rho} + e^{2(Z(\zeta) + \delta - 2i\theta_\rho)}. \quad (5.5)$$

Setting $r = 2$ we work out the determinant of the respective 2×2 matrix explicitly given above and thus obtain for the two-soliton solution the expression

$$\begin{aligned} \det T_{\alpha+1} &= 1 + 2 \frac{\cos(2\alpha - 1)\theta_{\rho_1}}{\cos \theta_{\rho_1}} e^{\tilde{Z}_1} + 2 \frac{\cos(2\alpha - 1)\theta_{\rho_2}}{\cos \theta_{\rho_2}} e^{\tilde{Z}_2} + e^{2\tilde{Z}_1} + e^{2\tilde{Z}_2} \\ &\quad + \left(2\eta_{12}^+ \frac{\cos(2\alpha - 1)(\theta_{\rho_1} - \theta_{\rho_2})}{\cos \theta_{\rho_1} \cos \theta_{\rho_2}} + 2\eta_{12}^- \frac{\cos(2\alpha - 1)(\theta_{\rho_1} + \theta_{\rho_2})}{\cos \theta_{\rho_1} \cos \theta_{\rho_2}} \right) e^{\tilde{Z}_1 + \tilde{Z}_2} \\ &\quad + 2\eta_{12}^+ \eta_{12}^- \left(\frac{\cos(2\alpha - 1)\theta_{\rho_1}}{\cos \theta_{\rho_1}} e^{\tilde{Z}_1 + 2\tilde{Z}_2} + \frac{\cos(2\alpha - 1)\theta_{\rho_2}}{\cos \theta_{\rho_2}} e^{2\tilde{Z}_1 + \tilde{Z}_2} \right) \\ &\quad + (\eta_{12}^+ \eta_{12}^-)^2 e^{2(\tilde{Z}_1 + \tilde{Z}_2)}, \end{aligned} \quad (5.6)$$

with the ‘soliton interaction factors’

$$\eta_{12}^+ = \frac{(\zeta_1 \zeta_2^{-1} + \zeta_2 \zeta_1^{-1}) + 2 \cos(\theta_{\rho_1} + \theta_{\rho_2})}{(\zeta_1 \zeta_2^{-1} + \zeta_2 \zeta_1^{-1}) + 2 \cos(\theta_{\rho_1} - \theta_{\rho_2})}, \quad \eta_{12}^- = \frac{(\zeta_1 \zeta_2^{-1} + \zeta_2 \zeta_1^{-1}) - 2 \cos(\theta_{\rho_1} - \theta_{\rho_2})}{(\zeta_1 \zeta_2^{-1} + \zeta_2 \zeta_1^{-1}) - 2 \cos(\theta_{\rho_1} + \theta_{\rho_2})},$$

and the constant parameters

$$e^{\delta'_1} = \frac{(f_1 + f_2)(\tilde{f}_1 - f_2)}{(f_1 - f_2)(\tilde{f}_1 + f_2)}, \quad e^{\delta'_2} = \frac{(f_1 + f_2)(f_1 - \tilde{f}_2)}{(f_1 - f_2)(f_1 + \tilde{f}_2)}$$

giving rise to a shift in the exponents as

$$\tilde{Z}_i = Z_i(\zeta) + \delta_i + \delta'_i - 2i\theta_{\rho_i}.$$

It can also be shown that performing the corresponding change of variables as suggested in the paper [3], one can reach the same result along the lines of the Hirota’s approach. Here, the quantities $\det T_{\alpha+1}$ constructed by means of the rational dressing formalism, will coincide with the Hirota’s τ -functions τ_α , see the paper [3] where such correspondence was established for the untwisted case.

Now, considering $s = 2$, so that $N = 3$, we describe the Dodd-Bullough-Mikhailov equation from section 3.2. Here we have $\Gamma_1 = (-1)^r$, $\Gamma_3 = \Gamma_2^{-1}$, and so, the mapping γ is parameterized by the only nontrivial function Γ_2 , denoted here by Γ . The corresponding soliton solutions can easily be derived from the relations (5.5) and (5.6) putting $\alpha = 2$ and $\alpha = 3$ in order and taking into account that $\theta_\rho = \pi\rho/3$. Remember here that $\rho = J - K$, where J and K take values 1, 2 or 3 only. In particular, it is easy to see that the one-soliton solution can be written as

$$\Gamma = \frac{1 - 4e^{Z(\zeta)+\delta-2i\theta_\rho} + e^{2(Z(\zeta)+\delta-2i\theta_\rho)}}{(1 + e^{Z(\zeta)+\delta-2i\theta_\rho})^2}.$$

For the two-soliton solution, we should respectively simplify the expression (5.6). The corresponding expressions reproduce the one- and two-soliton solutions of the Dodd-Bullough-Mikhailov equation obtained in the paper [13] by means of the Hirota’s method.

5.2 Even-dimensional case

Here we consider the case of $n = 2s$, with $p = N = 2s - 1$. It means also that now we have $n_1 = 2$. The eigenvectors of the matrices ${}^t c_-$, ${}^t c_+$, c_- and c_+ are $2s$ -dimensional column vectors Ψ_ρ , $\rho = 1, \dots, 2s - 1$, satisfying the relations

$${}^t c_- \Psi_\rho = m \epsilon_{2N}^{s+2\rho} \Psi_\rho, \quad {}^t c_+ \Psi_\rho = m \epsilon_{2N}^{-s-2\rho} \Psi_\rho, \quad c_- \Psi_\rho = m \epsilon_{2N}^{-s-2\rho} \Psi_\rho, \quad c_+ \Psi_\rho = m \epsilon_{2N}^{s+2\rho} \Psi_\rho,$$

where we define the $2s$ components of Ψ_ρ as

$$(\Psi_\rho)_0 = (\Psi_\rho)_1 = \epsilon_{2N}^{\alpha(s+2\rho)}, \quad (\Psi_\rho)_\alpha = \sqrt{2} \epsilon_{2N}^{\alpha(s+2\rho)}, \quad \alpha = 2, \dots, s$$

and

$$(\Psi_\rho)_\alpha = (-1)^{\alpha-s-1} \sqrt{2} \epsilon_{2N}^{\alpha(s+2\rho)}, \quad \alpha = s + 1, \dots, 2s - 1.$$

Besides, respective to the only zero eigenvalue, c_- , c_+ and their transposed matrices have one and the same null-vector that can be defined as ${}^t\Psi_0 = (1, -1, 0, \dots, 0)$.

Consequently, the solution (4.28) takes the form

$$(u_{i,1})_0 = c_{i0} + \sum_{\rho=1}^{2s-1} c_{i\rho} \epsilon_{2N}^{s+2\rho} e^{-Z_\rho(\mu_i)}, \quad (u_{i,1})_1 = -c_{i0} + \sum_{\rho=1}^{2s-1} c_{i\rho} \epsilon_{2N}^{s+2\rho} e^{-Z_\rho(\mu_i)},$$

and

$$u_{i,\alpha} = \sum_{\rho=1}^{2s-1} c_{i\rho} \sqrt{2} \epsilon_{2N}^{\alpha(s+2\rho)} e^{-Z_\rho(\mu_i)}, \quad \alpha = 2, \dots, s$$

$$u_{i,\alpha} = \sum_{\rho=1}^{2s-1} c_{i\rho} (-1)^{\alpha-s-1} \sqrt{2} \epsilon_{2N}^{\alpha(s+2\rho)} e^{-Z_\rho(\mu_i)}, \quad \alpha = s+1, \dots, 2s-1,$$

where c_{i0} and $c_{i\rho}$ are arbitrary constants and, as usual, we have introduced the notation

$$Z_\rho(\mu_i) = m(\epsilon_{2N}^{-s-2\rho} \mu_i^{-1} z^- + \epsilon_{2N}^{s+2\rho} \mu_i z^+).$$

Note that $u_{i,1}$ is now a 2-dimensional column vector with the components $(u_{i,1})_0$ and $(u_{i,1})_1$ given in order.

For the quasi-periodic quantities \tilde{R}_α introduced in section 4 we obtain the expressions

$$(\tilde{R}_1)_{ij} = -\frac{2\mu_i^{2s} \mu_j^{2s}}{\mu_i^{2s-1} + \mu_j^{2s-1}} c_{i0} c_{j0}$$

$$+ 2\mu_i \mu_j^{2s} \sum_{\rho,\sigma=1}^{2s-1} c_{i\rho} c_{j\sigma} \frac{\epsilon_{2N}^{2(s+\rho+\sigma)}}{1 + \mu_j \mu_i^{-1} \epsilon_{2N}^{-2(\rho-\sigma)}} e^{-Z_\rho(\mu_i) - Z_\sigma(\mu_j)},$$

$$(\tilde{R}_\alpha)_{ij} = \frac{2\mu_i^{2s} \mu_j^{2s}}{\mu_i^{2s-1} + \mu_j^{2s-1}} c_{i0} c_{j0}$$

$$+ 2(-1)^\alpha \mu_i^{2s+1-\alpha} \mu_j^\alpha \sum_{\rho,\sigma=1}^{2s-1} c_{i\rho} c_{j\sigma} \frac{\epsilon_{2N}^{4\rho+1-2(\rho-\sigma)\alpha}}{1 + \mu_j \mu_i^{-1} \epsilon_{2N}^{-2(\rho-\sigma)}} e^{-Z_\rho(\mu_i) - Z_\sigma(\mu_j)},$$

that are to be used in the determinant representation derived earlier for constructing the soliton solutions.

It is easy to see that if we set here $c_{i0} = 0$, then we come to the same solutions given for $n = 2s - 1$ by the relations (5.4)–(5.6), with the only unessential difference that for $n = 2s$ the rational dressing gives $\Gamma_1 = (-1)^r J_2^r$. Therefore, to obtain new solutions we consider that in what follows c_{i0} does not vanish.

To construct such new simplest soliton solutions we thus assume that for each value of the index i only one arbitrary constant $c_{i\rho}$, apart from the c_{i0} , is different from zero. To keep up with the notations used in the preceding section, we denote such nonvanishing coefficients by C_{I_i} and C_{0_i} . Then we can write for the above $r \times r$ matrices \tilde{R}_1 and \tilde{R}_α

$$(\tilde{R}_1)_{ij} = -2\mu_i C_{0_i} (\tilde{R}'_1)_{ij} C_{0_j} \mu_j^{2s}, \quad (\tilde{R}_\alpha)_{ij} = 2\mu_i C_{0_i} (\tilde{R}'_\alpha)_{ij} C_{0_j} \mu_j^{2s},$$

where \tilde{R}'_1 and \tilde{R}'_α can be represented as

$$\begin{aligned} (\tilde{R}'_1)_{ij} &= D_{ij}(\zeta^{2s-1}, \zeta^{2s-1}) - e^{-Z'_i} D_{ij}(\zeta, \zeta) e^{-Z'_j}, \\ (\tilde{R}'_\alpha)_{ij} &= D_{ij}(\zeta^{2s-1}, \zeta^{2s-1}) - (-1)^\alpha \zeta_i^{2s-\alpha} e^{-Z'_i} D_{ij}(\zeta, \zeta) e^{-Z'_j} \zeta_j^{\alpha-2s}. \end{aligned} \quad (5.7)$$

Here we use the same notation for the matrices $D(f, g)$ introduced in the preceding section 5.1, and besides,

$$Z'_i = Z_i(\zeta) - \delta_i - i\theta_{s+2I_i}, \quad Z_i(\zeta) = m(\zeta_i^{-1} z^- + \zeta_i z^+),$$

with the set of parameters

$$\zeta_i = \epsilon_{2N}^{s+2I_i} \mu_i, \quad e^{\delta_i} = \frac{C_{I_i}}{C_{0_i}}, \quad \theta_{s+2I_i} = \frac{\pi(s+2I_i)}{2s-1}.$$

We also rewrite the explicit forms of the components of the 2-dimensional column vector $\tilde{u}_{i,1}$ in terms of the notations introduced above. We have

$$(\tilde{u}_{i,1})_0 = \mu_i C_{0_i} (1 + \exp(-Z'_i)), \quad (\tilde{u}_{i,1})_1 = -\mu_i C_{0_i} (1 - \exp(-Z'_i)).$$

Hence, according to the general relations derived in section 4, we can take the matrices $T_\alpha = D^{-1}(\zeta^{2s-1}, \zeta^{2s-1}) \tilde{R}'_\alpha$ instead of \tilde{R}_α and write for the solutions of the Toda equations (3.13) the following expressions:

$$\Gamma_1 = I_2 + \sum_{i,j=1}^r v_i (\tilde{R}'_1)^{-1}_{ij} {}^t v_j J_2,$$

where v_i are 2-dimensional column vectors with the components

$$v_{i,0} = \frac{1}{\sqrt{2}}(1 + \exp(-Z'_i)), \quad v_{i,1} = -\frac{1}{\sqrt{2}}(1 - \exp(-Z'_i)),$$

and

$$\Gamma_\alpha = \frac{\det T_{\alpha+1}}{\det T_\alpha}, \quad \alpha = 2, \dots, s.$$

To obtain a one-soliton solution of the type under consideration, we put $r = 1$, for which T_α are ordinary functions. It is easy to show that in this case we have

$$\Gamma_1 = \begin{pmatrix} 0 & \Gamma \\ \Gamma^{-1} & 0 \end{pmatrix}, \quad \Gamma = \frac{1 + \exp(-Z')}{1 - \exp(-Z')},$$

and

$$\Gamma_\alpha = \frac{1 + (-1)^\alpha \exp(-2Z')}{1 - (-1)^\alpha \exp(-2Z')}, \quad \alpha = 2, \dots, s.$$

Note that apart from the relation $\Gamma_{2s+1-\alpha} = \Gamma_\alpha^{-1}$ here we also have $\Gamma_{\alpha+1} = \Gamma_\alpha^{-1}$. It is clear that to have a mapping γ belonging to G_0 we should take $\Gamma_1 J_2$ instead of the above Γ_1 .

Setting $r = 2$ we work out the corresponding 2×2 matrices and thus obtain new two-soliton solutions to (3.13). The calculations lead to the expressions

$$\Gamma_1 = \begin{pmatrix} \Gamma & 0 \\ 0 & \Gamma^{-1} \end{pmatrix}, \quad \Gamma = \frac{1 + e^{-\tilde{Z}_1} - e^{-\tilde{Z}_2} - \eta_{12} e^{-(\tilde{Z}_1 + \tilde{Z}_2)}}{1 - e^{-\tilde{Z}_1} + e^{-\tilde{Z}_2} - \eta_{12} e^{-(\tilde{Z}_1 + \tilde{Z}_2)}},$$

where the ‘soliton interaction factor’ is now

$$\eta_{12} = \frac{\zeta_1 - \zeta_2}{\zeta_1 + \zeta_2} \cdot \frac{\zeta_1^{2s-1} - \zeta_2^{2s-1}}{\zeta_1^{2s-1} + \zeta_2^{2s-1}},$$

and we have introduced a new parameter δ' defined by

$$e^{\delta'} = \frac{\zeta_1^{2s-1} + \zeta_2^{2s-1}}{\zeta_1^{2s-1} - \zeta_2^{2s-1}}$$

and producing a shift in the exponents,

$$\tilde{Z}_i = Z'_i - \delta' = Z_i(\zeta) - \delta_i - \delta' - i\theta_{s+2I_i}.$$

We also have

$$\begin{aligned} \det T_{\alpha+1} &= 1 + (-1)^\alpha (e^{-2\tilde{Z}_1} + e^{-2\tilde{Z}_2}) \\ &- 4(-1)^\alpha \frac{\zeta_1^\alpha \zeta_2^{2s-\alpha} + \zeta_2^\alpha \zeta_1^{2s-\alpha}}{(\zeta_1 + \zeta_2)(\zeta_1^{2s-1} + \zeta_2^{2s-1})} e^{-(\tilde{Z}_1 + \tilde{Z}_2)} + \eta_{12}^2 e^{-2(\tilde{Z}_1 + \tilde{Z}_2)}. \end{aligned} \tag{5.8}$$

Note finally that under the permutation of the parameters ζ_1 and ζ_2 the function Γ transforms into Γ^{-1} , thus Γ_1 goes to Γ_1^{-1} , while Γ_α for the other values of α all stay invariant.

6. Conclusion

We have considered the abelian Toda systems associated with the loop groups of the complex general linear groups. Using the method of rational dressing, along the lines of [3], we have constructed soliton solutions to these equations in the twisted cases, that is, when the gradations are generated by outer automorphisms of the structure Lie algebras. In fact, we have constructed solutions which obviously form wider classes than the solitons. The soliton solutions can be obtained from the more general solutions constructed by means of the rational dressing by setting the initial data of the problems under consideration to specific values. And in this sense the soliton solutions that can be found also by the Hirota’s approach, form a subset of the solutions constructed by the rational dressing method. We have demonstrated this, in particular, by the solutions to the famous Dodd-Bullough-Mikhailov equation.

Our consideration can be generalized to Toda systems connected with other loop groups, such as twisted and untwisted loop groups of the complex orthogonal and symplectic groups. It is worth noting that, as we have already observed here, the pole positions of the dressing meromorphic mappings and their inverse ones turn out to be bound up with each other because of the specific structure of the outer automorphism leading to the twisted cases. This circumstance made part of the formulae more intricate than in the untwisted general linear case considered in the preceding paper [3]. Actually, similar problems of coinciding pole positions arise also due to the specific group conditions. We will address to this problem and present our respective results in some future publications.

Note finally that Toda systems based on twisted affine Kac-Moody algebras were approached in [24] by means of vertex operators, by constructing twisted algebras from untwisted ones with the help of relevant folding techniques. Although the constructions of [24] do not present explicit forms of the solutions, however, it is not difficult to see that they reproduce the soliton solutions that can be obtained by the Hirota's approach. It should be an interesting problem to make a comparative analysis, from a group-algebraic point of view, of the rational dressing and vertex operators methods.

Acknowledgments

This work was supported in part by the Russian Foundation for Basic Research under grant #07-01-00234 and by the joint DFG-RFBR grant #08-01-91953. One of the authors (A.V.R.) wishes to acknowledge the warm hospitality of the Erwin Schrödinger International Institute for Mathematical Physics where a part of this work was carried out.

References

- [1] A.N. Leznov and M.V. Saveliev, *Group-theoretical methods for the integration of nonlinear dynamical systems*, Birkhauser, Basel Switzerland (1992).
- [2] A.V. Razumov and M.V. Saveliev, *Lie algebras, geometry and Toda-type systems*, Cambridge University Press, Cambridge U.K. (1997).
- [3] Kh.S. Nirov and A.V. Razumov, *Abelian Toda solitons revisited*, *Rev. Math. Phys.* **20** (2008) 1209 [[arXiv:0802.0593](#)].
- [4] R. Hirota, *The direct method in soliton theory*, Cambridge University Press, Cambridge U.K. (2004).
- [5] T.J. Hollowood, *Solitons in affine Toda field theories*, *Nucl. Phys.* **B 384** (1992) 523.
- [6] C.P. Constantinidis, L.A. Ferreira, J.F. Gomes and A.H. Zimerman, *Connection between affine and conformal affine Toda models and their Hirota's solution*, *Phys. Lett.* **B 298** (1993) 88 [[hep-th/9207061](#)].
- [7] N.J. MacKay and W.A. McGhee, *Affine Toda solitons and automorphisms of Dynkin diagrams*, *Int. J. Mod. Phys.* **A 8** (1993) 2791 [*Erratum ibid.* **8** (1993) 3830] [[hep-th/9208057](#)].
- [8] H. Aratyn, C.P. Constantinidis, L.A. Ferreira, J.F. Gomes and A.H. Zimerman, *Hirota's solitons in the affine and the conformal affine Toda models*, *Nucl. Phys.* **B 406** (1993) 727 [[hep-th/9212086](#)].
- [9] Z. Zhu and D.G. Caldi, *Multi-soliton solutions of affine Toda models*, *Nucl. Phys.* **B 436** (1995) 659 [[hep-th/9307175](#)].
- [10] V.E. Zakharov and A.B. Shabat, *Integration of nonlinear equations of mathematical physics by the method of inverse scattering. II*, *Func. Anal. Appl.* **13** (1979) 166.
- [11] A.V. Mikhailov, *The reduction problem and the inverse scattering method*, *Physica* **3D** (1981) 73.

- [12] A.G. Bueno, L.A. Ferreira and A.V. Razumov, *Confinement and soliton solutions in the SL(3) Toda model coupled to matter fields*, *Nucl. Phys. B* **626** (2002) 463 [[hep-th/0105078](#)].
- [13] P.E.G. Assis and L.A. Ferreira, *The Bullough-Dodd model coupled to matter fields*, *Nucl. Phys. B* **800** (2008) 409 [[arXiv:0708.1342](#)].
- [14] Kh.S. Nirov and A.V. Razumov, *Toda equations associated with loop groups of complex classical Lie groups*, *Nucl. Phys. B* **782** (2007) 241 [[math-ph/0612054](#)].
- [15] Kh.S. Nirov and A.V. Razumov, *On Z-graded loop Lie algebras, loop groups and Toda equations*, *Theor. Math. Phys.* **154** (2008) 385 [*Teor. Mat. Fiz.* **154** (2008) 451] [[arXiv:0705.2681](#)].
- [16] A.V. Razumov and M.V. Savelev, *Differential geometry of Toda systems*, *Commun. Anal. Geom.* **2** (1994) 461 [[hep-th/9311167](#)].
- [17] A.V. Razumov and M.V. Savelev, *Multidimensional Toda type systems*, *Teor. Mat. Fiz.* **112** (1997) 254 [*Theor. Math. Phys.* **112** (1997) 999] [[hep-th/9609031](#)].
- [18] Kh.S. Nirov and A.V. Razumov, *On Z-gradations of twisted loop Lie algebras of complex simple Lie algebras*, *Commun. Math. Phys.* **267** (2006) 587 [[math-ph/0504038](#)].
- [19] A. Kriegl and P. Michor, *Aspects of the theory of infinite dimensional manifolds*, *Diff. Geom. Appl.* **1** (1991) 159 [[math.DG/9202206](#)].
- [20] A. Kriegl and P. Michor, *The convenient setting of global analysis*, *Mathematical Surveys and Monographs*, vol. 53, American Mathematical Society, Providence U.S.A. (1997).
- [21] A.V. Mikhailov, M.A. Olshanetsky and A.M. Perelomov, *Two-dimensional generalized Toda lattice*, *Commun. Math. Phys.* **79** (1981) 473.
- [22] R.K. Dodd and R.K. Bullough, *Polynomial conserved densities for the sine-Gordon equations*, *Proc. Roy. Soc. Lond. A* **352** (1977) 481.
- [23] G. Tzitzéica, *Sur une nouvelle classe de surfaces*, *Rend. Circ. Mat. Palermo* **25** (1908) 180.
- [24] M.A.C. Kneipp and D.I. Olive, *Solitons and vertex operators in twisted affine Toda field theories*, *Commun. Math. Phys.* **177** (1996) 561 [[hep-th/9404030](#)].