

## Manifest spin 2 duality with electric and magnetic sources

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

JHEP01(2009)030

(<http://iopscience.iop.org/1126-6708/2009/01/030>)

[The Table of Contents](#) and [more related content](#) is available

Download details:

IP Address: 80.92.225.132

The article was downloaded on 03/04/2010 at 11:36

Please note that [terms and conditions apply](#).

## Manifest spin 2 duality with electric and magnetic sources

---

Glenn Barnich\* and Cédric Troessaert†

*Physique Théorique et Mathématique, Université Libre de Bruxelles and  
International Solvay Institutes,  
Campus Plaine C.P. 231, B-1050 Bruxelles, Belgium  
E-mail: gbarnich@ulb.ac.be, Cedric.Troessaert@ulb.ac.be*

**ABSTRACT:** We extend the formulation of spin 2 fields on Minkowski space which makes the action manifestly invariant under duality rotations to the case of interactions with external electric and magnetic sources by adding suitable potentials for the longitudinal and trace parts. In this framework, the string singularity of the linearized Taub-NUT solution is resolved into a Coulomb-like solution. Suitable surface charges to measure energy-momentum and angular momentum of both electric and magnetic type are constructed.

**KEYWORDS:** Duality in Gauge Field Theories, Solitons Monopoles and Instantons, Gauge Symmetry, Classical Theories of Gravity.

---

\*Research Director of the Fund for Scientific Research-FNRS (Belgium). Also at Laboratoire de Mathématiques et Physique Théorique, Unité Mixte de Recherche du CNRS, Fédération Denis Poisson, Université François Rabelais, Parc de Grandmont, 37200 Tours, France.

†Research Fellow of the Fund for Scientific Research-FNRS (Belgium).

## Contents

<b>1. Introduction</b>	<b>2</b>
<b>2. Preliminaries</b>	<b>3</b>
2.1 Canonical formulation of Pauli-Fierz theory	3
2.2 Decomposition of symmetric rank two tensors	3
2.3 Reduced phase space for linearized gravity	5
<b>3. Action and symmetries</b>	<b>6</b>
3.1 Degree of freedom count	6
3.2 Change of variables and duality rotations	6
3.3 Action principle and locality	7
3.4 Canonical structure	7
3.5 Gauge structure	8
3.6 Duality generator	9
3.7 Hamiltonian	9
3.8 Poincaré generators	10
<b>4. Coupling to conserved electric and magnetic sources</b>	<b>11</b>
4.1 Interacting variational principle	11
4.2 Linearized Taub-NUT solution	11
<b>5. Surface charges</b>	<b>12</b>
5.1 Regge-Teitelboim revisited	12
5.2 Linear theories	13
5.3 Electric and magnetic energy-momentum and angular momentum surface charges	14
5.4 Poincaré transformations of surface charges	17
<b>A. Poincaré generators for Pauli-Fierz theory</b>	<b>18</b>
<b>B. Riemann tensor and canonical variables</b>	<b>21</b>
B.1 Covariant equations in the presence of magnetic sources	21
B.2 Canonical expressions	24
B.3 Riemann tensor for linearized Taub-NUT	26

## 1. Introduction

Duality rotations for massless spin 2 fields have recently been shown to be symmetries of the action in the context of a Hamiltonian formulation involving suitable potentials that arise when solving the constraints [1]. This symmetry can be extended to higher spin fields [2] and also to the case of spin 2 fields propagating on an (A)dS background [3, 4], but does not survive gravitational self-interactions [5].

The spin 2 result generalizes the so-called double potential formalism for spin 1 fields [6, 7], which has been extended so as to include couplings to dynamical dyons by using Dirac strings [8, 9]. In the original double potential formalism, Gauss's constraint is solved in terms of new transverse vector potentials for the electric field so that electromagnetism is effectively formulated on a reduced phase space with all gauge invariance eliminated. Alternatively, one may choose [10] to double the gauge redundancy of standard electromagnetism by using a description with independent vector and longitudinal potentials for the magnetic and electric fields and 2 scalar potentials that appear as Lagrange multipliers for the electric and magnetic Gauss constraints. In this framework, the string-singularity of the solution describing a static dyon is resolved into a Coulomb-like solution. Furthermore, magnetic charge no longer appears as a topological conservation law but as a surface charge on a par with electric charge.

The aim of the present work is to apply the same strategy to the spin 2 case. Doubling the gauge invariance by keeping all degrees of freedom of symmetric tensors now leads to a second copy of linearized lapse and shifts as Lagrange multipliers for the new magnetic constraints. As a consequence, the string singularity of the gravitational dyon, the linearized Taub-NUT solution is resolved and becomes Coulomb-like exactly as the purely electric linearized Schwarzschild solution. Furthermore, as required by manifest duality, magnetic mass, momentum and Lorentz charges also appear as surface integrals.

Our work thus presents a manifestly duality invariant alternative to [11] where the coupling of spin 2 fields to conserved electric and magnetic sources has been achieved in a manifestly Poincaré invariant way through the introduction of Dirac strings.

Recent and not so recent related work includes for instance [12–24] and references therein.

In section 2, we briefly recall the two ingredients needed for our formulation: the Hamiltonian description of spin 2 fields propagating on Minkowski spacetime and the decomposition of symmetric tensors into their irreducible components, giving rise to the reduced phase space description of linearized gravity.

Our analysis starts in section 3 with a degree of freedom count that shows that the phase space of duality invariant spin 2 fields with doubled gauge invariance can be taken to consist of 2 symmetric tensors, 2 vectors and 2 scalars in 3 dimensions. We then define the metric, extrinsic curvature and their duals in terms of the phase space variables and propose our duality invariant action principle with enhanced gauge invariance. We proceed by identifying the canonically conjugate pairs and discuss the gauge structure, Hamiltonian and duality generators of the theory. In the absence of sources, we then show how the

generators for global Poincaré transformations of Pauli-Fierz theory, reviewed in detail in appendix A, can be extended to the duality invariant theory.

The coupling to external electric and magnetic sources is discussed in section 4. The equations of motion are first solved in the simplest case of a point-particle dyon sitting at the origin. They are Coulomb-like without string singularities. By identifying the Riemann tensor in terms of the canonical variables and computing it for this case, we show in appendix B that this solution indeed describes the linearized Taub-NUT solution.

In section 5 we discuss the surface charges of the theory and show that they include electric and magnetic energy-momentum and angular momentum. Because of the non-locality of the Poisson structure, we proceed indirectly and show that the expressions obtained by generalizing the surface charges of Pauli-Fierz theory in a duality invariant way fulfill the standard properties. Finally, we investigate how the surface charges transform under a global Poincaré transformation of the sources.

## 2. Preliminaries

### 2.1 Canonical formulation of Pauli-Fierz theory

The Hamiltonian formulation of general relativity linearized around flat spacetime is

$$S_{\text{PF}}[h_{mn}, \pi^{mn}, n_m, n] = \int dt \left[ \int d^3x (\pi^{mn} \dot{h}_{mn} - n^m \mathcal{H}_m - n \mathcal{H}) - H_{\text{PF}} \right], \quad (2.1)$$

with

$$H_{\text{PF}}[h_{mn}, \pi^{mn}] = \int d^3x \left( \pi^{mn} \pi_{mn} - \frac{1}{2} \pi^2 + \frac{1}{4} \partial^r h^{mn} \partial_r h_{mn} - \frac{1}{2} \partial_m h^{mn} \partial^r h_{rn} + \frac{1}{2} \partial^m h \partial^n h_{mn} - \frac{1}{4} \partial^m h \partial_m h \right), \quad (2.2)$$

and

$$\mathcal{H}_m = -2\partial^n \pi_{mn}, \quad \mathcal{H}_\perp = \Delta h - \partial^m \partial^n h_{mn}. \quad (2.3)$$

Here, indices are lowered and raised with the flat space metric  $\delta_{mn}$  and its inverse,  $h = h^m_m$ ,  $\pi = \pi^m_m$  and  $\Delta = \partial_m \partial^m$  is the Laplacian in flat space. The linearized 4 metric is reconstructed using  $h_{00} = -2n$  and  $h_{0i} = n_i$ .

### 2.2 Decomposition of symmetric rank two tensors

Symmetric rank two tensors  $\phi_{mn}$  decompose as [25, 26]

$$\phi_{mn} = \phi_{mn}^{TT} + \phi_{mn}^T + \phi_{mn}^L, \quad (2.4)$$

$$\phi_{mn}^L = \partial_m \psi_n + \partial_n \psi_m, \quad (2.5)$$

$$\phi_{mn}^T = \frac{1}{2} (\delta_{mn} \Delta - \partial_m \partial_n) \psi^T. \quad (2.6)$$

Here  $\phi_{mn}^{TT}$  is the transverse-traceless part, containing two independent components. The tensor  $\phi_{mn}^T$  contains the trace of the transverse part of  $\phi_{mn}$  and only one independent

component. The last three components are the longitudinal part contained in  $\phi_{mn}^L$ . In terms of the original tensor  $\phi_{mn}$  the potentials for the longitudinal part and the trace are given by

$$\psi_m = \Delta^{-1} \left( \partial^n \phi_{mn} - \frac{1}{2} \Delta^{-1} \partial_m \partial^k \partial^l \phi_{kl} \right), \quad (2.7)$$

$$\psi^T = \Delta^{-1} (\phi - \Delta^{-1} \partial^m \partial^n \phi_{mn}), \quad (2.8)$$

while the transverse-traceless part is then defined as the remainder,

$$\phi_{mn}^{TT} = \phi_{mn} - \phi_{mn}^T - \phi_{mn}^L. \quad (2.9)$$

This implies

$$\begin{aligned} \Delta^2 \phi_{mn}^{TT} = & \Delta^2 \phi_{mn} - \Delta \partial_m \partial^k \phi_{kn} - \Delta \partial_n \partial^k \phi_{km} \\ & - \frac{1}{2} \Delta (\delta_{mn} \Delta - \partial_m \partial_n) \phi + \frac{1}{2} (\delta_{mn} \Delta + \partial_m \partial_n) \partial^k \partial^l \phi_{kl}, \end{aligned} \quad (2.10)$$

$$\begin{aligned} \int d^3x \phi^{mn} \Delta^2 \phi_{mn}^{TT} = & \int d^3x \left( \Delta \phi^{mn} \Delta \phi_{mn} + 2 \partial_m \phi^{mn} \Delta \partial^k \phi_{kn} - \frac{1}{2} (\Delta \phi)^2 \right. \\ & \left. + \partial_m \partial_n \phi^{mn} \Delta \phi + \frac{1}{2} \partial_m \partial_n \phi^{mn} \partial_k \partial_l \phi^{kl} \right). \end{aligned} \quad (2.11)$$

Alternatively, one can introduce the local operator  $\mathcal{P}^{TT}$

$$(\mathcal{P}^{TT} \phi)_{mn} = \frac{1}{2} [\epsilon_{mpq} \partial^p (\Delta \phi_n^q - \partial_n \partial_r \phi^{qr}) + \epsilon_{npq} \partial^p (\Delta \phi_m^q - \partial_m \partial_r \phi^{qr})], \quad (2.12)$$

which projects out the longitudinal and trace parts and onto a transverse-traceless tensor,

$$(\mathcal{P}^{TT} \phi)_{mn} = \mathcal{P}^{TT} (\phi^{TT})_{mn} = (\mathcal{P}^{TT} \phi)_{mn}^{TT}. \quad (2.13)$$

In addition,

$$(\mathcal{P}^{TT} (\mathcal{P}^{TT} \phi))_{mn} = -\Delta^3 \phi_{mn}^{TT}. \quad (2.14)$$

As a consequence, the transverse-traceless tensor  $\phi_{mn}^{TT}$  can be written as  $\mathcal{P}^{TT}$  acting on a suitable potential  $\psi_{mn}^{TT}$ ,

$$\phi_{mn}^{TT} = (\mathcal{P}^{TT} \psi^{TT})_{mn}, \quad \psi_{mn}^{TT} = -\Delta^{-3} (\mathcal{P}^{TT} \phi)_{mn}. \quad (2.15)$$

The operator  $\mathcal{P}^{TT}$  is related to the way the Hamiltonian constraint  $\mathcal{H} = 0$  is solved by expressing the metric  $h_{mn}$  in terms of superpotentials in [1]. When acting on a transverse-traceless tensor, the two terms of (2.12) involving  $\partial_r \phi^{qr}$  can be dropped. In this case,  $\mathcal{P}^{TT}$  is related to the generalized curl [2, 5],

$$(\mathcal{O} \phi)_{mn} = \frac{1}{2} (\epsilon_{mpq} \partial^p \phi_n^q + \epsilon_{npq} \partial^p \phi_m^q), \quad (2.16)$$

$$(\mathcal{P}^{TT} \phi^{TT})_{mn} = \Delta (\mathcal{O} \phi^{TT})_{mn}. \quad (2.17)$$

A second operator that projects out the longitudinal and trace parts and onto a transverse-traceless tensor is  $\mathcal{Q}^{TT}$ ,

$$(\mathcal{Q}^{TT}\phi)_{mn} = \epsilon_{mpq}\epsilon_{nrs}\partial^p\partial^r\Delta\phi^{qs} - \frac{1}{2}(\delta_{mn}\Delta - \partial_m\partial_n)(\Delta\phi - \partial^p\partial^r\phi_{pr}). \quad (2.18)$$

In this case,

$$(\mathcal{Q}^{TT}(\mathcal{Q}^{TT}\phi))_{mn} = \Delta^4\phi_{mn}^{TT}, \quad (2.19)$$

so that the transverse-traceless tensor  $\phi_{mn}^{TT}$  can be written as  $\mathcal{Q}^{TT}$  acting on another potential  $\chi_{mn}^{TT}$ ,

$$\phi_{mn}^{TT} = (\mathcal{Q}^{TT}\chi^{TT})_{mn}, \quad \chi_{mn}^{TT} = \Delta^{-4}(\mathcal{Q}^{TT}\phi)_{mn}. \quad (2.20)$$

In turn this operator is related to the way the constraints  $\mathcal{H}_m = 0$  are solved by expressing the momenta  $\pi^{mn}$  in terms of superpotentials in [1]. When acting on a transverse-traceless tensor, the last term can again be dropped and it is related to the square of the generalized curl,

$$(\mathcal{Q}^{TT}\phi^{TT})_{mn} = \Delta(\mathcal{O}(\mathcal{O}\phi^{TT}))_{mn} = -\Delta^2\phi_{mn}^{TT}. \quad (2.21)$$

The elements of the decomposition are orthogonal under integration if boundary terms can be neglected,

$$\int d^3x \phi^{mn}\varphi_{mn} = \int d^3x (\phi^{TTmn}\varphi_{mn}^{TT} + \phi^{Lmn}\varphi_{mn}^L + \phi^{Tmn}\varphi_{mn}^T). \quad (2.22)$$

and the operators  $\mathcal{P}^{TT}$ ,  $\mathcal{Q}^{TT}$ ,  $\mathcal{O}$  are self-ajoint, e.g.,

$$\int d^3x (\mathcal{P}^{TT}\phi)^{mn}\varphi_{mn} = \int d^3x \phi^{mn}(\mathcal{P}^{TT}\varphi)_{mn}. \quad (2.23)$$

### 2.3 Reduced phase space for linearized gravity

For completeness, let us briefly recall [25] the reduced phase space associated with the Pauli-Fierz action. Because of the orthogonality of the decomposition, the canonically conjugate pairs can be directly read off from the kinetic term and are given by

$$(h_{mn}^{TT}(x), \pi_{TT}^{kl}(\vec{y})), \quad (h_{mn}^L(\vec{x}), \pi_L^{kl}(y)), \quad (h_{mn}^T(x), \pi_T^{kl}(y)). \quad (2.24)$$

The first class constraints  $\mathcal{H}_m = 0 = \mathcal{H}$  are equivalent to  $\pi_L^{kl} = 0 = h_{mn}^T$ . They can be gauge fixed through the conditions  $h_{mn}^L = 0 = \pi_T^{kl}$ . The reduced theory only depends on 2 degrees of freedom (per spacetime point), the transverse-traceless components  $(h_{mn}^{TT}(\vec{x}), \pi_{TT}^{kl}(y))$  and the reduced Hamiltonian simplifies to

$$H^R = \int d^3x \left( \pi_{TT}^{mn}\pi_{mn}^{TT} + \frac{1}{4}\partial_r h_{mn}^{TT}\partial^r h_{TT}^{mn} \right). \quad (2.25)$$

### 3. Action and symmetries

#### 3.1 Degree of freedom count

In order to be able to couple to sources of both electric and magnetic type in a duality invariant way, we want to keep all components and double the gauge invariance of the theory. With 2 degrees of freedom,  $\# \text{ dof} = 2$ , and 8 first class constraints,  $\# \text{ fcc} = 8$ , we thus need 10 canonical pairs,  $\# \text{ cp} = 10$ , according to the degree of freedom count [27]

$$2 * (\# \text{ cp}) = 2 * (\# \text{ dof}) + 2 * (\# \text{ fcc}). \quad (3.1)$$

This can be done by taking 2 symmetric tensors, 2 vectors and 2 scalars as fundamental canonical variables,

$$z^A = (H_{mn}^a, A_m^a, C^a). \quad (3.2)$$

#### 3.2 Change of variables and duality rotations

For  $a = 1, 2$ , consider  $h_{mn}^a = (h_{mn}, h_{mn}^D)$  and  $\pi_a^{mn} = (\pi_D^{mn}, \pi^{mn})$  and the definitions

$$\begin{aligned} h_{mn}^a &= \epsilon_{mpq} \partial^p H^{aq}{}_n + \epsilon_{npq} \partial^p H^{aq}{}_m + \partial_m A_n^a + \partial_n A_m^a + \frac{1}{2} (\delta_{mn} \Delta - \partial_m \partial_n) C^a \\ &= 2\Delta^{-1} (\mathcal{P}^{TT} H^a)_{mn} + \partial_m (\Delta^{-1} \epsilon_{npq} \partial^p \partial_r H^{aqr} + A_n^a) \\ &\quad + \partial_n (\Delta^{-1} \epsilon_{mpq} \partial^p \partial_r H^{aqr} + A_m^a) + \frac{1}{2} (\delta_{mn} \Delta - \partial_m \partial_n) C^a, \end{aligned} \quad (3.3)$$

$$\begin{aligned} \pi_{mn}^a &= \epsilon_{mpq} \epsilon_{nrs} \partial^p \partial^r H^{aqs} - \partial_m \partial^r H_{rn}^a - \partial_n \partial^r H_{rm}^a - (\delta_{mn} \Delta - \partial_m \partial_n) H^a + \delta_{mn} \partial^k \partial^l H_{kl}^a \\ &= \Delta^{-1} (\mathcal{Q}^{TT} H^a)_{mn} - \partial_m \partial^r H_{rn}^a - \partial_n \partial^r H_{rm}^a - \frac{1}{2} (\delta_{mn} \Delta - \partial_m \partial_n) H^a \\ &\quad + \frac{1}{2} \Delta^{-1} (\delta_{mn} \Delta + \partial_m \partial_n) \partial^p \partial^r H_{pr}^a \\ &= -\Delta H_{mn}^a. \end{aligned} \quad (3.4)$$

The relations for  $h_{mn}[H^1, A^1, C^1]$  and  $\pi^{mn}[H^2]$  are the local change of coordinates from the standard canonical variables of linearized gravity to the new variables. They are invertible and, as usual, the inverse is not local. The relations for  $h_{mn}^2 = h_{mn}^D$ ,  $\pi_1^{mn} = \pi_D^{mn}$  serve to denote convenient combinations of the new variables in terms of which expressions below will simplify. As indicated by the notation, the infinitesimal duality rotations among the fundamental variables are

$$\delta_D H_{mn}^a = \epsilon^{ab} H_{bmn}, \quad \delta_D A_m^a = \epsilon^{ab} A_{bm}, \quad \delta_D C^a = \epsilon^{ab} C_b. \quad (3.5)$$

Here,  $\epsilon_{ab}$  is skew-symmetric with  $\epsilon_{12} = 1$  and indices are lowered and raised with  $\delta_{ab}$  and its inverse. Duality invariance will be manifest if all the internal indices  $a$  are contracted with the invariant tensors  $\delta_{ab}, \epsilon_{ab}$ . Since  $h_{mn}^a, \pi_a^{mn}$  are linear combinations of the fundamental variables, they are rotated in exactly the same way. We can thus consider  $h_{mn}^2 = h_{mn}^D$ ,  $\pi_1^{mn} = \pi_D^{mn}$  as the dual spatial metric and the dual extrinsic curvature in the linearized theory.



### 3.3 Action principle and locality

The duality invariant local action principle that we propose is of the form

$$S_G[z^A, u^\alpha] = \int d^4x (a_A[z] \dot{z}^A - u^\alpha \gamma_\alpha[z]) - \int dt H[z], \quad (3.6)$$

where  $u^\alpha$  denote the 8 Lagrange multipliers and  $\gamma_\alpha$  the constraints.

Let us stress here that we use the assumption that the flat space Laplacian  $\Delta$  is invertible in order to show equivalence with the usual Hamiltonian or covariant formulation of Pauli-Fierz theory and also to disentangle the canonical structure. The action principle (3.6) itself and the associated equations of motion will be local both in space and in time independently of this assumption. The theory itself is not local as a Hamiltonian gauge theory (see e.g. [28], section 12) because the Poisson brackets among the canonical variables will not be local.

### 3.4 Canonical structure

The explicit expression for the kinetic term is

$$a_A \dot{z}^A = \epsilon_{ab} H^{amn} \left( (\mathcal{P}^{TT} \dot{H}^b)_{mn} + \partial_m \Delta \dot{A}_n^b + \partial_n \Delta \dot{A}_m^b + \frac{1}{2} (\delta^{mn} \Delta - \partial^m \partial^n) \Delta \dot{C}^b \right). \quad (3.7)$$

The canonically conjugate pairs are identified by writing the integrated kinetic term as

$$\int d^4x a_A \dot{z}^A = \int d^4x \left( -2\Delta (\mathcal{O}H_{TT}^2)^{mn} \dot{H}_{mn}^{1TT} + 2\Delta \partial_m H_L^{2mn} \dot{A}_n^1 - 2\Delta \partial_m \dot{H}_L^{1mn} A_n^2 - \frac{1}{2} \Delta (\Delta H_T^2 - \partial_p \partial_q H_T^{2pq}) \dot{C}^1 + \frac{1}{2} \Delta (\Delta H_T^1 - \partial_p \partial_q H_T^{1pq}) \dot{C}^2 \right). \quad (3.8)$$

This means that the usual canonical pairs of linearized gravity can be chosen in terms of the new variables as

$$\left( H_{mn}^{1TT}(x), -2\Delta (\mathcal{O}H_{TT}^2)^{kl}(y) \right), \left( C^1(x), -\frac{1}{2} \Delta (\Delta H_T^2 - \partial_p \partial_q H_T^{2pq})(y) \right), \\ \left( A_m^1(x), 2\Delta \partial_r H_L^{2rn}(y) \right), \quad (3.9)$$

The 4 additional canonical pairs are

$$\left( A_m^2(x), -2\Delta \partial_r H_L^{1rn}(y) \right), \left( C^2(x), \frac{1}{2} \Delta (\Delta H_T^1 - \partial_p \partial_q H_T^{1pq})(y) \right). \quad (3.10)$$

In particular, it follows that

$$\{h_{mn}^a(x), \pi^{bkl}(y)\} = \epsilon^{ab} \frac{1}{2} (\delta_m^k \delta_n^l + \delta_m^l \delta_n^k) \delta^{(3)}(x, y). \quad (3.11)$$

### 3.5 Gauge structure

The constraints  $\gamma_\alpha \equiv (\mathcal{H}_{am}, \mathcal{H}_{a\perp})$  are chosen as

$$\mathcal{H}_{am} = -2\epsilon_{ab}\partial^n \pi_{mn}^b = 2\epsilon_{ab}\Delta\partial^n H_{mn}^b, \quad (3.12)$$

$$\mathcal{H}_{a\perp} = \Delta h_a - \partial_m \partial_n h_a^{mn} = \Delta^2 C_a. \quad (3.13)$$

They are first class and abelian

$$\{\gamma_\alpha, \gamma_\beta\} = 0. \quad (3.14)$$

The constraints  $\mathcal{H}_{1m}, \mathcal{H}_{1\perp}$  are those of the standard Hamiltonian formulation of Pauli-Fierz theory expressed in terms of the new variables. The new constraints  $\gamma_\Delta^N = 0$  are

$$\mathcal{H}_{2m} = 0 = \mathcal{H}_{2\perp}. \quad (3.15)$$

They are equivalent to  $\partial^r H_{rm}^1 = 0 = C^2$  and are gauge fixed through the conditions  $A_m^2 = 0 = H_{mn}^{1T}$ . This does not affect  $\pi^{2kl}$ , while  $h_{mn}^1$  is changed by a gauge transformation. The partially gauge fixed theory corresponds to the usual Pauli-Fierz theory in Hamiltonian form as described in section 2.1.

More precisely, the observables of a Hamiltonian field theory with constraints are defined as equivalence classes of functionals that have weakly vanishing Dirac brackets with the constraints and where two functionals are considered as equivalent if they agree on the surface defined by the constraints (see e.g. [28]). The new constraints together with the gauge fixing conditions form second class constraints. The Dirac bracket algebra of observables of this (partially) gauge fixed formulation is isomorphic to the Poisson bracket algebra of observables of the extended formulation on the one hand, and to the Poisson bracket algebra of observables of Pauli-Fierz theory on the other hand.

In the same way, the original constraints  $\mathcal{H}_{1m} = 0 = \mathcal{H}_{1\perp}$  are equivalent to  $\partial^r H_{rm}^2 = 0 = C^1$  and are gauge fixed through  $A_m^2 = 0 = H_{mn}^{2T}$ , leading to the completely reduced theory in terms of the 2 transverse-traceless physical degrees of freedom.

If  $\epsilon^\alpha = (\xi^{am}, \xi^{a\perp})$  collectively denote the gauge parameters, the gauge symmetries are canonically generated by the smeared constraints,

$$\delta_\epsilon z^A = \{z^A, \Gamma[\epsilon]\}, \quad \Gamma[\epsilon] = \int d^3x \gamma_\alpha \epsilon^\alpha, \quad (3.16)$$

so that

$$\delta_\epsilon H_{mn}^a = -\Delta^{-1}\epsilon^{ab}(\delta_{mn}\Delta - \partial_m\partial_n)\xi_b^\perp, \quad \delta_\epsilon A_m^a = \xi_m^a, \quad \delta_\epsilon C^a = 0, \quad (3.17)$$

which implies in particular

$$\delta_\epsilon h_{mn}^a = \partial_m \xi_n^a + \partial_n \xi_m^a, \quad \delta_\epsilon \pi_{mn}^a = \epsilon^{ab}(\delta^{mn}\Delta - \partial^m\partial^n)\xi_b^\perp. \quad (3.18)$$

Note that a way to get local gauge transformations for the fundamental variables is to multiply the constraints by  $\Delta$ , which is allowed when the flat space Laplacian is invertible. This amounts to introducing suitable potentials for the gauge parameters and Lagrange multipliers.

### 3.6 Duality generator

The canonical generator for the infinitesimal duality rotations (3.5) is

$$\begin{aligned}
 D &= \int d^3x \left( -(\mathcal{P}^{TT} H^a)_{mn} H_a^{mn} + 2\Delta \partial_r H_a^{rm} A_m^a \right. \\
 &\quad \left. - \frac{1}{2} \Delta (\Delta H^a - \partial^m \partial^n H_{mn}^a) C_a \right) \\
 &\approx - \int d^3x \mathcal{P}^{TT} (H^a)_{mn} H_a^{mn}.
 \end{aligned} \tag{3.19}$$

The duality generator is only weakly gauge invariant,

$$\{\mathcal{H}_{am}, D\} = \epsilon_{ab} \mathcal{H}_m^b \quad \{\mathcal{H}_{a\perp}, D\} = \epsilon_{ab} \mathcal{H}_\perp^b. \tag{3.20}$$

On the constraint surface, it coincides with the generator found in [1] up to normalisation, where it has been cast in the form of a Chern-Simons term.

### 3.7 Hamiltonian

In terms of the new variables (3.3)–(3.4), the Pauli-Fierz Hamiltonian reads

$$\begin{aligned}
 H_{\text{PF}} &= \int d^3x \left( H^{amn} \Delta^2 H_{amn}^{TT} - 2\Delta \partial^r H_{rn}^2 \partial_s H^{2sn} - \right. \\
 &\quad \left. - \partial^r \partial^s H_{rs}^2 \Delta H^2 - \frac{1}{2} (\partial^r \partial^s H_{rs}^2)^2 + \frac{1}{8} \Delta C^1 \Delta^2 C^1 \right),
 \end{aligned} \tag{3.21}$$

where one can use (2.10) to expand the first term as a local functional of  $H_{mn}^a$ .

The local Hamiltonian  $H = \int d^3x h$  of the manifestly duality invariant action principle (3.6) is

$$\begin{aligned}
 H &= \int d^3x \left( H^{amn} \Delta^2 H_{amn}^{TT} - 2\Delta \partial^r H_{rn}^a \partial_s H_a^{sn} - \right. \\
 &\quad \left. - \partial^r \partial^s H_{rs}^a \Delta H_a - \frac{1}{2} \partial^r \partial^s H_{rs}^a \partial_k \partial_l H_a^{kl} + \frac{1}{8} \Delta C^a \Delta^2 C_a \right) \\
 &= \int d^3x \left( \Delta H_{mn}^a \Delta H_a^{mn} - \frac{1}{2} \Delta H^a \Delta H_a + \frac{1}{8} \Delta C^a \Delta^2 C_a \right).
 \end{aligned} \tag{3.22}$$

It is equivalent to the Pauli-Fierz Hamiltonian since it reduces to the latter when the additional constraints  $\partial^r H_{rm}^1 = 0 = C^2$  hold. Note that the terms proportional to  $\partial^r H_{rm}^a$  and  $C^a$  may be dropped since they vanish on the constraint surface,  $H \approx \int d^3x H^{amn} \Delta^2 H_{amn}^{TT}$ .

The Hamiltonian is gauge invariant on the constraint surface,

$$\{H, \Gamma[\xi]\} = \int d^3x \mathcal{H}_m^a \partial^m \xi_a^\perp. \tag{3.23}$$

In order for the action (3.6) to be gauge invariant, it follows from (3.23) that the Lagrange multipliers  $u^\alpha$  need to transform as

$$\delta_\xi u^{am} = \dot{\xi}^{am} - \partial^m \xi^{a\perp}, \quad \delta_\xi u^{a\perp} = \dot{\xi}^{a\perp}. \tag{3.24}$$

### 3.8 Poincaré generators

Consider now a symmetry generator of Pauli-Fierz theory. It is defined by an observable  $K[h^1, \pi^2]$  whose representative is weakly conserved in time,

$$\frac{\partial}{\partial t}K + \{K, H_{\text{PF}}\} \approx 0. \quad (3.25)$$

Since the new Hamiltonian differs from the Pauli-Fierz one by terms proportional to the new constraints  $\gamma_{\Delta}^N = 0$  given explicitly in (3.15), we have

$$H = H_{\text{PF}} + \int d^3x \gamma_{\Delta}^N k^{\Delta}. \quad (3.26)$$

Furthermore, since  $K$ , when expressed in terms of the new variables, does not depend on  $H_L^2, H_T^1$ , so that  $\{K, \int d^3x \gamma_{\Delta}^N k^{\Delta}\} \approx 0$  in the extended theory, it follows that  $K$  is also weakly conserved and thus a symmetry generator of the extended theory,

$$\frac{\partial}{\partial t}K + \{K, H\} \approx 0. \quad (3.27)$$

Consider then the Poincaré generators  $Q_G(\omega, a)$  of Pauli-Fierz theory as described in appendix A. When expressed in terms of the new variables, they are representatives for the Poincaré generators of the extended theory. Indeed, we just have shown that they are symmetry generators, while we have argued in section 3.5 that their Poisson algebra is isomorphic when restricted to their respective constraint surfaces.

Since symmetry generators form a Lie algebra with respect to the Poisson bracket,  $\{Q_G(\omega, a), D\}$  is a symmetry generator. In much the same way as for the Hamiltonian, we now want to show that one can find representatives for the Poincaré generators that are duality invariant,

$$\{Q_G^D(\omega, a), D\} = 0, \quad (3.28)$$

by adding terms proportional to the new constraints.

The first step in the proof consists in showing that the reduced phase space generators, i.e., the generators  $Q_G(\omega, a)$  for which all variables except for the physical  $H_{TT}^a$  have been set to zero, are duality invariant. All other contributions to  $Q_G(\omega, a)$  are then shown to be proportional to the constraints of Pauli-Fierz theory. Both these steps follow from straightforward but slightly tedious computations. For the generators of rotations and boosts for instance the computation is more involved because the explicit  $x^i$  dependence has to be taken into account when performing integrations by parts.

In terms of the new variables, the terms proportional to the constraints are bilinear in  $(h^1, A^2)$ ,  $(\pi^2, A^2)$ ,  $(h^1, C^1)$  and  $(\pi^2, C^1)$ . The duality invariant generators  $Q_G^D(\omega, a)$  are then obtained by adding the same terms with the substitution  $h^1 \rightarrow h^2$ ,  $A^2 \rightarrow -A^1$ ,  $\pi^2 \rightarrow -\pi^1$  and  $C^1 \rightarrow C^2$ , while keeping unchanged the terms involving only the physical variables  $H_{TT}^a$ .

As a consequence, the duality invariant Poincaré transformations of  $h^1, \pi^2$  are unchanged on the extended constraint surface. They are given by (A.31)–(A.32) where

$\xi^\perp = -\omega^0{}_\nu x^\nu + a^0$  and  $\xi^i = -\omega^i{}_\nu x^\nu + a^i$ . Because of (3.28), those of  $h^2, -\pi^1$  are obtained, on the constraint surface, by applying a duality rotation to the right hand-sides of (A.31)–(A.32).

An open question that we plan to address elsewhere is the construction of the canonical generators for global Poincaré transformations in the presence of both types of sources that will be introduced in the next section.

## 4. Coupling to conserved electric and magnetic sources

### 4.1 Interacting variational principle

We define

$$h_{0m}^a = n_m^a = h_{m0}^a, \quad h_{00}^a = -2n^a, \quad (4.1)$$

and consider the action

$$S_T[z^A, u^\alpha; T^{a\mu\nu}] = \frac{1}{16\pi G} S_G + S^J, \quad (4.2)$$

with  $S_G$  given in (3.6) and the gauge invariant interaction term

$$S^J = \int d^4x \frac{1}{2} h_{\mu\nu}^a T_a^{\mu\nu}, \quad \partial_\mu T_a^{\mu\nu} = 0, \quad (4.3)$$

where  $T_a^{\mu\nu} \equiv (T^{\mu\nu}, \Theta^{\mu\nu})$  are external, conserved electric and magnetic energy-momentum tensors.

### 4.2 Linearized Taub-NUT solution

We start by considering the sources corresponding to a point-particle gravitational dyon with electric mass  $M$  and magnetic mass  $N$  at rest at the origin of the coordinate system, for which

$$T_a^{\mu\nu}(x) = \delta_0^\mu \delta_0^\nu M_a \delta^{(3)}(x^i), \quad M_a = (M, N). \quad (4.4)$$

In this case, only the constraints (3.13) are affected by the interaction and become

$$\mathcal{H}_{a\perp} = -16\pi G M_a \delta^{(3)}(x). \quad (4.5)$$

They are solved by

$$\Delta C^a = GM^a \left( \frac{4}{r} \right), \quad (4.6)$$

where  $r = \sqrt{x^i x_i}$ . It is then straightforward to check that all equations of motions are solved by

$$\begin{aligned} C^a &= GM^a(2r), & n^a &= GM^a \left( -\frac{1}{r} \right), & A_m^a &= n^{am} = H_{mn}^a = 0, \\ h_{mn}^a &= GM^a \left( \delta_{mn} + \frac{x_m x_n}{r^3} \right), & \pi_a^{mn} &= 0. \end{aligned} \quad (4.7)$$

The usual Schwarzschild form is obtained after a gauge transformation with parameter  $\xi^{am} = GM^a(-\frac{1}{2}\frac{x^m}{r})$ ,  $\xi^{a\perp} = 0$ . The solution then reads

$$\begin{aligned} C^a &= GM^a(2r), & n^a &= GM^a\left(-\frac{1}{r}\right), & A_m^a &= GM^a\left(-\frac{1}{2}\frac{x_m}{r}\right), & n^{am} &= H_{mn}^a = 0, \\ h_{mn}^a &= GM^a\left(\frac{2x_mx_n}{r^3}\right), & \pi_a^{mn} &= 0. \end{aligned} \tag{4.8}$$

By computing the Riemann tensor in terms of the canonical variables in appendix B, we show that this solution describes the linearized Taub-NUT solution. It resolves the string singularity of the linearized Taub-NUT solution in the standard Pauli-Fierz formulation. In spherical coordinates, the latter can for instance be described by

$$h_{rr} = \frac{2GM}{r} = h_{00}, \quad h_{0\varphi} = -2N(1 - \cos\theta), \tag{4.9}$$

and all other components vanishing, with a string-singularity along the negative  $z$ -axis.

## 5. Surface charges

### 5.1 Regge-Teitelboim revisited

Because the theory is not local as a Hamiltonian gauge theory, the analysis of surface charges cannot directly be performed as in [29, 30]. We thus revert to the original Hamiltonian method of [31, 32] and adapt it to the present situation of exact solutions, where there is no need to discuss fall-off conditions.

Let  $\mathcal{L}_H = a_A \dot{z}^A - h - \gamma_\alpha u^\alpha$ , with  $h$  a first class Hamiltonian density and  $\gamma_\alpha$  first class constraints and define  $\phi^i = (z^A, u^\alpha)$ . Even though it is not so for our theory, let us first run through the arguments in the case where one has Darboux coordinates for the symplectic structure, i.e., when  $\sigma_{AB} = \frac{\partial a_B}{\partial z^A} - \frac{\partial a_A}{\partial z^B}$  is the constant symplectic matrix. We furthermore suppose that we are in a source-free region of spacetime. In this case one can show that

$$\delta_\varepsilon z^A \frac{\delta \mathcal{L}_H}{\delta z^A} + \delta_\varepsilon u^\alpha \frac{\delta \mathcal{L}_H}{\delta u^\alpha} = -\partial_0(\gamma_\alpha \varepsilon^\alpha) - \partial_i s_\varepsilon^i, \tag{5.1}$$

where  $s_\varepsilon^i = s_\varepsilon^i[z, u]$  vanishes when the Hamiltonian equations of motion, including constraints, are satisfied,  $s_\varepsilon^i \approx 0$ . This identity merely expresses the general fact that the Noether current  $s_\varepsilon^\mu$  associated to a gauge symmetry can be taken to vanish when the equations of motions hold (see e.g. [28], section 3),  $s_\varepsilon^\mu \approx 0$ , and that the integrand of the generator is given by (minus) the constraints contracted with the gauge parameters in the Hamiltonian formalism,  $s_\varepsilon^0 = -\gamma_\alpha \varepsilon^\alpha$ . An explicit expression for  $s_\varepsilon^i$  in terms of the structure functions can for instance be found in appendix D of [30]. Using integrations by parts, one can write the variations of the constraints under a change of the canonical coordinates  $z^A$  as an Euler-Lagrange derivative, up to a total derivative,

$$\delta_z(\gamma_\alpha \varepsilon^\alpha) = \delta z^A \frac{\delta(\gamma_\alpha \varepsilon^\alpha)}{\delta z^A} - \partial_i k_\varepsilon^i. \tag{5.2}$$

where  $k_\varepsilon^i = k_\varepsilon^i[\delta z, z]$  depends linearly on  $\delta z^A$  and its spatial derivatives. Taking the time derivative of (5.2) and using a variation  $\delta\phi$  of (5.1) to eliminate  $\partial_0\delta z(\gamma_\alpha\varepsilon^\alpha)$ , one finds

$$\partial_i\left(\partial_0 k_\varepsilon^i - \delta\phi s_\varepsilon^i\right) = \partial_0\left(\delta z^A \frac{\delta(\gamma_\alpha\varepsilon^\alpha)}{\delta z^A}\right) + \delta\phi\left(\delta_\varepsilon z^A \frac{\delta\mathcal{L}_H}{\delta z^A} + \delta_\varepsilon u^\alpha \frac{\delta\mathcal{L}_H}{\delta u^\alpha}\right). \quad (5.3)$$

One now takes  $\varepsilon_s^\alpha$  to satisfy  $\delta_{\varepsilon_s} z_s^A = 0 = \delta_{\varepsilon_s} u_s^\alpha$ . Note that in the case of Darboux coordinates, this also implies that  $\frac{\delta(\gamma_\alpha\varepsilon_s^\alpha)}{\delta z^A} = 0$ . If furthermore  $z_s^A, u_s^\alpha$  is a solution of the Hamiltonian equations of motion, the r.h.s. of (5.3) also vanishes. By using a contracting homotopy with respect to  $\delta\phi^i$  and their spatial derivatives, one deduces that

$$\partial_0 k_{\varepsilon_s}^i[\delta z, z_s] = (\delta\phi s_{\varepsilon_s}^i)[z_s] - \partial_j k_{\varepsilon_s}^{[ij]}, \quad (5.4)$$

where  $k_{\varepsilon_s}^{[ij]} = k_{\varepsilon_s}^{[ij]}[\delta\phi, \phi_s]$  depends linearly on  $\delta\phi^i$  and their spatial derivatives. Finally, when  $\delta z_s^A, \delta u_s^\alpha$  satisfy the linearized Hamiltonian equations of motion, including constraints, we find from (5.2) and (5.4) that

$$\partial_i k_{\varepsilon_s}^i[\delta z_s, z_s] = 0, \quad \partial_0 k_{\varepsilon_s}^i[\delta z_s, z_s] - \partial_j t_{\varepsilon_s}^{[ij]} = 0. \quad (5.5)$$

At a fixed time  $t = x^0$ , consider a closed 2 dimensional surface  $S$ ,  $\partial S = 0$ , for instance a sphere with radius  $r$  and define the surface charge 1-forms by

$$\oint_S \mathcal{Q}_{\varepsilon_s}[\delta z_s, z_s] = \oint_S d^2 x_i k_{\varepsilon_s}^i[\delta z_s, z_s], \quad (5.6)$$

where  $d^2 x_i = \frac{1}{2}\epsilon_{ijk} dx^j \wedge dx^k$ . The first relation of (5.5) implies that the surface charge 1-form only depends on the homology class of the closed surface  $S$ ,

$$\oint_{S_1} d^2 x_m k_{\varepsilon_s}^m[\delta z_s, z_s] = \oint_{S_2} d^2 x_m k_{\varepsilon_s}^m[\delta z_s, z_s]. \quad (5.7)$$

Here  $S_1 - S_2 = \partial\Sigma$ , where  $\Sigma$  is a three-dimensional volume at fixed time  $t$  containing no sources. For instance, the surface charge 1-form does not depend on  $r$ . The second relation of (5.5) implies that it is conserved in time and so does not depend on  $t$  either,

$$\frac{d}{dt} \oint_S \mathcal{Q}_{\varepsilon_s}[\delta z_s, z_s] = 0. \quad (5.8)$$

The question is then whether these charge 1-forms are integrable, see e.g. [33, 34, 30] for a discussion.

## 5.2 Linear theories

In the case of linear theories, the latter problem does not arise and the whole analysis simplifies. One can replace (5.2) by

$$\gamma_\alpha\varepsilon^\alpha = z^A \frac{\delta(\gamma_\alpha\varepsilon^\alpha)}{\delta z^A} - \partial_i k_\varepsilon^i[z], \quad (5.9)$$

where  $\delta/\delta z^A$  are the (spatial) Euler-Lagrange derivatives and  $k_\varepsilon^i[z]$  depends linearly both on the phase space variables  $z^A$  and their spatial derivatives and on the gauge parameters. One then uses (5.1) directly to eliminate  $\partial_0(\gamma_\alpha \varepsilon^\alpha)$  from the time derivative of (5.9), to get

$$\partial_i [\partial_0 k_\varepsilon^i - s_\varepsilon^i] = \partial_0 \left[ z^A \frac{\delta(\gamma_\alpha \varepsilon^\alpha)}{\delta z^A} \right] + \delta_\varepsilon z^A \frac{\delta \mathcal{L}_H}{\delta z^A} + \delta_\varepsilon u^\alpha \frac{\delta \mathcal{L}_H}{\delta u^\alpha}. \quad (5.10)$$

For gauge parameters  $\varepsilon_s^\alpha$  that satisfy

$$\delta_{\varepsilon_s} z^A = 0 = \delta_{\varepsilon_s} u^\alpha, \quad (5.11)$$

one then arrives at

$$\partial_i k_{\varepsilon_s}^i [z] = -\gamma_\alpha \varepsilon^\alpha, \quad \partial_0 k_{\varepsilon_s}^i [z] = s_{\varepsilon_s}^i [z, u] - \partial_j k_{\varepsilon_s}^{[ij]}. \quad (5.12)$$

For a solution  $z_s^a, u_s^\alpha$ , the surface charges

$$\mathcal{Q}_{\varepsilon_s} [z_s] = \oint_S d^2 x_i k_{\varepsilon_s}^i [z_s], \quad (5.13)$$

are again independent of  $r$  and  $t$ .

When this analysis is applied to the Hamiltonian formulation of Pauli-Fierz theory, one finds the standard expressions

$$k_\varepsilon^i [z] = 2\xi_m \pi^{mi} - \xi^\perp (\delta^{mn} \partial^i - \delta^{mi} \partial^n) h_{mn} + h_{mn} (\delta^{mn} \partial^i - \delta^{ni} \partial^m) \xi^\perp, \quad (5.14)$$

while the only solutions to (5.11) are  $\xi_{\mu s} = -\omega_{[\mu\nu]} x^\nu + a_\mu$ , for some constants  $a_\mu$ ,  $\omega_{[\mu\nu]} = -\omega_{[\nu\mu]}$ . In this context of flat space, Greek indices take values from 0 to 3 with  $\mu = (\perp, i)$ . Indices  $\mu$  are lowered and raised with  $\eta_{\mu\nu} = \text{diag}(-1, 1, 1, 1)$ .

### 5.3 Electric and magnetic energy-momentum and angular momentum surface charges

The previous analysis is not directly applicable to our case since we do not have Darboux coordinates and the Poisson brackets of the fundamental variables are non-local. In particular, the gauge transformations (3.17) do not allow for non trivial solutions to  $\delta_{\varepsilon_s} z^A = 0$ . We also have to keep the sources explicitly throughout the argument, because  $\Delta^{-1}$  applied to localized sources will spread them out throughout space and we need to check that we are only dropping terms that indeed vanish outside of the sources.

In the presence of the sources, the constraints  $\gamma_\alpha^J = (\mathcal{H}_{am}^J, \mathcal{H}_{a\perp}^J)$  are determined

$$\mathcal{H}_{am}^J = \mathcal{H}_{am} - (16\pi G) T_{am}^0, \quad \mathcal{H}_{a\perp}^J = \mathcal{H}_{a\perp} - (16\pi G) T_{a0}^0. \quad (5.15)$$

Instead of (5.9), we can write

$$\begin{aligned} \gamma_\alpha^J \varepsilon^\alpha &= (\partial^m \xi^{an} + \partial^n \xi^{am}) \epsilon_{ab} \pi_{mn}^b + (\delta^{mn} \Delta - \partial^m \partial^n) \xi^{a\perp} h_{amn} - \partial_i \tilde{k}_\varepsilon^i [z] \\ &\quad - (16\pi G) (T_{am}^0 \xi^{am} + T_{a0}^0 \xi^{a\perp}), \end{aligned} \quad (5.16)$$



where

$$\tilde{k}_\varepsilon^i[z] = 2\xi_m^a \epsilon_{ab} \pi^{bmi} - \xi^{a\perp} (\delta^{mn} \partial^i - \delta^{mi} \partial^n) h_{amn} + h_{amn} (\delta^{mn} \partial^i - \delta^{ni} \partial^m) \xi^{a\perp}. \quad (5.17)$$

Consider now gauge parameters  $\epsilon_s^\alpha(x)$  satisfying the conditions

$$\begin{cases} \partial^m \xi_s^{an} + \partial^n \xi_s^{am} = 0 = \partial_0 \xi_s^{am} - \partial^m \xi_s^{a\perp}, \\ (\delta^{mn} \Delta - \partial^m \partial^n) \xi_s^{a\perp} = 0 = \partial_0 \xi_s^{a\perp}, \end{cases} \quad (5.18)$$

The general solution to conditions (5.18) can be written as

$$\xi_{\mu s}^a = -\omega_{[\mu\nu]}^a x^\nu + a_\mu^a, \quad (5.19)$$

for some constants  $a_\mu^a$ ,  $\omega_{[\mu\nu]}^a = -\omega_{[\nu\mu]}^a$ . It follows in particular that the surface charges

$$\mathcal{Q}_{\varepsilon_s}[z_s] = \frac{1}{16\pi G} \oint_S d^2 x_i \tilde{k}_{\varepsilon_s}^i[z_s], \quad (5.20)$$

do not depend on the homology class of  $S$  outside of sources.

Assuming  $\Delta$  invertible, the equations of motion associated to  $\mathcal{L}_T = \frac{1}{16\pi G} \mathcal{L}_H + \mathcal{L}^J$  imply in particular that

$$\begin{aligned} \partial_0 h_{mn}^a &= \partial_m n_n^a + \partial_n n_m^a - 2\epsilon^{ab} \Delta H_{bmn} + \epsilon^{ab} \delta_{mn} \Delta H_b \\ &\quad + (16\pi G) \epsilon^{ab} \left( \Delta^{-1} (\mathcal{O}T_b)_{mn} + \frac{1}{2} \Delta^{-2} \partial_m \epsilon_{npq} \partial^p \partial_k T_b^{kq} + \frac{1}{2} \Delta^{-2} \partial_n \epsilon_{mpq} \partial^p \partial_k T_b^{kq} \right), \\ \epsilon_{ab} \partial_0 \pi_{mn}^b &= (\mathcal{P}^{TT} H_a)_{mn} + (8\pi G) T_{amn} - \\ &\quad - \frac{1}{2} (\delta^{mn} \Delta - \partial^m \partial^n) \left( 2n_a + \frac{1}{2} \Delta C_a \right). \end{aligned} \quad (5.21)$$

By direct computation using the equations of motion, one then finds

$$\partial_0 \tilde{k}_{\varepsilon_s}^i[z_s] = (16\pi G) (\xi_{\mu s}^a T_a^{\mu i}) - \partial_j k_{\varepsilon_s}^{[ij]}[z_s, u_s], \quad (5.22)$$

with

$$\begin{aligned} k_{\varepsilon_s}^{[ij]}[z, u] &= \left( 2n_a^i \partial^j \xi_s^{a\perp} + \xi_s^{a\perp} \partial^i n_a^j + \xi_s^{ai} \partial^j \left( 2n_a + \frac{1}{2} \Delta C_a \right) + \xi_{sm}^a \epsilon^{mpq} \partial_p \partial^i H_{aq}^j \right. \\ &\quad \left. + \omega^{aj} \partial^k H_{ak}^i + \omega^{ai} \partial^j H_a + 2\omega^{ak} \partial^i H_{ak}^j + 16\pi G \epsilon^{ab} \epsilon^{imq} \Delta^{-1} T_{bq}^j \partial_m \xi_{as}^\perp \right. \\ &\quad \left. + 8\pi G \epsilon^{ab} \epsilon^{mpq} \partial_p \Delta^{-2} \partial^i T_{bq}^j \partial_m \xi_{as}^\perp - (i \longleftrightarrow j) \right) \\ &\quad + \epsilon^{ijk} \left[ \omega_k^a \left( 2n_a + \frac{1}{2} \Delta C_a \right) - \xi_s^{am} (\Delta H_{amk} - \partial_m \partial^r H_{ark}) \right. \\ &\quad \left. - 16\pi G \epsilon^{ab} \Delta^{-1} \partial^r T_{brk} \xi_{as}^\perp + 8\pi G \epsilon^{ab} (\Delta^{-1} T_{bk}^m + \Delta^{-2} \partial^m \partial^r T_{brk}) \partial_m \xi_{as}^\perp \right], \end{aligned} \quad (5.23)$$

where  $\omega_{mn}^a = \omega^{ak} \epsilon_{kmn}$ . The surfaces charges (5.20) are thus also time-independent outside of sources.

Finally, the surface charges are gauge invariant,

$$\tilde{k}_{\varepsilon_s}^i[\delta\eta z] = \partial_j r_{\varepsilon_s, \eta}^{[ij]}, \quad (5.24)$$

$$r_{\varepsilon_s, \eta}^{[ij]} = \left( 2\xi_s^{aj} \partial^i \eta_a^\perp + 2\eta_a^j \partial^i \xi_s^{a\perp} + \xi_s^{a\perp} \partial^j \eta_a^i - (i \longleftrightarrow j) \right) - 2\epsilon^{ijk} \omega_k^a \eta_a^\perp. \quad (5.25)$$

Defining

$$Q_{\varepsilon_s}[z] = \frac{1}{2} \omega_{\mu\nu}^a J_a^{\mu\nu} - a_\mu^a P_a^\mu, \quad (5.26)$$

we get for the individual generators

$$(16\pi G)P_a^\perp = - \oint_{S^\infty} d^2x_m \partial^m \Delta C_a = \oint_{S^\infty} d^2x_m (\partial_n h_a^{mn} - \partial^m h_a), \quad (5.27)$$

$$(16\pi G)P_a^n = 2 \oint_{S^\infty} d^2x_m \epsilon_{ab} \Delta H^{bnm} = -2 \oint_{S^\infty} d^2x_m \epsilon_{ab} \pi^{bmn}, \quad (5.28)$$

$$(16\pi G)J_a^{kl} = 2 \oint_{S^\infty} d^2x_m \epsilon_{ab} \left( \Delta H^{bmk} x^l - \Delta H^{bml} x^k \right) \quad (5.29)$$

$$= -2 \oint_{S^\infty} d^2x_m \epsilon_{ab} \left( \pi^{bmk} x^l - \pi^{bml} x^k \right), \quad (5.30)$$

$$(16\pi G)J_a^{\perp k} = \oint_{S^\infty} d^2x_m \left( \Delta C_a \delta^{mk} - \partial^m \Delta C_a x^k \right) \quad (5.31)$$

$$= \oint_{S^\infty} d^2x_m \left[ (\partial_n h_a^{mn} - \partial^m h_a) x^k - h_a^{mk} + h_a \delta^{mk} \right]. \quad (5.32)$$

The only non-vanishing surface charges of the dyon sitting at the origin are

$$P_a^\perp = M_a. \quad (5.33)$$

As expected, they measure the electric and magnetic mass of the dyon.

For later use, we combine  $\tilde{k}_\varepsilon^i, k_\varepsilon^{[ij]}$  into the  $n - 2$  forms  $k_\varepsilon[z, u]$  through the following expressions in Cartesian coordinates,

$$k_\varepsilon = k_\varepsilon^{[\mu\nu]} d^{n-2} x_{\mu\nu}, \quad k_\varepsilon^{[0i]} = \tilde{k}_\varepsilon^i, \quad (5.34)$$

$$d^{n-k} x_{\mu_1 \dots \mu_k} = \frac{1}{k!(4-k)!} \epsilon_{\mu_1 \dots \mu_k \nu_{k+1} \dots \nu_4} dx^{\nu_{k+1}} \dots dx^{\nu_4}, \quad (5.35)$$

where  $\epsilon_{\alpha\beta\gamma\delta}$  is completely skew-symmetric with  $\epsilon_{0123} = 1$  and the wedge product between the differentials is understood. Equations (5.16) and (5.22) can then be summarized by

$$dk_{\varepsilon_s} \approx -(16\pi G)T_{\varepsilon_s}, \quad T_{\varepsilon_s} = T_{\alpha\nu}^\mu \xi_s^{\alpha\nu} d^3 x_\mu, \quad dT_{\varepsilon_s} = 0, \quad (5.36)$$

where closure of the  $n - 1$ -forms  $T_{\varepsilon_s}$  follows from the conservation of the sources, the symmetry of the energy-momentum tensor and (5.19).

**Remark.** In fact we have checked here that the standard expressions for surface charges in Pauli-Fierz theory, when extended in a duality invariant way, have all the expected properties. More interesting would be to develop the theory of surface charges from scratch in theories where the Poisson brackets among the fundamental variables are not local to see if the ones we have found exhaust all possibilities. From the preceding discussion we see that pseudo-differential operators will play a crucial role for a discussion of these generalized conservation laws, as they do in the discussion of ordinary conservation laws for evolution equations of the Korteweg-de Vries type for instance.

### 5.4 Poincaré transformations of surface charges

Suppose now that  $z_s^A, u_s^\alpha$  solve the equations of motions for the conserved sources  $T_a^{\mu\nu}(x)$ . Let  $z_s'^A, u_s'^\alpha$  be the solution associated to new sources  $T_a'^{\mu\nu}(x)$  related to  $T_a^{\mu\nu}(x)$  through a (proper) Poincaré transformation,  $x'^\mu = \Lambda^\mu{}_\nu x^\nu + b^\mu$  with  $|\Lambda| = 1$ ,

$$T_a'^{\mu\nu}(x') = \Lambda^\mu{}_\alpha \Lambda^\nu{}_\beta T_a^{\alpha\beta}(x). \quad (5.37)$$

For instance, starting from the conserved energy-momentum tensors (4.4) of a dyon sitting at the origin with world-line  $z^\mu = \delta_0^\mu s$ , one can obtain in this way the conserved energy-momentum tensors of a dyon moving along a straight line,  $z'^\mu = u^\mu s + a^\mu$  with  $u^\mu, a^\mu$  constant,  $u^\mu u_\mu = -1$  and  $s$  the proper time,

$$T_a'^{\mu\nu}(x') = M_a u^\nu \int d\lambda \delta^{(4)}(x' - z'(\lambda)) \frac{dz'^\mu}{d\lambda} = M_a \frac{u^\mu u^\nu}{u^0} \delta^{(3)}(x'^i - z'^i(x^0)). \quad (5.38)$$

when  $\Lambda^\mu{}_0 = u^\mu$ .

Assume then that the  $\xi_{as}^\mu(x)$  transform like vectors

$$\xi_{as}'^\nu(x') = \Lambda^\nu{}_\alpha \xi_{as}^\alpha(x) = -(\Lambda \omega_a \Lambda^{-1} x')^\nu + (\Lambda \omega_a \Lambda^{-1} b + \Lambda a_a)^\nu, \quad (5.39)$$

which implies that the  $T_{\varepsilon_s}$  are closed Poincaré invariant  $n - 1$  forms,

$$T_{\varepsilon_s}'(x', dx') = T_{\varepsilon_s}(x, dx). \quad (5.40)$$

We can then use the following variant of the tube lemma. Suppose that at fixed time  $x^0$ ,  $T_a^{0\nu}(x) \xi_s^{a\nu}(x)$  has compact support and that there exists a tube, i.e., a space-time volume  $\mathcal{W}$  connecting the hypersurfaces  $\Omega : K = x^0$  and  $\Omega' : K = x'^0 = \Lambda^0{}_\nu x^\nu + b^0$  with  $K$  a constant such that,  $\partial\mathcal{W} = \Omega' - \Omega + \mathcal{T}$ . If nothing flows out through  $\mathcal{T}$ ,  $\int_{\mathcal{T}} T_{\varepsilon_s} = 0$ , it follows from Stokes' theorem and (5.40) that

$$\int_{\Omega} T_{\varepsilon_s} = \int_{\Omega'} T_{\varepsilon_s} = \int_{\Omega'} T_{\varepsilon_s}'. \quad (5.41)$$

If we now compute the surface charges for a large enough sphere  $S$  at fixed  $x^0$  containing both  $T_a^{0\nu}(x) \xi_s^{a\nu}(x)$  and  $T_a'^{0\nu}(x) \xi_s'^{a\nu}(x)$ , it finally follows from (5.16) or (5.36) that the surface charges evaluated for the new solutions  $z'^A$  are obtained from those of the old solutions  $z^A$  through

$$\mathcal{Q}_{\varepsilon_s}'[z'_s] = \mathcal{Q}_{\varepsilon_s}[z_s]. \quad (5.42)$$

### Acknowledgments

The authors thank R. Argurio, X. Bekaert, F. Dehouck, A. Gomberoff, and L. Houart for useful discussions. This work is supported in part by a ‘‘Pôle d’Attraction Interuniversitaire’’ (Belgium), by IISN-Belgium, convention 4.4505.86, by the Fund for Scientific Research-FNRS (Belgium), and by the European Commission programme MRTN-CT-2004-005104, in which the authors are associated to V.U. Brussel.

## A. Poincaré generators for Pauli-Fierz theory

In this appendix, we assume that the canonical variables vanish sufficiently fast at the boundary so that integrations by parts can be used even if the gauge parameters do not vanish at the boundary.

In the Hamiltonian formulation of general relativity [25], the canonically conjugate variables are the spatial 3 metric  $g_{ij}$  and the extrinsic curvature  $\pi^{ij}$ . The constraints are explicitly given by

$$\mathcal{H}_\perp = \frac{1}{\sqrt{g}} \left( \pi^{mn} \pi_{mn} - \frac{1}{2} \pi^2 \right) - \sqrt{g} R, \quad \mathcal{H}_i = -2 \nabla_j \pi^j_i. \quad (\text{A.1})$$

The associated generators of gauge transformation  $H[\xi] = \int d^3x (\mathcal{H}_\perp \xi^\perp + \mathcal{H}_i \xi^i)$  satisfy the so-called surface deformation algebra [35, 36],

$$\{H[\xi], H[\eta]\} = H[[\xi, \eta]_{\text{SD}}], \quad (\text{A.2})$$

$$[\xi, \eta]_{\text{SD}}^\perp = \xi^i \partial_i \eta^\perp - \eta^i \partial_i \xi^\perp, \quad (\text{A.3})$$

$$[\xi, \eta]_{\text{SD}}^i = g^{ij} \left( \xi^\perp \partial_j \eta^\perp - \eta^\perp \partial_j \xi^\perp \right) + \xi^j \partial_j \eta^i - \eta^j \partial_j \xi^i. \quad (\text{A.4})$$

When the parameters  $f, g$  of gauge transformations depend on the canonical variables, (A.2) is replaced by [37]

$$\{H[f], H[g]\} = H[k], \quad (\text{A.5})$$

$$k = [f, g]_{\text{SD}} + \delta_g f - \delta_f g - \bar{m}, \quad (\text{A.6})$$

$$m^\perp = \int d^3x' \left[ \{f^\perp, g^\perp(x')\} \mathcal{H}_\perp(x') + \{f^\perp, g^j(x')\} \mathcal{H}_j(x') \right], \quad (\text{A.7})$$

$$m^i = \int d^3x' \left[ \{f^i, g^\perp(x')\} \mathcal{H}_\perp(x') + \{f^i, g^j(x')\} \mathcal{H}_j(x') \right], \quad (\text{A.8})$$

where

$$\delta_\xi g_{ij} = \nabla_i \xi_j + \nabla_j \xi_i + 2D_{ijkl} \pi^{kl} \xi^\perp, \quad (\text{A.9})$$

$$D_{ijkl} = \frac{1}{2\sqrt{g}} (g_{ik} g_{jl} + g_{jk} g_{il} - g_{ij} g_{kl}), \quad (\text{A.10})$$

$$\begin{aligned} \delta_\xi \pi^{ij} = & -\xi^\perp \sqrt{g} \left( R^{ij} - \frac{1}{2} g^{ij} R \right) + \frac{\xi^\perp}{2\sqrt{g}} g^{ij} \left( \pi^{kl} \pi_{kl} - \frac{1}{2} \pi^2 \right) \\ & - 2 \frac{\xi^\perp}{\sqrt{g}} \left( \pi^{im} \pi_m^j - \frac{1}{2} \pi^{ij} \pi \right) + \sqrt{g} (\nabla^j \nabla^i \xi^\perp - g^{ij} \nabla_m \nabla^m \xi^\perp) \\ & + \nabla_m (\pi^{ij} \xi^m) - \nabla_m \xi^i \pi^{mj} - \nabla_m \xi^j \pi^{mi}. \end{aligned} \quad (\text{A.11})$$

Let  $g_{ij} = \delta_{ij} + h_{ij}$  and consider the canonical change of variables from  $g_{ij}, \pi^{kl}$  to  $z^A = (h_{ij}, \pi^{kl})$ . We will expand in terms of the homogeneity in the new variables and use the flat metric  $\delta_{ij}$  to raise and lower indices in the remainder of this appendix. Furthermore, Greek indices take values from 0 to 3 with  $\mu = (\perp, i)$ . Indices are lowered and raised with  $\eta_{\mu\nu} = \text{diag}(-1, 1, 1, 1)$  and its inverse. Let  $\tilde{\omega}_{\mu\nu} = -\tilde{\omega}_{\nu\mu}$ .

To lowest order, i.e., when  $g_{ij} = \delta_{ij}$ , the vector fields

$$\xi_P(\tilde{\omega}, \tilde{a})^\mu = -\tilde{\omega}^\mu{}_i x^i + \tilde{a}^\mu, \quad (\text{A.12})$$

equipped with the surface deformation bracket form a representation of the Poincaré algebra [31],

$$[\xi_P(\tilde{\omega}_1, \tilde{a}_1), \xi_P(\tilde{\omega}_2, \tilde{a}_2)]_{\text{SD}}^{(0)} = \xi_P([\tilde{\omega}_1, \tilde{\omega}_2], \tilde{\omega}_1 \tilde{a}_2 - \tilde{\omega}_2 \tilde{a}_1). \quad (\text{A.13})$$

For the gauge generators, we find  $H[\xi] = H^{(1)}[\xi] + H^{(2)}[\xi] + H^{(3)}[\xi] + \dots$ , where

$$H^{(1)}[\xi] = \int d^3x \left( -2\partial^j \pi_{ij} \xi^i + (\partial^i \partial^j h_{ij} - \Delta h) \xi^\perp \right) \quad (\text{A.14})$$

$$= \int d^3x \left( \mathcal{H}_m^{(1)} \xi^m + \mathcal{H}_\perp^{(1)} \xi^\perp \right) \quad (\text{A.15})$$

are the gauge generators associated to the constraints (2.3) of the Pauli-Fierz theory. Because

$$H[[\xi, \eta]_{\text{SD}}] = H^{(1)}[[\xi, \eta]_{\text{SD}}^{(0)}] + H^{(2)}[[\xi, \eta]_{\text{SD}}^{(0)}] + H^{(1)}[[\xi, \eta]_{\text{SD}}^{(1)}] + O(z^3), \quad (\text{A.16})$$

we have to lowest non trivial order

$$\left\{ H^{(1)}[\xi], H^{(2)}[\eta] \right\} = H^{(1)}[[\xi, \eta]_{\text{SD}}^{(0)}]. \quad (\text{A.17})$$

This means that  $H^{(2)}[\eta]$  are observables, i.e., weakly gauge invariant functionals.

One can use integrations by parts to show that  $H^{(1)}[\xi_P] = 0$ . It then follows that

$$\left\{ H[\xi_P], H[\eta_P] \right\} = \left\{ H^{(2)}[\xi_P], H^{(2)}[\eta_P] \right\} + O(z^3). \quad (\text{A.18})$$

For vectors  $\xi_P(\tilde{\omega}, \tilde{a}), \eta_P(\tilde{\theta}, \tilde{b})$  of the form (A.12), the first term on the r.h.s. of (A.16) vanishes on account of (A.13). To lowest non trivial order, (A.2) then implies

$$\left\{ H^{(2)}[\xi_P], H^{(2)}[\eta_P] \right\} = H^{(2)}[[\xi_P, \eta_P]_{\text{SD}}^{(0)}] + H^{(1)}[[\xi_P, \eta_P]_{\text{SD}}^{(1)}]. \quad (\text{A.19})$$

The generators  $H^{(2)}[\xi_P]$  equipped with the Poisson bracket thus form a representation of the Poincaré algebra when the constraints of the Pauli-Fierz theory are satisfied. Explicitly, the term proportional to the constraints is

$$H^{(1)}[[\xi, \eta]_{\text{SD}}^{(1)}] = -2 \int d^3x \partial^j \pi_{ji} h^{ik} (\xi_P^\perp \theta^\perp{}_k - \eta_P^\perp \omega^\perp{}_k), \quad (\text{A.20})$$

while

$$\mathcal{H}_i^{(2)} = -2\partial_j \left( \pi^{jk} h_{ik} \right) + \pi^{jk} \partial_i h_{jk} \quad (\text{A.21})$$

$$\begin{aligned} \mathcal{H}_\perp^{(2)} = & \pi^{ij} \pi_{ij} - \frac{1}{2} \pi^2 \\ & + \frac{1}{4} \partial_k h_{ij} \partial^k h^{ij} - \frac{1}{2} \partial_k h^{ki} \partial^j h_{ij} + \frac{1}{2} \partial_i h \partial_j h^{ij} - \frac{1}{4} \partial_i h \partial^i h \\ & + \partial_l \left( \frac{1}{2} h \partial^l h - h^{ij} \partial^l h_{ij} - \frac{1}{2} h \partial_i h^{il} - h^{il} \partial_i h + \frac{3}{2} h^{lj} \partial^i h_{ij} + \frac{1}{2} h_{ij} \partial^i h^{jl} \right). \end{aligned} \quad (\text{A.22})$$

Isolating terms proportional to the constraints, we find

$$H^{(2)}[\xi] = \int d^3x \left( \mathcal{H}_m h^{mi} \xi_i + \frac{1}{2} \mathcal{H} h \xi^\perp \right) + \bar{H}^{(2)}[\xi], \quad (\text{A.23})$$

$$\bar{\mathcal{H}}_i^{(2)} = -\pi^{jk} (\partial_j h_{ki} + \partial_k h_{ji} - \partial_i h_{jk}), \quad (\text{A.24})$$

$$\begin{aligned} \bar{\mathcal{H}}_\perp^{(2)} = & \pi^{ij} \pi_{ij} - \frac{1}{2} \pi^2 \\ & + \frac{1}{4} \partial_k h_{ij} \partial^k h^{ij} - \frac{1}{2} \partial_k h^{ki} \partial^j h_{ij} + \frac{1}{4} \partial_i h \partial^i h \\ & + \partial_l \left( -h^{ij} \partial^l h_{ij} - h^{il} \partial_i h + \frac{3}{2} h^{lj} \partial^i h_{ij} + \frac{1}{2} h_{ij} \partial^i h^{jl} \right), \end{aligned} \quad (\text{A.25})$$

with  $\bar{H}^{(2)}[\xi] = \int d^3x \left( \bar{\mathcal{H}}_i^{(2)} \xi^i + \bar{\mathcal{H}}_\perp^{(2)} \xi^\perp \right)$ . On account of (A.17) and the analog of (A.5) for  $H^{(1)}[f]$ , it follows that

$$\left\{ \bar{H}^{(2)}[\xi_P], \bar{H}^{(2)}[\eta_P] \right\} \approx \bar{H}^{(2)}[[\xi_P, \eta_P]_{\text{SD}}^{(0)}], \quad (\text{A.26})$$

where  $\approx$  means an equality up to terms proportional to the constraints  $\mathcal{H}_m, \mathcal{H}_\perp$  of Pauli-Fierz theory. Note that the functionals  $H^{(2)}[\xi_P]$  and  $\bar{H}^{(2)}[\xi_P]$  generate transformations of the canonical variables that are equivalent because they differ at most by a gauge transformation of the Pauli-Fierz theory when restricted to the constraint surface.

The generators for global Poincaré transformations of Pauli-Fierz theory can then be identified as

$$\begin{aligned} Q_G(\omega, a) &= \frac{1}{2} \omega_{\mu\nu} J_G^{\mu\nu} - a_\mu P_G^\mu = \bar{H}^{(2)}[\xi_P(\tilde{\omega}, \tilde{a})] \\ \tilde{\omega}_{\mu\nu} &= \omega_{\mu\nu}, \quad \tilde{a}_\perp = a_\perp, \quad \tilde{a}_i = a_i + \omega_{\perp i} x^0. \end{aligned} \quad (\text{A.27})$$

Indeed, differentiating (A.19) with respect to  $b_\perp$  gives

$$\{H, Q_G(\omega, a)\} = \frac{\partial}{\partial t} Q_G(\omega, a) + 2 \int d^3x \partial^j \pi_{ji} h^{ik} \omega_{\perp k}. \quad (\text{A.28})$$

When combined with (A.19) and (A.27), this shows that, on the constraint surface, the generators  $Q_G(\omega, a)$  are conserved and satisfy the Poincaré algebra.

Finally, we can further simplify the explicit expression for  $\bar{H}^{(2)}[\xi_P]$  by using linearity of  $\xi_P$  in  $x^i$  and integrations by parts to show that

$$\begin{aligned} \int d^3x \bar{\mathcal{H}}_\perp^{(2)} \xi_P^\perp = & \int d^3x \left[ \pi^{ij} \pi_{ij} - \frac{1}{2} \pi^2 + \frac{1}{4} \partial_k h_{ij} \partial^k h^{ij} - \frac{1}{2} \partial_k h^{ki} \partial^j h_{ij} \right. \\ & \left. + \frac{1}{4} \partial_i h \partial^i h + \partial_l (h \partial_i h^{il} + h^{lj} \partial^i h_{ij}) \right] \xi_P^\perp. \end{aligned} \quad (\text{A.29})$$

The expansion of the gauge transformations (A.9), (A.11) gives to first order:

$$\delta_\xi^{(0)} h_{ij} = \partial_i \xi_j + \partial_j \xi_i, \quad \delta_\xi^{(0)} \pi^{ij} = (\partial^i \partial^j - \delta^{ij} \Delta) \xi^\perp, \quad (\text{A.30})$$

$$\delta_\xi^{(1)} h_{ij} = \xi^k \partial_k h_{ij} + \partial_i \xi^k h_{kj} + \partial_j \xi^k h_{ik} + 2\pi_{ij} \xi^\perp - \delta_{ij} \pi \xi^\perp, \quad (\text{A.31})$$

$$\begin{aligned}
 \delta_\xi^{(1)} \pi^{ij} &= \frac{1}{2} h (\partial^i \partial^j - \delta^{ij} \Delta) \xi^\perp - h^{im} \partial_m \partial^j \xi^\perp - h^{jm} \partial_m \partial^i \xi^\perp + h^{ij} \Delta \xi^\perp \\
 &+ \delta^{ij} h^{mn} \partial_m \partial_n \xi^\perp + \partial_m (\pi^{ij} \xi^m) - \pi^{mj} \partial_m \xi^i - \pi^{mi} \partial_m \xi^j \\
 &+ \frac{1}{2} \partial_k \xi^\perp \left[ -\partial^j h^{ki} - \partial^i h^{kj} + \partial^k h^{ij} + \delta^{ij} (2\partial_l h^{kl} - \partial^k h) \right] \\
 &+ \frac{1}{2} \xi^\perp \left[ \partial^i \partial^j h + \Delta h^{ij} - \partial_k \partial^i h^{jk} - \partial_k \partial^j h^{ik} - \delta^{ij} (\Delta h - \partial_k \partial_l h^{kl}) \right]. \quad (\text{A.32})
 \end{aligned}$$

## B. Riemann tensor and canonical variables

By following [1, 11] (up to conventions), we show in this appendix that the duality rotations that we have defined coincide on-shell with the standard simultaneous duality rotations among the (linearized) Riemann tensor and its dual together with those for the electric and magnetic conserved sources. We do this by showing how the covariant Riemann tensor is expressed in terms of the canonical variables. This gives us the appropriate generalization of the Gauss-Codazzi relations in the case of both electric and magnetic sources.

### B.1 Covariant equations in the presence of magnetic sources

Our conventions are as follows. Define  $\epsilon_{a_1 \dots a_n} = \epsilon^{a_1 \dots a_n}$  to be totally skew-symmetric with  $\epsilon_{1 \dots n} = 1$ . The Levi-Civita tensor is  $\epsilon_{a_1 \dots a_n} = \sqrt{|g|} \epsilon_{a_1 \dots a_n}$ . Indices on this tensor are raised with the metric, which implies that  $\epsilon^{a_1 \dots a_n} = \frac{(-)^\sigma}{\sqrt{|g|}} \epsilon_{a_1 \dots a_n}$  where  $\sigma$  is the signature of the metric. Our convention for the dual is  ${}^* \omega_{a_1 \dots a_{n-p}} = \frac{1}{p!} \omega^{b_1 \dots b_p} \epsilon_{b_1 \dots b_p a_1 \dots a_{n-p}}$ .

In flat Minkowski spacetime, traces are taken with the flat Minkowski metric  $\eta_{\mu\nu} = \text{diag}(-1, 1, 1, 1)$ . Start with the “(linearized) Riemann tensor”  $R_{\mu\nu\rho\sigma}^1 \equiv R_{\mu\nu\rho\sigma}$ , with only symmetry properties skew-symmetry in the first and last pairs of indices,

$$R_{\mu\nu\rho\sigma} = -R_{\nu\mu\rho\sigma} = -R_{\mu\nu\sigma\rho}. \quad (\text{B.1})$$

Its double dual (cf. MTW [38])

$${}^{\mathcal{G}}_{\mu\nu\rho\sigma}^1 \equiv {}^{\mathcal{G}}_{\mu\nu\rho\sigma} = \frac{1}{4} \epsilon_{\mu\nu}{}^{\alpha\beta} R_{\alpha\beta}{}^{\gamma\delta} \epsilon_{\gamma\delta\rho\sigma}, \quad (\text{B.2})$$

has the same symmetry properties. If ordinary duals of  $R_{\mu\nu\rho\sigma}$  and  ${}^{\mathcal{G}}_{\mu\nu\rho\sigma}$  are taken with respect to the last pair of indices, and we define

$$R_{\mu\nu\rho\sigma}^2 \equiv -{}^* R_{\mu\nu\rho\sigma} = -\frac{1}{2} R_{\mu\nu}{}^{\alpha\beta} \epsilon_{\alpha\beta\rho\sigma}, \quad (\text{B.3})$$

$${}^{\mathcal{G}}_{\mu\nu\rho\sigma}^2 \equiv -{}^* {}^{\mathcal{G}}_{\mu\nu\rho\sigma} = \frac{1}{2} \epsilon_{\mu\nu}{}^{\alpha\beta} R_{\alpha\beta\rho\sigma}, \quad (\text{B.4})$$

we have

$$R_{\mu\nu\rho\sigma}^a = \epsilon^{ab} ({}^* R)_{b\mu\nu\rho\sigma}, \quad {}^{\mathcal{G}}_{\mu\nu\rho\sigma}^a = \epsilon^{ab} {}^* {}^{\mathcal{G}}_{b\mu\nu\rho\sigma}. \quad (\text{B.5})$$

The 36 independent components of the Riemann tensor can be encoded in

$$R_{0m0n}^a, \quad {}^{\mathcal{G}}_{0m0n}^a. \quad (\text{B.6})$$

The Ricci and Einstein tensors are defined as

$$R_{\mu\nu}^a = R^{a\alpha}{}_{\mu\alpha\nu}, \quad G_{\mu\nu}^a = G^{a\alpha}{}_{\mu\alpha\nu} = R_{\nu\mu}^a - \frac{1}{2}\eta_{\mu\nu}R^a. \quad (\text{B.7})$$

If electric and magnetic conserved sources are  $T_a^{\mu\nu} \equiv (T^{\mu\nu}, \Theta^{\mu\nu})$ , with  $T_a^{\mu\nu} = T_a^{\nu\mu}$  symmetric,  $\partial_\mu T_a^{\mu\nu} = 0$ , the duality rotations are defined by

$$\begin{aligned} R_{\mu\nu\rho\sigma}^{a'} &= M^a{}_b R_{\mu\nu\rho\sigma}^b, & T_{\mu\nu}^{a'} &= M^a{}_b T_{\mu\nu}^b, \\ M_{ca}M^c{}_b &= \delta_{ab}. \end{aligned} \quad (\text{B.8})$$

For a tensor  $K^{\mu\nu}$ , let  $\bar{K}^{\mu\nu} = K^{\mu\nu} - \frac{1}{2}\eta^{\mu\nu}K$ .

It has been shown in [11] that the duality invariant equations of motion are

$$G_a^{\mu\nu} = 8\pi G T_a^{\mu\nu} \iff R_{\mu\nu\rho\sigma}^a + R_{\mu\sigma\nu\rho}^a + R_{\mu\rho\sigma\nu}^a = 8\pi G \epsilon^{ab} \epsilon_{\delta\nu\rho\sigma} \bar{T}_{b\mu}^\delta. \quad (\text{B.9})$$

They imply in particular that, on-shell, the tensors  $R_{\mu\nu\rho\sigma}^a$  are symmetric in the exchange of pairs of indices and that  $R_{\mu\nu}^a, G_{\mu\nu}^a$  are symmetric. Furthermore, the Bianchi “identities” read

$$\begin{aligned} \partial_\epsilon R_{\gamma\delta\alpha\beta}^a + \partial_\beta R_{\gamma\delta\epsilon\alpha}^a + \partial_\alpha R_{\gamma\delta\beta\epsilon}^a &= 8\pi G \epsilon^{ab} \epsilon_{\epsilon\alpha\beta\rho} (\partial_\gamma \bar{T}_{b\delta}^\rho - \partial_\delta \bar{T}_{b\gamma}^\rho) \\ &\iff \partial_\mu R_a^{\gamma\delta\rho\mu} = 8\pi G (\partial^\delta \bar{T}_a^{\rho\gamma} - \partial^\gamma \bar{T}_a^{\rho\delta}), \end{aligned} \quad (\text{B.10})$$

while the contracted Bianchi identities are

$$\partial_\nu G_a^{\mu\nu} = 0. \quad (\text{B.11})$$

Let

$$\begin{aligned} K^{\lambda\tau}{}_{\mu\nu\rho\sigma} [R_{\lambda\tau}^a] &= \frac{1}{2} [\eta_{\mu\rho} R_{\nu\sigma}^a + \eta_{\nu\sigma} R_{\mu\rho}^a - \eta_{\mu\sigma} R_{\nu\rho}^a - \eta_{\nu\rho} R_{\mu\sigma}^a] - \\ &\quad - \frac{R^a}{6} [\eta_{\mu\rho} \eta_{\nu\sigma} - \eta_{\mu\sigma} \eta_{\nu\rho}]. \end{aligned} \quad (\text{B.12})$$

Defining

$$\tilde{R}_{\mu\nu\rho\sigma}^a = R_{\mu\nu\rho\sigma}^a - \frac{1}{2}\epsilon^{ab} \epsilon_{\rho\sigma\alpha\beta} K^{\lambda\tau}{}_{\mu\nu}{}^{\alpha\beta} [R_{b\lambda\tau}^a], \quad (\text{B.13})$$

the tensor  $\tilde{R}_{\mu\nu\rho\sigma}^a$  is skew in the first and last pairs of indices, satisfies the cyclic identity because  $\epsilon^{\gamma\nu\rho\sigma} R_{\mu\nu\rho\sigma}^a = \epsilon^{\gamma\nu\rho\sigma} \frac{1}{2}\epsilon^{ab} \epsilon_{\rho\sigma\alpha\beta} K^{\lambda\tau}{}_{\mu\nu}{}^{\alpha\beta} [R_{b\lambda\tau}^a]$  and, as a consequence, is also symmetric in the exchange of the first and last pair of indices,  $\tilde{R}_{\mu\nu\rho\sigma}^a = \tilde{R}_{\rho\sigma\mu\nu}^a$ . The associated Ricci tensors  $\tilde{R}_{\nu\sigma}^a = R_{\nu\sigma}^a - \frac{1}{2}\epsilon^{ab} \epsilon_{\nu\sigma\mu\alpha} R_b^{\mu\alpha}$  is then symmetric,  $\tilde{R}_{\nu\sigma}^a = \tilde{R}_{\sigma\nu}^a$ . It follows that  $\tilde{R}_{\nu\sigma}^a = R_{(\nu\sigma)}^a$  and  $R_{[\nu\sigma]}^a = \frac{1}{2}\epsilon^{ab} \epsilon_{\nu\sigma\mu\alpha} R_b^{\mu\alpha}$ . The Weyl tensors are then defined as usual in terms of  $\tilde{R}_{\mu\nu\rho\sigma}^a$ ,

$$C_{\mu\nu\rho\sigma}^a = \tilde{R}_{\mu\nu\rho\sigma}^a - K^{\lambda\tau}{}_{\mu\nu\rho\sigma} [\tilde{R}_{\lambda\tau}^a], \quad (\text{B.14})$$

and satisfy all standard symmetry properties: skew-symmetry in the first and last pairs of indices, tracelessness (because  $\tilde{R}_{\nu\sigma}^a = K^{\lambda\tau}{}_{\nu\mu\sigma} [\tilde{R}_{\lambda\tau}^a]$ ), the cyclic identity (because



$\epsilon^{\gamma\nu\rho\sigma} K^{\lambda\tau}{}_{\mu\nu\rho\sigma} [\tilde{R}^a_{\lambda\tau}] = 0$ ), which implies also symmetry in the exchange of the first and last pair of indices,

$$C^a_{\mu\nu\rho\sigma} = -C^a_{\nu\mu\rho\sigma} = -C^a_{\mu\nu\sigma\rho}, \quad (\text{B.15})$$

$$C^{\mu a}_{\nu\mu\sigma} = 0, \quad \epsilon^{\gamma\nu\rho\sigma} C^a_{\mu\nu\rho\sigma} = 0, \quad C^a_{\mu\nu\rho\sigma} = C^a_{\rho\sigma\mu\nu}. \quad (\text{B.16})$$

As usual, the 10 independent components of the Weyl tensor can be parametrized by the electric and magnetic components  $E^a_{mn} \equiv (E_{mn}, B_{mn})$ , symmetric and traceless tensors defined by

$$E^a_{mn} = C^a_{0m0n} = \frac{1}{2} \epsilon_{nj k} \epsilon^{ab} C^j{}_{b0m}. \quad (\text{B.17})$$

Putting all definitions together, the relation between the Riemann and Weyl tensors is

$$R^a_{\mu\nu\rho\sigma} = C^a_{\mu\nu\rho\sigma} + K^{\lambda\tau}{}_{\mu\nu\rho\sigma} [R^a_{\lambda\tau}] + \frac{1}{2} \epsilon^{ab} \epsilon_{\rho\sigma\alpha\beta} K^{\lambda\tau}{}_{\mu\nu}{}^{\alpha\beta} [R_{b(\lambda\tau)}] \quad (\text{B.18})$$

$$= C^a_{\mu\nu\rho\sigma} + K^{\lambda\tau}{}_{\mu\nu\rho\sigma} [R^a_{(\lambda\tau)}] + \frac{1}{2} \epsilon^{ab} \epsilon_{\rho\sigma\alpha\beta} K^{\lambda\tau}{}_{\mu\nu}{}^{\alpha\beta} [R_{b\lambda\tau}]. \quad (\text{B.19})$$

In particular, it follows that the 36 independent components of the Riemann tensor  $R^1_{\mu\nu\rho\sigma}$  can be parameterized by the 10 independent components of the Weyl tensor  $C^1_{\mu\nu\rho\sigma}$ , the 16 components of the Ricci tensor  $R^1_{\lambda\tau}$ , and the 10 components of  $R^2_{(\lambda\tau)}$ .

If we define

$$\mathcal{E}^a_{mn} = R^a_{0(m|0|n)}, \quad \mathcal{F}^{am} = \frac{1}{2} \epsilon^{mj k} R^a_{0[j|0|k]}, \quad \mathcal{R}^a_{mn} = R^a_{(mn)} + \mathcal{E}^a_{mn} \quad (\text{B.20})$$

the parameterization consisting in choosing the symmetric tensors  $\mathcal{E}^a_{mn}, \mathcal{R}^a_{mn}$  (24 components),  $\mathcal{F}^a_m$ , (6 components), and  $R^1_{[\mu\nu]} (= {}^*R^2_{[\mu\nu]})$  (6 components) is more useful for our purpose. That all tensors can be reconstructed from these variables follows from the fact that

$$R^a_{0m} = -2\epsilon^{ab} \mathcal{F}_{bm}, \quad R^a_{00} = \mathcal{E}^a, \quad R^a_{(mn)} = \mathcal{R}^a_{mn} - \mathcal{E}^a_{mn}. \quad (\text{B.21})$$

This means that the symmetric part of the Ricci tensors can be reconstructed from the variables. Since the antisymmetric parts belong to the variables, so can the complete Ricci tensors  $R^a_{\mu\nu}$ . Using now (B.18) and definitions (B.17), (B.20), (B.12), we find

$$E^a_{mn} = \frac{1}{2} (\mathcal{E}^a_{mn} + \mathcal{R}^a_{mn}) - \frac{\delta_{mn}}{6} (\mathcal{E}^a + \mathcal{R}^a). \quad (\text{B.22})$$

It follows that the Weyl tensors and then, using again (B.18), the Riemann tensors can be reconstructed.

In terms of the new parameterization, the equations of motion (B.9) read  $R^a_{[\mu\nu]} = 0$  and

$$-2\epsilon^{ab} \mathcal{F}_{bm} = 8\pi G T^a_{0m}, \quad (\text{B.23})$$

$$\frac{1}{2} \mathcal{R}^a = 8\pi G T^a_{00}, \quad (\text{B.24})$$

$$\mathcal{R}^a_{mn} - \mathcal{E}^a_{mn} + \delta_{mn} \left( \mathcal{E}^a - \frac{1}{2} \mathcal{R}^a \right) = 8\pi G T^a_{mn}. \quad (\text{B.25})$$

Using these equations of motion, the Bianchi identities (B.10) are equivalent to

$$\partial^k(\epsilon_{ikm}\mathcal{F}^{am} + \mathcal{R}_{ik}^a) = \frac{1}{2}\partial_i\mathcal{R}^a, \quad (\text{B.26})$$

$$2\epsilon^{ab}\partial_0\mathcal{F}_{bm} = \partial^n(\mathcal{E}_{mn}^a + \epsilon_{mnk}\mathcal{F}^{ak}) - \partial_m\mathcal{E}^a, \quad (\text{B.27})$$

$$\begin{aligned} \partial_0\mathcal{R}_{ik}^a &= \frac{1}{2}\epsilon^{ab}[\epsilon_{kjl}\partial^j\mathcal{E}_{bi}^l + \epsilon_{ijl}\partial^j\mathcal{E}_{bk}^l - 2\delta_{ik}\partial_j\mathcal{F}_b^j - \partial_i\mathcal{F}_{bk} - \partial_k\mathcal{F}_{bi}] \iff \\ \epsilon^{ab}\partial_0\left(\mathcal{R}_b^{ik} - \frac{1}{2}\delta^{ik}\mathcal{R}_b\right) &= -\frac{1}{2}[\epsilon^{klm}\partial_l\mathcal{E}_m^a + \epsilon^{ilm}\partial_l\mathcal{E}_m^k + 2\delta^{ik}\partial^j\mathcal{F}_j^a - \partial^i\mathcal{F}^{ak} - \partial^k\mathcal{F}^{ai}]. \end{aligned} \quad (\text{B.28})$$

## B.2 Canonical expressions

We will now express the Riemann tensor in terms of the canonical variables in such a way that the covariant equations (B.23)–(B.28) coincide with the Hamiltonian equations deriving from (4.2).

From the constraints with sources, we find

$$\mathcal{R}^a = \partial^m\partial^n h_{mn}^a - \Delta h^a = -\Delta^2 C^a, \quad (\text{B.29})$$

$$\mathcal{F}_m^a = \frac{1}{2}\Delta\partial^n H_{mn}^a. \quad (\text{B.30})$$

Assuming  $\Delta$  to be invertible, which we do in the rest of this appendix,  $\mathcal{R}^a$  and  $C^a$ , respectively  $\mathcal{F}_m^a$  and  $\partial^n H_{mn}^a$  determine each other. By taking the divergence, the Bianchi identity (B.26) implies that

$$\partial^m\partial^n\mathcal{R}_{mn}^a = -\frac{1}{2}\Delta^3 C^a.$$

Similarly, the Bianchi identity (B.27) implies in particular that  $\Delta\mathcal{E}^a - \partial^m\partial^n\mathcal{E}_{mn}^a = \epsilon^{ab}\partial_0\Delta\partial^m\partial^n H_{bmn}$ . When combined with (B.25), the equations of motion following from variation with respect to  $C^a$  read

$$\frac{1}{2}\Delta^3 C^a + \epsilon^{ab}\Delta\partial_0(\Delta H_b - \partial^m\partial^n H_{bmn}) + 2\Delta^2 n^a = \Delta\mathcal{E}^a - \partial^m\partial^n(\mathcal{R}_{mn}^a - \mathcal{E}_{mn}^a).$$

When combined with the previous relations, they imply that

$$\begin{aligned} \mathcal{E}^a &= -\frac{1}{2}\epsilon^{ab}\partial_0\Delta H_b + \Delta n^a, \\ \partial^m\partial^n\mathcal{E}_{mn}^a &= -\frac{1}{2}\epsilon^{ab}\partial_0\Delta(\Delta H_b - 2\partial^m\partial^n H_{bmn}) + \Delta^2 n^a. \end{aligned}$$

The rest of the Bianchi identities (B.26), (B.27) are taken into account by applying a curl. This gives  $\epsilon^{rsi}\partial_s\partial^k\mathcal{R}_{ik}^a = \frac{1}{2}\Delta(\Delta\partial^k H_k^{ar} - \partial^r\partial^m\partial^n H_{mn}^a)$  and  $\epsilon^{rsi}\partial_s\partial^k\mathcal{E}_{ik}^a = \epsilon^{rsi}2\epsilon^{ab}\partial_0\partial_s\mathcal{F}_{bi} - \partial^r\partial^k\mathcal{F}_k^a + \Delta\mathcal{F}^{ar}$ . Yet another curl gives  $\partial_k\partial^m\partial^n\mathcal{R}_{mn}^a - \Delta\partial^n\mathcal{R}_{kn}^a = \frac{1}{2}\epsilon_{klr}\partial^l\Delta^2\partial^n H_n^{ar}$  and  $\partial_k\partial^m\partial^n\mathcal{E}_{mn}^a - \Delta\partial^n\mathcal{E}_{kn}^a = 2\epsilon^{ab}\partial_0(\partial_k\partial^n\mathcal{F}_{bn} - \Delta\mathcal{F}_{bk}) + \epsilon_{klr}\partial^l\Delta\mathcal{F}^{ar}$ . Using the previous relations we then get

$$\begin{aligned} \partial^n\mathcal{R}_{kn}^a &= -\frac{1}{2}\partial_k\Delta^2 C^a - \frac{1}{2}\epsilon_{klr}\partial^l\Delta\partial^n H_n^{ar}, \\ \partial^n\mathcal{E}_{kn}^a &= \epsilon^{ab}\partial_0\Delta\left(-\frac{1}{2}\partial_k H_b + \partial^n H_{bkn}\right) + \partial_k\Delta n^a - \frac{1}{2}\epsilon_{klr}\partial^l\Delta\partial^n H_n^{ar}. \end{aligned}$$

The equations of motion following from variation with respect to  $A_m^a$  are then identically satisfied.

Defining  $\mathcal{D}_{mn}^a = \mathcal{R}_{mn}^a - \mathcal{E}_{mn}^a$  and using definition (2.12) of  $\mathcal{P}^{TT}$  combined with (B.25), the equations of motion following from variation with respect to  $H_{mn}^a$  read

$$\begin{aligned} \epsilon_{ab}\partial_0 \left[ 2 \left( \mathcal{P}^{TT} H^b \right)_{mn} + \partial_m \Delta A_n^b + \partial_n \Delta A_m^b + \frac{1}{2} (\delta_{mn} \Delta - \partial_m \partial_n) C^b \right] - \epsilon_{ab} \Delta (\partial_m n_n^b + \partial_n n_m^b) - \\ - 2 \Delta^2 H_{mn}^a + \delta_{mn} \Delta^2 H^a = -\epsilon_{mpq} \partial^p \mathcal{D}_{an}^q - \epsilon_{npq} \partial^p \mathcal{D}_{am}^q. \end{aligned} \quad (\text{B.31})$$

Taking into account definition (2.12) and previous relations, we can extract

$$\begin{aligned} -\Delta^{-1} (\mathcal{P}^{TT} \mathcal{D}^a)_{mn} = \frac{1}{2} \epsilon_{ab} \partial_0 \left[ 2 \left( \mathcal{P}^{TT} H^b \right)_{mn} - \epsilon_{mpq} \partial_n \partial^p \partial^r H_r^{bq} - \epsilon_{npq} \partial_m \partial^p \partial^r H_r^{bq} + \right. \\ \left. + \partial_m \Delta A_n^b + \partial_n \Delta A_m^b + \frac{1}{2} (\delta_{mn} \Delta - \partial_m \partial_n) C^b \right] - \epsilon_{ab} \Delta (\partial_m n_n^b + \partial_n n_m^b) - \\ - \Delta^2 H_{mn}^a + \frac{1}{2} \delta_{mn} \Delta^2 H^a. \end{aligned} \quad (\text{B.32})$$

In order to extract the remaining information from (B.31), we first apply  $\delta^{mn} \Delta - \partial^m \partial^n$  to get

$$\epsilon_{ab} \partial_0 \Delta C^b + 2 \partial^m \partial^n H_{am} = 0, \quad (\text{B.33})$$

and then a divergence  $\partial^m$  giving

$$\epsilon_{ab} \partial_0 (\Delta A_n^b - \epsilon_{npq} \partial^p \partial^k H_k^{bq}) = \epsilon_{ab} \Delta n_n^b + 2 \Delta \partial^k H_{an}^k - \frac{1}{2} \partial_n \Delta H^a - \partial_n \partial^k \partial^l H_{kl}^a. \quad (\text{B.34})$$

We can now inject the latter relations into (B.31) and use (2.14), (2.6) to get

$$\mathcal{D}_{mn}^{aTT} = -\epsilon^{ab} \partial_0 \Delta H_{bmn}^{TT} - (\mathcal{P}^{TT} H^a)_{mn}, \quad (\text{B.35})$$

$$\begin{aligned} \mathcal{D}_{mn}^a = -\epsilon^{ab} \partial_0 \Delta \left[ H_{bmn} - \frac{1}{2} \delta_{mn} H_b \right] - (\mathcal{P}^{TT} H^a)_{mn} - \partial_m \partial_n n^a - \\ - \frac{1}{4} (\delta_{mn} \Delta + \partial_m \partial_n) \Delta C^a. \end{aligned} \quad (\text{B.36})$$

Injecting into the second form of the last Bianchi identity (B.28) and using previous relations gives

$$\begin{aligned} \epsilon_{ab} \partial_0 \mathcal{R}_{ij}^b = -(\mathcal{O}\mathcal{R}_a)_{ij} + \Delta^2 H_{aij}^{TT} + \frac{1}{4} \Delta \partial_i \partial^k H_{akj} + \frac{1}{4} \Delta \partial_j \partial^k H_{aki} - \frac{1}{2} \partial_i \partial_j \partial^k \partial^l H_{akl} \\ - \frac{1}{2} \epsilon_{ab} \partial_0 \left[ \epsilon_{iqn} \partial^q \Delta H_j^{bn} + \epsilon_{jqn} \partial^q \Delta H_i^{bn} + \frac{1}{2} (\delta_{ij} \Delta + \partial_i \partial_j) \Delta C^b \right]. \end{aligned} \quad (\text{B.37})$$

Identifying the terms with time derivatives gives

$$\begin{aligned} \mathcal{R}_{ij}^a = -\frac{1}{2} \left[ \epsilon_{iqn} \partial^q \Delta H_j^{an} + \epsilon_{jqn} \partial^q \Delta H_i^{an} + \frac{1}{2} (\delta_{ij} \Delta + \partial_i \partial_j) \Delta C^a \right] \\ = \frac{1}{2} \left[ \partial_i \partial^k h_{kj}^a + \partial_j \partial^k h_{ki}^a - \partial_i \partial_j h^a - \Delta h_{ij}^a - \epsilon_{ikl} \partial^k \partial^p \partial_j H_p^{al} - \epsilon_{jkl} \partial^k \partial^p \partial_i H_p^{al} \right]. \end{aligned} \quad (\text{B.38})$$

The terms without time derivatives in (B.37) then cancel identically. Together with (B.36) this then finally gives

$$\begin{aligned}\mathcal{E}_a^{ij} &= \epsilon_{ab}\partial_0\Delta\left[H^{bij} - \frac{1}{2}\delta^{ij}H^b\right] + \partial^i\partial^j n_a - \frac{1}{2}\epsilon^{ikl}\partial_k\partial^j\partial^p H_{alp} - \frac{1}{2}\epsilon^{jkl}\partial_k\partial^i\partial^p H_{alp} \\ &= -\epsilon_{ab}\partial_0\left(\pi^{bij} - \frac{1}{2}\delta^{ij}\pi^b\right) + \partial^i\partial^j n_a - \frac{1}{2}\epsilon^{ikl}\partial_k\partial^j\partial^p H_{alp} - \frac{1}{2}\epsilon^{jkl}\partial_k\partial^i\partial^p H_{alp}.\end{aligned}\quad (\text{B.39})$$

### B.3 Riemann tensor for linearized Taub-NUT

Following for instance [39] section A<sub>1</sub>.2 and using a regularization in Fourier space, we find for the gravitational dyon at rest at the origin discussed in section 4.2,

$$\mathcal{R}_{ij}^a = GM^a\left[\frac{16\pi}{3}\delta_{ij}\delta^3(x) + \frac{\eta(r)}{r^3}\left(\delta_{ij} - \frac{3x_ix_j}{r^2}\right)\right],\quad (\text{B.40})$$

$$\mathcal{E}_{ij}^a = GM^a\left[\frac{4\pi}{3}\delta_{ij}\delta^3(x) + \frac{\eta(r)}{r^3}\left(\delta_{ij} - \frac{3x_ix_j}{r^2}\right)\right],\quad (\text{B.41})$$

where  $\eta(r)$  is a regularizing function that suppresses the divergence at the origin and is 1 away from the origin. We then find

$$R_{00}^a = GM^a 4\pi\delta^3(x),\quad R_{ij}^a = GM^a 4\pi\delta_{ij}\delta^3(x),\quad (\text{B.42})$$

$$E_{ij}^a = GM^a\frac{\eta(x)}{r^3}\left(\delta_{ij} - \frac{3x_ix_j}{r^2}\right),\quad (\text{B.43})$$

and all other components of  $R_{\mu\nu}^a$  vanishing. For the Riemann tensor, this implies

$$R_{0i0j}^a = GM^a\left[\frac{4\pi}{3}\delta_{ij}\delta^3(x) + \frac{\eta(x)}{r^3}\left(\delta_{ij} - \frac{3x_ix_j}{r^2}\right)\right],\quad (\text{B.44})$$

$$R_{0ijk}^a = -\epsilon^{ab}\epsilon_{jk}{}^l GM^a\left[\frac{4\pi}{3}\delta_{il}\delta^3(x) + \frac{\eta(x)}{r^3}\left(\delta_{il} - \frac{3x_ix_l}{r^2}\right)\right],\quad (\text{B.45})$$

with all other components obtained through the on-shell symmetries of the Riemann tensor. This is the usual Riemann tensor for the linearized Taub-NUT solution.

### References

- [1] M. Henneaux and C. Teitelboim, *Duality in linearized gravity*, *Phys. Rev.* **D 71** (2005) 024018 [[gr-qc/0408101](#)].
- [2] S. Deser and D. Seminara, *Duality invariance of all free bosonic and fermionic gauge fields*, *Phys. Lett.* **B 607** (2005) 317 [[hep-th/0411169](#)].
- [3] B. Julia, J. Levie and S. Ray, *Gravitational duality near de Sitter space*, *JHEP* **11** (2005) 025 [[hep-th/0507262](#)].
- [4] R.G. Leigh and A.C. Petkou, *Gravitational duality transformations on (A)dS<sub>4</sub>*, *JHEP* **11** (2007) 079 [[arXiv:0704.0531](#)].
- [5] S. Deser and D. Seminara, *Free spin 2 duality invariance cannot be extended to GR*, *Phys. Rev.* **D 71** (2005) 081502 [[hep-th/0503030](#)].

- [6] S. Deser and C. Teitelboim, *Duality transformations of abelian and nonabelian gauge fields*, *Phys. Rev. D* **13** (1976) 1592.
- [7] J.H. Schwarz and A. Sen, *Duality symmetric actions*, *Nucl. Phys. B* **411** (1994) 35 [[hep-th/9304154](#)].
- [8] S. Deser, A. Gomberoff, M. Henneaux and C. Teitelboim, *Duality, self-duality, sources and charge quantization in abelian N-form theories*, *Phys. Lett. B* **400** (1997) 80 [[hep-th/9702184](#)].
- [9] S. Deser, A. Gomberoff, M. Henneaux and C. Teitelboim, *p-brane dyons and electric-magnetic duality*, *Nucl. Phys. B* **520** (1998) 179 [[hep-th/9712189](#)].
- [10] G. Barnich and A. Gomberoff, *Dyons with potentials: duality and black hole thermodynamics*, *Phys. Rev. D* **78** (2008) 025025 [[arXiv:0705.0632](#)].
- [11] C.W. Bunster, S. Cnockaert, M. Henneaux and R. Portugues, *Monopoles for gravitation and for higher spin fields*, *Phys. Rev. D* **73** (2006) 105014 [[hep-th/0601222](#)].
- [12] T. Curtright, *Generalized gauge fields*, *Phys. Lett. B* **165** (1985) 304.
- [13] J.A. Nieto, *S-duality for linearized gravity*, *Phys. Lett. A* **262** (1999) 274 [[hep-th/9910049](#)].
- [14] C.M. Hull, *Duality in gravity and higher spin gauge fields*, *JHEP* **09** (2001) 027 [[hep-th/0107149](#)].
- [15] X. Bekaert and N. Boulanger, *Tensor gauge fields in arbitrary representations of  $GL(D,R)$ : duality and Poincaré lemma*, *Commun. Math. Phys.* **245** (2004) 27 [[hep-th/0208058](#)].
- [16] X. Bekaert and N. Boulanger, *Massless spin-two field S-duality*, *Class. and Quant. Grav.* **20** (2003) S417 [[hep-th/0212131](#)].
- [17] U. Ellwanger, *S-dual gravity in the axial gauge*, *Class. and Quant. Grav.* **24** (2007) 785 [[hep-th/0610206](#)].
- [18] A.J. Nurbagametov, *Duality-symmetric approach to general relativity and supergravity*, *SIGMA* **2** (2006) 020 [[hep-th/0602145](#)].
- [19] S. de Haro, *Dual gravitons in  $AdS_4/CFT_3$  and the holographic Cotton tensor*, [arXiv:0808.2054](#).
- [20] E.A. Bergshoeff, M. de Roo, S.F. Kerstan, A. Kleinschmidt and F. Riccioni, *Dual gravity and matter*, [arXiv:0803.1963](#).
- [21] N. Boulanger and O. Hohm, *Non-linear parent action and dual gravity*, *Phys. Rev. D* **78** (2008) 064027 [[arXiv:0806.2775](#)].
- [22] R. Argurio, F. Dehouck and L. Houart, *Supersymmetry and gravitational duality*, [arXiv:0810.4999](#).
- [23] R. Argurio, F. Dehouck and L. Houart, *Boosting Taub-NUT to a BPS NUT-wave*, [arXiv:0811.0538](#).
- [24] I. Bakas, *Duality in linearized gravity and holography*, [arXiv:0812.0152](#).
- [25] R. Arnowitt, S. Deser and C. Misner, *Gravitation, an introduction to current research*, chapter 7 “The dynamics of general relativity”, Wiley, New York U.S.A. (1962), pg. 227
- [26] S. Deser, *Covariant decomposition of symmetric tensors and the gravitational Cauchy problem*, section A, *Ann. Inst. Henri Poincaré* **7** (1967) 149.

- [27] M. Henneaux, C. Teitelboim and J. Zanelli, *Gauge invariance and degree of freedom count*, *Nucl. Phys. B* **332** (1990) 169.
- [28] M. Henneaux and C. Teitelboim, *Quantization of gauge systems*, Princeton University Press, Princeton U.S.A. (1992).
- [29] G. Barnich and F. Brandt, *Covariant theory of asymptotic symmetries, conservation laws and central charges*, *Nucl. Phys. B* **633** (2002) 3 [[hep-th/0111246](#)].
- [30] G. Barnich and G. Compere, *Surface charge algebra in gauge theories and thermodynamic integrability*, *J. Math. Phys.* **49** (2008) 042901 [[arXiv:0708.2378](#)].
- [31] T. Regge and C. Teitelboim, *Role of surface integrals in the Hamiltonian formulation of general relativity*, *Ann. Phys. (NY)* **88** (1974) 286.
- [32] R. Benguria, P. Cordero and C. Teitelboim, *Aspects of the Hamiltonian dynamics of interacting gravitational gauge and Higgs fields with applications to spherical symmetry*, *Nucl. Phys. B* **122** (1977) 61.
- [33] V. Iyer and R.M. Wald, *Some properties of Nöther charge and a proposal for dynamical black hole entropy*, *Phys. Rev. D* **50** (1994) 846 [[gr-qc/9403028](#)].
- [34] G. Barnich, *Boundary charges in gauge theories: using Stokes theorem in the bulk*, *Class. and Quant. Grav.* **20** (2003) 3685 [[hep-th/0301039](#)].
- [35] C. Teitelboim, *How commutators of constraints reflect the space-time structure*, *Ann. Phys. (NY)* **79** (1973) 542.
- [36] S.A. Hojman, K. Kuchar and C. Teitelboim, *Geometrodynamics regained*, *Ann. Phys. (NY)* **96** (1976) 88.
- [37] J.D. Brown and M. Henneaux, *Central charges in the canonical realization of asymptotic symmetries: an example from three-dimensional gravity*, *Commun. Math. Phys.* **104** (1986) 207.
- [38] C. Misner, K. Thorne and J. Wheeler, *Gravitation*, W.H. Freeman, New York U.S.A. (1973).
- [39] C. Cohen-Tannoudji, J. Dupont-Roc and G. Grynberg, *Photons and atoms: introduction to quantum electrodynamics*, Wiley-Interscience, New York U.S.A. (1989).