

$\mathcal{N} = 8$ superconformal gauge theories and $M2$ branes

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

JHEP01(2009)078

(<http://iopscience.iop.org/1126-6708/2009/01/078>)

[The Table of Contents](#) and [more related content](#) is available

Download details:

IP Address: 80.92.225.132

The article was downloaded on 03/04/2010 at 11:34

Please note that [terms and conditions apply](#).

$\mathcal{N} = 8$ superconformal gauge theories and $M2$ branes

Sergio Benvenuti,^a Diego Rodríguez-Gómez,^{ab} Erik Tonni^c and Herman Verlinde^a

^a*Department of Physics, Princeton University,
Princeton, NJ 08544, U.S.A.*

^b*Center for Research in String Theory, Queen Mary University of London,
Mile End Road, London, E1 4NS, U.K.*

^c*Dipartimento di Fisica, Università di Pisa and INFN, sezione di Pisa,
Largo Bruno Pontecorvo 3, 56127 Pisa, Italy*

*E-mail: sbenvenu@princeton.edu, drodrigu@princeton.edu, tonni@df.unipi.it,
verlinde@princeton.edu*

ABSTRACT: Based on recent developments, in this letter we find $2 + 1$ dimensional gauge theories with scale invariance and $\mathcal{N} = 8$ supersymmetry. The gauge theories are defined by a Lagrangian and are based on an infinite set of 3-algebras, constructed as an extension of ordinary Lie algebras. Recent no-go theorems on the existence of 3-algebras are circumvented by relaxing the assumption that the invariant metric is positive definite. The gauge group is non compact, and its maximally compact subgroup can be chosen to be any ordinary Lie group, under which the matter fields are adjoints or singlets. Interestingly, the theories are parity invariant and do not admit any tunable coupling constant.

KEYWORDS: Brane Dynamics in Gauge Theories, Conformal Field Models in String Theory, M-Theory.

Contents

1.	Introduction	1
2.	New $\mathcal{N} = 8$ superconformal gauge theories	3
2.1	Mini-review of the Bagger-Lambert framework	3
2.2	The special case of the 3-algebra (1.1)	4
2.3	The Lagrangian	6
2.4	Absence of coupling constant	7
2.5	Comments on the physical spectrum	7
2.6	Moduli space of vacua for $\mathcal{G} = \text{SU}(N)$	8
2.6.1	Massive excitations on the abelian branch	10
3.	Comments	11
A.	Check of the fundamental identity	12

1. Introduction

In a series of papers by Bagger and Lambert [1–3] and Gustavsson [4, 5], a new set of 2+1 dimensional theories enjoying $\mathcal{N} = 8$ supersymmetry, $\text{SO}(8)$ global symmetry and scale invariance has been proposed. These theories are supposed to describe multiple coincident M2 branes; a Chern-Simons term and a sextic potential for the scalars are present in the Lagrangian, as expected [6]. This framework involves an unusual algebraic structure called 3-algebra: a vector space endowed with a set of structure constants with four indices $f^{abc}{}_d$. The structure constants satisfy antisymmetry in the upper three indices and a fundamental identity analogue to the Jacobi identity satisfied by the structure constants $f^{\alpha\beta}{}_{\gamma}$ of ordinary Lie algebras.

The requirement that the 3-algebra has a positive definite metric is very strong: it was recently proven in [7, 8] that there is only one such 3-algebra, called \mathcal{A}_4 (or linear sums thereof). \mathcal{A}_4 is 4-dimensional and $f^{abcd} = \epsilon^{abcd}$. The corresponding $\mathcal{N} = 8$ theory has gauge symmetry $\text{SO}(4)$ and is expected to describe two M2 branes sitting at the origin of $\mathbb{R}^8/\mathbb{Z}_2$ [9–11]. It was shown in [12] that, upon giving a VEV to a scalar, it is possible to recover the theory on coincident D2 branes via a novel type of Higgs mechanism. For additional results see [13–21].

The problem of describing N M2 branes in flat space is the main motivation for our study. It seems reasonable to relax some of the constraints in order to evade the no-go theorems, and maybe describe N M2's. This direction has been pursued in [15, 16]. In [16] Gran, Nilsson and Petersson proposed to focus only on the equations of motion, relying

on the fact that many 3-algebras exist if we don't require the existence of a metric. It's important to remark that in the Bagger-Lambert work at the level of the equation of motion the metric is not used. The metric is needed in order to have a Lagrangian and gauge invariant local operators, such as the energy-momentum tensor.

In this letter we study the structure of the theory that arises if one relaxes the assumption that the metric on the 3-algebra is positive definite. This allows us to find an infinite set of 3-algebras. The construction starts from any ordinary Lie algebra \mathcal{G} and consists in adding two directions to \mathcal{G} as a vector space, which we call $+$ and $-$, thus the 3-algebra has dimension $\dim(\mathcal{G}) + 2$. Using indices $a, b, \dots = \{+, -, \alpha\}$, the structure constants are given in terms of the \mathcal{G} -structure constants $f^{\alpha\beta}{}_{\gamma}$ as

$$f^{+\alpha\beta}{}_{\gamma} = -f^{\alpha+\beta}{}_{\gamma} = f^{\alpha\beta+}{}_{\gamma} = f^{\alpha\beta}{}_{\gamma} \qquad f^{\alpha\beta\gamma}{}_{-} = f^{\alpha\beta\gamma} \qquad (1.1)$$

all other possible components of $f^{abc}{}_{d}$ simply vanish. This 3-algebra is related to a similar algebra proposed in [5, 16, 22].¹ One nice feature of (1.1) is the existence of an invariant metric, given in terms of the standard metric on \mathcal{G}

$$\left(\begin{array}{cc|ccc} 0 & -1 & 0 & \dots & 0 \\ -1 & 0 & 0 & \dots & 0 \\ \hline 0 & 0 & & & \\ \vdots & \vdots & & h_{\mathcal{G}} & \\ 0 & 0 & & & \end{array} \right) \qquad (1.2)$$

This metric is clearly non positive definite, having signature $(\dim(\mathcal{G}) + 1, 1)$ if \mathcal{G} is compact and semisimple. The metric is invariant under the symmetry group of the equations of motion, which turns out to be a Inonu-Wigner contraction of $\mathcal{G} \otimes \mathcal{G}$. Our theory has thus a non compact gauge group of dimension $2 \dim(\mathcal{G})$, that can be embedded in $SO(\dim(\mathcal{G}) + 1, 1)$. The scalar fields X^I and fermion fields Ψ transform in the 8 of $SO(8)$. Two fields (X^I_{\pm} and Ψ_{\pm}) are singlets of \mathcal{G} (the maximally compact subgroup of the full non compact gauge group), while the other fields $X^I_{\alpha}, \Psi_{\alpha}$ transform in the adjoint representation. Since the gauge theory discussed in this paper can be recast in the Bagger-Lambert framework, it automatically enjoys $\mathcal{N} = 8$ supersymmetry.

One new feature is that the gauge interactions turns out to be of the BF-type. BF theories do not admit a tunable coupling constant, and this property extends to the full superconformal theory. This is expected for the gauge theory living on M2 branes. Also parity invariance is preserved.

In the main body of this letter, section 2, we discuss in detail the classical aspects of the theory based on the 3-algebra (1.1), finding the gauge symmetries and an explicit Lagrangian. We show how the overall coupling in front of the Lagrangian can be reabsorbed via rescaling of the fields. We also comment on the structure of the moduli space of vacua and the mass of the low energy fluctuations. We conclude in section 3 with some speculative remarks about the quantization and the unitarity of the theory, and we comment on the possible relation to M theory.

¹More precisely if we project (1.1) along the $\dim(\mathcal{G}) + 1$ dimensional subspace generated by $\{+, \alpha\}$ we obtain the 3-algebra discussed in [5, 16, 22]. We can also project along the subspace generated by $\{-, \alpha\}$.

Note added: at the same time when this letter was first posted on the arXiv, two other preprints [23, 25] appeared with substantial overlap with our results. Since then, several new papers have appeared with results that clarify the interpretation of the class of theories, that we introduce here. We briefly comment on these developments at the end of this letter.

2. New $\mathcal{N} = 8$ superconformal gauge theories

2.1 Mini-review of the Bagger-Lambert framework

The theory on coincident M2 branes should involve 8 real scalar fields X_a^I , $I = 1, \dots, 8$. Then, $\mathcal{N} = 8$ supersymmetry requires a 16 component spinor Ψ_a^I , which we can take a chiral spinor of $SO(8)$. The fields carry an internal index a running from 1 to D , where D is the dimension of the 3-algebra. With these ingredients, Bagger and Lambert proposed the following $\mathcal{N} = 8$ SUSY transformations, consistent with classical scale invariance:

$$\begin{aligned} \delta X_a^I &= i\bar{\epsilon}\Gamma^I\Psi_a, \\ \delta\Psi_a &= D_\mu X_a^I\Gamma^\mu\Gamma_I\epsilon - \frac{1}{6}X_b^IX_c^JX_d^Kf^{bcd}{}_a\Gamma_{IJK}\epsilon, \\ \delta(\tilde{A}_\mu)_b^a &= i\bar{\epsilon}\Gamma_\mu\Gamma_I X_c^IX_d^J\Psi_d f^{cdab}; \end{aligned} \tag{2.1}$$

where $f^{abc}{}_d$ are the structure constants of the 3-algebra and are completely antisymmetric in the 3 upper indices. The closure of the SUSY transformations implies the equations of motion

$$\begin{aligned} \Gamma^\mu D_\mu\Psi_a + \frac{1}{2}\Gamma_{IJ}X_c^IX_d^J\Psi_b f^{cdb}{}_a &= 0, \\ D^2X_a^I - \frac{i}{2}\bar{\Psi}_c\Gamma_J^IX_d^J\Psi_b f^{cdb}{}_a + \frac{1}{2}f^{bcd}{}_af^{efg}{}_dX_b^JX_c^KX_e^IX_f^JX_g^K &= 0, \\ (\tilde{F}_{\mu\nu})_a^b + \epsilon_{\mu\nu\lambda}\left(X_c^JD^\lambda X_d^J + \frac{i}{2}\bar{\Psi}_c\Gamma^\lambda\Psi_d\right) f^{cdb}{}_a &= 0. \end{aligned} \tag{2.2}$$

The fields transform in the following way under gauge transformation

$$\delta X_a^I = \tilde{\Lambda}_a^b X_b^I, \quad \delta\Psi_a = \tilde{\Lambda}_a^b \Psi_b, \quad \delta(\tilde{A}_\mu)_a^b = D_\mu\tilde{\Lambda}_a^b. \tag{2.3}$$

where $\tilde{\Lambda}_b^a = \Lambda_{mn}f^{mna}{}_b$ and $(\tilde{A}_\mu)_b^a = (A_\mu)_{mn}f^{mna}{}_b$. The gauge group is generated by the $\tilde{\Lambda}_b^a$, while the antisymmetric Λ_{mn} are auxiliary parameters. The gauge group is thus a subgroup of $GL(D)$. (If we add a metric of signature $(D - k, k)$ on the 3-algebra, then we can say that the gauge group is a subgroup of $SO(D - k, k)$). In order for the equations of motion to be consistent with gauge symmetry and supersymmetry, a constraint on the structure constants has to be satisfied:

$$f^{efg}{}_df^{abc}{}_g - f^{efa}{}_gf^{bcg}{}_d - f^{efb}{}_gf^{cag}{}_d - f^{efc}{}_gf^{abg}{}_d = 0. \tag{2.4}$$

This is known as the fundamental identity. At the level of equations of motion the only constraints on the structure constants come from the fundamental identity and antisymmetry in the upper 3 indices.

2.2 The special case of the 3-algebra (1.1)

As a first step we determine the gauge group for the special choice of structure constants (1.1)

$$f^{+\alpha\beta}{}_{\gamma} = -f^{\alpha+\beta}{}_{\gamma} = f^{\alpha\beta+}{}_{\gamma} = f^{\alpha\beta}{}_{\gamma} \qquad f^{\alpha\beta\gamma}{}_{-} = f^{\alpha\beta\gamma} \qquad (2.5)$$

Since the fields X_{α} transform in the adjoint of \mathcal{G} we can introduce the matrices T^{α} such that

$$X^I = X_{\alpha}^I T^{\alpha} \qquad (2.6)$$

$$[T^{\alpha}, T^{\beta}] = f^{\alpha\beta}{}_{\gamma} T^{\gamma} \qquad (2.7)$$

$$\text{Tr}(T^{\alpha} T^{\beta}) = \delta^{\alpha\beta} \qquad (2.8)$$

We focus on the equations of motion (2.2) but, for the sake of simplicity, we set to zero the fermions and the gauge fields. (2.2) becomes

$$\begin{aligned} \partial^2 X_{+}^I &= 0, \\ D^2 X_{-}^I &= \frac{1}{2} X_{+}^I \text{Tr}([X^J, X^K]^2) - X_{+}^J \text{Tr}([X^J, X^K][X^I, X^K]) \\ D^2 X^I &= (X_{+}^J)^2 [X^K, [X^I, X^K]] - X_{+}^J X_{+}^I [X^K, [X^J, X^K]] - X_{+}^J X_{+}^K [X^K, [X^I, X^J]], \end{aligned} \qquad (2.9)$$

where for the moment the precise definition of the covariant derivative D^{μ} is not important. We now want to study the symmetries of (2.9) under global transformations of the fields (X_{\pm}^I, X^I) . The symmetry generators act on the matter field through the matrix

$$\tilde{\Lambda}_b^a = \Lambda_{mn} f^{mna}{}_b. \qquad (2.10)$$

From (1.1) it is easy to see that

$$\tilde{\Lambda}_{+}^{-} = \tilde{\Lambda}_{+}^{\alpha} = \tilde{\Lambda}_{-}^{+} = 0 \qquad (2.11)$$

while $\tilde{\Lambda}_{\beta}^{\alpha}$, $\tilde{\Lambda}_{\alpha}^{+}$ and $\tilde{\Lambda}_{-}^{\alpha}$ are non-zero. The $\tilde{\Lambda}_{\beta}^{\alpha}$ transformations are in one-to-one correspondence with the generators of the compact group \mathcal{G} . These are obviously symmetries of (2.9). The corresponding generators will be denoted by T_c^{α} (where the subscript c stands for ‘compact’).

An additional set of $\dim(\mathcal{G})$ symmetries are present, corresponding to $\tilde{\Lambda}_{\alpha}^{+}$ and $\tilde{\Lambda}_{-}^{\alpha}$, which come from $\Lambda_{\alpha\beta}$. These additional symmetries are non-compact, and we denote their generators by T_{nc}^{α} . They act on the fields as

$$\begin{aligned} \delta X_{+}^I &= 0 \\ \delta X_{-}^I &= \text{Tr}(M X^I) + \frac{1}{2} \text{Tr}(M^2) X_{+}^I \\ \delta X^I &= M X_{+}^I \end{aligned} \qquad (2.12)$$

Where M is a matrix in the adjoint of \mathcal{G} which is not necessarily infinitesimal. It is clear that at the infinitesimal level the transformations T_{nc}^{α} commute among themselves.

In order to find the full set of commutation relations it is useful to write the generators of the symmetry group as $(\dim(\mathcal{G}) + 2) \times (\dim(\mathcal{G}) + 2)$ matrices.

$$T_c^\alpha = \left(\begin{array}{c|ccc} 0 & 0 & \dots & 0 \\ 0 & 0 & 0 & \dots & 0 \\ \hline 0 & 0 & & & \\ \vdots & \vdots & & & T^\alpha \\ 0 & 0 & & & \end{array} \right) \qquad T_{nc}^\alpha = \left(\begin{array}{c|ccc} 0 & 0 & 0 & \dots & 0 & \dots & 0 \\ 0 & 0 & 0 & \dots & 1 & \dots & 0 \\ \hline 0 & 0 & & & & & \\ \vdots & \vdots & & & & & \\ \alpha \rightarrow 1 & 0 & & & 0 & & \\ \vdots & \vdots & & & & & \\ 0 & 0 & & & & & \end{array} \right) \quad (2.13)$$

It is straightforward to check that the algebra is

$$[T_c^\alpha, T_c^\beta] = f^{\alpha\beta}{}_\gamma T_c^\gamma, \quad [T_{nc}^\alpha, T_{nc}^\beta] = 0, \quad [T_{nc}^\alpha, T_c^\beta] = f^{\alpha\beta}{}_\gamma T_{nc}^\gamma. \quad (2.14)$$

To gain further insight we modify this algebra with a parameter ϵ

$$[T_c^\alpha, T_c^\beta] = f^{\alpha\beta}{}_\gamma T_c^\gamma, \quad [T_{nc}^\alpha, T_{nc}^\beta] = \epsilon f^{\alpha\beta}{}_\gamma T_c^\gamma, \quad [T_{nc}^\alpha, T_c^\beta] = f^{\alpha\beta}{}_\gamma T_{nc}^\gamma. \quad (2.15)$$

so that in the limit $\epsilon \rightarrow 0$ we recover (2.14). It is easy to see² that (2.15) is the Lie algebra of $\mathcal{G} \otimes \mathcal{G}$. The symmetry algebra of the ungauged theory is thus a non compact Inonu-Wigner contraction of $\mathcal{G} \times \mathcal{G}$. This analysis has been carried out just at the level of scalar fields, but it's easy to see that it extends to the full set of equation of motion. Notice that the theory classically have a shift symmetry acting on the 8 scalars X_-^I . The role of this symmetry is not clear.

The theory therefore has $2 \dim(\mathcal{G})$ gauge fields. We denote the fields associated the compact part \mathcal{G} and the non compact part by \mathcal{A}^μ and \mathcal{B}^μ , respectively.

$$(\mathcal{A}_\mu)_\alpha = (\mathcal{A}_\mu)_{+\alpha} \qquad (\mathcal{B}_\mu)_\alpha = (\mathcal{A}_\mu)_{\beta\gamma} f^{\beta\gamma}{}_\alpha \quad (2.16)$$

and $\mathcal{A}_\mu = (\mathcal{A}_\mu)_\alpha T^\alpha$, $\mathcal{B}_\mu = (\mathcal{B}_\mu)_\alpha T^\alpha$. So \mathcal{A}_μ and \mathcal{B}_μ are matrices in the adjoint of \mathcal{G} . The covariant derivatives are defined as

$$D_\mu X^I = \partial_\mu X^I - 2[\mathcal{A}_\mu, X^I] - \mathcal{B}_\mu X_+^I; \quad (2.17)$$

$$D_\mu X_-^I = \partial_\mu X_-^I - \text{Tr}(\mathcal{B}_\mu X^I); \quad (2.18)$$

$$D_\mu X_+^I = \partial_\mu X_+^I, \quad (2.19)$$

and similarly for the fermions. X^I , Ψ are fields in the adjoint of \mathcal{G} and X_\pm^I , Ψ_\pm are singlets. The gauge parameters can also be assembled in matrices of the adjoint of \mathcal{G} as

$$\Lambda = \Lambda_{+\alpha} T^\alpha \qquad M = \Lambda_{\beta\gamma} f^{\beta\gamma}{}_\alpha T^\alpha, \quad (2.20)$$

²We can now re-scale the generators and combine them into two sets $\{T_\pm^\alpha\}$ in such a way that the deformed algebra is $[T_+^\alpha, T_-^\beta] = 0$, $[T_+^\alpha, T_+^\beta] = f^{\alpha\beta}{}_\gamma T_+^\gamma$, $[T_-^\alpha, T_-^\beta] = f^{\alpha\beta}{}_\gamma T_-^\gamma$

under which the gauge fields transform as

$$\delta\mathcal{A}_\mu = \partial_\mu\Lambda - 2[\mathcal{A}_\mu, \Lambda] \tag{2.21}$$

$$\delta\mathcal{B}_\mu = \partial_\mu M - 2[\mathcal{A}_\mu, M] - 2[\mathcal{B}_\mu, \Lambda] \tag{2.22}$$

The equations of motion for the gauge fields, in absence of matter fields, are simply $(\tilde{F}_{\mu\nu})^b_a = 0$. The non trivial ones are

$$(\tilde{F}_{\mu\nu})^+_\alpha = 0 \qquad (\tilde{F}_{\mu\nu})^\beta_\alpha = 0, \tag{2.23}$$

After contracting with $\epsilon^{\lambda\mu\nu}$, in terms of \mathcal{A}_μ and \mathcal{B}_μ (2.23) imply

$$0 = \epsilon^{\lambda\mu\nu}(\partial_\mu\mathcal{B}_\nu - [\mathcal{A}_\mu, \mathcal{B}_\nu]) \tag{2.24}$$

$$0 = \epsilon^{\lambda\mu\nu}(\partial_\mu\mathcal{A}_\nu - [\mathcal{A}_\mu, \mathcal{A}_\nu]) \tag{2.25}$$

The action for the gauge fields \mathcal{A} and \mathcal{B} , from which the above equations can be derived, is of the BF -type:

$$S_{gauge} = \int d^3x \epsilon^{\lambda\mu\nu} \text{Tr}\left(\mathcal{B}_\lambda(\partial_\mu\mathcal{A}_\nu - [\mathcal{A}_\mu, \mathcal{A}_\nu])\right) \tag{2.26}$$

This action is parity invariant, if we define parity to act on the gauge field \mathcal{B}^μ by flipping its sign. Notice the no tunable coupling constant can appear in (2.26).

2.3 The Lagrangian

The metric associated to our choice of structure constants is

$$h^{ab} = \left(\begin{array}{cc|ccc} 0 & -1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & & & \\ 0 & 0 & & \mathcal{I}_G & \\ 0 & 0 & & & \end{array} \right)$$

It is easy to check that this metric is invariant under the transformations (2.13). Using this metric it is possible to write down a Lagrangian. As a first illustrative step we consider just the scalar part. The equations of motion (2.9) are derived from the following ungauged Lagrangian:

$$\mathcal{L}_S = -\frac{1}{2}\text{Tr}\left(\partial_\mu X^I \partial_\mu X^I\right) + \partial_\mu X^I_+ \partial_\mu X^I_- - \frac{1}{12}\text{Tr}\left(X^I_+[X^J, X^K] + X^J_+[X^K, X^I] + X^K_+[X^I, X^J]\right)^2 \tag{2.27}$$

It is important that this Lagrangian does not have a coupling constant. Indeed, the same is true for the gauge Lagrangian (2.26).³

³This 3D theory is of the form $\partial X_+ \partial X_- - (\partial X)^2 + X_+^2 X^4$, which is scale invariant and does not admit a tunable parameter but still seems non trivial. The analogous scale invariant Lagrangian in 4D, $\partial X_+ \partial X_- - (\partial X)^2 + X_+^2 X^2$, is instead much simpler, since the equations of motions are all linear if solved in the right order: first the one coming from X_- ($\partial^2 X_+ = 0$), then the one coming from X ($\partial^2 X = X_+^2 X$) and finally the equation coming from X_+ ($\partial^2 X_- = -2X_+ X^2$).

We finally consider the complete gauged Lagrangian, with $\mathcal{N} = 8$ SUSY, SO(8) global symmetry, from which the equations of motion follow.⁴ The full Lagrangian contains standard kinetic terms with covariant derivatives, the gauge term (2.26), a sextic potential and Yukawa couplings:

$$\begin{aligned} \mathcal{L} = & -\frac{1}{2}h^{ab}D_\mu X_a^I D_\mu X_b^I + \frac{i}{2}h^{ab}\bar{\Psi}_a\Gamma^\mu D_\mu\Psi_b + \epsilon^{\mu\nu\lambda}\text{Tr}\left(\mathcal{B}_\lambda(\partial_\mu\mathcal{A}_\nu - [\mathcal{A}_\mu, \mathcal{A}_\nu])\right) \\ & + \frac{1}{12}h^{mn}f^{abc}{}_m f^{efg}{}_n X_a^I X_b^J X_c^K X_e^I X_f^J X_g^K + \frac{i}{4}h^{de}f^{abc}{}_e X_a^I X_b^J \bar{\Psi}_c\Gamma_{IJ}\Psi_d \end{aligned} \quad (2.28)$$

In this form the invariance under the non compact gauge group is manifest, moreover from the results of Bagger and Lambert it is clear that (2.28) is $\mathcal{N} = 8$ supersymmetric.

We can rewrite the Lagrangian in a \mathcal{G} -invariant notation:

$$\begin{aligned} \mathcal{L} = & -\frac{1}{2}\text{Tr}\left(D_\mu X^I D_\mu X^I\right) + D_\mu X_+^I D_\mu X_-^I + \frac{i}{2}\text{Tr}\left(\bar{\Psi}\Gamma^\mu D_\mu\Psi\right) - \frac{i}{2}\bar{\Psi}_+\Gamma^\mu D_\mu\Psi_- \\ & - \frac{i}{2}\bar{\Psi}_-\Gamma^\mu D_\mu\Psi_+ + \epsilon^{\mu\nu\lambda}\text{Tr}\left(\mathcal{B}_\lambda(\partial_\mu\mathcal{A}_\nu - [\mathcal{A}_\mu, \mathcal{A}_\nu])\right) \\ & - \frac{1}{12}\text{Tr}\left(X_+^I[X^J, X^K] + X_+^J[X^K, X^I] + X_+^K[X^I, X^J]\right)^2 + \frac{i}{2}\text{Tr}\left(\bar{\Psi}\Gamma_{IJ}X_+^I[X^J, \Psi]\right) \\ & + \frac{i}{4}\text{Tr}\left(\bar{\Psi}\Gamma_{IJ}[X^I, X^J]\Psi_+\right) - \frac{i}{4}\text{Tr}\left(\bar{\Psi}_+\Gamma_{IJ}[X^I, X^J]\Psi\right), \end{aligned} \quad (2.29)$$

3-dimensional parity is preserved if the fields \mathcal{B}_μ , X^I and Ψ are parity-odd.

2.4 Absence of coupling constant

As for the previously studied truncations, the Lagrangian (2.29) does not admit any tunable coupling constant. Indeed, had we considered including a coupling constant $\frac{1}{g^2}\mathcal{L}(X_\pm, X, \mathcal{B}, \mathcal{A})$, we could redefine

$$X^I = gY^I \quad X_+^I = g^{-1}Y_+^I \quad X_-^I = g^3Y_-^I \quad \mathcal{B} = g^2\tilde{\mathcal{B}}, \quad (2.30)$$

(and consistently the same for the fermions) in such a way that we have

$$\frac{1}{g^2}\mathcal{L}(X_\pm, X, \mathcal{B}, \mathcal{A}) = \mathcal{L}(Y_\pm, Y, \tilde{\mathcal{B}}, \mathcal{A}), \quad (2.31)$$

i.e. the coupling constant can be always reabsorbed. This is a highly non-trivial hint that (2.29) is related to M2 branes.

2.5 Comments on the physical spectrum

Given that the metric in (2.29) is not positive definite, one could worry about the presence of negative norm states in the quantum theory. Since this problem is already present in the ungauged theory, we can start considering just (2.27). The field X_-^I appears only through $\partial X_+^I \partial X_-^I$, we could perform the functional integral over it, which would lead to

⁴Notice that in [2] the Lagrangian was derived by the eqs. of motion under the assumption that the metric is positive definite, so we cannot use directly their results. At the end however we get the same form of the Lagrangian, defining $f^{abcd} = h^{de}f^{abc}{}_e$.

a functional delta localizing the X_+^I integral on the solutions to $\partial^2 X_+^I = 0$ (this is the quantum-mechanical counterpart of the classical observation that we can regard X_-^I as a Lagrange multiplier, thus enforcing a constraint). We would be left with an effective theory whose partition function is

$$\mathcal{Z} = \int \mathcal{D}X_+^I \mathcal{D}X^I \delta(\partial^2 X_+^I) \exp\left\{i \int -\frac{1}{2} \text{Tr}(\partial_\mu X^I \partial_\mu X^I) - \frac{1}{12} \text{Tr}(X_+^I [X^J, X^K] + X_+^J [X^K, X^I] + X_+^K [X^I, X^J])^2\right\}. \quad (2.32)$$

This theory can be regarded as a version of $\lambda\phi^4$ theory in 3 dimensions where we integrate over all the possible λ (which in general are space-time dependent, since they will be harmonic functions in 3 dimensions with suitable boundary conditions). However, the coupling λ is schematically $(X_+^I)^2$, which ensures that the theory (2.32) does not contain any negative norm states. Since the X_-^I integral is exact, we believe that this hints that the negative norm states can be consistently decoupled from the physical Hilbert space.⁵ We have not performed the analysis in the fully gauged Lagrangian (2.29), but we expect that, along the same lines as in the ungauged case, it should be possible to consistently decouple negative norm states even-though the presence of a gauge field (the \mathcal{B} field) of a non-compact gauge symmetry.

One important aspect of having a metric on the space of fields invariant under the symmetry transformations is that it's possible to construct local gauge invariant observables as

$$\mathcal{O}^{IJ}(x) = h^{ab} X_a^I(x) X_b^J(x) = \text{Tr}(X^I X^J) - X_-^I X_+^J - X_-^J X_+^I \quad (2.33)$$

This scalar operator transforms in the $35 \oplus 1$ of $\text{SO}(8)$. The 35 should be BPS and should have a dual in the graviton supermultiplet of 11-dimensional supergravity reduced on $AdS_4 \times S^7$. We can proceed to construct higher order operators as $X_+^{I_1} \dots X_+^{I_n} \mathcal{O}^{IJ}$ and decompose them into $\text{SO}(8)$ irreps.

Note however that this class operators are not invariant under the X_- shift symmetry. Deforming the Lagrangian by adding these operators would translate in a different constraint of the form $\partial^2 X_+ = f(X_+)$. Therefore, it remains to be seen whether this operators survive as physical operators after the negative norm states have been eliminated.

Similarly there are the $\text{SO}(8)$ currents:

$$\mathcal{K}_\mu^{IJ}(x) = h^{ab} (X_a^I(x) \partial_\mu X_b^J(x) - X_a^J(x) \partial_\mu X_b^I(x) + \text{fermions}) \quad (2.34)$$

The energy momentum tensor should have a very similar structure.

2.6 Moduli space of vacua for $\mathcal{G} = \text{SU}(N)$

In this section we perform a preliminary study of the moduli space of vacua of the theory.

⁵The constraint $\partial^2 X_+$ imposed by the delta function in (2.32) can be regarded as the condition that the current associated to the X_- shift symmetry is divergence-free. Therefore, operators that act within the physical Hilbert space should be invariant under the shift symmetry.

Setting the fermions and the gauge fields to zero, and the scalar fields to be constant, we can look for the moduli space of supersymmetric vacua as those configurations preserving supersymmetry. This requires that

$$X_c^I X_d^J X_b^K f^{cd}{}_a = 0 . \tag{2.35}$$

In our case these equations are equivalent to

$$0 = X_+^I [X^J, X^K] + X_+^J [X^K, X^I] + X_+^K [X^I, X^J], \tag{2.36}$$

$$0 = \text{Tr} \left(X^I [X^J, X^K] \right) \tag{2.37}$$

Notice that (2.36) is a set of matrix equations, while (2.37) is just a set of scalar equations. The set of solutions divides into two branches: on one branch $X_+^I X_+^I = 0$, on the other branch $X_+^I X_+^I > 0$.

If $X_+^I X_+^I > 0$ we can use $\text{SO}(8)$ rotations to set $X_+^I = (X_+^1, 0, \dots, 0)$. It is then easy to see that the (2.36) imply $[X^I, X^J] = 0$ for any $I, J = \{1 \dots 8\}$, so that (2.37) is automatically satisfied. In the case $\mathcal{G} = \text{SU}(N)$, the solutions of (2.36) and (2.37) are of the form

$$X^I = U^{-1} (D^I + M X_+^I) U \tag{2.38}$$

where M is an adjoint matrix and U is unitary. X_-^I is unconstrained. We can at this point quotient by global gauge transformations. Setting $M = -D^1/X_+^1$ we can set $X^1 = 0$, and also X_-^1 is fixed. The quotient by the compact part can be used to set X^I to be diagonal and non vanishing for $I = 2 \dots 8$, so it appears that the moduli space has dimension $7(N - 1) + 16$. In a generic point in this moduli space, the surviving gauge symmetry is $\text{U}(1)^{N-1}$. We can now dualize these $N - 1$ low energy gauge fields into $N - 1$ scalars $\Phi_i, i = 1 \dots N - 1$ such that $d\Phi_i = \star F_i$. A priori it looks like these Φ_i will be periodically identified moduli 2π , since they come from dualizing a $\text{U}(1)$ gauge field. Nevertheless, if one looks at the equation of motion for the gauge field B associated with the non-compact gauge symmetry, it demands that $B \sim \star F + \dots$. So the dual scalar Φ is related to the gauge field B via $d\Phi_i \sim B$. Since B is a non-compact gauge field, it is tempting to speculate that the quantum theory allows for a choice of boundary conditions such that Φ is not compact. Assuming this is the case,⁶ and keeping track of the shift symmetry of X_-^I , this abelian branch takes the form $\mathcal{M} = \frac{(\mathbb{R}^8)^N}{S_N}$, where S_N is the discrete group of permutations of the eigenvalues of the X^I matrices, which is the moduli space which one would expect for M2 branes in flat space.

On the other branch $X_+^I = 0$, the eqs. (2.36) are trivially satisfied, while the equations (2.37) impose a set of at most $\mathcal{C}(8, 3) = 56$ independent constraints on the 8 matrices X^I . The non compact part of the gauge symmetry can be used to gauge fix X_-^I , thus we are led to the set of solutions to $\text{Tr}(X^I [X^J, X^K]) = 0$ modulo \mathcal{G} transformations. The dimension of this branch is of order N^2 .

⁶We return to this issue in the note added at the end of this letter, where we briefly address the interpretation of our model in the light of subsequent developments.

2.6.1 Massive excitations on the abelian branch

Let us now examine the effective theory at a generic point on the moduli space \bar{X}^I , where $[\bar{X}^I, \bar{X}^J] = 0$. We can take

$$\bar{X}^I = \text{diag}(a_1^I, \dots, a_N^I) = a^I(\alpha) \delta_\beta^\alpha, \quad \sum_\alpha a^I(\alpha) = 0; \quad (2.39)$$

and then consider linearized fluctuations around this vacuum $X^I = \bar{X}^I + \epsilon^I$. Upon defining

$$L^I(\alpha, \beta) = a^I(\alpha) - a^I(\beta), \quad (2.40)$$

it is easy to see that the linearized equations of motion for ϵ^I become

$$\partial^2(\epsilon^I)_\beta^\alpha - [M^2(\alpha, \beta)]_J^I(\epsilon^J)_\beta^\alpha = 0. \quad (2.41)$$

The mass-squared matrix, in SO(8) space, reads

$$[M^2]_J^I = \left(\bar{L}^2 \bar{X}_+^2 - (\bar{X}_+ \cdot \bar{L})^2 \right) \delta_J^I - \bar{L}^2 X_+^I X_+^J - \bar{X}_+^2 L^I L^J + (\bar{X}_+ \cdot \bar{L}) \left(X_+^I L^J + X_+^J L^I \right), \quad (2.42)$$

where $L^I = L^I(\alpha, \beta)$. It is clear that, for $\alpha = \beta$, since $L^I = 0$, the $(\epsilon^I)_\alpha^\alpha$ will remain massless. However, the shift symmetry $X^I \rightarrow X^I + M X_+^I$ allows to gauge-fix one of the X^I directions. Therefore, the massless fields will be the $(\epsilon^I)_\alpha^\alpha$, with $I = 1, \dots, 7$. Given that the matrix ϵ^I has to be traceless, we are left with $7(N - 1)$ real scalar fields parametrizing the moduli space. In addition, the from the covariant derivatives expanded around this vacuum, it is easy to see that the gauge field B_μ acquires a mass given by $(X_+^I)^2$, while the SU(N) part, corresponding to A_μ , gets broken to $U(1)^{N-1}$. We can now dualize these $N - 1$ abelian gauge fields to obtain $N - 1$ scalars, in such a way that, assuming that there is a choice of boundary conditions where these scalars are not periodic due to the identification $B \sim d\Phi$ imposed by the equation of motion for B , we are left with the $(\mathbb{R}^8)^N / S_N$. On the other hand, hadn't we dualized the abelian gauge fields, we would have $7(N - 1)$ real scalars and $N - 1$ gauge fields, which is the expected field content for D2 branes in flat space. In this case, for a large value of X_+^I , as shown in [16] along the lines of [12], the theory precisely approaches the SYM in 3 dimensions corresponding to a stack of D2 branes.

The massive excitations should carry information about the degrees of freedom through which the membranes interact. With no loss of generality we can choose X_+^I to be aligned with the direction 1, i.e. $X_+^I = (X_+^1, \dots, 0)$. Then, the mass-squared matrix can be easily diagonalized. The squared masses are

$$m^2 = (X_+^1)^2 (\bar{L}^2 - (L^1)^2), \quad (2.43)$$

in such a way that $m = X_+^1 \sqrt{(\bar{L}^2)^2 + \dots + (L^8)^2}$. Note that the L^1 does not appear in the mass expression, which is, as expected, an area. However, at this point it is unclear the meaning of this area, which, however, seems to have one of its vertices trapped at X_+^1 .

3. Comments

In this letter we proposed a large set of superconformal gauge theories described by a Lagrangian with $\mathcal{N} = 8$ supersymmetry which fits into the class of models introduced by Bagger, Lambert and Gustavsson. Part of the field content transforms in the adjoint of an $SU(N)$ gauge symmetry, which appears in the Lagrangian through a BF theory. The theory has a non-compact gauge symmetry, and — under the assumption explained in section 2.6. — allows for a moduli space consistent with a potential interpretation in terms of M2 branes in flat space-time. We have shown that the Lagrangian does not have a tunable coupling, but that the strength of interactions is set by the field X^+ .

It is not evident that the theory is unitary, since some matter fields appear with the wrong sign in the Lagrangian. We argued that the negative norm states can be eliminated via a suitable projection onto physical states. However, a more careful analysis should be undertaken. A related issue is the proper treatment of the shift symmetry of the fields X^I .

Note added. Since this paper first appeared on the arXiv, the understanding of the M2-brane world-volume theory has much improved. Here we briefly comment on the interpretation of our results in the light of these later developments. The most notable progress was made by Aharony, Bergman, Jafferis and Maldacena, who in [24] proposed a description of the M2-brane theory, based on an $\mathcal{N} = 6$ Chern-Simons gauge theory. This ABJM model lacks manifest $SO(8)$ covariance, but (for CS level $k = 1$) appears to possess all the essential properties to describe the worldvolume of coincident M2 branes on \mathbb{R}^8 . It has been shown that the BF membrane theory introduced here and in [23, 25], can be obtained as a limit of the ABJM theory in which the CS level k is sent to infinity [26, 27]. The ABJM theory at level k appears to be related to M2-branes on a \mathbb{Z}_k orbifold of \mathbb{R}^8 , and therefore sending $k \rightarrow \infty$ effectively compactifies one of the directions of \mathbb{R}^8 . When obtained via this limit, the moduli space for the lorentzian BLG theory is therefore $\mathbb{R}^7 \times S^1$, suggesting a more direct relationship with D2-branes. The behavior of the solitons of the BF theory indeed resemble more that of D-branes than M-branes [28].⁷

Our theory as it stands has negative norm states in the spectrum linked to the lightcone-like fields X_{\pm} . It has been argued in [31] that a suitable $\mathcal{N} = 8$ ghost sector can be added to the theory in such a way that the negative norm states decouple. The most simple way of adding such a ghost sector effectively forces X_+ to be constant, and projects the theory onto that of D2 branes. In this case, the BF theory and maximally SUSY 3d SYM are equivalent via the de Wit-Nicolai-Santleben duality [32]. However, there exist alternative ways to eliminate the negative norm states, that still keeps some dynamics that is distinct from that of the pure D2 brane theory, [33].

Acknowledgments

We would like to thank Juan Maldacena and Nathan Seiberg for important suggestions. We

⁷Along a somewhat different line, in [29, 30] it was argued that the moduli space of the Lorentzian BLG theory should be viewed as \mathbb{R}^8 , but written in polar coordinates in a plane, with X^+ as the radial center of mass coordinate. However, as of today, the understanding of this proposal is not entirely satisfactory.

are also grateful to Marcus Benna, Seba Franco, Michele Papucci and Antonello Scardicchio for discussions. E.T. is grateful to Princeton University for the warm hospitality during the development of this paper. The research of SB and HV is supported in part by the National Science Foundation under Grant No. PHY-0756966. D. R-G. acknowledges financial support from the European Commission through Marie Curie OIF grant contract no. MOIF-CT-2006-38381.

A. Check of the fundamental identity

In this appendix we check that the fundamental identity (2.4) is satisfied by the choice (1.1) for the structure constants $f^{abc}{}_d$.

As for the free low index d , when $d = +$ all the 4 terms in the l.h.s. of (2.4) vanish, therefore we are left with $d = -$ and $d = \delta$. From (1.1), the $-$ cannot appear among the upper indices and the low index cannot be $+$ to give non trivial result, therefore from the repeated index we have $g = \rho$.

Starting with $d = -$, we get

$$f^{ef\rho}{}_-\!f^{abc}{}_\rho - f^{efa}{}_\rho f^{bc\rho}{}_-\! - f^{efb}{}_\rho f^{ca\rho}{}_-\! - f^{efc}{}_\rho f^{ab\rho}{}_-\! = 0 . \quad (\text{A.1})$$

When either $e = +$ or $f = +$ and the remaining indices are greek, the (A.1) reduces to the Jacobi identity

$$f^{\phi\alpha}{}_\rho f^{\beta\gamma\rho} + f^{\phi\beta}{}_\rho f^{\gamma\alpha\rho} + f^{\phi\gamma}{}_\rho f^{\alpha\beta\rho} = 0 . \quad (\text{A.2})$$

Instead, when one index among $\{a, b, c\}$ is $+$ and all the other ones (included also e and f) are greek, the two remaining terms in the l.h.s. of (2.4) cancel between themselves. In all the other cases with $d = -$, the four terms in the r.h.s. of (2.4) simply vanish.

When $d = \delta$, then only one index $+$ must occur in the corresponding f

$$f^{ef\rho}{}_\delta f^{abc}{}_\rho - f^{efa}{}_\rho f^{bc\rho}{}_\delta - f^{efb}{}_\rho f^{ca\rho}{}_\delta - f^{efc}{}_\rho f^{ab\rho}{}_\delta = 0 . \quad (\text{A.3})$$

References

- [1] J. Bagger and N. Lambert, *Modeling multiple M2's*, *Phys. Rev. D* **75** (2007) 045020 [[hep-th/0611108](#)].
- [2] J. Bagger and N. Lambert, *Gauge symmetry and supersymmetry of multiple M2-branes*, *Phys. Rev. D* **77** (2008) 065008 [[arXiv:0711.0955](#)].
- [3] J. Bagger and N. Lambert, *Comments on multiple M2-branes*, *JHEP* **02** (2008) 105 [[arXiv:0712.3738](#)].
- [4] A. Gustavsson, *Algebraic structures on parallel M2-branes*, [arXiv:0709.1260](#).
- [5] A. Gustavsson, *Selfdual strings and loop space Nahm equations*, *JHEP* **04** (2008) 083 [[arXiv:0802.3456](#)].
- [6] J.H. Schwarz, *Superconformal Chern-Simons theories*, *JHEP* **11** (2004) 078 [[hep-th/0411077](#)].

- [7] G. Papadopoulos, *M2-branes, 3-Lie algebras and Plücker relations*, *JHEP* **05** (2008) 054 [[arXiv:0804.2662](#)].
- [8] J.P. Gauntlett and J.B. Gutowski, *Constraining maximally supersymmetric membrane actions*, [arXiv:0804.3078](#).
- [9] M. Van Raamsdonk, *Comments on the Bagger-Lambert theory and multiple M2-branes*, *JHEP* **05** (2008) 105 [[arXiv:0803.3803](#)].
- [10] N. Lambert and D. Tong, *Membranes on an orbifold*, *Phys. Rev. Lett.* **101** (2008) 041602 [[arXiv:0804.1114](#)].
- [11] J. Distler, S. Mukhi, C. Papageorgakis and M. Van Raamsdonk, *M2-branes on M-folds*, *JHEP* **05** (2008) 038 [[arXiv:0804.1256](#)].
- [12] S. Mukhi and C. Papageorgakis, *M2 to D2*, *JHEP* **05** (2008) 085 [[arXiv:0803.3218](#)].
- [13] M.A. Bandres, A.E. Lipstein and J.H. Schwarz, *$N = 8$ superconformal Chern-Simons theories*, *JHEP* **05** (2008) 025 [[arXiv:0803.3242](#)].
- [14] D.S. Berman, L.C. Tadrowski and D.C. Thompson, *Aspects of multiple membranes*, *Nucl. Phys. B* **802** (2008) 106 [[arXiv:0803.3611](#)].
- [15] A. Morozov, *On the problem of multiple M2 branes*, *JHEP* **05** (2008) 076 [[arXiv:0804.0913](#)].
- [16] U. Gran, B.E.W. Nilsson and C. Petersson, *On relating multiple M2 and D2-branes*, *JHEP* **10** (2008) 067 [[arXiv:0804.1784](#)].
- [17] P.-M. Ho, R.-C. Hou and Y. Matsuo, *Lie 3-algebra and multiple M2-branes*, *JHEP* **06** (2008) 020 [[arXiv:0804.2110](#)].
- [18] J. Gomis, A.J. Salim and F. Passerini, *Matrix theory of type IIB plane wave from membranes*, *JHEP* **08** (2008) 002 [[arXiv:0804.2186](#)].
- [19] E.A. Bergshoeff, M. de Roo and O. Hohm, *Multiple M2-branes and the embedding tensor*, *Class. and Quant. Grav.* **25** (2008) 142001 [[arXiv:0804.2201](#)].
- [20] K. Hosomichi, K.-M. Lee and S. Lee, *Mass-deformed Bagger-Lambert theory and its BPS objects*, *Phys. Rev. D* **78** (2008) 066015 [[arXiv:0804.2519](#)].
- [21] P.-M. Ho and Y. Matsuo, *M5 from M2*, *JHEP* **06** (2008) 105 [[arXiv:0804.3629](#)].
- [22] H. Awata, M. Li, D. Minic and T. Yoneya, *On the quantization of Nambu brackets*, *JHEP* **02** (2001) 013 [[hep-th/9906248](#)].
- [23] J. Gomis, G. Milanesi and J.G. Russo, *Bagger-Lambert theory for general Lie algebras*, *JHEP* **06** (2008) 075 [[arXiv:0805.1012](#)].
- [24] O. Aharony, O. Bergman, D.L. Jafferis and J. Maldacena, *$N = 6$ superconformal Chern-Simons-matter theories, M2-branes and their gravity duals*, *JHEP* **10** (2008) 091 [[arXiv:0806.1218](#)].
- [25] P.-M. Ho, Y. Imamura and Y. Matsuo, *M2 to D2 revisited*, *JHEP* **07** (2008) 003 [[arXiv:0805.1202](#)].
- [26] Y. Honma, S. Iso, Y. Sumitomo and S. Zhang, *Scaling limit of $N = 6$ superconformal Chern-Simons theories and lorentzian Bagger-Lambert theories*, *Phys. Rev. D* **78** (2008) 105011 [[arXiv:0806.3498](#)].

- [27] E. Antonyan and A.A. Tseytlin, *On 3D $N = 8$ lorentzian BLG theory as a scaling limit of 3D superconformal $N = 6$ ABJM theory*, [arXiv:0811.1540](#).
- [28] R. Iengo and J.G. Russo, *Non-linear theory for multiple M2 branes*, *JHEP* **10** (2008) 030 [[arXiv:0808.2473](#)].
- [29] S. Banerjee and A. Sen, *Interpreting the M2-brane action*, [arXiv:0805.3930](#).
- [30] S. Cecotti and A. Sen, *Coulomb branch of the lorentzian three algebra theory*, [arXiv:0806.1990](#).
- [31] J. Gomis, D. Rodriguez-Gomez, M. Van Raamsdonk and H. Verlinde, *Supersymmetric Yang-Mills theory from lorentzian three-algebras*, *JHEP* **08** (2008) 094 [[arXiv:0806.0738](#)].
- [32] B. Ezhuthachan, S. Mukhi and C. Papageorgakis, *D2 to D2*, *JHEP* **07** (2008) 041 [[arXiv:0806.1639](#)].
- [33] H. Verlinde, *D2 or M2? A note on membrane scattering*, [arXiv:0807.2121](#).