

Torsion and the Gravity Dual of Parity Breaking in AdS_4/CFT_3 Holography

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

JHEP03(2009)033

(<http://iopscience.iop.org/1126-6708/2009/03/033>)

[The Table of Contents](#) and [more related content](#) is available

Download details:

IP Address: 80.92.225.132

The article was downloaded on 03/04/2010 at 10:41

Please note that [terms and conditions apply](#).

Torsion and the Gravity Dual of Parity Breaking in $\text{AdS}_4/\text{CFT}_3$ Holography

Robert G. Leigh,^a Nam Nguyen Hoang^a Anastasios C. Petkou^b

^a*Department of Physics, University of Illinois,
1110 W. Green St., IL 61801-3080 Urbana, U.S.A.*

^b*Department of Physics, University of Crete,
71003 Heraklion, Greece*

E-mail: rgleigh@gmail.com, nnguyen2@illinois.edu,
petkou@physics.uoc.gr

ABSTRACT: We study four dimensional gravity with a negative cosmological constant deformed by the Nieh-Yan torsional topological invariant with a spacetime-dependent coefficient. We find an exact solution of the Euclidean system, which we call the torsion vortex, having two asymptotic AdS_4 regimes supported by a pseudoscalar with a kink profile. We propose that the torsion vortex is the holographic dual of a three dimensional system that exhibits distinct parity breaking vacua. The torsion vortex represents a (holographic) transition between these distinct vacua. We expect that from the boundary point of view, the torsion vortex represents a “domain wall” between the two distinct vacua.

From a bulk point of view, we point out an intriguing identification of the parameters of the torsion vortex with those of an Abrikosov vortex in a Type I superconductor. Following the analogy, we find that external Kalb-Ramond flux then appears to support bubbles of flat spacetime within an asymptotically AdS geometry.

KEYWORDS: Field Theories in Lower Dimensions, AdS-CFT Correspondence, Classical Theories of Gravity, Discrete and Finite Symmetries

ARXIV EPRINT: [0809.5258](https://arxiv.org/abs/0809.5258)

Contents

1	Introduction and summary of the results	1
2	Torsional degrees of freedom in gravity	3
2.1	Preliminaries	3
2.2	Torsion and the magnetic field of gravity	4
3	The Nieh-Yan models	6
3.1	General aspects	6
3.2	The 3+1-split of the pseudoscalar Nieh-Yan model	7
4	The torsion vortex	8
5	The torsion vortex as the gravity dual of parity symmetry breaking	11
6	Physics in the bulk: the superconductor analogy	12
6.1	Gravity vortex as Abrikosov vortex	13
6.2	Multi-vortices and vortex condensation	14
7	Conclusions	16
A	The 3 + 1 split in gravity	17
B	Parity breaking in three dimensions	18

1 Introduction and summary of the results

AdS₄/CFT₃ is currently emerging as a novel paradigm of holography that has qualitatively different properties from the more familiar AdS₅/CFT₄ correspondence. Particularly intriguing is the recent accumulation of evidence that AdS₄/CFT₃ can be used to describe a plethora of phenomena in 2 + 1 dimensional condensed matter systems, such as quantum criticality [1, 2], Quantum Hall transitions [3–6], superconductivity [7–11], superfluidity [12, 13] and spontaneous symmetry breaking [14–16]. Furthermore, AdS₄/CFT₃ is the appropriate setup to study the holographic consequences of generalized electric-magnetic duality of gravity and higher-spin gauge fields [17–21].

In the absence of an explicit AdS₄/CFT₃ correspondence example,¹ various toy models have been used to study its general qualitative aspects. One of the aims of the present

¹The recently suggested field theoretic models for M2 branes [22–27] are important steps towards the understanding of the boundary side of AdS₄/CFT₃.

work it to provide yet another model that can be used to unveil some salient and intriguing properties of AdS₄/CFT₃ holography. However, this is not our only aim. We also wish to study here the relevance of torsion to four dimensional gravity from a holographic point of view. The study of torsion is an interesting subject in itself that poses formal and phenomenological challenges.² In the context of a string theory description of gravity, torsion is omnipresent through antisymmetric tensor fields. AdS₄/CFT₃ provides the basic setup where four dimensional torsion can be holographically investigated.

We consider a simple toy model where torsion is introduced via the topological Nieh-Yan class. In particular, we consider the modification of the Einstein-Hilbert action with a negative cosmological constant by the Nieh-Yan class, the latter having a spacetime-dependent coefficient. In the context of the 3+1-split formalism for gravity [17] we point out that the torsional degrees of freedom are carried by the “gravitational magnetic field”. In pure gravity the magnetic field is fully determined by the frame field and torsion vanishes. In our model, the spacetime dependence of the Nieh-Yan coefficient makes some of the components of the magnetic field dynamical and as a consequence torsional degrees of freedom enter the theory. Our toy model is simple enough such that only one of the torsional degrees of freedom becomes dynamical. This degree of freedom can be either carried by a pseudoscalar, in which case our model is equivalent to a massless pseudoscalar coupled to gravity, or by a two-form gauge potential. In the latter case our model becomes equivalent to a Kalb-Ramond field coupled to gravity.

Next, we find an exact solution of the equations of motion in Euclidean signature. Our metric ansatz is that of a bulk domain wall. The solution, the *torsion vortex*, has two distinct asymptotically AdS₄ regimes along the “radial” coordinate. The pseudoscalar has a kink profile and it is finite at both of the asymptotic regimes. Our torsion vortex can be viewed as a generalization of the axionic wormhole solution of [33] in the case of non-zero cosmological constant. See also [34] for recent work on AdS wormholes. Having in mind the holographic interpretation of our model we focus mainly on the case where the torsional degree of freedom is carried by a pseudoscalar field. Following standard holographic recipes we find that the torsion vortex is the gravity dual of a three dimensional system that possesses two distinct parity breaking vacua. The two vacua are distinguished by the relative sign of the pseudoscalar order parameter. Our bulk picture suggests that the transition from one vacuum to the other can be done by a marginal deformation of the boundary theory. In appendix B we suggest that the above qualitative properties can be realized in the boundary by the three dimensional Gross-Neveu model coupled to U(1) gauge fields.

Finally, we point out that the bulk physics of our vortex solution bears some resemblance to the Abrikosov vortex of superconducting systems. There is a natural mapping of the parameters of the torsion vortex to those of the Abrikosov vortex. We show that the gravitational parameter that is interpreted as an order parameter satisfies a ϕ^4 -like equation and this motivates us to suggest that the cosmological constant is related to the “critical temperature” as $\Lambda \sim T - T_c$. We end with a discussion of multi-vortex config-

²See [28–30] for recent reviews and [31, 32] for other recent works.

urations and vortex condensation. The outcome of this analysis is that H -flux supports bubbles of flat spacetime.

The paper is organized as follows. In section 2 we discuss the torsional degrees of freedom in gravity and their relation with the gravitational magnetic field. In section 3 we present our toy model and its various equivalent manifestations and discuss its 3+1-split formalism of [17]. In section 4 we present the explicit torsion vortex solution of our model. In section 5 we discuss the holography of the torsion vortex. Section 6 contains the bulk physics of the vortex and its relationship to the Abrikosov vortex. It also contains the discussion of multi-vortices and vortex condensation. Technical details and the discussion of the three-dimensional Gross-Neveu model coupled to U(1) gauge fields are contained in the appendices.

2 Torsional degrees of freedom in gravity

2.1 Preliminaries

In this paper we will consider a four dimensional spacetime with a negative cosmological constant. The Einstein-Hilbert action may be written as³

$$I_{\text{EH}} = \int_M \left(\epsilon_{abcd} e^a \wedge e^b \wedge R^{cd} - \frac{1}{6} \Lambda \epsilon_{abcd} e^a \wedge e^b \wedge e^c \wedge e^d \right), \quad (2.1)$$

where e^a denote the one-form frame fields, while ω^a_b are the connection one-forms with curvature $R^a_b = d\omega^a_b + \omega^a_c \wedge \omega^c_b$. As is well-known the variation of (2.1) gives the Einstein equations and also the zero torsion constraint $T^a = de^a + \omega^a_b \wedge e^b = 0$. By virtue of the latter this action can be written entirely in terms of metric variables.

There are also a number of other terms that one may consider. These are all of potential interest to holography because being total derivatives they may induce interesting boundary effects. We may parameterize these terms as follows (writing all possible SO(3,1)-invariant 4-forms constructed from e^a, R^a_b, T^a):

$$I_{\text{top}} = n \int_M C_{\text{NY}} + 2\gamma^{-1} \int_M C_{\text{Im}} + p \int_M P_4 + q \int_M E_4, \quad (2.2)$$

where $C_{\text{NY}} = T^a \wedge T_a - R_{ab} \wedge e^a \wedge e^b = d(T^a \wedge e_a)$ is the Nieh-Yan form, γ is often referred to as the Immirzi parameter with $C_{\text{Im}} = R^a_b \wedge e^b \wedge e_a$, $P_4 = -\frac{1}{8\pi^2} R^a_b \wedge R^b_a = -\frac{1}{8\pi^2} d(\omega^a_b \wedge R^b_a - \frac{1}{3} \omega^a_b \wedge \omega^b_c \wedge \omega^c_a)$ is the Pontryagin form and $E_4 = -\frac{1}{32\pi^2} \epsilon_{abcd} R^{ab} \wedge R^{cd}$ is the Euler form. We note that $P_4 + \frac{\sigma_{\perp} a^2}{4\pi^2} C_{\text{NY}}$ and $C_{\text{NY}} - C_{\text{Im}}$ are actually SO(3,2) invariants [29]. These terms become of more interest, even in gravity, if we allow the coefficients to become fields. Although we will not consider this problem here in full generality, we will consider a particular example. We note that there is older literature, principally by d'Auria and Regge [35] that also considered some such cases (usually in asymptotically Minkowski geometries). In the course of the paper, we will review what is known from those older works. The purpose of our work, amongst other things, is to bring this up to date, and in particular focus on aspects of holography.

³We use $I_{\text{EH}} = -16\pi G_4 S_{\text{EH}}$ where S_{EH} is the usually normalized gravitational action. To fix notation we note that the Einstein equations that follow from S_{EH} are $G_{\mu\nu} + \Lambda g_{\mu\nu} = 0$. We will also write $\Lambda = -3\sigma_{\perp}/L^2$.

2.2 Torsion and the magnetic field of gravity

Our simple model involves only the Nieh-Yan (NY) term. It is interesting to discuss the physics of this topological invariant before we embark on detailed calculations. We will see below that the NY term in gravity plays a role similar to that of the θ -angle in gauge theories.

To see this, we explain below the relationship between the gravitational magnetic field B^α and torsion. Consider the 3+1 split⁴ of the Einstein-Hilbert action (1) with the addition of the usual gravitational Gibbons-Hawking boundary term I_{GH} [17, 19, 20]

$$\begin{aligned}
 I_{\text{EH}} + I_{\text{GH}} = \int dt \wedge & \left(\dot{\tilde{e}}^\alpha \wedge (-4\sigma_\perp \epsilon_{\alpha\beta\gamma} \tilde{e}^\beta \wedge K^\gamma) \right. \\
 & + 2\sigma_\perp N \left\{ 2\tilde{d}(B^\alpha \wedge \tilde{e}_\alpha) + 2B^\alpha \wedge \tilde{T}_\alpha \right. \\
 & + \epsilon_{\alpha\beta\gamma} \left(\sigma B^\alpha \wedge B^\beta - K^\alpha \wedge K^\beta - \frac{\sigma_\perp \Lambda}{3} \tilde{e}^\alpha \wedge \tilde{e}^\beta \right) \wedge \tilde{e}^\gamma \left. \right\} \\
 & - 4\sigma_\perp N^\alpha \epsilon_{\alpha\beta\gamma} (\tilde{D}K)^\beta \wedge \tilde{e}^\gamma + 4Q^\alpha (K_\beta \wedge \tilde{e}^\beta) \wedge \tilde{e}_\alpha \\
 & \left. + 4q^0_\alpha \left\{ \epsilon^\alpha_{\beta\gamma} \tilde{T}^\beta \wedge \tilde{e}^\gamma \right\} \right). \tag{2.3}
 \end{aligned}$$

In the 3+1 split formalism the dynamical variables in (2.3) are the “spatial”⁵ one-forms \tilde{e}^α , K^α and B^α . The first two are canonically conjugate variables. The magnetic field B^α carries the torsional degrees of freedom as it can be seen for example if we write the definition of the non-trivial “spatial” torsion as

$$\tilde{T}^\alpha = \tilde{d}\tilde{e}^\alpha - \sigma \epsilon^{\alpha\beta\gamma} B_\beta \wedge \tilde{e}_\gamma. \tag{2.4}$$

It is easily seen that the radial component of torsion T^0 is determined by \tilde{e}^α and K^α . Notice that (2.4) implies that the tensor $B_{\alpha\beta}$ is odd under “spatial” parity, hence its trace B^α_α is a pseudoscalar. Although a priori the torsional degrees of freedom are not connected with the pair of conjugate variables \tilde{e}^α and K^α , they are not dynamical as there is no kinetic term for B^α . Rather, they enter (2.3) algebraically and as such they give the algebraic zero torsion condition by virtue of which the magnetic field is related to the frame field. Indeed, as discussed in ref. [17], the q^0_α constraint sets to zero the antisymmetric part of B^α in deDonder gauge, such that the first term in the second line of (2.3) vanishes. Then, the variation of (2.3) with respect to B^α yields $\tilde{T}^\alpha = 0$, leaving as true dynamical variables \tilde{e}^α and K^α . This is the gravitational analogue of the electromagnetic case where the magnetic field is related to the gauge potential via the Bianchi identity.

Consider now adding to the Einstein-Hilbert action the Nieh-Yan class C_{NY} with a constant coefficient θ . Over a compact manifold, the NY class is a topological invariant

⁴In appendix A we present a brief review of the 3+1 split formalism where the definitions of the various relevant quantities appear and notation is explained. We note here that σ is the overall signature of the spacetime, σ_\perp is the signature of the “radial” direction and σ_3 the signature of the boundary.

⁵By spatial, we will mean orthogonal to the “radial” coordinate t . In the case of AdS_4 , this radial coordinate is spacelike, and thus $\sigma_\perp = +1$. Then, Lorentzian AdS_4 corresponds to $\sigma_3 = -1$ while Euclidean AdS_4 corresponds to $\sigma_3 = 1$.

and takes integer values⁶ [29]. Having in mind holography, we are interested here in manifolds with boundary. In particular, the 3+1 split has been set up so that the boundary is a constant- t slice. The NY term reduces to a boundary contribution. The explicit calculation yields

$$\mathcal{I}_{\text{NY}} \equiv -2\sigma_{\perp}\theta \int C_{\text{NY}} = 2\sigma_{\perp}\theta \int dt \wedge \left[2\epsilon_{\alpha\beta\gamma}\dot{\tilde{e}}^{\alpha} \wedge \tilde{e}^{\beta} \wedge B^{\gamma} + \epsilon_{\alpha\beta\gamma}\dot{B}^{\alpha} \wedge \tilde{e}^{\beta} \wedge \tilde{e}^{\gamma} \right]. \quad (2.5)$$

Adding (2.5) to (2.3) we obtain

$$I_{\text{EH}} + I_{\text{GH}} + \mathcal{I}_{\text{NY}} = \int dt \wedge \left(\dot{\tilde{e}}^{\alpha} \wedge (-4\sigma_{\perp}\epsilon_{\alpha\beta\gamma}\tilde{e}^{\beta} \wedge [K^{\gamma} - \theta B^{\gamma}]) + 2\sigma_{\perp}\theta\epsilon_{\alpha\beta\gamma}\dot{B}^{\alpha} \wedge \tilde{e}^{\beta} \wedge \tilde{e}^{\gamma} + \text{constraint terms} \right). \quad (2.6)$$

Notice that the \mathcal{I}_{NY} term has two effects. One is to modify the canonical momentum variable $K^{\alpha} \mapsto K^{\alpha} - \theta B^{\alpha}$. This is analogous to the effect of the θ -angle in the canonical description of electromagnetism [36]. The other is to provide a kinetic term for the singlet component of the magnetic field (one easily verifies that only B^{α}_{α} contributes in the second term in the first line of (2.6)). This second effect has no analogue in electromagnetism. Taking the variation of (2.6) with respect to B^{α} , one finds that the zero torsion condition still holds. This is expected of course since the \mathcal{I}_{NY} term is purely a boundary term. As a consequence, the true dynamical variables remain \tilde{e}^{α} and K^{α} . However, the holography is slightly modified. The variation of (2.6) gives on-shell

$$\delta (I_{\text{EH}} + I_{\text{GH}} + \mathcal{I}_{\text{NY}})_{\text{on shell}} = \int_{\partial\mathcal{M}} \delta\tilde{e}^{\alpha} \wedge \left(-4\sigma_{\perp}\epsilon_{\alpha\beta\gamma}\tilde{e}^{\beta} \wedge [K^{\gamma} - \theta B^{\gamma}] \right)_{\text{on shell}}. \quad (2.7)$$

After the appropriate subtraction of divergences [19, 20], (2.7) yields a modified boundary energy momentum tensor. The modification is due to the term $4\sigma_{\perp}\theta\epsilon_{\alpha\beta\gamma}\tilde{e}^{\beta} \wedge B^{\gamma}$ which is parity odd and corresponds to the unique symmetric, conserved and traceless tensor of rank two and scaling dimension three that can be constructed from the three-dimensional metric [37]. It is the exact analogue of the topological spin-1 current constructed from the three dimensional gauge potential.

The form of the action (2.6) unveils an intriguing possibility. The above holographic interpretation was based on the zero torsion condition that connects B^{α} to the frame field. However, to get the zero torsion condition from (2.6) we needed to integrate by parts the last term in the first line. Hence, if θ were t -dependent, the torsion would no longer be zero and the trace B^{α}_{α} would become a proper dynamical degree of freedom independent of \tilde{e}^{α} . In such a case the holographic interpretation of (2.6) would change. The new bulk degree of freedom would couple to a new pseudoscalar boundary operator. As a consequence, we have the possibility to probe additional aspects of the boundary physics and describe new 2+1 dimensional phenomena. That we do in the next section.

⁶More precisely, $C_{\text{NY}}/(2\pi L)^2$ is integral, as it is equal to the difference of two Pontryagin forms, one $\text{SO}(3,2)$ and one $\text{SO}(3,1)$.

3 The Nieh-Yan models

3.1 General aspects

In the previous section we sketched a mechanism by which torsional degrees of freedom become dynamical. In particular, we have argued that the addition of the Nieh-Yan class with a space-time-dependent coefficient in the Einstein-Hilbert action makes dynamical one pseudoscalar degree of freedom which is connected to the trace of the gravitational magnetic field. Adding boundary terms to the bulk action corresponds to a canonical transformation. Consequently, by adding boundary terms we can change the canonical interpretation and the variational principle. Consider first the action

$$I'_{\text{NY}} = I_{\text{EH}}[e, \omega] + I_{\text{GH}}[e, \omega] + 2 \int_M F(x) C_{\text{NY}}, \quad (3.1)$$

where F is a pseudoscalar “axion” field with no kinetic term. If $F \equiv -\sigma_{\perp} \theta$ were a constant, this theory would be equivalent to that studied in the last section. With $F = F(x)$, we have additional terms in the action involving gradients of F . If we perform the 3 + 1 split on this action, we will find that \tilde{e}^{α} and B^{α} are canonical coordinates, and their conjugate momenta will depend on F .

The action as given may be supplemented by additional boundary terms. Such boundary terms are analogous to the Gibbons-Hawking term in gravity, but here involve the torsional degrees of freedom. In particular, we can replace I'_{NY} by

$$I_{\text{NY}} = I_{\text{EH}}[e, \omega] + I_{\text{GH}}[e, \omega] - 2 \int_M dF \wedge T_a \wedge e^a. \quad (3.2)$$

This action is such that \tilde{e}^{α} and F are canonical coordinates with appropriate boundary conditions, while B^{α} appears in the momentum conjugate to F . To investigate this theory, we note that the variation of the action takes the form

$$\begin{aligned} \delta I_{\text{NY}} = & 2 \int_M \delta e^d \wedge \left[\epsilon_{abcd} e^b \wedge \left(R^{cd} - \frac{1}{3} \Lambda e^c \wedge e^d \right) + 2dF \wedge T_d \right] \\ & + 2 \int_M \delta \omega^{ab} \wedge \left[\epsilon_{abcd} T^c \wedge e^d + dF \wedge e_b \wedge e_a \right] + 2 \int_M \delta F C_{\text{NY}} \\ & + 2 \int_M d \left[\delta e^a \wedge \left(\epsilon_{abcd} e^b \wedge \omega^{cd} - dF \wedge e_a \right) - T_a \wedge e^a \delta F \right]. \end{aligned} \quad (3.3)$$

A non-trivial configuration of F sources a particular component of the torsion. Indeed the classical equations of motion can be manipulated to yield in the bulk

$$T^a \wedge e_a = 3 *_4 dF, \quad (3.4)$$

where $*_4$ denotes the Hodge-* operation. However, as d’Auria and Regge [35] showed, this classical system is equivalent to a pseudoscalar coupled to torsionless gravity.

$$I_{\text{PS}} = I_{\text{EH}}[e, \mathring{\omega}] + I_{\text{GH}}[e, \mathring{\omega}] - 3 \int_M dF \wedge *_4 dF. \quad (3.5)$$

This comes about as follows. We write the connection as $\omega = \overset{\circ}{\omega} + \Omega$, where $\overset{\circ}{\omega}$ is torsionless, and insert the equation of motion (3.4). The latter becomes an equation⁷ for Ω , and we obtain (3.5).

A massless pseudoscalar field coupled to torsionless gravity is holographically dual to composite pseudoscalar operators of dimension $\Delta = 3, 0$ in the boundary. The usual holographic dictionary then says that only the $\Delta = 3$ operator appears in the boundary theory since only this is above the unitarity bound of the three dimensional conformal group $SO(3,2)$. A scalar operator with dimension $\Delta = 0$ would simply correspond to a constant in the boundary. Hence, the sensible holographic interpretation of the massless bulk pseudoscalar is that its leading behaviour determines the marginal coupling of a $\Delta = 3$ operator; the expectation value of the operator itself is determined by the subleading behaviour of the bulk pseudoscalar.

Another equivalent formulation of this bulk theory is obtained by writing

$$*_4 dF = \frac{1}{3}H. \tag{3.6}$$

with H a 3-form field. This is the parameterization that would be most familiar from string theory, as the system simply corresponds to an antisymmetric 2-form field. In this formulation, we write

$$\begin{aligned} I_{\text{KR}} &= I_{\text{EH}}[e, \overset{\circ}{\omega}] + I_{\text{GH}}[e, \overset{\circ}{\omega}] + \frac{1}{3} \int_M H \wedge *_4 H + \sqrt{\frac{2}{3}} \int_M C \wedge d *_4 H \\ &= I_{\text{EH}}[e, \overset{\circ}{\omega}] + I_{\text{GH}}[e, \overset{\circ}{\omega}] - \frac{1}{2} \int_M dC \wedge *_4 dC + \int_M d(C \wedge *_4 dC). \end{aligned} \tag{3.7}$$

In the first equation, C appears as a Lagrange multiplier for the ‘‘Gauss constraint’’ and in the second expression, we have solved for the H equation of motion in the bulk, which is just $H = \sqrt{\frac{3}{2}}dC$.

3.2 The 3+1-split of the pseudoscalar Nieh-Yan model

To investigate the holographic aspects of our model it is most useful to use the ‘‘radial quantization’’ in which we think of the radial coordinate as ‘‘time’’ t . We have derived the radial 3+1 split in the first order formalism in [17], and this is summarized with explanations of notation in appendix A. Here we update that calculation to include torsional terms. The Nieh-Yan deformation gives

$$\begin{aligned} -2 \int dF \wedge T^\alpha \wedge e_\alpha &= 2 \int dt \wedge \left\{ -\dot{F} \tilde{T}_\alpha \wedge \tilde{e}^\alpha - \dot{\tilde{e}}_\alpha \wedge \tilde{d}F \wedge \tilde{e}^\alpha + N[2\tilde{d}F \wedge K_\alpha \wedge \tilde{e}^\alpha] \right. \\ &\quad \left. + N^\alpha[2\tilde{d}F \wedge \tilde{T}_\alpha] + Q^\alpha[-\sigma \epsilon_{\alpha\beta\gamma} \tilde{d}F \wedge \tilde{e}^\beta \wedge \tilde{e}^\gamma] \right\}. \end{aligned} \tag{3.8}$$

We see that the F field makes a contribution to the constraints, and has a conjugate momentum proportional to the scalar part of the torsion (the part transverse to the radial

⁷Explicitly this is $\Omega^a{}_b = \frac{\sigma}{4} \epsilon^{acd} \partial_c F e_d$.

direction). The full bulk action becomes

$$\begin{aligned}
I = \int dt \wedge & \left(\dot{\tilde{e}}^\alpha \wedge (4\sigma_\perp \epsilon_{\alpha\beta\gamma} K^\gamma \wedge \tilde{e}^\beta - 2\tilde{d}F \wedge \tilde{e}_\alpha) - 2\dot{F}(\tilde{e}^\alpha \wedge \tilde{T}_\alpha) \right. \\
& + N \left\{ 2\epsilon_{\alpha\beta\gamma} \left({}^{(3)}R^{\alpha\beta} - \sigma_\perp K^\alpha \wedge K^\beta - \frac{\Lambda}{3} \tilde{e}^\alpha \wedge \tilde{e}^\beta \right) \wedge \tilde{e}^\gamma + 4\tilde{d}F \wedge K_\alpha \wedge \tilde{e}^\alpha \right\} \\
& + 4N^\alpha \left\{ -\sigma_\perp \epsilon_{\alpha\beta\gamma} (\tilde{D}K)^\beta \wedge \tilde{e}^\gamma + \tilde{d}F \wedge \tilde{T}_\alpha \right\} \\
& + 4Q^\alpha \left\{ (K_\beta \wedge \tilde{e}^\beta) \wedge \tilde{e}_\alpha - \frac{1}{2} \sigma \epsilon_{\alpha\beta\gamma} \tilde{d}F \wedge \tilde{e}^\beta \wedge \tilde{e}^\gamma \right\} \\
& \left. + 4q^0_\alpha \left\{ \epsilon^\alpha_{\beta\gamma} \tilde{T}^\beta \wedge \tilde{e}^\gamma \right\} \right). \tag{3.9}
\end{aligned}$$

We notice that the Q -constraint term can be written in the form

$$4Q_\alpha \tilde{e}^\alpha \wedge (K_\beta \wedge \tilde{e}^\beta - \sigma *_3 \tilde{d}F). \tag{3.10}$$

Because of this constraint (which relates the antisymmetric part of the extrinsic curvature to the vorticity of F), the momentum conjugate to \tilde{e}^α is symmetric, i.e.

$$\begin{aligned}
\Pi_\alpha &= 4\sigma_\perp \epsilon_{\alpha\beta\gamma} K^\gamma \wedge \tilde{e}^\beta - 2\tilde{d}F \wedge \tilde{e}_\alpha \\
&= 4\sigma_\perp \left(\epsilon_{\alpha\beta\gamma} K^\gamma \wedge \tilde{e}^\beta - \frac{1}{2} \sigma *_3 (K_\beta \wedge \tilde{e}^\beta) \wedge \tilde{e}_\alpha \right). \tag{3.11}
\end{aligned}$$

When written out in components, one finds that the antisymmetric part $K_{[\alpha\beta]}$ cancels

$$\Pi_\alpha = 4\sigma_\perp (K_{(\beta\alpha)} - tr K \eta_{\beta\alpha}) \tilde{e}^\beta. \tag{3.12}$$

This result is consistent with the fact noted above, that the system may be equivalently described as a pseudoscalar field coupled to torsionless gravity. Moreover, if we take the deDonder gauge $d^\dagger \tilde{e}^\alpha = 0$, the torsion constraint implies that B is symmetric.

The q^0_α constraint yields $\tilde{T}^\beta_{\alpha\beta} = 0$. Out of the nine components of \tilde{T} , which transform as $\mathbf{5} + \mathbf{3} + \mathbf{1}$ under $SO(3)$ (or $SO(2,1)$), this sets the triplet to zero (the $\mathbf{5}$ also vanishes on an equation of motion). The momentum conjugate to F is given by

$$\Pi_F = -2\epsilon^{\alpha\beta\gamma} \tilde{T}_{\alpha\beta\gamma}. \tag{3.13}$$

This is the singlet part of the torsion, which has become dynamical in this description of the theory, in the sense that it is canonically conjugate to F .

4 The torsion vortex

We will now simplify the analysis by taking a coordinate basis and looking for solutions of the form

$$\tilde{e}^\alpha = e^{A(t)} dx^\alpha, \quad N = 1, \quad N^\alpha = 0, \tag{4.1}$$

and we will further suppose that $F = F(t)$. In this case K^α and B^α reduce to one degree of freedom each as a result of the constraints

$$K_\alpha = k \tilde{e}_\alpha, \quad B_\alpha = b \tilde{e}_\alpha, \tag{4.2}$$

and one finds $\Pi_A = -4\sigma_\perp k$ and $\Pi_F = 2\sigma b$. The action then takes the following relatively simple Hamiltonian form

$$I_{\text{NY}} \propto \int dt d^3x e^{3A(t)} \left[\dot{A}\Pi_A + \dot{F}\Pi_F - \left(\frac{1}{2}\sigma_3\Pi_F^2 + \frac{1}{8}\sigma_\perp\Pi_A^2 + \frac{2}{3}\Lambda \right) \right]. \quad (4.3)$$

and the equations of motion give

$$\dot{\Pi}_A = 3\dot{F}\Pi_F, \quad \dot{\Pi}_F + 3\Pi_F\dot{A} = 0, \quad \Pi_A = 4\sigma_\perp\dot{A}, \quad \Pi_F = \sigma_3\dot{F}, \quad (4.4)$$

$$\Pi_A^2 + 4\sigma\Pi_F^2 + \frac{16}{3}\sigma_\perp\Lambda = 0. \quad (4.5)$$

These equations of motion could of course alternatively be obtained by considering the theory in the form (3.5). It is convenient to rescale $F(t) = \frac{1}{3}\Theta(t)$. Then the equations of motion can be put in the form

$$\ddot{A} + 3\dot{A}^2 - 3a^2 = 0, \quad \ddot{A} = \frac{1}{12}\sigma\dot{\Theta}^2, \quad \ddot{\Theta} + 3\dot{\Theta}\dot{A} = 0. \quad (4.6)$$

where we have set $\Lambda = -3\sigma_\perp a^2$ with $a = 1/L$. These are of the standard form of domain wall equations that have appeared numerous times in the AdS/CFT literature. However, there is a crucial difference. Notice that the first two of (4.6) imply

$$\dot{A}^2 + \frac{1}{36}\sigma\dot{\Theta}^2 - a^2 = 0. \quad (4.7)$$

For Euclidean signature ($\sigma = \sigma_3 = 1$) the second term in (4.7) has *positive* sign in contrast to most of the other holographic studies. This is due to the fact that in passing from Lorentzian to Euclidean signature the pseudoscalar kinetic term acquires the “wrong sign” [38]. This property allows for a remarkable exact solution to the above system of non-linear equations in Euclidean signature, which we refer to as the *torsion vortex*. To obtain it we define

$$h(t) = \dot{A}(t), \quad (4.8)$$

at which point we have

$$\dot{h} = \frac{1}{12}\dot{\Theta}^2, \quad \dot{h} + 3(h^2 - a^2) = 0. \quad (4.9)$$

The general solution is of the form

$$h(t) = a \tanh 3a(t - t_0) \quad (4.10)$$

and we then have

$$\Pi_F = \dot{F} = \pm 2\sqrt{a^2 - h^2(t)} = \pm 2a \operatorname{sech} 3a(t - t_0) \quad (4.11)$$

which gives

$$\Theta(t) = \Theta_0 \pm 4 \arctan \left(e^{3a(t-t_0)} \right). \quad (4.12)$$




Figure 1. Plot of the torsion vortex solution vs. t . The blue dashed line is $e^{A(t)}$ while the red solid line is $\Theta(t)$. To make the plot, we have chosen $\Theta_0 = 0$, $t_0 = 0$ and $\alpha = 1$.

The \pm sign corresponds to kink/antikink and we will without loss of generality choose the $+$ sign. We may also solve for

$$e^{A(t)} = \alpha(2 \cosh 3a(t - t_0))^{1/3} \tag{4.13}$$

The parameter α is an arbitrary positive integration constant that sets the overall scale of the spatial part of the metric. t_0 may be interpreted as the position of the vortex; when $t_0 = 0$ the torsion vortex sits in the middle between the two asymptotically AdS_4 regimes. Below, we will discuss the interesting holographic interpretation of the torsion vortex.

Note the curvature and torsion of this solution:

$$R^\alpha{}_\beta = -\dot{F}\dot{A} \epsilon^\alpha{}_{\beta\gamma} dt \wedge e^\gamma - a^2 e^\alpha \wedge e_\beta, \tag{4.14}$$

$$R^\alpha{}_0 = (\dot{h} + h^2) dt \wedge e^\alpha - \frac{1}{2} \dot{F}\dot{A} \epsilon^\alpha{}_{\beta\gamma} e^\beta \wedge e^\gamma, \tag{4.15}$$

$$T^\alpha = -\frac{1}{2} \dot{F} \epsilon^\alpha{}_{\beta\gamma} e^\beta \wedge e^\gamma, \tag{4.16}$$

$$T^0 = 0. \tag{4.17}$$

These are non-singular for all $t \in (-\infty, \infty)$. The torsion vortex solution has divergent action, but this divergence is cancelled by boundary counterterms, the same counterterms which render the action of AdS_4 finite. To see this, the energy of the torsion vortex can be computed by evaluating the Euclidean action on the solution. Introducing a cutoff at $t = \pm L$, we find

$$I_{\text{tv, on-shell}} = 4a^2 \int \epsilon_{\alpha\beta\gamma} dx^\alpha \wedge dx^\beta \wedge dx^\gamma \int dt e^{3A(t)} \tag{4.18}$$

$$= (6 \int \widehat{Vol}_3) \cdot \left(\frac{4}{3} a \alpha^3 e^{3aL} + \dots \right), \tag{4.19}$$

where the ellipsis contains terms that vanish when the cutoff is removed. As in pure AdS_4 , an appropriate counterterm is of the form [39, 40]

$$I_{\text{c.t.}} = -\frac{4a}{3} \int_{\partial M} \epsilon_{\alpha\beta\gamma} \tilde{e}^\alpha \wedge \tilde{e}^\beta \wedge \tilde{e}^\gamma. \tag{4.20}$$

In the present case, we have such a counterterm on *each* asymptotic boundary, and thus we find

$$I_{\text{c.t.}} = -2\frac{2a}{3}\alpha^3 e^{3aL} \cdot (6 \int \widehat{Vol}_3), \quad (4.21)$$

which exactly cancels the divergent energy of the torsion vortex.

Furthermore, we note that in the Kalb-Ramond representation, the solution has

$$H = \dot{\Theta} Vol_3 = \pm 6a\alpha^3 \widehat{Vol}_3 \equiv \hat{H} \widehat{Vol}_3, \quad (4.22)$$

where $\widehat{Vol}_3 = \frac{1}{6}\epsilon_{\alpha\beta\gamma} dx^\alpha \wedge dx^\beta \wedge dx^\gamma$. This corresponds to a “topological quantum number” of the kink

$$\int *_4 H = \pm \Delta \Theta = \pm 2\pi. \quad (4.23)$$

5 The torsion vortex as the gravity dual of parity symmetry breaking

The holographic interpretation of the torsion vortex is rather interesting. To study this, we set to zero without loss of generality the integration constant $\Theta_0 = 0$ and pick the plus sign in (4.11), (4.12). Next we need the asymptotic expansion of the vierbein which reads

$$\bar{e}^\alpha = 2^{-1/3} \alpha e^{\pm a(t-t_0)} \left(1 + \frac{1}{3} e^{\mp 6a(t-t_0)} + \dots \right) dx^\alpha \text{ for } t \rightarrow \pm\infty. \quad (5.1)$$

This shows that our solution is asymptotically anti-de Sitter for both $t \rightarrow \pm\infty$. The two asymptotic AdS spaces have the same cosmological constant. From this expansion we could read the expectation value of the renormalized boundary energy momentum tensor which would be given by the coefficient of the $e^{\pm 3at}$ term (see e.g. [19, 20]). Such a term is missing in (5.1), hence the expectation value of the boundary energy momentum tensor is zero.

It is not immediately apparent how to interpret these two asymptotic regimes. Are they truly distinct, or should they be identified in some way? We note that the pseudoscalar behaves in these asymptotic regimes as

$$\Theta(t) \rightarrow 4e^{-3a(t-t_0)} - \frac{4}{3}e^{-9a(t-t_0)} + \dots \text{ for } t \rightarrow -\infty, \quad (5.2)$$

$$\Theta(t) \rightarrow 2\pi - 4e^{3a(t-t_0)} + \frac{4}{3}e^{9a(t-t_0)} + \dots \text{ for } t \rightarrow +\infty. \quad (5.3)$$

From the above we confirm that $\Theta(t)$ is dual to a dimension $\Delta = 3$ boundary pseudoscalar that we denote \mathcal{O}_3 . In each one of the asymptotically AdS regimes, the leading constant behavior of $\Theta(t)$ corresponds to the source (i.e., coupling constant) for \mathcal{O}_3 and the sub-leading term proportional to $e^{\mp 3a(t-t_0)}$ to the expectation value $\langle \mathcal{O}_3 \rangle$. The two asymptotic regimes are distinguished by the behavior of Θ . In fact, the essential difference is *parity*.

We can now describe the holography of our torsion vortex. In the $t \rightarrow -\infty$ boundary sits a three-dimensional CFT at a parity breaking vacuum state. The order parameter is the expectation value of the pseudoscalar which is $\langle \mathcal{O}_3 \rangle = 4$ in units of the AdS radius. The expectation value breaks of course the conformal invariance of the boundary theory. Then, the theory is deformed by the same pseudoscalar operator $g\mathcal{O}_3$ where g is a marginal

coupling. The torsion vortex provides the holographic description of that deformation. Nevertheless, our solution should not be interpreted in terms of the usual holographic renormalization group flow. In our case, at $t \rightarrow +\infty$ the space becomes AdS with the *same* radius as at $t \rightarrow -\infty$. Hence, the two boundary theories have the same “central charges”.⁸

We suggest that instead of interpreting the solution in terms of an RG flow, we should think of it as a transition between two inequivalent vacua of a single boundary theory. This statement is supported by the behavior of $\Theta(t)$ in the two asymptotic regimes. For $t \rightarrow \infty$ the pseudoscalar asymptotes to the configuration (5.3). The interpretation is now that when the marginal coupling takes the fixed value $g_* = 2\pi$ we are back to the *same* CFT (i.e. having the same central charge) however in a distinct parity breaking vacuum such that $\langle \mathcal{O}_3 \rangle = -4$. In others words, the two asymptotic AdS regimes seem to describe two distinct parity breaking vacua of the same theory. The two vacua are distinguished by the expectation value of the parity breaking order parameter being $\langle \mathcal{O}_3 \rangle = \pm 4$. Quite remarkably, we also seem to find that starting in one of the two vacua, we can reach the other by a marginal deformation with a *fixed* value of the deformation parameter.

Since the marginal operator is of dimension $\Delta = 3$ and parity odd, we tentatively identify it with a Chern-Simons operator of a boundary gauge field. In this case the torsion vortex induces the T-transformation in the boundary CFT [37, 41]. In appendix B we will argue that the three dimensional Gross-Neveu model coupled to abelian gauge fields exhibits a large- N vacuum structure that matches our holographic findings. Although our bulk model is extremely simple to provide details for its possible holographic dual, we regard this remarkable similarity as strong qualitative evidence that our torsion vortex is the gravity dual of the “tunneling” between different parity breaking vacua in three dimensions. However, in a three-dimensional quantum field theory, we do not expect that tunneling can occur because of large volume effects, and distinct vacua remain orthogonal. Thus, referring to the torsion vortex as a tunneling event should be taken figuratively. We leave to future work a more careful study of the boundary interpretation of the torsion vortex solution. An interpretation will depend on the precise topology of the boundary. [42] Moreover, embedding our model into M-theory could provide additional clues regarding its holography.

6 Physics in the bulk: the superconductor analogy

The bulk interpretation of the exact solution is also interesting. Because the pseudoscalar field undergoes $\Theta(t) \rightarrow \Theta(t) + 2\pi$ under t goes from $-\infty$ to $+\infty$, the exact solution corresponds to a topological kink. It satisfies

$$\int dt \dot{\Theta} = 2\pi .$$

In figure 2, we plot the solution.

⁸We use “central charge” in $d = 3$ for a quantity that counts the massless degrees of freedom at the fixed point. Such a quantity may be taken to be the coefficient in the two-point function of the energy momentum tensor or the coefficient of the free energy density. There is no conformal anomaly in $d = 3$.




Figure 2. The blue dashed line is $|h(t)|$, resembling the order parameter of a superconductor, while the solid red line is Π_F , analogous to the magnetic induction of an Abrikosov vortex.

Abrikosov vortex	Torsion vortex
order parameter Φ	order parameter $ h = \dot{A} $
$T - T_c$	Λ
magnetic induction B	Π_F
magnetic field H	\hat{H}
\mathbb{Z} -quantized magnetic flux	\mathbb{Z}_2 -quantized electric flux

Table 1. Abrikosov vortex v.s. Torsion vortex.

6.1 Gravity vortex as Abrikosov vortex

The gravity vortex solution (4.9), (4.12) bears some resemblance to the Abrikosov vortex of superconducting systems. To avoid confusion, we emphasise here that this is *not* a holographic interpretation i.e. the Abrikosov vortex (or more precisely, domain wall) is in the bulk. In this section, we will explore this and point out some possibly interesting features. The first thing to notice is that the plot in figure 2 is identical to the profile of an Abrikosov vortex (see for example figure 5.1 in ref. [43].) The codimension differs,⁹ but there is a correspondence between our radial t -direction and the radial direction in the Abrikosov vortex, and $|h|$ and Π_F correspond to the condensate and magnetic induction of the superconductor, respectively. Table 1 summarizes the correspondence. In this correspondence, since the order parameter is $h = \dot{A}$, the superconducting phase (constant order parameter) corresponds to AdS_4 , while the normal phase corresponds to flat space ($h = 0$). Far away from the core of the torsion vortex, the geometry is asymptotically AdS, but at the core the spatial slice (at $t \rightarrow t_0$) becomes flat. To see this, note that if we think of the system as a pseudoscalar coupled to torsionless gravity, the torsion vortex has $\hat{\omega}^{\alpha\beta} = 0$

⁹The difference in dimensionality of the core is what we expect, since it supports a 3-form field strength in contrast to a 2-form field strength in superconductivity.

and $\overset{\circ}{\omega}{}^\alpha{}_0 = \dot{A}\tilde{e}^\alpha$, and so

$$\hat{R}^\alpha{}_\beta = -h^2\tilde{e}^\alpha \wedge \tilde{e}_\beta, \tag{6.1}$$

$$\hat{R}^\alpha{}_0 = (\dot{h} + h^2)dt \wedge \tilde{e}^\alpha, \tag{6.2}$$

$$\hat{T}^\alpha = 0. \tag{6.3}$$

Thus, at the core, we find that the Riemann tensor has components

$$R^\alpha{}_{0\alpha 0} \rightarrow -3a^2\alpha, \tag{6.4}$$

$$R^\alpha{}_{\beta\alpha\beta} \rightarrow 0. \tag{6.5}$$

This behavior is in line with an Abrikosov vortex in which there is normal phase at the core and superconducting phase away from the core.

The analogue of the magnetic field is what we have called \hat{H} , proportional to the constant α^3 . In the vortex, the magnetic induction, analogous to Π_F , has a penetration length $\lambda \sim 1/3a$, and the coherence length of the order parameter is $\xi \sim 1/6a$. The penetration and coherence lengths are obtained by looking at the exponential fall-off of these quantities in the core of the vortex, away from their values in the superconducting phase.

The torsion vortex also has a quantized flux $\int *_4H = \Delta\Theta = 2\pi$. This flux is independent of any parameters of the solution and of any rescaling of fields in the theory. Thus, this is an analogue of the quantized magnetic flux in superconductivity.

Finally, note the following interesting feature. If we take a derivative of the second equation in (4.9), we arrive at

$$\ddot{h} - 6\Lambda h - 18h^3 = 0. \tag{6.6}$$

This looks like a Landau-Ginzburg equation of motion of an effective ϕ^4 theory. This leads us to interpret $\Lambda \sim T - T_c$. Of course, there is no real temperature in the case of the torsion vortex, but we note that this implies that the penetration and coherence lengths diverge as $T \rightarrow T_c$ with exponent 1/2, as in superconductivity.

6.2 Multi-vortices and vortex condensation

In the last section, we noted that there is a strong analogue between the torsion vortex solution and superconductivity. It is intriguing to carry the analogy further and consider multi-vortex configurations. We have noted that at the core of the torsion vortex, the spatial sections are flat. Thus, one might imagine that if it was favourable for torsion vortices to condense, as vortices do in Type I superconductors, then finite regions of normal phase (corresponding to $\Lambda = 0$) would be obtained. We will argue below that this can in fact occur, although the system appears not to be unstable.

To understand the physics involved, the first step is to consider a configuration of two vortices. In the superconductivity literature, this is a standard computation. One takes two vortices separated by a distance ℓ and computes the Euclidean action. More precisely, we will treat this here as follows. Denoting the torsion vortex schematically as $\Phi(t_0)$, we take a configuration

$$\begin{cases} \Phi(\ell/2), & t > 0 \\ \Phi(-\ell/2), & t < 0. \end{cases} \tag{6.7}$$




Figure 3. Size of normal state droplet vs. n for multi-vortex.




Figure 4. Energy vs. n for multi-vortex.

We have taken a piecewise solution, because solutions of non-linear equations cannot be simply superimposed. The result is not quite a solution to the equations of motion of course, failing at the midpoint between the vortices. However, if we simply evaluate the Euclidean action, we find

$$S_E(\ell) = 4a\alpha^3 \sinh(3a\ell/2). \tag{6.8}$$

Note that this is positive, so one might naively conclude that the vortices repel each other. However, recall that the vortex profile exists not in flat space-time, but in the metric given by (4.13), which rises asymptotically. As a result, as we move the vortices further apart, there is a corresponding rise in the metric between the vortices. So, we should directly evaluate the force

$$F = -\frac{dS_E}{d\ell} = -6a\alpha^3 \cosh(3a\ell/2) < 0. \tag{6.9}$$

and thus we conclude that the vortices in fact *attract* each other. In the superconducting analogue, this implies that we have a *Type I superconductor*. In such a superconductor, the number of vortices is determined by the total magnetic flux, and the vortices tend to clump together forming (potentially) finite regions of normal phase within the superconductor.

We now describe the analogous situation in our gravitational system. We have noted that the constant \hat{H} plays the role of the external magnetic induction, while H is the magnetic field, varying within the vortex, with $\Delta\Theta = \int *_4 H$. Following the superconducting analogue, if we put the system in a box of size $2L$ (that is we impose a cutoff on each AdS asymptotic) the flux conservation equation is of the form

$$\Delta\Theta = 2L\hat{H}. \tag{6.10}$$

The vortices carry the flux in the superconductor, and so it is natural to ask what is the lowest energy configuration satisfying (6.10)? To analyze this, consider an array of n vortices in a region of size L_0 . We take the vortices to be equally spaced, as one can show that deviating from such a configuration causes a rise in energy. For such a configuration, the flux quantization condition (6.10) gives a relation between n, L_0 and \hat{H} . Such a representative curve is shown in figure 3. If we solve this equation for L_0 as a function of n

and \hat{H} , we can then compute the energy as a function of n . One obtains a curve as in figure 4. One notes that the energy is minimized for large n , and in that case, the size L_0 asymptotes to a fixed value, which is found to be

$$L_0 = \frac{\hat{H}}{6a} \cdot 2L = \alpha^3 \cdot 2L. \tag{6.11}$$

We conclude that the preferred configuration, given a fixed external flux, is a continuum of vortices arrayed over a finite size region. Within this droplet, the system is in the normal phase. We have noted that the vortex core is spatially flat, and so we surmise that within the droplet, the space-time is flat. The asymptotic value of energy in figure 4 is precisely minus that contributed by the cosmological constant. Again, the size of the droplet is set by the value of the external H -flux, and the boundary conditions are AdS. Note that for a fixed cutoff, there is a critical field (given by $\hat{H} = 6a$) for which the entire spacetime is flat.

7 Conclusions

In this work we have presented in detail a simple toy model, the Nieh-Yan model, where torsion enters through the spacetime dependence of the coupling constant of the Nieh-Yan topological invariant. Although we have discussed the model directly in terms of torsion, it can classically be put into equivalent forms as either a massless pseudoscalar or a Kalb-Ramond field coupled to gravity. The model has an interesting and non-trivial holographic interpretation. In particular, we have shown that it possesses an exact bulk solution in Euclidean signature, termed the torsion vortex, having two asymptotically AdS₄ regimes, while the pseudoscalar acquires a kink profile. We have argued then that the holographic interpretation of this torsion vortex is a three-dimensional CFT with two distinct parity breaking vacua. Moreover, our bulk solution may imply that the deformation by a classically marginal pseudoscalar with a fixed coupling constant induces a transition between the two parity breaking vacua separated by a domain wall, which would be at infinity in the boundary components. [42] Remarkably, this qualitative behaviour is seen already in the three-dimensional Gross-Neveu model coupled to U(1) gauge fields. The economy of our bulk model does not allow a detailed identification of the bulk and boundary theories, nevertheless we believe that our results provide a strong base where an exact bulk/boundary dictionary for AdS₄/CFT₃ can be based. A further rather intriguing property of the torsion vortex is that it can be mapped into the standard Abrikosov vortex of superconductivity. Such a map identifies flat spacetime with a superconductor's normal phase, while AdS is identified with a superconducting phase. The cosmological constant would then measure the deviation from the "critical temperature". A phenomenon of vortex condensation is found, similar to the analogous case in type I superconductors.

The upshot of our results is that the torsional degrees of freedom of four dimensional gravity can provide holographic descriptions for a number of interesting properties of three dimensional critical systems. It would be interesting to extend our analysis to more elaborate models where more torsional degrees of freedom become dynamical. It is also of interest to discuss whether our simple model can be embedded into M-theory.

Acknowledgments

We thank F. Larsen, D. Minic and N. Papanicolaou for discussions. ACP is partially supported by the European RTN Program MRTN-CT-2004-512194, and RGL receives partial support from the US Department of Energy, under contract DE-FG02-91ER40709.

A The 3 + 1 split in gravity

We recall the 3+1 split formalism. We will refer to the radial coordinate as t , although its signature will be left arbitrary, and indicated by $\sigma_\perp = \pm 1$ ($\sigma_\perp = +1$ for AdS_4). The overall signature of the 4-manifold is denoted $\sigma = \det \eta$. In addition, we note the notation

$$*_4(e^a \wedge e^b \wedge e^c) = \epsilon^{abcd} e_d \quad (\text{A.1})$$

$$*_4 e_d = -\frac{\sigma}{6} \epsilon_{defg} e^e \wedge e^f \wedge e^g \quad (\text{A.2})$$

and for example

$$*_3 \tilde{e}^\alpha = \frac{1}{2} \epsilon^{\alpha\beta\gamma} \tilde{e}^\beta \wedge \tilde{e}^\gamma \quad (\text{A.3})$$

where \tilde{e}^α is defined as follows. As described in [17], we split the 1-forms via

$$e^0 = N dt \quad (\text{A.4})$$

$$e^\alpha = \tilde{e}^\alpha + N^\alpha dt \quad (\text{A.5})$$

$$\omega^0_\alpha = \sigma_\perp K_\alpha + q^0_\alpha dt \quad (\text{A.6})$$

$$\omega^\alpha_\beta = \sigma \epsilon^{\alpha\beta\gamma} (B^\gamma + Q^\gamma dt). \quad (\text{A.7})$$

We then find

$$T^\alpha = \tilde{T}^\alpha + dt \wedge \left\{ \dot{\tilde{e}}^\alpha - \tilde{d}N^\alpha + NK^\alpha - \sigma \epsilon^{\alpha\beta\gamma} Q^\beta e^\gamma - \sigma \epsilon^{\alpha\beta\gamma} N^\beta B^\gamma \right\} \quad (\text{A.8})$$

$$T^0 = \sigma_\perp K_\alpha \wedge \tilde{e}^\alpha + dt \wedge \left\{ -\tilde{d}N - \sigma_\perp N_\alpha K^\alpha + q^0_\beta \tilde{e}^\beta \right\} \quad (\text{A.9})$$

and we write

$$R^a_b = \tilde{R}^a_b + dt \wedge r^a_b \quad (\text{A.10})$$

$$\tilde{R}^0_\alpha = \sigma_\perp (\tilde{d}K_\alpha + K_\beta \wedge \tilde{\omega}^\beta_\alpha) \equiv \sigma_\perp (\tilde{D}K)_\alpha. \quad (\text{A.11})$$

$$\tilde{R}^\alpha_\beta = {}^{(3)}R^\alpha_\beta - \sigma_\perp K^\alpha \wedge K_\beta, \quad (\text{A.12})$$

with

$${}^{(3)}R^{\alpha\beta} = \sigma \left[\epsilon^{\alpha\beta\gamma} dB_\gamma - \sigma_\perp B^\alpha \wedge B^\beta \right]. \quad (\text{A.13})$$

and

$$2\epsilon_{\alpha\beta\gamma} r^{0\alpha} \wedge \tilde{e}^\beta \wedge \tilde{e}^\gamma = 2\sigma_\perp \epsilon_{\alpha\beta\gamma} \dot{K}^\alpha \wedge \tilde{e}^\beta \wedge \tilde{e}^\gamma + 4Q_\alpha K_\beta \wedge \tilde{e}^\beta \wedge \tilde{e}^\alpha + 4q^{0\alpha} \left[\epsilon_{\alpha\beta\gamma} \tilde{T}^\beta \wedge \tilde{e}^\gamma \right].$$

Including the Gibbons-Hawking term, which is of the form $2\sigma_\perp \int_{\partial M} \epsilon_{\alpha\beta\gamma} K^\alpha \wedge \tilde{e}^\beta \wedge \tilde{e}^\gamma$, we find

$$I_{\text{EH}} = \int dt \wedge \left\{ \dot{\tilde{e}}^\alpha \wedge (4\sigma_\perp \epsilon_{\alpha\beta\gamma} K^\gamma \wedge \tilde{e}^\beta) + 2N \epsilon_{\alpha\beta\gamma} \left({}^{(3)}R^{\alpha\beta} - \sigma_\perp K^\alpha \wedge K^\beta - \frac{\Lambda}{3} \tilde{e}^\alpha \wedge \tilde{e}^\beta \right) \wedge \tilde{e}^\gamma \right. \\ \left. - 4\sigma_\perp N^\alpha \epsilon_{\alpha\beta\gamma} (\tilde{D}K)^\beta \wedge \tilde{e}^\gamma - 4q^{0\alpha} \epsilon_{\alpha\beta\gamma} \tilde{T}^\beta \wedge \tilde{e}^\gamma + 4Q_\alpha \tilde{e}^\alpha \wedge K_\beta \wedge \tilde{e}^\beta \right\} \quad (\text{A.14})$$

Here, N and N^α are the usual Lagrange multiplier fields for the lapse and shift constraints, while $q_{0\alpha}$ and Q_α are Lagrange multipliers that, in the pure gravity case, set the torsion and the antisymmetric part of the extrinsic curvature to zero.

B Parity breaking in three dimensions

Consider the three dimensional Gross-Neveu model coupled to abelian gauge fields. The Euclidean action is¹⁰

$$I = - \int d^3x \left[\bar{\psi}^a (\not{\partial} - ie\mathcal{A}) \psi^a + \frac{G}{2N} (\bar{\psi}^a \psi^a)^2 + \frac{1}{4M} F_{\mu\nu} F_{\mu\nu} \right]. \quad (\text{B.1})$$

M is an UV mass scale. Introducing the usual Lagrange multiplier field σ , whose equation of motion is $\sigma = \frac{-2G}{N} \bar{\psi}^a \psi^a$ we can make the action quadratic in the fermions

$$I = - \int d^3x \left[\bar{\psi}^a (\not{\partial} + \sigma - ie\mathcal{A}) \psi^a - \frac{N}{2G} \sigma^2 - \frac{1}{4M} F^{\mu\nu} F_{\mu\nu} \right]. \quad (\text{B.2})$$

The model possesses two parity breaking vacua distinguished by the value of the pseudoscalar order parameter $\langle \sigma \rangle$. This is seen as follows: switching off the gauge fields momentarily one integrates over the fermions to produce a large- N effective action as

$$\mathcal{Z} = \int (\mathcal{D}\sigma) e^{N [\text{Tr} \log(\not{\partial} + \sigma) - \frac{1}{2G} \int d^3x \sigma^2]}. \quad (\text{B.3})$$

The path integral has a non-zero large- N extremum σ_* found by setting $\sigma = \sigma_* + \frac{1}{\sqrt{N}} \lambda$

$$\mathcal{Z} = \int (\mathcal{D}\lambda) e^{N [\text{Tr} \log(\not{\partial} + \sigma_*) - \frac{1}{2G} \int d^3x \sigma_*^2 + \frac{1}{\sqrt{N}} \{ \text{Tr} \frac{\lambda}{\not{\partial} + \sigma_*} - \frac{\sigma_*}{G} \int d^3x \lambda \} + O(1/N)]}. \quad (\text{B.4})$$

The term in the curly brackets is the gap equation. To study it one considers a uniform momentum cutoff Λ to obtain

$$\frac{1}{G} = \int^\Lambda \frac{d^3p}{(2\pi)^3} \frac{2}{p^2 + \sigma_*^2} = (\text{Tr}1) \left[\frac{\Lambda}{\pi^2} - \frac{|\sigma_*|}{\pi^2} \arctan \frac{\Lambda}{|\sigma_*|} \right]. \quad (\text{B.5})$$

Defining the critical coupling as

$$\frac{1}{G_*} = \frac{\Lambda}{\pi^2}, \quad (\text{B.6})$$

(B.5) possesses a non-zero solution for σ_* when $G > G_*$ given by

$$|\sigma_*| = \frac{2\pi}{G} \left(\frac{G}{G_*} - 1 \right) \equiv m. \quad (\text{B.7})$$

The two distinct parity breaking vacua then have

$$\sigma_* = -\frac{2G}{N} \langle \bar{\psi}^a \psi^a \rangle = \pm m. \quad (\text{B.8})$$

¹⁰We use $\bar{\psi}^i, \psi^i$ ($a = 1, 2, \dots, N$) two-component Dirac fermions. The γ -matrices are defined in terms of the usual Pauli matrices as $\gamma^i = \sigma^i$ $i = 1, 2, 3$.

Going back to (B.2) one can tune $G > G_*$ and start in any of the two parity breaking vacua. Suppose we start from $\sigma_* = +m$. To leading order in N we have

$$\mathcal{Z} = \int (\mathcal{D}A_\mu)(\mathcal{D}\bar{\psi}^a)(\mathcal{D}\psi^a) e^{\int d^3x [\bar{\psi}^a(\not{\partial} + m - ieA)\psi^a - \frac{N}{2G}m^2 + O(1/\sqrt{N}) - \frac{1}{4M}F^{\mu\nu}F_{\mu\nu}]} . \quad (\text{B.9})$$

As is well known [44, 45] for an odd number N of fermions the path integral (B.9) yields an effective action for the gauge fields which for low momenta is dominated by the Chern-Simons term i.e.

$$\mathcal{Z} \approx \int e^{S_{\text{CS}}} , \quad (\text{B.10})$$

with

$$S_{\text{CS}} = i \frac{ke^2}{4\pi} \int d^3x \epsilon^{\mu\nu\rho} A_\mu \partial_\nu A_\rho , \quad k = \frac{N}{2} . \quad (\text{B.11})$$

Had we started from the $\sigma_* = -m$ vacuum, we would have found again (B.10), however with $k = -\frac{N}{2}$, i.e. the vacuum with $\sigma_* = -m$ yields an effective Chern-Simons action with $k = -\frac{N}{2}$.

Consider now deforming the action (B.9) by the Chern-Simons term with a fixed coefficient as

$$\mathcal{Z} = \int (\mathcal{D}A_\mu)(\mathcal{D}\bar{\psi}^i)(\mathcal{D}\psi^i) e^{\int d^3x [\bar{\psi}^i(\not{\partial} + m - ieA)\psi^i - \frac{N}{2G}m^2 + O(1/\sqrt{N}) - \frac{1}{4M}F^{\mu\nu}F_{\mu\nu} - iq \int d^3x \epsilon^{\mu\nu\rho} A_\mu \partial_\nu A_\rho]} . \quad (\text{B.12})$$

If q is fixed to

$$q = \frac{Ne^2}{4\pi} , \quad (\text{B.13})$$

the effective action for the gauge fields resulting from the fermionic path integrals in (B.12) is going to be *exactly equal* the one obtained when we start at the $\sigma_* = -m$ vacuum. In other words, deforming the $\sigma_* = +m$ vacuum with a Chern-Simons term with a fixed coefficient is equivalent to being in the $\sigma_* = -m$ vacuum. This is exactly analogous to the holographic interpretation of our torsion vortex.

References

- [1] C.P. Herzog, P. Kovtun, S. Sachdev and D.T. Son, *Quantum critical transport, duality and M-theory*, *Phys. Rev. D* **75** (2007) 085020 [[hep-th/0701036](#)] [[SPIRES](#)].
- [2] S.A. Hartnoll, P.K. Kovtun, M. Muller and S. Sachdev, *Theory of the Nernst effect near quantum phase transitions in condensed matter and in dyonic black holes*, *Phys. Rev. B* **76** (2007) 144502 [[arXiv:0706.3215](#)] [[SPIRES](#)].
- [3] S.A. Hartnoll and P. Kovtun, *Hall conductivity from dyonic black holes*, *Phys. Rev. D* **76** (2007) 066001 [[arXiv:0704.1160](#)] [[SPIRES](#)].
- [4] S.A. Hartnoll and C.P. Herzog, *Ohm's Law at strong coupling: s duality and the cyclotron resonance*, *Phys. Rev. D* **76** (2007) 106012 [[arXiv:0706.3228](#)] [[SPIRES](#)].
- [5] E. Keski-Vakkuri and P. Kraus, *Quantum Hall Effect in AdS/CFT*, *JHEP* **09** (2008) 130 [[arXiv:0805.4643](#)] [[SPIRES](#)].

- [6] J.L. Davis, P. Kraus and A. Shah, *Gravity Dual of a Quantum Hall Plateau Transition*, *JHEP* **11** (2008) 020 [[arXiv:0809.1876](#)] [[SPIRES](#)].
- [7] S.A. Hartnoll, C.P. Herzog and G.T. Horowitz, *Building a Holographic Superconductor*, *Phys. Rev. Lett.* **101** (2008) 031601 [[arXiv:0803.3295](#)] [[SPIRES](#)].
- [8] D. Minic and J.J. Heremans, *High Temperature Superconductivity and Effective Gravity*, [arXiv:0804.2880](#) [[SPIRES](#)].
- [9] E. Nakano and W.-Y. Wen, *Critical magnetic field in a holographic superconductor*, *Phys. Rev. D* **78** (2008) 046004 [[arXiv:0804.3180](#)] [[SPIRES](#)].
- [10] T. Albash and C.V. Johnson, *A Holographic Superconductor in an External Magnetic Field*, *JHEP* **09** (2008) 121 [[arXiv:0804.3466](#)] [[SPIRES](#)].
- [11] S.S. Gubser and S.S. Pufu, *The gravity dual of a p-wave superconductor*, *JHEP* **11** (2008) 033 [[arXiv:0805.2960](#)] [[SPIRES](#)].
- [12] C.P. Herzog, P.K. Kovtun and D.T. Son, *Holographic model of superfluidity*, [arXiv:0809.4870](#) [[SPIRES](#)].
- [13] P. Basu, A. Mukherjee and H.-H. Shieh, *Supercurrent: vector Hair for an AdS Black Hole*, [arXiv:0809.4494](#) [[SPIRES](#)].
- [14] S.S. Gubser, *Breaking an Abelian gauge symmetry near a black hole horizon*, *Phys. Rev. D* **78** (2008) 065034 [[arXiv:0801.2977](#)] [[SPIRES](#)].
- [15] S.S. Gubser, *Colorful horizons with charge in anti-de Sitter space*, *Phys. Rev. Lett.* **101** (2008) 191601 [[arXiv:0803.3483](#)] [[SPIRES](#)].
- [16] S.S. Gubser and F.D. Rocha, *The gravity dual to a quantum critical point with spontaneous symmetry breaking*, *Phys. Rev. Lett.* **102** (2009) 061601 [[arXiv:0807.1737](#)] [[SPIRES](#)].
- [17] R.G. Leigh and A.C. Petkou, *Gravitational Duality Transformations on (A)dS₄*, *JHEP* **11** (2007) 079 [[arXiv:0704.0531](#)] [[SPIRES](#)].
- [18] S. de Haro and A.C. Petkou, *Holographic Aspects of Electric-Magnetic Dualities*, *J. Phys. Conf. Ser.* **110** (2008) 102003 [[arXiv:0710.0965](#)] [[SPIRES](#)].
- [19] D.S. Mansi, A.C. Petkou and G. Tagliabue, *Gravity in the 3+1-Split Formalism I: holography as an Initial Value Problem*, *Class. Quant. Grav.* **26** (2009) 045008 [[arXiv:0808.1212](#)] [[SPIRES](#)].
- [20] D.S. Mansi, A.C. Petkou and G. Tagliabue, *Gravity in the 3+1-Split Formalism II: self-Duality and the Emergence of the Gravitational Chern-Simons in the Boundary*, *Class. Quant. Grav.* **26** (2009) 045009 [[arXiv:0808.1213](#)] [[SPIRES](#)].
- [21] S. de Haro, *Dual Gravitons in AdS₄/CFT₃ and the Holographic Cotton Tensor*, *JHEP* **01** (2009) 042 [[arXiv:0808.2054](#)] [[SPIRES](#)].
- [22] J. Bagger and N. Lambert, *Modeling multiple M2's*, *Phys. Rev. D* **75** (2007) 045020 [[hep-th/0611108](#)] [[SPIRES](#)].
- [23] J. Bagger and N. Lambert, *Gauge Symmetry and Supersymmetry of Multiple M2-Branes*, *Phys. Rev. D* **77** (2008) 065008 [[arXiv:0711.0955](#)] [[SPIRES](#)].
- [24] J. Bagger and N. Lambert, *Comments On Multiple M2-branes*, *JHEP* **02** (2008) 105 [[arXiv:0712.3738](#)] [[SPIRES](#)].

- [25] A. Gustavsson, *Algebraic structures on parallel M2-branes*, *Nucl. Phys. B* **811** (2009) 66 [[arXiv:0709.1260](#)] [[SPIRES](#)].
- [26] K. Hosomichi, K.-M. Lee, S. Lee, S. Lee and J. Park, *N=4 Superconformal Chern-Simons Theories with Hyper and Twisted Hyper Multiplets*, *JHEP* **07** (2008) 091 [[arXiv:0805.3662](#)] [[SPIRES](#)].
- [27] O. Aharony, O. Bergman, D.L. Jafferis and J. Maldacena, *N=6 superconformal Chern-Simons-matter theories, M2-branes and their gravity duals*, *JHEP* **10** (2008) 091 [[arXiv:0806.1218](#)] [[SPIRES](#)].
- [28] I.L. Shapiro, *Physical aspects of the space-time torsion*, *Phys. Rept.* **357** (2002) 113 [[hep-th/0103093](#)] [[SPIRES](#)].
- [29] O. Chandía and J. Zanelli, *Torsional topological invariants (and their relevance for real life)*, [hep-th/9708138](#) [[SPIRES](#)].
- [30] L. Freidel, D. Minic and T. Takeuchi, *Quantum Gravity, Torsion, Parity Violation and all that*, *Phys. Rev. D* **72** (2005) 104002 [[hep-th/0507253](#)] [[SPIRES](#)].
- [31] S. Mercuri, *From the Einstein-Cartan to the Ashtekar-Barbero canonical constraints, passing through the Nieh-Yan functional*, *Phys. Rev. D* **77** (2008) 024036 [[arXiv:0708.0037](#)] [[SPIRES](#)].
- [32] F. Canfora, *Some solutions with torsion in Chern-Simons gravity and observable effects*, [arXiv:0706.3538](#) [[SPIRES](#)].
- [33] S.B. Giddings and A. Strominger, *Axion Induced Topology Change in Quantum Gravity and String Theory*, *Nucl. Phys. B* **306** (1988) 890 [[SPIRES](#)].
- [34] M. Gutperle and W. Sabra, *Instantons and wormholes in Minkowski and (A)dS spaces*, *Nucl. Phys. B* **647** (2002) 344 [[hep-th/0206153](#)] [[SPIRES](#)].
- [35] R. D'Auria and T. Regge, *Gravity theories with asymptotically flat instantons*, *Nucl. Phys. B* **195** (1982) 308 [[SPIRES](#)].
- [36] R. Jackiw, *Topological Investigations of Quantized Gauge Theories*, presented at *Les Houches Summer School on Theoretical Physics: Relativity Groups and Topology*, Les Houches, France, Jun 27 – Aug 4 1983.
- [37] R.G. Leigh and A.C. Petkou, *SL(2,Z) action on three-dimensional CFTs and holography*, *JHEP* **12** (2003) 020 [[hep-th/0309177](#)] [[SPIRES](#)].
- [38] G.W. Gibbons, M.B. Green and M.J. Perry, *Instantons and Seven-Branes in Type IIB Superstring Theory*, *Phys. Lett. B* **370** (1996) 37 [[hep-th/9511080](#)] [[SPIRES](#)].
- [39] V. Balasubramanian and P. Kraus, *A stress tensor for anti-de Sitter gravity*, *Commun. Math. Phys.* **208** (1999) 413 [[hep-th/9902121](#)] [[SPIRES](#)].
- [40] P. Kraus, F. Larsen and R. Siebelink, *The gravitational action in asymptotically AdS and flat spacetimes*, *Nucl. Phys. B* **563** (1999) 259 [[hep-th/9906127](#)] [[SPIRES](#)].
- [41] E. Witten, *SL(2,Z) action on three-dimensional conformal field theories with Abelian symmetry*, [hep-th/0307041](#) [[SPIRES](#)].
- [42] R.G. Leigh, N. Nguyen-Hoang and A.C. Petkou, to appear.

- [43] M. Tinkham, *Introduction to Superconductivity*, second edition, Dover Publications (1996) [ISBN: 0070648786].
- [44] A.J. Niemi and G.W. Semenoff, *Axial Anomaly Induced Fermion Fractionization and Effective Gauge Theory Actions in Odd Dimensional Space- Times*, *Phys. Rev. Lett.* **51** (1983) 2077 [SPIRES].
- [45] A.N. Redlich, *Parity Violation and Gauge Noninvariance of the Effective Gauge Field Action in Three-Dimensions*, *Phys. Rev. D* **29** (1984) 2366 [SPIRES].