

Lorentzian Lie (3-)algebra and toroidal compactification of M/string theory

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

JHEP03(2009)045

(<http://iopscience.iop.org/1126-6708/2009/03/045>)

[The Table of Contents](#) and [more related content](#) is available

Download details:

IP Address: 80.92.225.132

The article was downloaded on 03/04/2010 at 10:40

Please note that [terms and conditions apply](#).

Lorentzian Lie (3-)algebra and toroidal compactification of M/string theory

Pei-Ming Ho,^a Yutaka Matsuo^b and Shotaro Shiba^b

^a*Department of Physics, Center for Theoretical Sciences and Leung Center for Cosmology and Particle Astrophysics, National Taiwan University, Taipei 10617, Taiwan, R.O.C.*

^b*Department of Physics, Faculty of Science, University of Tokyo, Hongo 7-3-1, Bunkyo-ku, Tokyo 113-0033, Japan*

E-mail: pmho@phys.ntu.edu.tw, matsuo@phys.s.u-tokyo.ac.jp,
shiba@hep-th.phys.s.u-tokyo.ac.jp

ABSTRACT: We construct a class of Lie 3-algebras with an arbitrary number of pairs of generators with Lorentzian signature metric. Some examples are given and corresponding BLG models are studied. We show that such a system in general describes supersymmetric massive vector multiplets after the ghost fields are Higgsed. Simple systems with nontrivial interaction are realized by infinite dimensional Lie 3-algebras associated with the loop algebras. The massive fields are then naturally identified with the Kaluza-Klein modes by the toroidal compactification triggered by the ghost fields. For example, Dp -brane with an (infinite dimensional) affine Lie algebra symmetry \hat{g} can be identified with $D(p+1)$ -brane with gauge symmetry g .

KEYWORDS: D-branes, M-Theory

Contents

1	Introduction	1
2	Lorentzian BLG model	3
3	Analysis of Lie 3-algebra with two or more negative-norm generators	5
3.1	Fundamental identities	7
3.2	Lorentzian extension of Nambu bracket	7
3.3	Constraints from the fundamental identities for $F^{ijkl} = 0$	9
3.4	Solutions	10
3.5	Summary of the 3-algebra solutions	13
4	BLG model for Lorentzian 3-algebra with $J_{ab}^{ij} \neq 0$	14
4.1	Component expansion	15
4.2	Elimination of ghosts	16
4.3	Inclusion of $f_a^{ijk} \neq 0$	18
5	Application to toroidal compactification of M/string theories	22
5.1	D_p to $D(p+1)$ via Kac-Moody algebra	22
5.2	M2 to D_p via 3-algebra	26
5.3	M2 to M5 revisited	29
6	Conclusion and discussion	30

1 Introduction

Recently, Bagger, Lambert [1–3] and Gustavsson [4] constructed a three-dimensional superconformal field theory as a multiple-M2-brane world-volume theory in M-theory. This BLG model is characteristic of the novel feature that the gauge symmetry is based on a Lie 3-algebra, and thus various studies on this algebra have been undertaken [5, 6]. For the BLG model to work, the Lie 3-algebra needs to satisfy the fundamental identity (a generalization of Jacobi identity). If the positivity of the invariant metric is also imposed to avoid ghosts, the only non-trivial example of finite dimensional 3-algebra is \mathcal{A}_4 [7] and its direct sums.

If we relax the condition on dimensionality, Nambu-Poisson brackets give realizations of infinite dimensional Lie 3-algebra [8, 9]. The BLG model with this algebra realizes the world-volume theory of M5-branes in the C -field background on a 3-manifold where Nambu-Poisson bracket can be defined.

Similarly, when the requirement of a positive definite metric is given up, we also found physically meaningful models. Among the various examples, a Lie 3-algebra with a

negative-norm generator was constructed and was referred to as *Lorentzian* Lie 3-algebra.¹ The corresponding BLG model has ghosts, but they can be completely decoupled. It was realized that the inclusion of the Lorentzian generators is associated with the compactification of a spatial dimension, and this Lorentzian model reproduces the multiple- D2-brane world-volume theory in type IIA string theory.

In this paper, we study some generalizations of such Lorentzian 3-algebras for which ghost fields can still be decoupled. Such algebras have been considered extensively by de Medeiros *et. al* [11] when the number of Lorentzian pairs is two. Here we present more straightforward and explicit analysis in terms of the structure constants. We find it fruitful to consider generalizations with more Lorentzian pairs, as it gives us insight about how to circumvent the strict constraints from fundamental identities. We also study the BLG model associated with such 3-algebras. Our construction includes an interesting example which contains the massive Kaluza-Klein towers associated with additional compactified dimensions. This seems to be consistent with our expectation that adding Lorentzian pairs corresponds to additional compactifications. A typical feature of the generalized Lorentzian 3-algebra is indeed that we have a massive spectrum with $\mathcal{N} = 8$ SUSY in the BLG model, and we need an infinite dimensional realization to have nontrivial interacting models.

Our observation of the relation between the D-brane system with Lorentzian gauge symmetry and higher dimensional branes is not restricted to the context of BLG models. In fact, most of the examples considered here can be directly analyzed in the context of a Yang-Mills system whose gauge symmetry has Lorentzian signature. It was known that in some brane configurations (see for example [12]) we have to treat such an infinite dimensional gauge symmetry on D-branes. It was generally expected that the appearance of infinite dimensional symmetry should be related to closed string modes in a compactified space. However, the explicit analysis was not made because the Higgs mechanism which implement Kaluza-Klein mass was not known. Similar infinite dimensional symmetries were also studied in various contexts [13] in string/M theory and we hope that our method gives a simple direct interpretation to such systems.

This paper is organized as follows. In section 2, we first review the Lorentzian BLG model [14–16]. We describe the typical structure of the 3-algebra for which the removal of the ghost field [18, 19] is possible. In section 3, we give a detailed study of the constraint from the fundamental identity. Such study for two Lorentzian pairs was made in [11] but we generalize their result by considering an arbitrary number of Lorentzian pairs. We use a strategy to analyze the constraint for the structure constants directly. Although we do not claim that we could classify all possible algebras, we find a class of interesting 3-algebras through such analysis, with potential applications to string/M theory. We note that the many 3-algebras which we found can be realized by Lorentzian extension [14–16] of Lorentzian Lie algebras. It enables us to analyze some of the Lorentzian BLG models through gauge theories with Lorentzian Lie algebra symmetry. As we noted, such D-brane system is by itself an interesting object to study. In section 4, we derive the BLG

¹The Lie 3-algebra with zero-norm generators was also studied [10] to construct M2-brane model which produces the correct entropy $O(N^{3/2})$ in large N limit. It was suggested that we need 3-algebra instead of Lie algebra to have such scaling.

model associated with the simplest Lie 3-algebra with more than one Lorentzian pairs. We demonstrate that such system typically has massive vector fields where each gauge field absorbs two degrees of freedom from scalar fields. In section 5, we construct the BLG model (or super Yang-Mills theory) based on loop algebras which are the simplest nontrivial examples of generalized Lorentzian Lie (3-)algebra. Finally we comment that the description of M5-brane [8, 9] can be also regarded as the typical example of the compactification through the Lorentzian 3-algebra.

2 Lorentzian BLG model

In this section, we review the basic features of the *Lorentzian* BLG model [14–16]. The original BLG action for multiple M2-branes is

$$S = T_2 \int d^3x L = T_2 \int d^3x (L_X + L_\Psi + L_{\text{int}} + L_{\text{pot}} + L_{\text{CS}}), \quad (2.1)$$

$$L_X = -\frac{1}{2} \langle D_\mu X^I, D^\mu X^I \rangle, \quad (2.2)$$

$$L_\Psi = \frac{i}{2} \langle \bar{\Psi}, \Gamma^\mu D_\mu \Psi \rangle, \quad (2.3)$$

$$L_{\text{int}} = \frac{i}{4} \langle \bar{\Psi}, \Gamma_{IJ} [X^I, X^J, \Psi] \rangle, \quad (2.4)$$

$$L_{\text{pot}} = -\frac{1}{12} \langle [X^I, X^J, X^K], [X^I, X^J, X^K] \rangle, \quad (2.5)$$

$$L_{\text{CS}} = \frac{1}{2} f^{ABCD} A_{AB} \wedge dA_{CD} + \frac{1}{3} f^{CDA}{}_G f^{EFG}{}_B A_{AB} \wedge A_{CD} \wedge A_{EF}, \quad (2.6)$$

where T_2 is the M2-brane tension. The indices $\mu = 0, 1, 2$ specify the longitudinal directions of M2-branes; $I, J, K = 3, \dots, 10$ the transverse directions. The indices A, B, C, \dots denote components of Lie 3-algebra generators. The covariant derivative is

$$(D_\mu \Phi(x))_A = \partial_\mu \Phi_A - f^{CDB}{}_A A_{\mu CD}(x) \Phi_B \quad (2.7)$$

for $\Phi = X^I, \Psi$.

In order to define the BLG model action, the Lie 3-bracket

$$[T^A, T^B, T^C] = f^{ABC}{}_D T^D \quad (2.8)$$

for a Lie 3-algebra must satisfy the following constraints:

- Tri-linearity
- Skew symmetry
- Fundamental identity

$$f^{ABC}{}_F f^{FDE}{}_G + f^{ABD}{}_F f^{CFE}{}_G + f^{ABE}{}_F f^{CDF}{}_G = f^{CDE}{}_F f^{ABF}{}_G \quad (2.9)$$

- Invariant metric $\langle T^A, T^B \rangle = h^{AB}$:

$$f^{ABC}{}_E h^{ED} + f^{ABD}{}_E h^{CE} = 0. \quad (2.10)$$

The simplest Lorentzian Lie 3-algebra was defined as follows. Let \mathcal{G} be a given Lie algebra. We denote its generators as T^i , structure constants f^{ij}_k , and Killing form h^{ij} . Now we define a Lie 3-algebra whose generators are $T^A = \{u, v, T^i\}$ such that

$$\begin{aligned} [v, T^A, T^B] &= 0, & [u, T^i, T^j] &= f^{ij}_k T^k, & [T^i, T^j, T^k] &= -h^{kl} f^{ij}_l v, \\ \langle u, v \rangle &= 1, & \langle T^i, T^j \rangle &= h^{ij}, & \text{otherwise} &= 0. \end{aligned} \quad (2.11)$$

This 3-algebra satisfies the fundamental identities and the requirement of invariant metric, so we can use it as the gauge symmetry of BLG model. Since this algebra has a negative-norm generator $u - \alpha v$ (for $\alpha > 0$), BLG model has a ghost field. The mode expansion of the Langrangian becomes (up to total derivatives)

$$\begin{aligned} L &= \left\langle -\frac{1}{2}(\hat{D}_\mu \hat{X}^I - A'_\mu X_u^I)^2 + \frac{i}{2} \bar{\Psi} \Gamma^\mu \hat{D}_\mu \hat{\Psi} + \frac{i}{2} \bar{\Psi}_u \Gamma^\mu A'_\mu \hat{\Psi} \right. \\ &\quad + \frac{i}{2} \bar{\Psi} \Gamma_{IJ} X_u^I [\hat{X}^J, \hat{\Psi}] + \frac{1}{4} (X_u^K)^2 [\hat{X}^I, \hat{X}^J]^2 - \frac{1}{2} (X_u^I [\hat{X}^I, \hat{X}^J])^2 \\ &\quad \left. + \frac{1}{2} \epsilon^{\mu\nu\lambda} \hat{F}_{\mu\nu} A'_\lambda \right\rangle + L_{gh}, \end{aligned} \quad (2.12)$$

$$L_{gh} = - \left\langle \partial_\mu X_u^I A'_\mu \hat{X}^I + (\partial_\mu X_u^I) (\partial_\mu X_u^I) - \frac{i}{2} \bar{\Psi}_v \Gamma^\mu \partial_\mu \Psi_u \right\rangle, \quad (2.13)$$

where

$$\hat{D}_\mu \Phi := \partial_\mu \hat{\Phi} - [\hat{A}_\mu, \hat{\Phi}], \quad \hat{F}_{\mu\nu} := \partial_\mu \hat{A}_\nu - \partial_\nu \hat{A}_\mu - [\hat{A}_\mu, \hat{A}_\nu] \quad (2.14)$$

for $\Phi = X^I, \Psi$. As we see, fortunately, the ghost fields decouple, that is, they act only as Langrange multipliers. Their equations of motions are

$$\partial_\mu^2 X_u^I = 0, \quad \Gamma^\mu \partial_\mu \Psi_u = 0, \quad (2.15)$$

and we can set

$$X_u^I = \lambda^I := \lambda \delta_{10}^I, \quad \Psi_u^I = 0 \quad (2.16)$$

without breaking any supersymmetry or gauge symmetry [16]. This is motivated by the Higgs mechanism in BLG model first considered in [17]. The Lagrangian becomes, after integration over A' ,

$$L = -\frac{1}{2}(\hat{D}_\mu \hat{X}^I)^2 + \frac{i}{2} \bar{\Psi} \Gamma^\mu \hat{D}_\mu \hat{\Psi} + \frac{\lambda^2}{4} [\hat{X}^I, \hat{X}^J]^2 + \frac{i\lambda}{2} \bar{\Psi} \Gamma_I [X^I, \hat{\Psi}] - \frac{1}{4\lambda^2} \hat{F}_{\mu\nu}^2, \quad (2.17)$$

where $I, J = 3, \dots, 9$. This can be regarded as D2-branes theory in type IIA string theory which is the compactification of M-theory on a circle.

The origin of the decoupling of the ghost fields comes from the specific way that Lorentzian generators appear in the 3-algebra. Namely, the generator v is the center of the 3-algebra and u is not produced in any 3-commutators. This property ensures that the system is invariant under the translation of the scalar fields X_u^I . The decoupling of the

ghost fields can be made more rigorous [18, 19] by gauging this global symmetry. Namely by adding extra gauge fields C_μ, χ through

$$L_{\text{new}} = -\bar{\Psi}_u \chi + \partial^\mu X_u^I C_\mu^I, \quad (2.18)$$

we have an extra gauge symmetry:

$$\delta X_v^I = \Lambda^I, \quad \delta C_\mu^I = \partial_\mu \Lambda^I, \quad \delta \Psi_v = \eta, \quad \delta \chi = i\Gamma^\mu \partial_\mu \eta. \quad (2.19)$$

It enable us to put $X_v^I = \Psi_v = 0$. The equations of motion by variation of C_μ^I, χ give the assignment (2.16) correctly.

Another important feature of the Lorentzian BLG model is that the assignment of VEV to X_u^I triggers the compactification of 11 dimensional M-theory to 10 dimensional type IIA theory. The compactification radius of M-direction is given by [16]

$$\lambda = 2\pi R. \quad (2.20)$$

For various aspects of the Lorentzian model, see for example [20].

3 Analysis of Lie 3-algebra with two or more negative-norm generators

In the following, we consider some generalizations of the Lorentzian 3-algebra invented in [14–16] by adding pairs of generators with Lorentzian metric. Positive-norm generators are denoted as e^i ($i = 1, \dots, N$), and Lorentzian pairs as u_a, v_a ($a, b = 1, \dots, M$). We assume that the invariant metric for them is given by the following simple form

$$\langle e^i, e^j \rangle = \delta^{ij}, \quad \langle u_a, v_b \rangle = \delta_{ab}. \quad (3.1)$$

In terms of the four-tensor defined by

$$f^{ABCD} := f^{ABC}{}_E h^{ED}, \quad (3.2)$$

the invariance of the metric and the skew symmetry of the structure constant imply that the condition that this 4-tensor is anti-symmetric with respect to all indices.

We also assume that the generators v_a are in the center of the 3-algebra. This condition is necessary to apply the Higgs mechanism to get rid of the ghost fields as we have reviewed. In terms of the 4-tensor this condition is written as

$$f^{v_a BCD} = 0 \quad (3.3)$$

for arbitrary B, C, D . Therefore the index in the 4-tensor is limited to e^i and u_a . For the simplicity of the notation, we write i for e^i and a for u_a for indices of the 4-tensor, for example $f^{ijab} := f^{e^i e^j u_a u_b}$ and so on.

We note that there is some freedom in the choice of basis when keeping the metric (3.1) and the form of 4-tensor (3.3) invariant:

$$\tilde{e}^i = O_j^i e^j + P_a^i v^a, \quad \tilde{u}^a = Q_i^a e^i + R_b^a u^b + S_b^a v^b, \quad \tilde{v}^a = ((R^t)^{-1})_b^a v^b, \quad (3.4)$$

where

$$O^t O = 1, \quad Q = -R P^t O, \quad R^{-1} S + (R^{-1} S)^t = -P^t P. \quad (3.5)$$

The matrices O and R describe the usual rotations of the basis. The matrix P describes the mixing of the Lorentzian generators u_a, v_a with e^i .

We introduce some notation for the 4-tensor,

$$f^{ijkl} = F^{ijkl}, \quad f^{aijk} = f_a^{ijk}, \quad f^{abij} = J_{ab}^{ij}, \quad f^{abci} = K_{abc}^i, \quad f^{abcd} = L_{abcd}. \quad (3.6)$$

We rewrite the fundamental identity in terms of this notation below in section 3.1.

There are a few comments which can be made without detailed analysis:

- For lower M (i.e. smaller number of Lorentzian pairs (u_a, v_a)), some components of the structure constants (3.6) vanish identically due to the anti-symmetry of indices. For example, for $M = 1$, we need to put $J_{ab}^{ij} = K_{abc}^i = L_{abcd} = 0$. For $M = 2$, one may put J_{ab}^{ij} nonvanishing but we have to keep $K_{abc}^i = L_{abcd} = 0$ and so on.
- In the fundamental identity (3.11)–(3.24), there is no constraint on L_{abcd} . It comes from the fact that the contraction with respect to Lorentzian indices automatically vanishes due to the restriction of the structure constant (3.3). So it can take arbitrary value for $M \geq 4$. This term, however, is not physically relevant in BLG model, since they appear only in the interaction terms of the ghost fields which will be erased after Higgs mechanism.
- A constraint for F^{ijkl} (3.11) is identical to the fundamental identity of a 3-algebra with the structure constant F^{ijkl} . So if we assume positive definite metric for e^i , it automatically implies that F^{ijkl} is proportional to ϵ_{ijkl} or its direct sums [21].
- By a change of basis (3.4), various components of the structure constants (3.6) mix. For example, if we put $O = R = 1$ for simplicity and keep only the matrix P nontrivial (which implies $S = -\frac{1}{2}P^t P$), the structure constant in terms of the new basis $\{\tilde{e}^i, \tilde{u}^a, \tilde{v}^a\}$ are given as

$$\tilde{F}^{ijkl} = F^{ijkl}, \quad (3.7)$$

$$\tilde{f}_a^{jkl} = f_a^{jkl} + P_a^i F^{ijkl}, \quad (3.8)$$

$$\tilde{J}_{ab}^{ij} = J_{ab}^{ij} + P_a^k f_b^{ijk} - P_b^k f_a^{ijk} + F^{ijkl} P_a^k P_b^l, \quad (3.9)$$

$$\begin{aligned} \tilde{K}_{abc}^i &= K_{abc}^i + P_a^j J_{bc}^{ij} - P_b^j J_{ac}^{ij} + P_c^j J_{ab}^{ij} \\ &\quad + f_c^{ikl} P_a^k P_b^l - f_b^{ikl} P_a^k P_c^l + f_a^{ikl} P_b^k P_c^l + P_a^j P_b^k P_c^l F^{ijkl}. \end{aligned} \quad (3.10)$$

We will find that many solutions of the fundamental identities can indeed be identified with well-known 3-algebra after such redefinition of basis. In this sense, the classification of the Lorentzian 3-algebra has a character of cohomology, namely only solutions which can not reduce to known examples after all changes of basis give rise to physically new system.

In the following, we give a somewhat technical analysis of the fundamental identity (2.9). Solutions which we found are summarized in section 3.5. We do not claim that our analysis exhausts all the possible solutions. But as we will see in the later sections, they play an important physical role in string/M theory compactification.

3.1 Fundamental identities

We rewrite the fundamental identity (2.9) in the notation (3.6):

$$F^{ijkn} F^{nlmp} + F^{ijln} F^{knmp} + F^{ijmn} F^{klnp} - F^{klmn} F^{ijnp} = 0, \quad (3.11)$$

$$F^{ijkn} f_a^{nlm} + F^{ijln} f_a^{knm} + F^{ijmn} f_a^{kln} - F^{klmn} f_a^{ijn} = 0, \quad (3.12)$$

$$f_a^{ijn} F^{nklm} + f_a^{ikn} F^{jnlm} + f_a^{iln} F^{jknm} - f_a^{inm} F^{jklm} = 0, \quad (3.13)$$

$$(f_a^{ijn} f_b^{nkl} + f_a^{ikn} f_b^{jnl} + f_a^{iln} f_b^{jkn}) + F^{jklm} J_{ab}^{in} = 0, \quad (3.14)$$

$$J_{ab}^{im} F^{mjkl} + J_{ab}^{jm} F^{imkl} + J_{ab}^{km} F^{ijml} + J_{ab}^{lm} F^{ijkm} = 0, \quad (3.15)$$

$$(J_{ab}^{im} f_c^{mjk} + J_{ab}^{jm} f_c^{imk} + J_{ab}^{km} f_c^{ijm}) - F^{ijkm} K_{abc}^m = 0, \quad (3.16)$$

$$F^{ijkn} J_{ab}^{nl} - F^{ijln} J_{ab}^{nk} - f_a^{ijn} f_b^{nkl} + f_b^{ijn} f_a^{nkl} = 0, \quad (3.17)$$

$$(J_{ab}^{im} f_c^{mjk} - J_{ac}^{im} f_b^{mjk}) + (f_a^{ijm} J_{bc}^{mk} - f_a^{ikm} J_{bc}^{mj}) = 0, \quad (3.18)$$

$$-K_{abc}^l f_d^{lij} + K_{abd}^l f_c^{lij} + J_{ab}^{il} J_{cd}^{lj} - J_{cd}^{il} J_{ab}^{lj} = 0, \quad (3.19)$$

$$(f_a^{ikm} J_{bc}^{mi} + f_b^{jkm} J_{ca}^{mi} + f_c^{jkm} J_{ab}^{mi}) + K_{abc}^m F^{jkim} = 0, \quad (3.20)$$

$$(J_{ab}^{jl} J_{cd}^{li} + J_{ad}^{jl} J_{bc}^{li} - J_{ac}^{jl} J_{bd}^{li}) - f_c^{jil} K_{abd}^l = 0, \quad (3.21)$$

$$-J_{ab}^{ki} K_{cde}^k - J_{be}^{ki} K_{acd}^k + J_{ae}^{ki} K_{bcd}^k + J_{cd}^{ki} K_{abe}^k = 0, \quad (3.22)$$

$$f_a^{ijl} K_{bcd}^l - f_b^{ijl} K_{acd}^l + f_c^{ijl} K_{abd}^l - f_d^{ijl} K_{abc}^l = 0, \quad (3.23)$$

$$K_{abc}^i K_{def}^i - K_{ade}^i K_{bcf}^i + K_{acf}^i K_{bde}^i - K_{abf}^i K_{cde}^i = 0. \quad (3.24)$$

3.2 Lorentzian extension of Nambu bracket

Let us examine the case with $F^{ijkl} \neq 0$ first. As we already mentioned, eq. (3.11) implies that $F^{ijkl} \propto \epsilon_{ijkl}$ and its direct sum. So without losing generality, one may assume $N = 4$ and $F^{ijkl} = \epsilon_{ijkl}$ for the terms which include nontrivial contraction with F^{ijkl} .

Suppose $f_a^{ijk} \neq 0$ for some a . Then by the skew-symmetry of indices they can be written as $f_a^{ijk} = \epsilon_{ijkl} P_l^a$ for some P_l^a . This expression actually solves (3.12), (3.13). However, this form of f_a^{ijk} is exactly the same as the right hand side of (3.8). It implies that such f_a^{ijk} can be set to zero by a redefinition of basis.

Therefore, at least when the 3-algebra is finite dimensional, it is impossible to construct Lorentzian algebra with nontrivial $F^{ijkl} \neq 0$. The situation is totally different if the 3-algebra is infinite dimensional [8, 9] which is related to the description of M5-brane (for the various aspects of M5-brane in BLG context, see also [22] for example). The realization of the three-algebra was given as follows. We take \mathcal{N} as a compact three dimensional manifold where Nambu-Poisson bracket [23],

$$\{f_1, f_2, f_3\} = \sum_{a,b,c} \epsilon_{abc} \partial_a f_1 \partial_b f_2 \partial_c f_3 \quad (3.25)$$

is well defined. Namely \mathcal{N} is covered by the local coordinate patches where the coordinate transformation between the two patches keeps the 3-bracket (3.25) invariant. The simplest examples are T^3 and S^3 [8, 10]. If we take $\chi^i(y)$ as the basis of \mathcal{H} : the Hilbert space which consists of functions which are globally well-defined on \mathcal{N} , and one can choose a basis mutually orthonormal with respect to the inner product,

$$\langle \chi^i, \chi^j \rangle := \int_{\mathcal{N}} d^3y \chi^i(y) \chi^j(y) = \delta^{ij}. \tag{3.26}$$

It is known that the structure constant

$$F^{ijkl} = \langle \{ \chi^i, \chi^j, \chi^k \}, \chi^l \rangle \tag{3.27}$$

satisfies the fundamental identity (3.11).

We are going to show that it is possible to extend this 3-algebra with the additional generators with the Lorentzian signature. For simplicity, we consider the case $\mathcal{N} = T^3$. The Hilbert space \mathcal{H} is spanned by the periodic functions on T^3 . If we write the flat coordinates on T^3 as y^a ($a = 1, 2, 3$), where the periodicity is imposed as $y^a \sim y^a + p^a$, and $p^a \in \mathbf{Z}$. The basis of \mathcal{H} is then given by

$$\chi^{\vec{n}}(y) := e^{2\pi i n_a y^a}, \quad \vec{n} \in \mathbf{Z}^3, \tag{3.28}$$

with the invariant metric and the structure constant:

$$\langle \chi^{\vec{n}}, \chi^{\vec{m}} \rangle = \delta(\vec{n} + \vec{m}), \tag{3.29}$$

$$F^{\vec{n}\vec{m}\vec{l}\vec{p}} = (2\pi i)^3 \epsilon_{abc} n^a m^b l^c \delta(\vec{n} + \vec{m} + \vec{l} + \vec{p}). \tag{3.30}$$

The idea to extend the 3-algebra is to introduce the functions which are *not* well-defined on T^3 but the Nambu bracket among \mathcal{H} and these generators remains in \mathcal{H} . For T^3 , such generators are given by the functions $u_a = y^a$. The fundamental identity for the Nambu-bracket comes from the definition of derivative and it does not matter whether or not the functions in the bracket is well-defined globally. Therefore even if we include extra generators the analog of fundamental identity holds. More explicitly we define the extra structure constants as

$$f_a^{\vec{n}\vec{m}\vec{l}} := \langle \{ u^a, \chi^{\vec{n}}, \chi^{\vec{m}} \}, \chi^{\vec{l}} \rangle = (2\pi i)^2 \epsilon_{abc} n^b m^c \delta(\vec{n} + \vec{m} + \vec{l}), \tag{3.31}$$

$$J_{ab}^{\vec{n}\vec{m}} := \langle \{ u^a, u^b, \chi^{\vec{n}} \}, \chi^{\vec{m}} \rangle = (2\pi i) \epsilon_{abc} n^c \delta(\vec{n} + \vec{m}), \tag{3.32}$$

$$K_{abc}^{\vec{n}} := \langle \{ u^a, u^b, u^c \}, \chi^{\vec{n}} \rangle = \epsilon_{abc} \delta(\vec{n}). \tag{3.33}$$

It is not difficult to demonstrate explicitly that they satisfy all the fundamental identities (3.11)–(3.24).

We have to be careful in the treatment of the new generators. For example, the inner product (3.26) is not well-defined if the function is not globally well-defined on \mathcal{N} . The fact that the structure constants (3.30)–(3.33) satisfies the fundamental identities (3.11)–(3.24)

implies that we can define the inner product *abstractly* as (3.1). Namely we introduce extra generators v_a ($a = 1, 2, 3$) and define

$$\langle u_a, v_b \rangle = \delta_{ab}, \quad \langle u_a, \chi^{\vec{n}} \rangle = \langle v_a, \chi^{\vec{n}} \rangle = \langle u_a, u_b \rangle = \langle v_a, v_b \rangle = 0 \quad (3.34)$$

while keeping (3.29).

We also need to be careful in the definition of the three-bracket itself. The naive Nambu bracket needs to be modified to make the structure constant F^{ABCD} totally anti-symmetric in all four indices. This condition is broken in the original Nambu bracket after the introduction of the extra generators u^a . We have to come back to our original definition of 3-algebra where this symmetry is manifest. This implies the following redefinition of the 3-algebra:

$$[\chi^{\vec{n}}, \chi^{\vec{m}}, \chi^{\vec{l}}] = F^{\vec{n}\vec{m}\vec{l}} \bar{p} \chi^{\vec{p}} - f_a^{\vec{n}\vec{m}\vec{l}} v^a, \quad (3.35)$$

$$[u^a, \chi^{\vec{n}}, \chi^{\vec{m}}] = f_a^{\vec{n}\vec{m}} \bar{i} \chi^{\vec{i}} + J_{ab}^{\vec{n}\vec{m}} v^b, \quad (3.36)$$

$$[u^a, u^b, \chi^{\vec{n}}] = J_{ab\vec{m}}^{\vec{n}} \chi^{\vec{m}} - K_{abc}^{\vec{n}} v^c, \quad (3.37)$$

$$[u^a, u^b, u^c] = K_{abc\vec{n}} \chi^{\vec{n}}. \quad (3.38)$$

This 3-algebra may be regarded as the “central extension” of the Nambu-Poisson bracket. The additional factors which are proportional to v^a on the right hand side is necessary to make the metric invariant. One might worry if the fundamental identity may be violated by the redefinition of the algebra. In this example, fortunately this turns out not to be true. So we have a consistent 3-algebra with Lorentzian signature. It may be useful to repeat our emphasis that, although u^a was originally defined through ill-defined function y^a , we have to neglect this fact to define the metric and the 3-algebra.

While the 3-algebra (3.35)–(3.38) is new, we will see later in section 5.3 that the BLG model based on it turns out to be the same as the M5 models defined in [8, 9, 16] although it was not noticed explicitly. A glimpse of this fact appeared in §7 in [16] where a subalgebra of (3.35)–(3.38) appeared and the relation with the Lorentzian BLG model and M5 model was discussed. We will give more comments on this issue later in section 5.3.

It is straightforward to obtain similar Lorentzian extensions of Nambu-Poisson type Lie 3-algebras defined on different manifolds \mathcal{N} such as S^3 and $S^2 \times S^1$. So far, the only nontrivial Lie 3-algebra with positive definite metric are \mathcal{A}_4 and the Nambu-Poisson type 3-algebras. The examples we consider here would exhaust the Lorentzian extensions which can be obtained from them.

3.3 Constraints from the fundamental identities for $F^{ijkl} = 0$

In the following, we restrict ourselves to the case $F^{ijkl} = 0$. The fundamental identities (3.11)–(3.24) are now simplified to be the following:

$$f_a^{ni(j} f_b^{kl)n} = 0, \quad (3.39)$$

$$f_{(a}^{ijm} f_b^{mkl} = 0, \quad (3.40)$$

$$f_a^{ijm} f_b^{mkl} + f_a^{kim} f_b^{mjl} + f_b^{jkm} f_a^{mil} = 0, \quad (3.41)$$

$$J_{ab}^{l(i} f_c^{jk)l} = 0, \quad (3.42)$$

$$f_{(a}^{ijk} J_{bc)}^{kl} = 0, \quad (3.43)$$

$$J_{a(b}^{il} f_c^{ljk)} + J_{bc}^{l(j} f_a^{k)il} = 0, \quad (3.44)$$

$$2K_{ab(c}^k f_d^{kij)} = J_{ab}^{ik} J_{cd}^{kj} - J_{cd}^{ik} J_{ab}^{kj}, \quad (3.45)$$

$$f_a^{ijk} K_{bcd}^k = 3J_{a(b}^{ik} J_{cd)}^{kj}, \quad (3.46)$$

$$3K_{ab(c}^i J_{de)}^{ij} = K_{cde}^i J_{ab}^{ij}, \quad (3.47)$$

$$J_{a(b}^{ij} K_{cde)}^j = 0, \quad (3.48)$$

$$f_{(a}^{ijk} K_{bcd)}^k = 0, \quad (3.49)$$

$$3K_{ab(c}^i K_{de)f}^i = K_{cde}^i K_{abf}^i. \quad (3.50)$$

In the above, we used the notation that all indices in parentheses are fully antisymmetrized. For instance,

$$A_{a(b} B_{cd)e} := \frac{1}{6} (A_{ab} B_{cde} + A_{ac} B_{dbe} + A_{ad} B_{bce} - A_{ab} B_{dce} - A_{ac} B_{bde} - A_{ad} B_{cbe}). \quad (3.51)$$

The constraints above are not all independent. We can use (3.41) alone to derive (3.39) and (3.40) as follows. Taking (3.41) and replacing the indices as $(ijk) \rightarrow (jki)$ and $(ab) \rightarrow (ba)$ and subtracting the derived equation from (3.41), we get (3.40). It is also obvious that (3.40) and (3.41) implies (3.39).

Similarly, (3.44) can be easily derived from (3.42) and (3.43).

3.4 Solutions

In this subsection, we try to solve the fundamental identities displayed above and find a class of solutions.

First, a solution for (3.39) is to use a direct sum of Lie algebras $g = g_1 \oplus \dots \oplus g_n$, We divide the values of indices into n blocks $I = I_1 \cup \dots \cup I_n$ and let

$$f_a^{ijk} = \gamma_a^\alpha f_\alpha^{ijk}, \quad (3.52)$$

where f_α^{ijk} is defined by

$$f_\alpha^{ijk} = \begin{cases} f_{g_\alpha}^{ijk} & i, j, k \in I_\alpha, \\ 0 & \text{otherwise.} \end{cases} \quad (3.53)$$

Here $f_{g_\alpha}^{ijk}$ is the structure constant for g_α while γ_a^α is a real number.

Note that the number n does not have to equal M . It is possible to have some of the sets I_a empty. An example has $g = g_1$ and all $I_{a \neq 1}$ empty. In this case, for $\gamma_a^\alpha = \delta_a^\alpha$, we have $f_1^{ijk} = f_{g_1}^{ijk}$ and $f_a^{ijk} = 0$ for all $a \neq 1$.

If all the other components of the 3-algebra structure constant vanish, one obtains from (3.52) a set of solutions to the fundamental identity. The BLG model for this 3-algebra is not new, however. For each range of index, say I_α , we have

$$[e^i, e^j, e^k] = - \sum_a \gamma_a^\alpha v^a, \quad [u^a, e^i, e^j] = \sum_k \gamma_a^\alpha f_a^{ijk} e^k. \quad (3.54)$$

By a suitable rotation (3.4) with

$$v'^1 = \sum_a \gamma_a^\alpha v^a, \quad (3.55)$$

we always have

$$[e^i, e^j, e^k] = -v'^a, \quad [u'^a, e^i, e^j] = \delta_{a1} \sum_k f_a^{ijk} e^k. \quad (3.56)$$

Therefore it is reduced to the standard Lorentzian Lie 3-algebra for $M = 1$ after the restriction of indices to I_α .

In order to obtain something new, we have to allow other coefficients to be nonzero.

The simplest class of solutions can be found when $f_a^{ijk} = 0$ for $i, j, k \in I_a$. In this case, for this range I_a , arbitrary anti-symmetric matrix J^{ij} ($i, j \in I_a$) solves the constraints (this case is a special case of solutions in [11]). We will study the BLG model for this case in section 4. It demonstrates the essential feature that the supersymmetric system acquires mass proportional to eigenvalues of J . However, since we put $f_a^{ijk} = 0$, there is no interaction. In order to have the interacting system, we need nonvanishing f_a^{ijk} .

For simplicity, let us assume that there is a suitable basis of generators such that the solution (3.52) is simplified as

$$f_a^{ijk} = \begin{cases} f_a^{ijk} & i, j, k \in I_a, \\ 0 & \text{otherwise,} \end{cases} \quad (3.57)$$

where the indices are divided into n disjoint sets $I = I_1 \cup \dots \cup I_n$, and f_a^{ijk} is the structure constant for a Lie algebra g_a .

Starting with (3.57), we can solve all the constraints (3.39)–(3.50) as follows, while (3.57) already solves (3.39)–(3.41).

Eq. (3.43) is trivial if two of the indices a, b, c are identical. Assuming (3.57), eq. (3.43) imposes no constraint on J_{ab}^{ij} if $i \in I_a$ or $i \in I_b$. In general, if $f_c^{ijk} \neq 0$ for $c \neq a$ and $c \neq b$, then $J_{ab}^{ij} = 0$ if $i \in I_c$. Hence we consider the case

$$J_{ab}^{ij} \neq 0 \quad \text{only if} \quad i, j \in I_a \quad \text{or} \quad i, j \in I_b. \quad (3.58)$$

Eq. (3.44) is now trivial if all indices a, b, c are different. If two of the indices are the same, it is equivalent to (3.42).

According to (3.42), J_{ab} is a derivation for both Lie algebras g_a and g_b . A derivation \mathcal{D} is a map from g to g such that

$$\mathcal{D}([e^i, e^j]) = [\mathcal{D}(e^i), e^j] + [e^i, \mathcal{D}(e^j)]. \quad (3.59)$$

As a result of (3.42), one can define a derivations \mathcal{D}_{ab} by

$$\mathcal{D}_{ab}(e^i) = J_{ab}^{ij} e^j. \quad (3.60)$$

The simplest case is when J_{ab} corresponds to an inner automorphism, so

$$J_{ab}^{ij} = \Lambda_{ab}^k f_a^{ij} - \Lambda_{ba}^k f_b^{ij}, \quad (3.61)$$

where $\Lambda_{ab}^k = 0$ unless $k \in I_a$. (Note that the indices a, b are not summed over in (3.61).) In this case $\mathcal{D}_{ab}(\cdot) = [(\Lambda_{ab}^k - \Lambda_{ba}^k)e_k, \cdot]$. It will be more interesting if \mathcal{D}_{ab} instead corresponds to an infinitesimal outer automorphism (an *outer derivation*).²

If all indices a, b, c, d are all different, (3.45) is trivial due to (3.58). If $a = d \neq b \neq c$, (3.45) says that the Lie bracket $[J_{ab}, J_{ac}]$ is an inner automorphism. The solution of (3.45) is in general given by

$$K_{abc} := K_{abc}^i e^i = [\mathcal{D}_{ac}, \mathcal{D}_{bc}] + [\mathcal{D}_{ba}, \mathcal{D}_{ca}] + [\mathcal{D}_{cb}, \mathcal{D}_{ab}] + C_{abc}, \quad (3.62)$$

where the antisymmetric tensor $C_{abc} = C_{abc}^i$ is a central element in g . Since all derivations of a Lie algebra is always a Lie algebra, the Lie bracket $[\mathcal{D}_{ab}, \mathcal{D}_{cd}]$ satisfies the Jacobi identity.

For J_{ab} given by an inner automorphism (3.61), K_{abc}^i can be solved from (3.45) to be

$$K_{abc}^i = \Lambda_{ab}^j \Lambda_{ac}^k f_a^{ijk} + \Lambda_{bc}^j \Lambda_{ba}^k f_b^{ijk} + \Lambda_{ca}^j \Lambda_{cb}^k f_c^{ijk} + C_{abc}^i. \quad (3.63)$$

(Indices a, b, c are not summed over in this equation.) The term $\Lambda_{ab}^j \Lambda_{ac}^k f_a^{ijk}$ corresponds to the Lie bracket of the two automorphisms generated by Λ_{ab} and Λ_{ac} on g_a . However, the case of J_{ab} generating an inner automorphism is not interesting because J_{ab} and K_{abc}^i can be both set to zero after a change of basis (3.8), (3.9),

$$e'^i = e^i - \sum_b \Lambda_{ab}^i v^b \quad \text{for } i \in I_a, \quad (3.64)$$

$$u'_a = u_a - \sum_b \Lambda_{ba}^i e^i. \quad (3.65)$$

Therefore, in the following we will focus on the case when J_{ab} is an outer automorphism.

When all indices a, b, c, d, e are different, (3.47) can be easily satisfied if

$$C_{abc}^i = 0 \quad \text{unless } i \in I_a \cup I_b \cup I_c. \quad (3.66)$$

Together with (3.58), this implies that K_{abc}^i (3.62) vanishes unless $i \in I_a \cup I_b \cup I_c$.

Due to (3.58) and (3.66), eq. (3.47) is trivial if all indices a, b, c, d, e are different. If $e = a$, it is

$$K_{abc}^i J_{ad}^{ij} + K_{acd}^i J_{ab}^{ij} + K_{adb}^i J_{ac}^{ij} = 0. \quad (3.67)$$

One can then check that this follows from (3.62) and the constraint

$$\mathcal{D}_{ab}(C_{acd}) + \mathcal{D}_{ac}(C_{adb}) + \mathcal{D}_{ad}(C_{abc}) = 0 \quad (3.68)$$

²We have to keep in mind that the existence of such automorphisms is quite nontrivial. We will come back to this issue below.

as a result of the Jacobi identity of the Lie bracket of \mathcal{D}_{ab} 's. The same discussion applies to (3.46), (3.48), (3.49) and (3.50).

Before closing this subsection, let us comment on infinitesimal outer automorphisms. For finite dimensional Lie algebra, we have two examples. The first example is when the Lie algebra is Abelian, and any nontrivial linear map of the generators is an outer automorphism. The 2nd example is when the Lie algebra is that of matrices composed of upper triangular blocks

$$\begin{pmatrix} A & B \\ 0 & C \end{pmatrix}, \tag{3.69}$$

where A, B, C are $m \times m$, $m \times n$ and $n \times n$ matrices, respectively. An arbitrary scaling of the off-diagonal block B is an outer automorphism. In both of these examples, the coefficients of e_i in the expansion of X^I or Ψ do not participate in interactions in the BLG model, unless e^i is inert to the outer derivation. Hence the appearance of outer derivation in these cases is irrelevant to physics. A nontrivial example is found when g is an infinite dimensional Lie algebra. This example is studied in section 5.

3.5 Summary of the 3-algebra solutions

To summarize the result of our construction of a new 3-algebra, the general solution of the fundamental identity for our ansatz

$$[u_a, u_b, u_c] = K_{abc}^i e_i + L_{abcd} v^d, \tag{3.70}$$

$$[u_a, u_b, e^i] = J_{ab}^{ij} e_j - K_{abc}^i v^c, \tag{3.71}$$

$$[u_a, e^i, e^j] = J_{ab}^{ij} v^b + f_a^{ijk} e_k, \tag{3.72}$$

$$[e^i, e^j, e^k] = -f_a^{ijk} v^a, \tag{3.73}$$

is given by (3.57), (3.60) and (3.62), which are repeated here for the convenience of the reader,

$$f_a^{ijk} = \begin{cases} f_a^{ijk} & i, j, k \in I_a, \\ 0 & \text{otherwise,} \end{cases} \tag{3.74}$$

$$J_{ab}^{ij} e^j = \mathcal{D}_{ab}(e^i) \quad \text{for a derivation } \mathcal{D}_{ab}, \tag{3.75}$$

$$K_{abc} := K_{abc}^i e^i = [\mathcal{D}_{ac}, \mathcal{D}_{bc}] + [\mathcal{D}_{ba}, \mathcal{D}_{ca}] + [\mathcal{D}_{cb}, \mathcal{D}_{ab}] + C_{abc}, \tag{3.76}$$

where C_{abc} are central elements in g satisfying (3.66) and (3.68)

$$C_{abc}^i = 0 \quad \text{unless} \quad i \in I_a \cup I_b \cup I_c, \tag{3.77}$$

$$\mathcal{D}_{ab}(C_{acd}) + \mathcal{D}_{ac}(C_{adb}) + \mathcal{D}_{ad}(C_{abc}) = 0. \tag{3.78}$$

The nontrivial part of the metric is given by

$$\langle e^i, e^j \rangle = g^{ij}, \quad \langle u_a, v^b \rangle = \delta_a^b, \tag{3.79}$$

where g^{ij} is the Killing form of the Lie algebra g . Although we have assumed that g^{ij} is positive definite in the derivation above, it is obvious that the 3-algebra can be directly generalized to a generic Killing form which is not necessarily positive definite.

Compared with the 3-algebra discovered in [14, 16, 24], the 3-algebra constructed above contains more information. While e^i 's are generators of a Lie algebra $g = g_1 + \dots + g_n$, J_{ab} 's correspond to infinitesimal outer automorphisms (outer derivations), and K_{abc} encodes both the commutation relations among J_{ab} 's and choices of central elements in g .

Based on this analysis, we will analyze the BLG model for some examples of Lorentzian 3-algebras:

1. $M = 2$, $J_{ab}^{ij} = \epsilon_{ab} J^{ij}$ ($i, j = 1, \dots, n$), others = 0 (section 4.1): This is the simplest finite dimensional example where some character of the Lorentzian symmetry is displayed. Namely the BLG model defines the $\mathcal{N} = 8$ supersymmetric vector multiplets.
2. $M = 2$, $J_{ab}^{ij} = \epsilon_{ab} J^{ij}$, $f_1^{ijk} \neq 0$, others = 0 (section 4.3): This is the simplest nontrivial example which contains the interaction. We will present our result by studying the Yang-Mills system (2.17) where the gauge symmetry is defined by Lorentzian Lie algebra. This is possible since the 3-algebra can be written in the form (4.38). In such case, one can skip the discussion of eliminating one pair of ghost fields. It also illuminate the structure of the Yang-Mills system with Lorentzian Lie algebra.
3. Lie 3-algebra associated with affine Kac-Moody Lie algebra (section 5.1): This is the special case of above example where the Lorentzian Lie algebra is given by the affine Lie algebra. It illuminates how Kaluza-Klein mass is generated by the ghost fields.
4. Lie 3-algebra associated with general loop algebras (section 5.2): By this generalization we describe the compactification on general torus T^p with constant B field flux on it.
5. Lorentzian 3-algebra with $F^{ijkl} \neq 0$ (section 5.3): We give a brief explanation how construction of M5-brane [8, 9, 16] can be related to the Lorentzian 3-algebra (3.27), (3.31)–(3.33) and how the analysis in [8, 9, 16] can be related to the analysis in this paper.

4 BLG model for Lorentzian 3-algebra with $J_{ab}^{ij} \neq 0$

In this section, we describe generic features of BLG model when $J_{ab}^{ij} \neq 0$. We will first start with the “minimal” choice, namely we set other structure constants to zero,

$$F^{ijkl} = f_a^{ijk} = K_{abc}^i = 0. \tag{4.1}$$

We note that this is the simplest example considered in [11]. For this simplest choice, we see that BLG model gives rise to a free $\mathcal{N} = 8$ supersymmetric massive gauge theory after the Higgs mechanism is used to eliminate the negative-norm fields. After including other structure constants, we have an interacting theory. The direct analysis of interacting model from BLG model itself is somehow complicated and less illuminating, hence we will consider its equivalent version, the super Yang-Mills theory, in the following.

4.1 Component expansion

The BLG action is defined by (2.1)–(2.6), with the indices $A, B = (e^i, u_a, v_a)$. For simplicity, we first study the special case when the only nonvanishing part of the structure constant of the 3-algebra is

$$f^{u_a u_b i j} = \epsilon_{ab} J_{ij}, \quad (4.2)$$

where $a, b = 1, 2$ and $i, j = 1, \dots, n$.

As usual, we expand the relevant parts of the fields as

$$X^I = X_i^I e^i + X_a^I u_a + \underline{X}_a^I v_a, \quad (4.3)$$

$$\Psi = \Psi_i e^i + \Psi_a u_a + \underline{\Psi}_a v_a, \quad (4.4)$$

$$\sum_{ij} J_{ij} A_{\mu ij} =: A'_\mu, \quad (4.5)$$

$$A_{\mu i u_a} = -A_{\mu u_a i} =: \frac{1}{2} B_{\mu ia}, \quad (4.6)$$

$$A_{\mu u_a u_b} =: \frac{1}{2} C_\mu \epsilon_{ab}. \quad (4.7)$$

In terms of the modes, the covariant derivative (2.7) becomes

$$\begin{aligned} (D_\mu X^I)_i &= \partial_\mu X_i^I + \epsilon_{ab} J_{ij} B_{\mu ja} X_b^I + C_\mu J_{ij} X_j^I, \\ (D_\mu X^I)_{u_a} &= \partial_\mu X_a^I, \\ (D_\mu X^I)_{v_a} &= \partial_\mu \underline{X}_a^I + \epsilon_{ab} (A'_\mu X_b^I + J_{ij} B_{\mu bi} X_j^I), \end{aligned} \quad (4.8)$$

and similar expressions for Ψ .

The Chern-Simons action (2.6) can be rewritten in terms of the component gauge fields as

$$\begin{aligned} L_{\text{CS}} &= \epsilon^{\mu\nu\lambda} (A'_\mu \partial_\nu C_\lambda - \frac{1}{2} J_{ij} B_{\mu ia} (\partial_\nu B_{\lambda jb} + C_\nu J_{jk} B_{\lambda kb})) \\ &=: \epsilon^{\mu\nu\lambda} (A'_\mu \partial_\nu C_\lambda - \frac{1}{2} J_{ij} B_{\mu ia} \hat{D}_\nu B_{\lambda jb}). \end{aligned} \quad (4.9)$$

The gauge field A'_μ appears only in the Chern-Simons term. It does not participate in the dynamics but only imposes the flatness condition $\partial_{[\nu} C_{\mu]} = 0$ as the equation of motion.

In the original BLG model, the gauge symmetry transformations are

$$\begin{aligned} \delta X_A^I &= \tilde{\Lambda}^B{}_A X_B^I, \\ \delta \Psi_A &= \tilde{\Lambda}^B{}_A \Psi_B, \\ \delta \tilde{A}_\mu{}^B{}_A &= \partial_\mu \tilde{\Lambda}^B{}_A - \tilde{\Lambda}^B{}_C \tilde{A}_\mu{}^C{}_A + \tilde{A}_\mu{}^B{}_C \tilde{\Lambda}^C{}_A. \end{aligned} \quad (4.10)$$

We introduce the components of the gauge parameters as

$$\Lambda_{u_a u_b} =: \frac{1}{2} \gamma, \quad \Lambda_{i u_a} =: \frac{1}{2} \beta_{ia}, \quad J_{ij} \Lambda_{ij} =: \alpha. \quad (4.11)$$

Then the gauge symmetry transformation in terms of the modes becomes

$$\begin{aligned}\delta\Phi_i &= \epsilon_{ab}J_{ij}\beta_{jb}\Phi_a - \gamma J_{ij}\Phi_j, \\ \delta\Phi_a &= 0, \\ \delta\underline{\Phi}_a &= \alpha\epsilon_{ab}\Phi_b - J_{ij}\epsilon_{ab}\beta_{jb}\Phi_a,\end{aligned}\tag{4.12}$$

$$\begin{aligned}\delta A'_\mu &= \partial_\mu\alpha, \\ \delta B_{\mu ib} &= \partial_\mu\beta_{ib} + J_{ij}\beta_{jb}C_\mu - J_{ij}B_{\mu jb}\gamma, \\ \delta C_\mu &= \partial_\mu\gamma,\end{aligned}\tag{4.13}$$

where $\Phi = X^I, \Psi$. The gauge transformations for the gauge fields A'_μ, C_μ associated with the parameters α, γ are Abelian.

In the original BLG model, the supersymmetry transformations are

$$\begin{aligned}\delta X^I_A &= i\bar{\epsilon}\Gamma^I\Psi_A, \\ \delta\Psi_A &= D_\mu X^I_A\Gamma^\mu\Gamma^I\epsilon - \frac{1}{6}X^I_B X^J_C X^K_D f^{BCD}{}_A\Gamma^{IJK}\epsilon, \\ \delta\tilde{A}^B{}_{\mu A} &= i\bar{\epsilon}\Gamma_\mu\Gamma_I X^I_C\Psi_D f^{CDB}{}_A,\end{aligned}\tag{4.14}$$

where $\tilde{\Lambda}^B{}_A = \Lambda_{CD}f^{CDB}{}_A$. So, in terms of the components, the nontrivial parts of the supersymmetry transformation (namely, for Ψ and \tilde{A}_μ) become

$$\begin{aligned}\delta\Psi_i &= D_\mu X^I_i\Gamma^\mu\Gamma^I\epsilon + \frac{1}{2}\epsilon_{ab}J_{ij}X^I_a X^J_b X^K_j\Gamma^{IJK}\epsilon, \\ \delta\Psi_a &= \partial_\mu X^I_a\Gamma^\mu\Gamma^I\epsilon, \\ \delta\underline{\Psi}_a &= (D_\mu X)_{\nu a}\Gamma^\mu\Gamma^I\epsilon + \frac{1}{2}\epsilon_{ab}J_{ij}X^I_i X^J_j X^K_b\Gamma^{IJK}\epsilon, \\ \delta A'_\mu &= i\bar{\epsilon}\Gamma_\mu\Gamma_I X^I_j\Psi_j J_{ij}, \\ \delta B_{\mu ib} &= i\bar{\epsilon}\Gamma_\mu\Gamma_I X^I_{[i}\Psi_{b]}, \\ \delta C_\mu &= i\bar{\epsilon}\Gamma_\mu\Gamma_I X^I_a\Psi_b\epsilon_{ab}.\end{aligned}\tag{4.15}$$

By the definition of the BLG model, we obtain an $\mathcal{N} = 8$ SUSY system with ghost fields.

4.2 Elimination of ghosts

Variation of the Lagrangian by fields $\underline{X}^I, \underline{\Psi}$ gives

$$\partial^2 X^I_a = 0, \quad \Gamma^\mu\partial_\mu\Psi_a = 0.\tag{4.16}$$

As already reviewed in section 2, we solve them by the assignment [16] or the introduction of extra gauge symmetry [18, 19]:

$$X^I_a = \lambda^I_a, \quad \Psi_a = 0; \quad a = 1, 2.\tag{4.17}$$

It is clear that this choice does not break gauge symmetry nor supersymmetry, since the transformation of these fields is closed.

The Lagrangian is simplified considerably after inserting these VEV's:

$$\begin{aligned}
L = & -\frac{1}{2} \sum_i (\hat{D}_\mu X_i^I + \epsilon_{ab} J_{ij} B_{\mu ja} \lambda_b^I)^2 + \frac{i}{2} \bar{\Psi}_i \Gamma^\mu \hat{D}_\mu \Psi_i \\
& -\frac{1}{2} (J^2)_{ij} \Delta^2 X_i^I P_{IJ} X_j^J - \frac{i}{2} \Delta \bar{\Psi}_i \Gamma^\parallel J_{ij} \Psi_j \\
& + \epsilon^{\mu\nu\lambda} \left(A'_\mu \partial_\nu C_\lambda - \frac{1}{2} J_{ij} B_{\mu ia} \hat{D}_\nu B_{\lambda jb} \right), \tag{4.18}
\end{aligned}$$

where

$$\hat{D}_\mu \Phi_i := \partial_\mu \Phi_i + C_\mu J_{ij} \Phi_j, \tag{4.19}$$

$$\Delta^2 := |\vec{\lambda}_1|^2 |\vec{\lambda}_2|^2 - (\vec{\lambda}_1 \cdot \vec{\lambda}_2)^2, \tag{4.20}$$

$$P_{IJ} := \delta_{IJ} - \sum_{a=1,2} \lambda_a^I \pi_a^J, \tag{4.21}$$

$$\vec{\pi}_1 := \frac{1}{\Delta^2} (|\vec{\lambda}_2|^2 \vec{\lambda}_1 - (\vec{\lambda}_1 \cdot \vec{\lambda}_2) \vec{\lambda}_2), \quad \vec{\pi}_2 = (1 \leftrightarrow 2), \tag{4.22}$$

$$\Gamma^\parallel := \frac{1}{2\Delta} \Gamma_{IJ} \epsilon_{ab} \lambda_a^I \lambda_b^J, \quad (\Gamma^\parallel)^2 = 1. \tag{4.23}$$

The $\vec{\pi}_a$ ($a = 1, 2$) is the dual basis of $\vec{\lambda}_a$, namely $(\vec{\pi}_a, \vec{\lambda}_b) = \delta_{ab}$. The matrix P_{IJ} is a projector with codimension two which satisfies $P \vec{\lambda}_a = 0$ ($a = 1, 2$) and $P^2 = P$. The potential implies that six components of X^I become massive after putting VEV to X_a^I , while the two components in the plane spanned by $\vec{\lambda}_a$ remain massless. Actually the latter can be removed by redefinition of $B_{\mu ja}$,³

$$B'_{\mu ia} = B_{\mu ia} + \delta B_{\mu ia} \tag{4.24}$$

$$\delta B_{\mu ia} = \hat{D}_\mu \beta_{ia}, \quad \beta_{ia} := (J^{-1})_{ij} \epsilon_{ab} \pi_{bJ} X_j^J. \tag{4.25}$$

Since this redefinition takes the form of the gauge transformation for $B_{\mu ia}$, it does not change the form of Chern-Simons term. The gauge symmetry associated with β_{ia} is fixed by this manipulation and will not survive in the gauge fixed Lagrangian.

After this gauge transformation, the Chern-Simons Lagrangian L_{CS} remains the same while the kinetic term for X becomes

$$L_X = -\frac{1}{2} \sum_i (\hat{D}_\mu X^I)_i P_{IJ} (\hat{D}_\mu X^J)_i + \frac{1}{2} \sum B_{\mu ja} (J^2)_{jk} Q_{ab} B_{kb}^\mu, \tag{4.26}$$

$$Q_{ab} := \epsilon_{aa'} \epsilon_{bb'} (\vec{\lambda}_{a'}, \vec{\lambda}_{b'}). \tag{4.27}$$

The second term in L_X is the mass term for the gauge potential $B_{\mu ia}$.

To see the mass term for gauge fields more explicitly, we combine the relevant parts from L_{CS} and L_X to give the action for $B_{\mu ia}$,

$$L'_B = -\frac{1}{2} \epsilon^{\mu\nu\lambda} J_{ij} B_{\mu i1} (F_{\nu\lambda})_{j2} + \frac{1}{2} \sum B_{\mu ja} (J^2)_{jk} Q_{ab} B_{kb}^\mu, \tag{4.28}$$

³ If the matrix J_{ij} is not invertible, one can first decompose the linear space $\{e^i\}$ into two parts: the part on which J_{ij} is trivial and the part on which J is invertible. We focus our attention on the latter part.

where $(F_{\nu\lambda})_{ja} = (\hat{D}_\nu B_\lambda)_{ja} - (\hat{D}_\lambda B_\nu)_{ja}$. In the second term, we used partial integration. Since there are no derivatives of $B_{\mu i1}$, we integrate over them, and

$$L_B \rightarrow \frac{1}{Q_{11}} \left(-\frac{1}{4} \sum (F_{\nu\lambda})_{k2}^2 + \frac{1}{2} (J^2)_{ij} \Delta^2 B_{\mu i2} B^{\mu j2} \right). \quad (4.29)$$

The gauge symmetry is now reduced to Abelian transformations,

$$\delta\Phi_i = -\gamma J_{ij} \Phi_j, \quad \delta B_{\mu ia} = -J_{ij} B_{\mu ja} \gamma, \quad \delta A'_\mu = \partial_\mu \alpha, \quad \delta C_\mu = \partial_\mu \gamma. \quad (4.30)$$

They are, however, mostly trivial since the gauge field C_μ which appears in the covariant derivative is required to be flat by the equation of motion.

In the end, we find that we have n massive vector fields $B_{\mu i2}$, $6n$ massive scalars $P^{IJ} X_i^J =: (X^I)_i^I$ and $8n$ fermion fields Ψ_i . The mass spectrum of this supersymmetric system is given by

$$m^2 = \text{eigenvalues of } J^2 \Delta^2. \quad (4.31)$$

We note that this mass formula is invariant under $\text{SL}(2, \mathbf{R})$ transformations:

$$\vec{\lambda}'_a = g_{ab} \vec{\lambda}_b, \quad g_{ab} \in \text{SL}(2, \mathbf{R}). \quad (4.32)$$

This property is natural if we want to associate the system with T^2 compactification of M-theory, so that the mass spectrum corresponds to the Kaluza-Klein modes. This feature becomes more explicit in the example considered in the next section.

The original supersymmetry remains the same ($\mathcal{N} = 8$) after the Higgs mechanism,

$$\delta X^I_i = i\bar{\epsilon} P_{IJ} \Gamma^J \Psi_i, \quad (4.33)$$

$$\delta \Psi_i = D_\mu X_i^I P_{IJ} \Gamma^\mu \Gamma^J \epsilon + \Delta J_{ij} X_j^I P_{IJ} \Gamma^J \Gamma^\parallel \epsilon, \quad (4.34)$$

$$\delta A'_\mu = i\bar{\epsilon} \Gamma_\mu \Gamma^I P_{IJ} X_i^J \Psi_j J_{ij}, \quad (4.35)$$

$$\delta C_\mu = 0. \quad (4.36)$$

4.3 Inclusion of $f_a^{ijk} \neq 0$

By turning on $f_a^{ijk} \neq 0$, one may include interacting non-Abelian gauge symmetry in the action. For simplicity, we set

$$f_1^{ijk} \neq 0, \quad f_2^{ijk} = 0, \quad J^{ij} \neq 0. \quad (4.37)$$

In this case, we can rewrite it as

$$\begin{aligned} [u_1, T^A, T^B] &= f^{AB}{}_C T^C, \\ [v_1, T^A, T^B] &= 0, \\ [T^A, T^B, T^C] &= -h^{CD} f^{AB}{}_D v_1, \end{aligned} \quad (4.38)$$

where $A, B, \dots = \{u_2, v_2, i\}$, $f^{ijk} := f_1^{ijk}$ and $f^{u_2 ij} := J^{ij}$. This algebra is similar to that of [14–16], that is, a (u_1, v_1) -extension of Lie 3-algebra (2.11). A different point is that

this Lie 3-algebra $\{T^A\} = \{T^i, u_2, v_2\}$ has Lorentzian generators, while that of [14–16] is a standard (positive-definite) Lie algebra.

In this subsection, we denote generators of this algebra as $\{e^i, u, v\}$, instead of $\{T^i, u_2, v_2\}$. Then the metric (or Killing form) and structure constant is

$$\begin{aligned} \langle e^i, e^j \rangle &= \delta^{ij}, & \langle u, v \rangle &= 1; \\ f^{ijk}, \quad f^{uij} &= J^{ij}, & \text{otherwise} &= 0, \end{aligned} \quad (4.39)$$

where $i = 1, \dots, N$. The Jacobi identity is written as

$$f^{ijl} f^{lkm} + f^{jkl} f^{lim} + f^{kil} f^{ljm} = 0, \quad (4.40)$$

$$f^{ijl} J^{lk} + f^{jkl} J^{li} + f^{kil} J^{lj} = 0, \quad (4.41)$$

which are consistent with the fundamental identity for the Lie 3-algebra $\{T^i, u_{1,2}, v_{1,2}\}$. This is the simplest ‘‘Lorentzian extension’’ of Lie algebra,

$$[e^i, e^j] = f^{ij}_k e^k + J^{ij} v, \quad [u, e^i] = J^{ij} e^j. \quad (4.42)$$

This extension is trivial if J^{ij} is an inner automorphism

$$J^{ij} = f^{ij}_k \alpha^k \quad (4.43)$$

for some parameter α^k . One may then redefine the basis

$$e'^i = e^i + \alpha^i v, \quad u' = u - \alpha_i e^i, \quad v' = v, \quad (4.44)$$

such that the algebra becomes the direct sum of the original Lie algebra and Lorentzian pairs:

$$[e'^i, e'^j] = f^{ij}_k e'^k, \quad \text{other commutators} = 0; \quad (4.45)$$

$$\langle e'^i, e'^j \rangle = \delta^{ij}, \quad \langle u', v' \rangle = 1, \quad \text{other inner products} = 0. \quad (4.46)$$

In the following, we will focus on the nontrivial case where J gives an infinitesimal outer automorphism.

As we explained in section 2, (according to [16],) BLG model with Lorentzian Lie 3-algebra results in super Yang-Mills theory with Lie algebra. So, let us consider the Yang-Mills theory coupled with scalar fields X^I ($I = 1, \dots, n$) and spinor fields Ψ based on this extended algebra:

$$\begin{aligned} L &= -\frac{1}{2} \langle D_\mu X^I, D^\mu X^I \rangle + \frac{\lambda_1^2}{4} \langle [X^I, X^J], [X^I, X^J] \rangle \\ &\quad + \frac{i}{2} \langle \bar{\Psi}, \Gamma^\mu D_\mu \Psi \rangle + \frac{i\lambda_1}{2} \langle \bar{\Psi}, \Gamma_I [X^I, \Psi] \rangle - \frac{1}{4\lambda_1^2} \langle F_{\mu\nu} F^{\mu\nu} \rangle \end{aligned} \quad (4.47)$$

$$=: L_X + L_{\text{pot}} + L_\Psi + L_{\text{int}} + L_A, \quad (4.48)$$

where X^I takes the adjoint representation

$$X^I = X^I_i e^i + X^I_u u + X^I_v v, \quad (4.49)$$

$$\begin{aligned} (D_\mu X^I)_i &= \partial_\mu X^I_i - f^{jk}{}_i A_{\mu j} X^I_k - J^{ji} C_\mu X^I_j + J^{ji} A_{\mu j} X^I_u \\ &=: (\hat{D}_\mu X^I)_i + J^{ji} A_{\mu j} X^I_u, \end{aligned} \quad (4.50)$$

$$(D_\mu X^I)_u = \partial_\mu X^I_u, \quad (4.51)$$

$$(D_\mu X^I)_v = \partial_\mu X^I_v + J^{ij} A_{\mu i} X^I_j, \quad (4.52)$$

$$A_{\mu u} =: C_\mu, \quad A_{\mu v} =: B_\mu \quad (4.53)$$

and similar expressions for Ψ . The covariant derivative corresponding to the gauge symmetry generated by e^i should thus be defined as

$$\hat{D}_\mu = \partial_\mu - C_\mu \mathcal{D}_u - A_{\mu i} e^i, \quad (4.54)$$

where \mathcal{D}_u is the derivation defined by J :

$$\mathcal{D}_u(e^i) = J^{ij} e^j. \quad (4.55)$$

On the right hand side of (4.54), e^i is used to imply the adjoint action of e^i , namely $e^i(x) = [e^i, x]$. The gauge transformation is written as

$$\delta\Phi_i = f^{jk}{}_i \epsilon_j \Phi_k + J^{ki} \gamma \Phi_k - J^{ji} \epsilon_j \Phi_u, \quad (4.56)$$

$$\delta\Phi_u = 0, \quad (4.57)$$

$$\begin{aligned} \delta A_{\mu i} &= \partial_\mu \epsilon_i + f^{jk}{}_i \epsilon_j A_{\mu k} + J^{ki} \gamma A_{\mu k} - J^{ji} \epsilon_j C_\mu \\ &=: (\hat{D}_\mu \epsilon)_j + J^{ji} \gamma A_{\mu j} \end{aligned} \quad (4.58)$$

for $\Phi = X^I, \Psi$.

The kinetic term for X^I becomes

$$L_X = \frac{1}{2} (\hat{D}_\mu X^I_i + J^{ji} A_{\mu j} X^I_u)^2 + \partial^\mu X^I_u (\partial_\mu X^I_v - J^{ij} A_{\mu i} X^I_j). \quad (4.59)$$

The variation of X^I_v gives $\partial^2 X^I_u = 0$. So we take it as constant as before,

$$X^I_u = \lambda_2 \delta_{I1}. \quad (4.60)$$

After imposing this VEV,

$$L_X = -\frac{1}{2} \sum_{I'=2}^n (\hat{D}_\mu X^{I'})^2 - \frac{1}{2\lambda_1^2} F_{\mu u}^2, \quad (4.61)$$

where

$$F_{\mu u} := [\hat{D}_\mu, \hat{D}_u], \quad (4.62)$$

$$\hat{D}_u := \lambda_1 (\lambda_2 \mathcal{D}_u + X^1_i e^i). \quad (4.63)$$

We are thus led to interpret \mathcal{D}_u (or J) as the derivative of a certain noncommutative space in the direction of X_u . The situation here is reminiscent of the result of quotient conditions in the context of Matrix Models in dealing with orbifolds and orientifolds [25]. In analogy, since we have taken the VEV of X_u to be in the direction of X^1 , X^1_j plays the role of a gauge

potential and J_{ij} that of a covariant derivative on a noncommutative space, and thus \hat{D}_u mimics a covariant derivative. We will see in the next section that for the compactification on a circle, \hat{D}_u is indeed the covariant derivative in the compactified direction.

If we fix the gauge by $X_i^1 = 0$, the second term in (4.61) becomes

$$-\frac{\lambda_2^2}{2}(J^2)_{ij}A_{\mu i}A_{\mu j}. \tag{4.64}$$

This is the mass term for vector bosons.

The potential term is

$$L_{\text{pot}} = \frac{\lambda_1^2}{4} \sum_{I', J'=2}^n [X^{I'}, X^{J'}]^2 - \frac{1}{2} \sum_{J'=2}^n (\hat{D}_u X^{J'})^2. \tag{4.65}$$

If we gauge away X_i^1 using the gauge symmetry, the last term above is simply

$$-\frac{\lambda_1^2 \lambda_2^2}{2} \sum_{J'=2}^n (J^2)_{ij} X_i^{J'} X_j^{J'}. \tag{4.66}$$

It gives the mass term for $X^{J'}$ with exactly the same mass as eq. (4.31) with $\Delta = \lambda_1 \lambda_2$.⁴

The kinetic term for the gauge field becomes

$$-\frac{1}{4\lambda_1^2} \langle F_{\mu\nu}, F^{\mu\nu} \rangle = -\frac{1}{4\lambda_1^2} \{ (F_{\mu\nu i})^2 + F_{\mu\nu u} F^{\mu\nu}{}_v \}, \tag{4.67}$$

where

$$F_{\mu\nu i} = \partial_\mu A_{\nu i} - \partial_\nu A_{\mu i} - f^{jk}{}_i A_{\mu j} A_{\nu k} + J^{ij} (C_\mu A_{j\nu} - C_\nu A_{j\mu}), \tag{4.68}$$

$$F_{\mu\nu u} = \partial_\mu C_\nu - \partial_\nu C_\mu, \tag{4.69}$$

$$F_{\mu\nu v} = \partial_\mu B_\nu - \partial_\nu B_\mu - J^{ij} A_{\mu i} A_{\nu j}. \tag{4.70}$$

Variation of gauge field B_μ gives a free equation of motion for C_μ ,

$$\partial^\mu \partial_{[\mu} C_{\nu]} = 0. \tag{4.71}$$

If we start from the BLG action (4.18), we have slightly different Lagrangian,

$$L_{A'C} = \epsilon_{\mu\nu\lambda} A'_\mu \partial_\nu C_\lambda, \tag{4.72}$$

where A'_μ is an auxiliary field. From the viewpoint of the SYM, although it is not present from the beginning, one can add this term as a way to gauge the global symmetry of translation of C_μ , analogous to (2.18), where we gauged the translation of X_u and Ψ_u . By variation of A'_μ , C_μ becomes topological and pure gauge. Hence we should set C_ν to be a constant. It can be interpreted as the projection of the “ u ”-direction on the D-brane worldvolume, while X_u^I is the projection of the u -direction in the transverse directions.

⁴ If J is an inner automorphism, i.e. $J^{ki} = f^{jk}{}_i \mu_j$, one may shift $X_j^1 = -\mu_j$ to absorb J in X^1 . This is consistent with our comment above that J can be redefined away if it corresponds to an inner automorphism.

On the fermionic parts, after setting the VEV to $\Psi_u = 0$, they become

$$L_\Psi = \frac{i}{2} \langle \bar{\Psi}, \Gamma^\mu \hat{D}_\mu \Psi \rangle, \quad (4.73)$$

and

$$L_{\text{int}} = \sum_{I'=2}^n \frac{i\lambda_1}{2} \langle \bar{\Psi}_i, \Gamma_{I'} [X_j^{I'}, \Psi_k] \rangle + \frac{i}{2} \bar{\Psi}_i \Gamma_1 \hat{D}_u \Psi_i. \quad (4.74)$$

In the gauge $X_i^1 = 0$, the second term becomes the mass term for the fermions with their masses given by the matrix $\lambda_1 \lambda_2 J$.

To summarize, in the gauge $X^1 = 0$,

$$L = L_X + L_\Psi + L_{\text{int}} + L_A, \quad (4.75)$$

$$L_X = \sum_{I', J'=2}^n -\frac{1}{2} (\hat{D}_\mu X_i^{I'})^2 + \frac{\lambda_1^2 \lambda_2^2}{2} X_i^{I'} (J^2)_{ij} X_j^{I'}, \quad (4.76)$$

$$L_\Psi = \sum_{I'=2}^n \frac{i}{2} \bar{\Psi} \Gamma^\mu \hat{D}_\mu \Psi - \frac{\lambda_1 \lambda_2}{2} \bar{\Psi}_i (i\Gamma_1) J^{ij} \Psi_j, \quad (4.77)$$

$$L_{\text{int}} = \sum_{I', J'=2}^n \frac{\lambda_1^2}{4} [X^{I'}, X^{J'}]^2 + \frac{i\lambda_1}{2} \langle \bar{\Psi}, \Gamma_{I'} [X^{I'}, \Psi] \rangle, \quad (4.78)$$

$$L_A = -\frac{1}{4\lambda_1^2} F_{\mu\nu}^2 - \frac{\lambda_2^2}{2} (J^2)_{ij} A'_{\mu i} A'_{\mu j}, \quad (4.79)$$

which is of the form of a massive super Yang-Mills theory with the mass matrix $\lambda_1 \lambda_2 J_{ij}$.

5 Application to toroidal compactification of M/string theories

In this section we first consider an example of the general theory studied in section 4.3. We consider the Kac-Moody algebra as an example of the Lorentzian extension of a Lie algebra, and show in section 5.1 that the SYM theory with the gauge symmetry generated by the Kac-Moody algebra is equivalent to a SYM theory with a finite dimensional gauge group on a base space of higher dimensions. Finally, to be complete, in section 5.2 we consider the BLG model with the full 3-algebraic structure to describe M2-branes in flat spacetime compactified on a d -dimensional (noncommutative) torus with background fields.

5.1 D_p to $D(p+1)$ via Kac-Moody algebra

Before we go to the general discussion, let us briefly consider a simple case where Lie 3-algebra is defined as (2.11) where Lie algebra \mathcal{G} itself is a Lorentzian Lie algebra. The simplest example is when \mathcal{G} is the affine Lie algebra $\hat{\mathfrak{g}}$,

$$[u, T_m^a] = m T_m^a, \quad (5.1)$$

$$[T_m^a, T_n^b] = m v g^{ab} \delta_{m+n} + i f^{ab}{}_c T_{m+n}^c, \quad (5.2)$$

$$[v, u] = [v, T_m^a] = 0, \quad (5.3)$$

where $a, b, c = 1, \dots, \dim(g)$, $n, m \in \mathbf{Z}$ and g^{ab} is the Killing form of a compact Lie algebra g . This algebra has an invariant metric

$$\langle T_m^a, T_n^b \rangle = g^{ab} \delta_{m+n}, \quad \langle u, v \rangle = 1. \quad (5.4)$$

We note that the generator v is the center of Kac-Moody algebra and usually taken as a quantized c-number. Here we identify it as a nontrivial generator. On the other hand, the generator u gives the level (or $-L_0$ in the Virasoro algebra). While T_n^a has a positive-definite metric, the generators u, v have a negative-norm generator.⁵

We follow the method in section 4.3 where we use the super Yang-Mills system on D2 with gauge symmetry \hat{g} by using the Higgs mechanism for one Lorentzian pair.

In fact, the following analysis can be carried out for any Dp -brane system and provides a general mechanism of the gauge theory with affine gauge symmetry. What we are going to show is that the Dp -brane system whose gauge symmetry is \hat{g} can be identified with $D(p+1)$ -brane system with Lie algebra g .

If we start from the BLG model directly, we have a different perspective in which we will treat more general argument given in the next subsection.

We start from the action

$$\begin{aligned} L = & -\frac{1}{4\lambda^2} \langle F_{\mu\nu}, F^{\mu\nu} \rangle - \frac{1}{2} \langle D_\mu X^I, D^\mu X^I \rangle + \frac{\lambda^2}{4} \langle [X^I, X^J], [X^I, X^J] \rangle \\ & + \frac{i}{2} \bar{\Psi} \Gamma^\mu D_\mu \Psi + \frac{i\lambda}{2} \bar{\Psi} \Gamma_I [X^I, \Psi], \end{aligned} \quad (5.5)$$

where $X^I(x)$ ($I = 1, \dots, D$) are the scalar field and $\Psi(x)$ is the spinor field. Both are in the adjoint representation of g . The world volume index is given as $\mu, \nu = 0, \dots, p$. The covariant derivative and the field strength are defined (only in this subsection) as

$$D_\mu \Phi := \partial_\mu \Phi - i[A_\mu, \Phi], \quad F_{\mu\nu} := \partial_\mu A_\nu - \partial_\nu A_\mu - i[A_\mu, A_\nu] \quad (5.6)$$

for $\Phi = X^I, \Psi$. The convention here differs from that in section 2; here A_μ is Hermitian. We consider the following component expansion,

$$A_\mu = A_{\mu(a,n)} T_n^a + B_\mu v + C_\mu u, \quad (5.7)$$

$$X^I = X_{(a,n)}^I T_n^a + X_u^I u + X_v^I v, \quad (5.8)$$

$$\Psi = \Psi_{(a,n)} T_n^a + \Psi_u u + \Psi_v v. \quad (5.9)$$

Various components of the covariant derivative and the field strength are given as

$$\begin{aligned} (D_\mu X^I)_{(an)} &= \partial_\mu X_{an}^I + f^{bc}{}_a \sum_m A_{\mu(b,m)} X_{(c,n-m)}^I - n C_\mu X_{(a,n)}^I \\ &\quad + i n A_{\mu(a,n)} X_u^I \\ &=: (\hat{D}_\mu X^I)_{(a,n)} + i n A_{\mu(a,n)} X_u^I, \end{aligned} \quad (5.10)$$

$$(D_\mu X^I)_u = \partial_\mu X_u^I, \quad (5.11)$$

⁵We note that a different type of Lie 3-algebra based on Kac-Moody symmetry was obtained in [28].

$$(D_\mu X^I)_v = \partial_\mu X_v^I + \sum_m img^{ab} A_{\mu(a,m)} X_{(b,-m)}^I, \quad (5.12)$$

$$(F_{\mu\nu})_{(a,n)} = \partial_\mu A_{\nu(a,n)} - \partial_\nu A_{\mu(a,n)} + f^{bc}{}_a \sum_m A_{\mu(b,m)} A_{\nu(c,n-m)}, \quad (5.13)$$

$$(F_{\mu\nu})_u = \partial_\mu C_\nu - \partial_\nu C_\mu, \quad (5.14)$$

$$(F_{\mu\nu})_v = \partial_\mu B_\nu - \partial_\nu B_\mu + \sum_m img^{ab} A_{\mu(a,m)} A_{\nu(b,-m)}, \quad (5.15)$$

and similar expressions for $D_\mu \Psi$. From the kinetic part for u, v components, the equations of motion for X_u, Ψ_u and C_μ are free,

$$\partial^\mu \partial_\mu X_u^I = \Gamma^\mu \partial_\mu \Psi_u = \partial^\mu (\partial_\mu C_\nu - \partial_\nu C_\mu) = 0. \quad (5.16)$$

We fix their values as

$$X_u^I = \text{const.} =: \lambda' \delta^{ID}, \quad \Psi_u = 0, \quad \partial_\mu C_\nu - \partial_\nu C_\mu = 0. \quad (5.17)$$

For the first two relations, we need to use the method [18, 19] as reviewed in section 2. We need to introduce the extra gauge symmetry as commented in the paragraph after (4.71) to derive the last one. For general world volume dimensions, the additional action is

$$S_{\text{additional}} = -\frac{1}{4\lambda^2} D_{\mu\nu} (\partial_\mu C_\nu - \partial_\nu C_\mu), \quad (5.18)$$

where $D_{\mu\nu}$ is a new field. It gives rise to a new gauge symmetry,

$$\delta D_{\mu\nu} = \partial_\mu \Xi_\nu - \partial_\nu \Xi_\mu, \quad \delta B_\mu = -\Xi_\mu \quad (5.19)$$

by which we can gauge fix $B_\mu = 0$. The equation of motion by the variation of $D_{\mu\nu}$ gives the flatness condition of C_μ .

Since the gauge field C_μ is essentially flat, we can ignore it for simplicity (namely set $C_\mu = 0$). After this, the ghost fields $C_\mu, B_\mu, X_u^I, X_v^I, \Psi_u, \Psi_v$ disappear from the action, and the system is unitary.

We identify the infinite components of the scalar, spinor and gauge fields as fields in $p+2$ dimensions,

$$\begin{aligned} \tilde{X}_a^I(x, y) &= \sum_m X_{(a,n)}^I(x) e^{-iny/R}, & \tilde{\Psi}_a(x, y) &= \sum_m \Psi_{(a,n)}(x) e^{-iny/R}, \\ \tilde{A}_{\mu a}(x, y) &= \sum_m A_{\mu(a,n)}(x) e^{-iny/R}, \end{aligned} \quad (5.20)$$

where an extra coordinate y is introduced to parametrize S^1 with the radius R . We also rename

$$\tilde{X}_a^D(x, y) \rightarrow \frac{1}{\lambda} \tilde{A}_{ya}(x, y). \quad (5.21)$$

The kinetic term of the scalar field X^I can be rewritten as

$$-\frac{1}{2} \int \frac{dy}{2\pi R} \left[\sum_{I=1}^{D-1} (\partial_\mu \tilde{X}_a^I - f^{bc}{}_a \tilde{A}_{\mu b} \tilde{X}_c^I)^2 + \frac{1}{\lambda^2} \tilde{F}_{\mu ya}^2 \right], \quad (5.22)$$

where

$$\tilde{F}_{\mu ya} := \partial_\mu \tilde{A}_{ya} - \partial_y \tilde{A}_{\mu a} + f^{bc}{}_a \tilde{A}_{\mu b} \tilde{A}_{yc}. \quad (5.23)$$

Here the second term can be produced properly if we identify

$$R = 1/\lambda\lambda'. \quad (5.24)$$

This relation seems strange if we compare with (2.20). It can be fixed by applying the T-duality transformation [29].

The second term in (5.22), when combined with the kinetic term for gauge fields, properly reproduces the kinetic term for $p + 2$ dimensional world volume. The Kaluza-Klein mass from the compactification radius (5.24) is $n\lambda\lambda'$ which is consistent with the result (4.31).

Similarly, we can rewrite the commutator term,

$$\begin{aligned} \frac{\lambda^2}{4} \sum_{I,J=1}^D \langle [X^I, X^J], [X^I, X^J] \rangle &= \frac{\lambda^2}{4} \sum_{I,J=1}^{D-1} \int \frac{dy}{2\pi R} \langle [\tilde{X}^I, \tilde{X}^J], [\tilde{X}^I, \tilde{X}^J] \rangle \\ &\quad - \frac{1}{2} \sum_{I=1}^{D-1} \int \frac{dy}{2\pi R} (D_y \tilde{X}^I)^2. \end{aligned} \quad (5.25)$$

Here again the second term can be combined with the kinetic term for X^I to give the kinetic energy on $p + 2$ dimensional world volume.

Finally, we can rewrite the interaction term,

$$\frac{i\lambda}{2} \sum_{I=1}^D \tilde{\Psi} \Gamma_I [X^I, \Psi] = \frac{i\lambda}{2} \sum_{I=1}^{D-1} \int \frac{dy}{2\pi R} \tilde{\Psi} \Gamma_I [\tilde{X}^I, \tilde{\Psi}] + \frac{i}{2} \int \frac{dy}{2\pi R} \tilde{\Psi} \Gamma^y D_y \tilde{\Psi}. \quad (5.26)$$

Here, this time, the second term can be combined with the kinetic term for Ψ .⁶ In the end, the Lagrangian thus obtained is the same as the original Lagrangian (5.5) except that we change the dimension parameter $D \rightarrow D - 1$ and $p \rightarrow p + 1$ and the gauge symmetry $\mathcal{G} = \hat{g} \rightarrow g$:

$$L = L_A + L_X + L_\Psi + L_{\text{pot}} + L_{\text{int}}, \quad (5.27)$$

$$L_A = -\frac{1}{4\lambda^2} \int \frac{dy}{2\pi R} (\tilde{F}_{\mu\nu}^2 + 2\tilde{F}_{\mu y}^2), \quad (5.28)$$

$$L_X = -\frac{1}{2} \int \frac{dy}{2\pi R} \sum_{I=1}^{D-1} \left[(D_\mu \tilde{X}^I)^2 + (D_y \tilde{X}^I)^2 \right], \quad (5.29)$$

$$L_\Psi = \frac{i}{2} \int \frac{dy}{2\pi R} \tilde{\Psi} (\Gamma^\mu D_\mu + \Gamma^y D_y) \tilde{\Psi}, \quad (5.30)$$

⁶We should notice the definition of Γ_μ and Γ_I here. We see from the kinetic term of Ψ in the Lagrangian (5.5) that Γ_μ satisfies $\{\Gamma_\mu, \Gamma_\nu\} = \text{diag.}(+ - \dots -)$. On the other hand, Γ_I should satisfy $\{\Gamma_I, \Gamma_J\} = \delta_{IJ}$ as usual. So we choose $\Gamma^D = -i\Gamma^y$ and obtain (5.26).

$$L_{\text{pot}} = \frac{\lambda^2}{4} \sum_{I,J=1}^{D-1} \int \frac{dy}{2\pi R} \langle [\tilde{X}^I, \tilde{X}^J], [\tilde{X}^I, \tilde{X}^J] \rangle, \tag{5.31}$$

$$L_{\text{int}} = \frac{i\lambda}{2} \sum_{I=1}^{D-1} \int \frac{dy}{2\pi R} \tilde{\Psi} \Gamma_I [\tilde{X}^I, \tilde{\Psi}]. \tag{5.32}$$

5.2 M2 to Dp via 3-algebra

Here we consider essentially the same physical system as the previous subsection, namely the compactification of D2-branes on torus, but we start from the BLG model for multiple M2-branes corresponding to an example of the Lie 3-algebra summarized in (3.70)–(3.79). The formulation here will be more general than above as we will turn on noncommutativity and a gauge field background.

We start by defining a Lie algebra g_0 with generators $T_{\vec{m}}^i$, structure constants

$$f^{(i\vec{l})(j\vec{m})(k\vec{n})} = f_{\vec{l}\vec{m}}^{ijk} \delta_{\vec{0}}^{\vec{l}+\vec{m}+\vec{n}}, \tag{5.33}$$

and metric

$$g^{(i\vec{m})(j\vec{n})} = g_{\vec{m}}^{ij} \delta_{\vec{0}}^{\vec{m}+\vec{n}}. \tag{5.34}$$

Here \vec{m} is a d -dimensional vector of integers.

The simplest example of g_0 has

$$T_{\vec{m}}^i = T^i e^{i\vec{m}\cdot\vec{x}}, \tag{5.35}$$

where T^i is the generator for $U(N)$ and \vec{x} is the coordinate on a d -dimensional torus. More generally, one can consider a twisted bundle on a noncommutative torus T_θ^d . In this case

$$T_{\vec{m}}^i = T^i Z_1^{m_1} \dots Z_d^{m_d}, \tag{5.36}$$

where T^i denotes a generator of the $U(N)$ gauge group, and Z_i are noncommutative algebraic elements satisfying

$$Z_i Z_j = e^{i\theta'_{ij}} Z_j Z_i. \tag{5.37}$$

The parameter θ' is in general not the same as the noncommutative parameter θ of the noncommutative torus T_θ^d , and it depends on the rank of the gauge group and its twisting. Z_i maps a section of the twisted bundle to another section. For the trivial bundle, $Z_i = e^{ix_i}$ and (5.36) reduces to (5.35). The case of $d = 2$ was studied in [26, 27]. It is straightforward to generalize it to arbitrary dimensions.

Since the structure constant (5.33) of g_0 has the property

$$f^{(i\vec{l})(j\vec{m})(k\vec{n})} \propto \delta_{\vec{0}}^{\vec{l}+\vec{m}+\vec{n}}. \tag{5.38}$$

g_0 has derivations

$$J_{0a}^{(i\vec{m})(j\vec{n})} = m_a \delta^{(i\vec{m})(j\vec{n})}. \tag{5.39}$$

Now we consider the 3-algebra with the underlying Lie algebra $g = g_0$ and $I_{a \neq 0}$'s empty. We take $J_{ab} = 0$ if $a, b \neq 0$, and J_{0a} given by (5.39). It follows that the first 3 terms in (3.62) vanish, hence

$$K_{abc}^{(i\vec{m})} = \delta_0^i \delta_0^{\vec{m}} C_{abc}, \tag{5.40}$$

assuming that T^0 is the identity of $U(N)$, so that $T^{(0\vec{0})}$ is the identity of g_0 . In the following we choose

$$K_{0ab}^{i\vec{m}} = \delta_0^i \delta_0^{\vec{m}} C_{ab}, \tag{5.41}$$

$$K_{abc}^{i\vec{m}} = 0, \quad \text{otherwise.} \tag{5.42}$$

It will be shown below that the constants C_{ab} corresponds to a nontrivial gauge field background.

The 3-algebra is defined by the 3-brackets

$$[u_0, u_a, u_b] = C_{ab} T_0^0 + L_{0abc} v^c, \tag{5.43}$$

$$[u_0, u_a, T_{\vec{m}}^i] = m_a T_{\vec{m}}^i - \delta_0^i \delta_0^{\vec{m}} C_{ab} v^b, \tag{5.44}$$

$$[u_0, T_{\vec{m}}^i, T_{\vec{n}}^j] = m_a g_{\vec{m}}^{ij} \delta_{\vec{m}+\vec{n}}^0 v^a + f_{\vec{m}\vec{n}}^{ijk} T_{\vec{m}+\vec{n}}^k, \tag{5.45}$$

$$[T_{\vec{l}}^i, T_{\vec{m}}^j, T_{\vec{n}}^k] = -f_{\vec{l}\vec{m}}^{ijk} \delta_{\vec{l}+\vec{m}+\vec{n}}^0 v^0, \tag{5.46}$$

where $a, b, c = 0, 1, 2, \dots, d$ and $i, j, k = 1, 2, \dots, N$. (Note that we have changed the range of indices a, b, c from the convention used above.)

This 3-algebra is actually precisely the Lorentzian algebra discovered in [14, 16, 24] constructed from the (multiple) loop algebra defined by

$$[u_a, u_b] = C_{ab} T_0^0 + L_{0abc} v^c, \tag{5.47}$$

$$[u_a, T_{\vec{m}}^i] = m_a T_{\vec{m}}^i - K_{0ab}^i v^b, \tag{5.48}$$

$$[T_{\vec{m}}^i, T_{\vec{n}}^j] = m_a g_{\vec{m}}^{ij} \delta_{\vec{m}+\vec{n}}^0 v^a + f_{\vec{m}\vec{n}}^{ijk} T_{\vec{m}+\vec{n}}^k, \tag{5.49}$$

$$[v^a, T_{\vec{m}}^i] = 0, \tag{5.50}$$

where $(a, b = 1, \dots, d)$. In the sense that one can construct the 3-algebra (5.43)–(5.46) from a Lie algebra by adjoining two elements (u_0, v^0) , this 3-algebra is not a good representative of the new class of 3-algebras defined in (3.70)–(3.79). However, it is still a good example because it demonstrates the roles played by the new parameters J_{ab} and K_{abc} , which encode the information about derivatives of the Lie algebra g , which is a subalgebra of the loop algebra (5.47)–(5.50).

It follows from the result of [16] that the BLG model with the Lie 3-algebra (5.43)–(5.46) is exactly equivalent to the SYM theory defined with the Lie algebra (5.47)–(5.50). In section 5.1, we showed explicitly that for $d = 1$ the resulting SYM theory is the low energy theory for D3-branes. Now we briefly sketch the derivation for generic d to obtain the SYM theory for $D(d + 2)$ -branes.

Expanding the fields in the BLG model, we have

$$X^I = \sum_{a=0}^d X_a^I u_a + \hat{X}^I(Z) + Y_a^I v_a, \quad (5.51)$$

$$\Psi = \sum_{a=0}^d \Psi_a u_a + \hat{\Psi}(Z) + \Phi_a v_a, \quad (5.52)$$

$$A_\mu = \frac{1}{2} \sum_{a,b=0}^d A_{\mu ab} u_a \wedge u_b + \sum_{a=0}^d u_a \wedge \hat{A}_{\mu a}(Z) + \sum_{a=0}^d v^a \wedge \hat{A}'_{\mu a}(Z) \\ + \frac{1}{2} \sum_{a,b=0}^d A'_{\mu ab} v^a \wedge v^b + \frac{1}{2} \sum_{ij} A_{\mu(i\vec{m})(j\vec{n})} T_{\vec{m}}^i \wedge T_{\vec{n}}^j, \quad (5.53)$$

where we have used (5.36) and the notation

$$\hat{X}^I(Z) := \sum_{\vec{m}} X_{(i\vec{m})}^I T^i Z^{m_1} \dots Z^{m_d}, \quad (5.54)$$

$$\hat{\Psi}(Z) := \sum_{\vec{m}} \Psi_{(i\vec{m})} T^i Z^{m_1} \dots Z^{m_d}, \quad (5.55)$$

$$\hat{A}_{\mu a}(Z) := \sum_{\vec{m}} A_{\mu a(i\vec{m})} T^i Z^{m_1} \dots Z^{m_d}, \quad (5.56)$$

$$\hat{A}'_{\mu a}(Z) := \sum_{\vec{m}} A'_{\mu a(i\vec{m})} T^i Z^{m_1} \dots Z^{m_d}, \quad (5.57)$$

and $X_a^I(Z)$, $\Psi_a(Z)$, $\hat{A}_{\mu a}(Z)$ and $\hat{A}'_{\mu a}(Z)$ are sections of a twisted bundle on T_θ^d .

As we have done it many times already, we fix the coefficients of u_a as

$$X_a^I = \text{constant}, \quad \Psi_a = 0, \quad A_{\mu ab} = 0, \quad (a, b = 0, 1, \dots, d,) \quad (5.58)$$

and the coefficients of v_a can be ignored. Here $A_{\mu ab}$ is chosen to be zero for simplicity. If $A_{\mu ab}$'s are nonzero, it corresponds to turning on a constant background field strength with nonvanishing components of $F_{\mu I}$.

To proceed, we first define covariant derivatives \mathcal{D}_a on the noncommutative torus, such that

$$[\mathcal{D}_a, Z_1^{m_1} \dots Z_d^{m_d}] = m_a Z_1^{m_1} \dots Z_d^{m_d}, \quad (5.59)$$

$$[\mathcal{D}_a, \mathcal{D}_b] = C_{ab}, \quad (5.60)$$

where C_{ab} is the constant background field strength that determines the twisting of the bundle on T_θ^d .

The rest of the derivation is essentially the same as section 5.1. Finally, after integrating out the field \tilde{A} , the BL Lagrangian turns into that of a SYM theory

$$\mathcal{L} = -\frac{1}{4} \sum_{A,B=0}^9 \langle F_{AB}, F^{AB} \rangle + \frac{i}{2} \langle \bar{\Psi}, \Gamma^A \hat{D}_A \Psi \rangle, \quad (5.61)$$

where

$$F_{\mu\nu} := [\hat{D}_\mu, \hat{D}_\nu], \tag{5.62}$$

$$F_{\mu I} := [\hat{D}_\mu, \hat{D}_I], \tag{5.63}$$

$$F_{IJ} := [\hat{D}_I, \hat{D}_J] + C_{IJ}, \tag{5.64}$$

and

$$\hat{D}_\mu := \partial_\mu - \hat{A}_{\mu 0}(Z), \tag{5.65}$$

$$\hat{D}^I := X_a^I \mathcal{D}_a - \hat{X}^I(Z), \tag{5.66}$$

$$C^{IJ} := X_a^I X_b^J C_{ab}. \tag{5.67}$$

Roughly speaking, only d of the \hat{D}^I 's are covariant derivatives and the rest $7 - d$ are scalar fields. To turn on the background field $C_{\mu I}$, we can assign nonzero values to $A_{\mu 0a}$ and $A_{\mu ab}$.

5.3 M2 to M5 revisited

As we discussed in section 3.2, there is an interesting Lorentzian 3-algebra associated with the Nambu-Poisson bracket on T^3 defined through the structure constants (3.27), (3.31)–(3.33). We claim that the BLG model associated with this 3-algebra is exactly the description of M5-brane in [8, 9, 16] while it was not explicitly understood. We would like to give a brief sketch on this point.

The key observation to define 6-dimensional fields on M5 from BLG model is to use the “mode expansion” such as

$$X_i^I(x) T^i \rightarrow X_i^I(x) \chi^i(y) =: X^I(x, y). \tag{5.68}$$

If we add three pairs of Lorentzian generators (u^a, v^a) , we have to redefine the above expansion as

$$\tilde{X}^I(x) = X_i^I(x) \chi^i(y) + X_a^I(x) u^a + \underline{X}_a^I(x) v^a. \tag{5.69}$$

Here, the fields $X_a^I(x)$ and $\underline{X}_a^I(x) v^a$ are ghost fields. As we have seen repeatedly, one may put

$$\underline{X}_a^I(x) = 0, \quad X_a^I = \text{const.} \tag{5.70}$$

By change of basis in the transverse direction \mathbf{R}^8 , one may put

$$\begin{aligned} \tilde{X}^a(x, y) &= X_i^a(x) \chi^i(y) + \lambda_a u^a & (a = 1, 2, 3), \\ \tilde{X}^I(x, y) &= X_i^I(x) \chi^i(y) & (I = 4, \dots, 8). \end{aligned} \tag{5.71}$$

where λ_a are constant numbers. This is exactly the assignment by which we can reproduce the M5-brane action from BLG model (for example, eq. (30) in [8]). Various kinetic terms on M5 world volume such as $(\partial_{y^a} X^i)^2$ ($i = 4, \dots, 8$) are generated from the extra term in \tilde{X}^a . All the other analysis in [8, 9] remain the same and we have the same conclusion.

We note that if we do not include these extra terms, the BLG model associated with this 3-algebra would contain infinite number of massless mode even when we consider the compactification on T^3 . On the other hand, if we use the M5 action in [8, 9] on T^3 , we can produce the Kaluza-Klein mass correctly since we have the kinetic term as mentioned. Therefore, the generation of Kaluza-Klein mass on M5 can be again reduced to the inclusion of pairs of Lorentzian norm generators in the Nambu-Poisson 3-algebra.

6 Conclusion and discussion

In this paper, we considered some generalizations of the Lorentzian Lie 3-algebras and studied the BLG models based on the symmetry. In the examples we studied, we naturally obtain the string/M theory compactification on the torus. The mass term generated by the Higgs fields can be identified with the Kaluza-Klein mass in the toroidal compactification. The dimension of the torus can be identified with the number of negative-norm generators of the 3-algebra. We also argued that one may use our technique to consider the D-brane system where its gauge symmetry is described by infinite dimensional loop algebras.

We do not believe that our examples exhaust all possible 3-algebras which are relevant to M/string theories. For example, we did not fully examine the infinite dimensional case with $F^{ijkl} \neq 0$. Another interesting possibility is the description of more general background, such as orbifolds, through different choices of Lorentzian Lie (3-)algebras.

Acknowledgments

We appreciate partial financial support from Japan-Taiwan Joint Research Program provided by Interchange Association (Japan) by which this collaboration is made possible.

The authors thank Yosuke Imamura and Ta-sheng Tai for helpful discussions. Y. M. would like to thank the hospitality of the string theory group in Taiwan. The work of P.-M. H. is supported in part by the National Science Council, and the National Center for Theoretical Sciences, Taiwan, R.O.C. Y. M. is partially supported by KAKENHI (#20540253) from MEXT, Japan. S. S. is partially supported by Global COE Program “the Physical Sciences Frontier”, MEXT, Japan.

References

- [1] J. Bagger and N. Lambert, *Modeling multiple M2's*, *Phys. Rev. D* **75** (2007) 045020 [[hep-th/0611108](#)] [[SPIRES](#)].
- [2] J. Bagger and N. Lambert, *Gauge symmetry and supersymmetry of multiple M2-branes*, *Phys. Rev. D* **77** (2008) 065008 [[arXiv:0711.0955](#)] [[SPIRES](#)].
- [3] J. Bagger and N. Lambert, *Comments on multiple M2-branes*, *JHEP* **02** (2008) 105 [[arXiv:0712.3738](#)] [[SPIRES](#)].
- [4] A. Gustavsson, *Algebraic structures on parallel M2-branes*, *Nucl. Phys. B* **811** (2009) 66 [[arXiv:0709.1260](#)] [[SPIRES](#)].

- [5] P.-M. Ho, R.-C. Hou and Y. Matsuo, *Lie 3-algebra and multiple M2-branes*, *JHEP* **06** (2008) 020 [[arXiv:0804.2110](#)] [[SPIRES](#)].
- [6] S.A. Cherkis and C. Sämann, *Multiple M2-branes and generalized 3-Lie algebras*, *Phys. Rev. D* **78** (2008) 066019 [[arXiv:0807.0808](#)] [[SPIRES](#)];
M. Ali-Akbari, M.M. Sheikh-Jabbari and J. Simon, *Relaxed three-algebras: their matrix representations and implications for multi M2-brane theory*, *JHEP* **12** (2008) 037 [[arXiv:0807.1570](#)] [[SPIRES](#)];
M. Alishahiha and S. Mukhi, *Higher-derivative 3-algebras*, *JHEP* **10** (2008) 032 [[arXiv:0808.3067](#)] [[SPIRES](#)];
M. Axenides and E. Floratos, *Nambu-Lie 3-algebras on fuzzy 3-manifolds*, *JHEP* **02** (2009) 039 [[arXiv:0809.3493](#)] [[SPIRES](#)];
J.M. Figueroa-O'Farrill, *Three lectures on 3-algebras*, [arXiv:0812.2865](#) [[SPIRES](#)];
S.A. Cherkis, V. Dotsenko and C. Sämann, *On superspace actions for multiple M2-branes, metric 3-algebras and their classification*, [arXiv:0812.3127](#) [[SPIRES](#)].
- [7] Y. Kawamura, *Cubic matrix, generalized spin algebra and uncertainty relation*, *Prog. Theor. Phys.* **110** (2003) 579 [[hep-th/0304149](#)] [[SPIRES](#)].
- [8] P.-M. Ho and Y. Matsuo, *M5 from M2*, *JHEP* **06** (2008) 105 [[arXiv:0804.3629](#)] [[SPIRES](#)].
- [9] P.-M. Ho, Y. Imamura, Y. Matsuo and S. Shiba, *M5-brane in three-form flux and multiple M2-branes*, *JHEP* **08** (2008) 014 [[arXiv:0805.2898](#)] [[SPIRES](#)].
- [10] C.-S. Chu, P.-M. Ho, Y. Matsuo and S. Shiba, *Truncated Nambu-Poisson bracket and entropy formula for multiple membranes*, *JHEP* **08** (2008) 076 [[arXiv:0807.0812](#)] [[SPIRES](#)].
- [11] P. de Medeiros, J.M. Figueroa-O'Farrill and E. Méndez-Escobar, *Metric Lie 3-algebras in Bagger-Lambert theory*, *JHEP* **08** (2008) 045 [[arXiv:0806.3242](#)] [[SPIRES](#)].
- [12] O. DeWolfe, T. Hauer, A. Iqbal and B. Zwiebach, *Uncovering infinite symmetries on (p,q) 7-branes: Kac-Moody algebras and beyond*, *Adv. Theor. Math. Phys.* **3** (1999) 1835 [[hep-th/9812209](#)] [[SPIRES](#)].
- [13] R.W. Gebert and H. Nicolai, *On E_{10} and the DDF construction*, *Commun. Math. Phys.* **172** (1995) 571 [[hep-th/9406175](#)] [[SPIRES](#)];
J.A. Harvey and G.W. Moore, *On the algebras of BPS states*, *Commun. Math. Phys.* **197** (1998) 489 [[hep-th/9609017](#)] [[SPIRES](#)].
- [14] J. Gomis, G. Milanesi and J.G. Russo, *Bagger-Lambert theory for general Lie algebras*, *JHEP* **06** (2008) 075 [[arXiv:0805.1012](#)] [[SPIRES](#)].
- [15] S. Benvenuti, D. Rodriguez-Gomez, E. Tonni and H. Verlinde, *$N = 8$ superconformal gauge theories and M2 branes*, *JHEP* **01** (2009) 078 [[arXiv:0805.1087](#)] [[SPIRES](#)].
- [16] P.-M. Ho, Y. Imamura and Y. Matsuo, *M2 to D2 revisited*, *JHEP* **07** (2008) 003 [[arXiv:0805.1202](#)] [[SPIRES](#)].
- [17] S. Mukhi and C. Papageorgakis, *M2 to D2*, *JHEP* **05** (2008) 085 [[arXiv:0803.3218](#)] [[SPIRES](#)].
- [18] M.A. Bandres, A.E. Lipstein and J.H. Schwarz, *Ghost-free superconformal action for multiple M2-branes*, *JHEP* **07** (2008) 117 [[arXiv:0806.0054](#)] [[SPIRES](#)].

- [19] J. Gomis, D. Rodriguez-Gomez, M. Van Raamsdonk and H. Verlinde, *Supersymmetric Yang-Mills theory from Lorentzian three-algebras*, *JHEP* **08** (2008) 094 [[arXiv:0806.0738](#)] [[SPIRES](#)].
- [20] P. de Medeiros, J. Figueroa-O'Farrill, E. Méndez-Escobar and P. Ritter, *On the Lie-algebraic origin of metric 3-algebras*, [arXiv:0809.1086](#) [[SPIRES](#)];
 B. Ezhuthachan, S. Mukhi and C. Papageorgakis, *D2 to D2*, *JHEP* **07** (2008) 041 [[arXiv:0806.1639](#)] [[SPIRES](#)];
 Y. Honma, S. Iso, Y. Sumitomo and S. Zhang, *Scaling limit of $N = 6$ superconformal Chern-Simons theories and Lorentzian Bagger-Lambert theories*, *Phys. Rev. D* **78** (2008) 105011 [[arXiv:0806.3498](#)] [[SPIRES](#)];
 J. Gomis, D. Rodriguez-Gomez, M. Van Raamsdonk and H. Verlinde, *A massive study of M2-brane proposals*, *JHEP* **09** (2008) 113 [[arXiv:0807.1074](#)] [[SPIRES](#)];
 H. Verlinde, *D2 or M2? A note on membrane scattering*, [arXiv:0807.2121](#) [[SPIRES](#)];
 S. Banerjee and A. Sen, *Interpreting the M2-brane action*, [arXiv:0805.3930](#) [[SPIRES](#)];
 S. Cecotti and A. Sen, *Coulomb branch of the Lorentzian three algebra theory*, [arXiv:0806.1990](#) [[SPIRES](#)];
 E. Antonyan and A.A. Tseytlin, *On 3D $N = 8$ Lorentzian BLG theory as a scaling limit of 3D superconformal $N = 6$ ABJM theory*, *Phys. Rev. D* **79** (2009) 046002 [[arXiv:0811.1540](#)] [[SPIRES](#)].
- [21] J.P. Gauntlett and J.B. Gutowski, *Constraining maximally supersymmetric membrane actions*, [arXiv:0804.3078](#) [[SPIRES](#)];
 G. Papadopoulos, *M2-branes, 3-Lie algebras and Plucker relations*, *JHEP* **05** (2008) 054 [[arXiv:0804.2662](#)] [[SPIRES](#)].
- [22] M. Li and T. Wang, *M2-branes coupled to antisymmetric fluxes*, *JHEP* **07** (2008) 093 [[arXiv:0805.3427](#)] [[SPIRES](#)];
 J.-H. Park and C. Sochichiu, *Single M5 to multiple M2: taking off the square root of Nambu-Goto action*, [arXiv:0806.0335](#) [[SPIRES](#)];
 I.A. Bandos and P.K. Townsend, *Light-cone M5 and multiple M2-branes*, *Class. Quant. Grav.* **25** (2008) 245003 [[arXiv:0806.4777](#)] [[SPIRES](#)];
 K. Furuuchi, S.-Y.D. Shih and T. Takimi, *M-theory superalgebra from multiple membranes*, *JHEP* **08** (2008) 072 [[arXiv:0806.4044](#)] [[SPIRES](#)];
 G. Bonelli, A. Tanzini and M. Zabzine, *Topological branes, p-algebras and generalized Nahm equations*, [arXiv:0807.5113](#) [[SPIRES](#)];
 I.A. Bandos and P.K. Townsend, *SDiff gauge theory and the M2 condensate*, *JHEP* **02** (2009) 013 [[arXiv:0808.1583](#)] [[SPIRES](#)];
 C.-S. Chu and D.J. Smith, *Towards the quantum geometry of the M5-brane in a constant C-field from multiple membranes*, [arXiv:0901.1847](#) [[SPIRES](#)].
- [23] Y. Nambu, *Generalized Hamiltonian dynamics*, *Phys. Rev. D* **7** (1973) 2405 [[SPIRES](#)];
 L. Takhtajan, *On foundation of the generalized Nambu mechanics* (second version), *Commun. Math. Phys.* **160** (1994) 295 [[hep-th/9301111](#)] [[SPIRES](#)];
 I. Vaisman, *A survey on Nambu-Poisson brackets*, *Acta. Math. Univ. Comenianae* **2** (1999) 213 [[math.DG/9901047](#)].
- [24] U. Gran, B.E.W. Nilsson and C. Petersson, *On relating multiple M2 and D2-branes*, *JHEP* **10** (2008) 067 [[arXiv:0804.1784](#)] [[SPIRES](#)].

- [25] P.-M. Ho, Y.-Y. Wu and Y.-S. Wu, *Towards a noncommutative geometric approach to matrix compactification*, *Phys. Rev. D* **58** (1998) 026006 [[hep-th/9712201](#)] [[SPIRES](#)];
P.-M. Ho and Y.-S. Wu, *Noncommutative gauge theories in matrix theory*,
Phys. Rev. D **58** (1998) 066003 [[hep-th/9801147](#)] [[SPIRES](#)]; *Matrix compactification on orientifolds*, *Phys. Rev. D* **60** (1999) 026002 [[hep-th/9812143](#)] [[SPIRES](#)].
- [26] P.-M. Ho, *Twisted bundle on quantum torus and BPS states in matrix theory*,
Phys. Lett. B **434** (1998) 41 [[hep-th/9803166](#)] [[SPIRES](#)].
- [27] D. Brace, B. Morariu and B. Zumino, *Dualities of the matrix model from T-duality of the type-II string*, *Nucl. Phys. B* **545** (1999) 192 [[hep-th/9810099](#)] [[SPIRES](#)].
- [28] H. Lin, *Kac-Moody extensions of 3-algebras and M2-branes*, *JHEP* **07** (2008) 136 [[arXiv:0805.4003](#)] [[SPIRES](#)].
- [29] W. Taylor, *D-brane field theory on compact spaces*, *Phys. Lett. B* **394** (1997) 283 [[hep-th/9611042](#)] [[SPIRES](#)].