

Twist-three at five loops, Bethe ansatz and wrapping

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

JHEP03(2009)129

(<http://iopscience.iop.org/1126-6708/2009/03/129>)

[The Table of Contents](#) and [more related content](#) is available

Download details:

IP Address: 80.92.225.132

The article was downloaded on 03/04/2010 at 10:37

Please note that [terms and conditions apply](#).

Twist-three at five loops, Bethe ansatz and wrapping

Matteo Beccaria,^a Valentina Forini,^b Tomasz Łukowski^c and Stefan Zieme^{d,e}

^a*Physics Department and Sezione INFN, Salento University,
Via Arnesano, 73100 Lecce, Italy*

^b*Max-Planck-Institut für Gravitationsphysik, Albert-Einstein-Institut,
Am Mühlenberg 1, D-14476 Potsdam, Germany*

^c*Institute of Physics, Jagellonian University,
ul. Reymonta 4, 30-059 Kraków, Poland*

^d*Max-Planck-Institut für Gravitationsphysik, Albert-Einstein-Institut,
Am Mühlenberg 1, D-14476 Potsdam, Germany*

^e*Kavli Institute for Theoretical Physics, University of California,
Santa Barbara, CA 93106, U.S.A.*

E-mail: matteo.beccaria@le.infn.it, valentina.forini@aei.mpg.de,
tomaszlukowski@gmail.com, stefan.zieme@aei.mpg.de

ABSTRACT: We present a formula for the five-loop anomalous dimension of $\mathcal{N} = 4$ SYM twist-three operators in the $\mathfrak{sl}(2)$ sector. We obtain its asymptotic part from the Bethe Ansatz and finite volume corrections from the generalized Lüscher formalism, considering scattering processes of spin chain magnons with virtual particles that travel along the cylinder. The complete result respects the expected large spin scaling properties and passes non-trivial tests including reciprocity constraints. We analyze the pole structure and find agreement with a conjectured resummation formula. In analogy with the twist-two anomalous dimension at four-loops wrapping effects are of order $(\log^2 M/M^2)$ for large values of the spin.

KEYWORDS: AdS-CFT Correspondence, Integrable Field Theories, Bethe Ansatz

ARXIV EPRINT: [0901.4864](https://arxiv.org/abs/0901.4864)

Contents

1	Introduction and discussion	1
2	Asymptotic Bethe equations	3
3	Parity invariance of $\gamma(M)$	5
3.1	Harmonic combinations with definite parity	5
4	Determination of γ_{10}^{ABA}	7
5	The wrapping contribution	9
6	Analytic continuation	11
A	Harmonic sums	13
B	Analysis of the asymptotic structure of γ and P	14

1 Introduction and discussion

Recent developments in computing finite size effects on the asymptotic spectrum of $\mathcal{N} = 4$ SYM twist-two operators are very promising in order to ultimately find the complete spectral equations of the dilatation generator.

The revealing of integrable structures on both sides of the AdS/CFT correspondence [1] gradually led to powerful tools for computing anomalous dimension of gauge invariant operators by means of the Bethe Ansatz [2]. The factorized two-body S-Matrix [3] that governs scattering processes of the spin chain particles and excitations of the $AdS_5 \times S^5$ string worldsheet is determined by the $\mathfrak{psu}(2, 2|4)$ symmetry of $\mathcal{N} = 4$ up to a phase factor [4]. In order to also determine this algebraic ambiguity a crossing-like equation for the dressing phase has been derived [5]. It allows for multiple solutions [6], one of which gets singled out [7] by reconciliation with an explicit diagrammatic calculation of the four-loop anomalous dimensions of twist-two operators in the large spin limit [8]. It still remains an open problem to explicitly show the crossing invariance of the dressed asymptotic Bethe equations. For this purpose the representations of the dressing factor in [9] might prove useful.

It was shown that these equations are asymptotic in nature, and need to be corrected by wrapping effects [10]. An explicit calculation of the anomalous dimension of twist-two operators from the asymptotic Bethe Ansatz at four-loops unequivocally showed that the pole prescribed by BFKL physics can not be fulfilled [11]. The Bethe Ansatz therefore does not produce the correct result at and beyond wrapping order.

However, for exactly these operators complete results have been obtained for the first time. For the simplest representative of twist-two operators, the Konishi-field, a field theoretical calculation starting from the asymptotic dilatation generator [12], finite-size corrections to the Bethe Ansatz using Lüscher formulas [13], and finally a full-fledged Feynman calculus [14] identically determined the complete anomalous dimension.

The successful application of the Lüscher formalism relies on a generalization of the Lüscher formulas [15] to both non-relativistic models as well as multi-particle states, which had been conjectured in [13, 16]. With this formalism applied to the 2d worldsheet QFT of the $AdS_5 \times S^5$ superstring it has been possible to compute the four-loop anomalous dimension of twist-two operators at general values of the spin [17]. The result passed several non-trivial tests from BFKL and reciprocity constraints [17, 18]. The leading transcendental part had been confirmed in an impressive field theory computation [19].

For the complete spectral equations of $\mathcal{N} = 4$ SYM, however, thermodynamic Bethe Ansatz methods ought to be applied, as has been initiated for string and gauge theory in [20, 21]. A Y-system, which is believed to yield anomalous dimensions of arbitrary local operators of planar $\mathcal{N} = 4$ SYM has been recently conjectured in [22].

The aim of this work is to continue the application of the Bethe Ansatz and the Lüscher formalism to the next operators in reach, namely twist-three operators. The leading wrapping contribution to the anomalous dimension of twist-three operators will appear at five-loops. In order to compute the complete five-loop anomalous dimension of the ground state we start from an ansatz based on the maximum transcendentality principle [23] for both the asymptotic and wrapping contributions. The asymptotic part can be determined from the Bethe equations after the initial ansatz has been upgraded with further constraints from reciprocity. To compute the wrapping contribution, we apply the generalized Lüscher formulas [17] to operators of twist-three.

Our result passes some important consistency tests. Its leading asymptotic behavior for large values of the spin reproduces the universal scaling function at five-loop order. The first subleading correction coincides with the results of [24, 25]. Contributions from finite-size effects start at order $(\log^2 M/M^2)$, as in the case of twist-two operators [18]. In contrast to the latter, there is no BFKL equation for twist-three operators, and therefore no prediction for the pole structure of our result. However, we analyze the behavior of the anomalous dimension at the singular value of spin $M = -2$. Interestingly, the pole structure agrees with the conjectured resummation formula of [11], once contributions from wrapping effects are taken into account. Additionally, the wrapping correction obtained from the Lüscher formula precisely matches with the computation from the Y-system¹ [22]. The complete result, including the wrapping contribution, is reciprocity respecting.

The main body of the paper follows to the above outlined procedure. Some basic definitions of harmonic sums are recalled in appendix A. Appendix B contains the analysis of the asymptotic structure of the anomalous dimensions and their corresponding P -kernels up to five loops. Speculations related to a pattern in the asymptotic structure of anomalous dimensions of twist operators are also given in appendix B.

¹We thank Pedro Vieira for his support in the numerical cross-check.

2 Asymptotic Bethe equations

We will start our analysis with the contribution to the final result stemming from the asymptotic Bethe Ansatz equations. Twist-three operators are embedded in the $\mathfrak{sl}(2)$ subsector of $\mathcal{N} = 4$ SYM. They can be represented by an insertion of M covariant derivatives \mathcal{D} into the protected half-BPS state $\text{Tr } \mathcal{Z}^3$

$$\text{Tr } (\mathcal{D}^{s_1} \mathcal{Z} \mathcal{D}^{s_2} \mathcal{Z} \mathcal{D}^{s_3} \mathcal{Z}) + \dots, \quad \text{with } M = s_1 + s_2 + s_3. \quad (2.1)$$

Their anomalous dimensions can be obtained from a non-compact, length-three $\mathfrak{sl}(2)$ spin chain with M excitations underlying a factorized two-body scattering [3]. However, the interaction range between scattering particles increases with orders of the coupling constant in perturbation theory. If it exceeds the length of the spin chain and *wraps* around it, the S-matrix picture [3, 4] loses its meaning, as no asymptotic region can be defined any longer. For twist L operators this effect, delayed by superconformal invariance, starts at order g^{2L+4} . Nevertheless, the Bethe Ansatz does not cease to work but gives an incomplete result, which does not incorporate these corrections [11].

It was shown that the Bethe Ansatz result for twist-two operators can be completed by considering additional scattering effects with virtual particles [17]. It passes non-trivial tests with BFKL [17], as well as reciprocity [18] constraints and reproduces the correct scaling behavior at large values of the spin M [7, 18], proposing a certain confidence. We compute these wrapping effects for twist $L = 3$ in section 5.

The Bethe Ansatz equations for the operators (2.1) with our choice of the coupling constant $g^2 = \frac{g_{\text{SYM}}^2 N}{16\pi^2}$ are given by

$$\left(\frac{x_k^+}{x_k^-}\right)^3 = \prod_{\substack{j=1 \\ j \neq k}}^M \frac{x_k^- - x_j^+}{x_k^+ - x_j^-} \frac{1 - g^2/x_k^+ x_j^-}{1 - g^2/x_k^- x_j^+} \exp(2i\theta(u_k, u_j)), \quad \prod_{k=1}^M \frac{x_k^+}{x_k^-} = 1. \quad (2.2)$$

The spectral parameters x^\pm are defined in terms of the rapidities u by [26]

$$x^\pm(u) = x(u \pm i/2), \quad x(u) = \frac{u}{2} \left(1 + \sqrt{1 - 4g^2/u^2}\right). \quad (2.3)$$

The phase shift $\theta(u_k, u_j)$ acquired by two particles with rapidities u_k, u_j passing each other is given to five-loop order by the expansion [7]

$$\theta(u_k, u_j) = (4\zeta_3 g^6 - 40\zeta_5 g^8) (q_2(u_k)q_3(u_j) - q_3(u_k)q_2(u_j)) + \mathcal{O}(g^{10}), \quad (2.4)$$

where the $q_r(u)$ correspond to the conserved magnon charges [26]

$$q_r(u) = \frac{i}{r-1} \left(\frac{1}{(x^+(u))^{r-1}} - \frac{1}{(x^-(u))^{r-1}} \right). \quad (2.5)$$

From the solution to (2.2) given in terms of the Bethe roots x_k^\pm one computes the anomalous dimension by

$$\gamma^{\text{ABA}}(g) = 2g^2 \sum_{k=1}^M q_2(u_k). \quad (2.6)$$

Due to the length $L = 3$ of the operators (2.1), wrapping effects are expected to contribute at order g^{10} , such that the anomalous dimension is written perturbatively as

$$\begin{aligned} \gamma(M) = & g^2 \gamma_2(M) + g^4 \gamma_4(M) + g^6 \gamma_6(M) + g^8 \gamma_8(M) \\ & + g^{10} (\gamma_{10}^{\text{ABA}}(M) + \gamma_{10}^{\text{wrapping}}(M)) + \dots, \end{aligned} \quad (2.7)$$

where we tacitly assumed that to order g^8 the complete result is identical to γ^{ABA} and therefore dropped its index.

At one-loop the Bethe roots u_k are given by zeros of the Wilson polynomial [11]

$$P_M(u) = {}_4F_3 \left(\begin{matrix} -\frac{M}{2}, \frac{M}{2} + 1, \frac{1}{2} + iu, \frac{1}{2} - iu \\ 1, 1, 1 \end{matrix} \middle| 1 \right). \quad (2.8)$$

Closed expressions for the corrections to the Bethe roots to three-loop order have also been obtained in [27] from the Baxter approach [28]. However, it is currently unclear if the asymptotic Baxter equation [28] reproduces the same result as the Bethe Ansatz at and beyond wrapping order.

In order to obtain closed expressions for the anomalous dimension we will therefore solve (2.2) perturbatively for fixed values of the spin M and match the coefficients in an appropriate ansatz which assumes the maximum transcendentality principle [23]. Up to four loops, these expressions have been derived in [11] and [29]. They are given by

$$\gamma_2 = 8 S_1 \quad (2.9)$$

$$\gamma_4 = -16 S_1 S_2 - 8 S_3 \quad (2.10)$$

$$\gamma_6 = 32 S_1 S_2^2 + 48 S_3 S_2 + 16 S_1 S_4 + 40 S_5 - 32 S_{2,3} + 64 S_1 S_{3,1} + 32 S_{4,1} - 64 S_{3,1,1} \quad (2.11)$$

$$\begin{aligned} \gamma_8 = & 8 S_7 + 112 S_{1,6} + 240 S_{2,5} - 80 S_{3,4} - 464 S_{4,3} - 336 S_{5,2} - 80 S_{6,1} - 640 S_{1,1,5} \\ & - 512 S_{1,2,4} + 384 S_{1,3,3} + 512 S_{1,4,2} - 512 S_{2,1,4} + 320 S_{2,2,3} + 640 S_{2,3,2} + 64 S_{2,4,1} \\ & + 384 S_{3,1,3} + 704 S_{3,2,2} + 384 S_{3,3,1} + 576 S_{4,1,2} + 576 S_{4,2,1} + 384 S_{5,1,1} + 1280 S_{1,1,1,4} \\ & - 256 S_{1,1,3,2} + 512 S_{1,1,4,1} - 384 S_{1,2,2,2} + 256 S_{1,2,3,1} - 384 S_{1,3,1,2} - 384 S_{1,3,2,1} \\ & - 384 S_{1,4,1,1} - 384 S_{2,1,2,2} + 256 S_{2,1,3,1} - 384 S_{2,2,1,2} - 384 S_{2,2,2,1} - 384 S_{2,3,1,1} \\ & - 384 S_{3,1,1,2} - 384 S_{3,1,2,1} - 384 S_{3,2,1,1} - 384 S_{4,1,1,1} - 1024 S_{1,1,1,3,1} - 128 S_1 S_3 \zeta_3. \end{aligned} \quad (2.12)$$

All sums are evaluated at argument $M/2$ and only positive indices appear. At four loop-order, the dressing phase of the Bethe equations starts to contribute to (2.12) with a term proportional to ζ_3 .

To determine the five-loop result in the same fashion implies a tremendous computational effort in view of the necessary precision.² We have obtained rational values for the anomalous dimension up to $M \approx 200$, which is however too far from the requirement to fit the coefficients of the corresponding ansatz vector of constant degree of transcendentality. In order to find a closed form for γ_{10}^{ABA} in terms of nested harmonic sums further constraints to reduce the number of coefficients are needed.

²See [30] for the five-loop anomalous dimension of a different class of operators (the field strength operators $\text{Tr } \mathcal{F}^L$), determined as a closed function of their length L .

3 Parity invariance of $\gamma(M)$

The multi-loop anomalous dimension $\gamma(M)$ is conjectured to obey a powerful constraint known as generalized Gribov-Lipatov *reciprocity*. This constraint, arising in the QCD context, has been presented in [31, 32] as a special (space-time symmetric) reformulation of the parton distribution functions evolution equation, and approached in [33] from the point of view of the large M expansion. In particular, in [33] such analysis has been generalised to anomalous dimensions of operators of arbitrary twist L , and reciprocity has been dubbed *parity invariance* in the sense clarified below. Reciprocity has been checked in various multi-loop calculations of weakly coupled $\mathcal{N} = 4$ gauge theory [18, 29, 34–37].

The reciprocity or parity invariance condition is easily expressed in terms of the P -function (kernel), depending on the Lorentz spin M , which is in one-to-one correspondence, at least perturbatively, with the anomalous dimension $\gamma(M)$ as follows from [31–33]

$$\gamma(M) = P\left(M + \frac{1}{2}\gamma(M)\right). \tag{3.1}$$

The parity invariance condition is a constraint that arises in the large M expansion of $P(M)$, which is expected to take the following form

$$P(M) = \sum_{\ell \geq 0} \frac{a_\ell (\log J^2)}{J^{2\ell}}, \quad J^2 = \frac{M}{2} \left(\frac{M}{2} + 1 \right), \tag{3.2}$$

where the a_ℓ are *coupling-dependent* polynomials. Eq. (3.2) implies an infinite set of constraints on the coefficients of the large M expansion of $P(M)$ organized in a standard $1/M$ power series. The name *parity-invariance* is related to the absence of terms of the form $1/J^{2n+1}$, odd under $J \rightarrow -J$.

In the following we will use the constraint eq. (3.2) as a guiding principle in order to obtain the five-loop expression γ_{10}^{ABA} . To this aim, we need to express it as a more practical test such that it can be applied to any proposed combination of harmonic sums. This task can be performed with a basic result of [18], which we recall in following.

3.1 Harmonic combinations with definite parity

The notation for complementary harmonic sums \underline{S}_a is recalled in appendix A. Let us introduce the map ω_a , $a \in \mathbb{N}$, which acts linearly on linear combinations of harmonic sums

$$\omega_a(S_{b,c}) = S_{a,b,c} - \frac{1}{2} S_{a+b,c}. \tag{3.3}$$

We also introduce a complementary map $\underline{\omega}_a$, acting in a similar way on complementary sums

$$\underline{\omega}_a(\underline{S}_{b,c}) = \underline{S}_{a,b,c} - \frac{1}{2} \underline{S}_{a+b,c}. \tag{3.4}$$

Finally, let us introduce the combinations Ω

$$\Omega_a = S_a, \tag{3.5}$$

$$\Omega_{a,b} = \omega_a(\Omega_b), \tag{3.6}$$

and the analogous complementary combinations $\underline{\Omega}_a$. It is of course possible to change the basis from S_a to $\underline{\Omega}_a$. For example, up to a degree of transcendentality three, we have

$$\begin{aligned} \underline{\Omega}_1 &= S_1, & \underline{\Omega}_2 &= S_2, & \underline{\Omega}_{1,1} &= S_{1,1} - \frac{S_2}{2}, & \underline{\Omega}_3 &= S_3, & \underline{\Omega}_{2,1} &= S_{2,1} - \frac{S_3}{2}, \\ \underline{\Omega}_{1,2} &= -\frac{1}{6}\pi^2 S_1 - \frac{S_3}{2} + S_{1,2}, & \underline{\Omega}_{1,1,1} &= \frac{S_3}{4} - \frac{S_{1,2}}{2} - \frac{S_{2,1}}{2} + S_{1,1,1}, \end{aligned} \quad (3.7)$$

which can be inverted in order to obtain

$$\begin{aligned} S_2 &= \underline{\Omega}_2, & S_{1,1} &= \frac{\underline{\Omega}_2}{2} + \underline{\Omega}_{1,1}, & S_3 &= \underline{\Omega}_3, & S_{2,1} &= \frac{\underline{\Omega}_3}{2} + \underline{\Omega}_{2,1}, \\ S_{1,2} &= \frac{\pi^2 \underline{\Omega}_1}{6} + \frac{\underline{\Omega}_3}{2} + \underline{\Omega}_{1,2}, & S_{1,1,1} &= \frac{\pi^2 \underline{\Omega}_1}{12} + \frac{\underline{\Omega}_3}{4} + \frac{\underline{\Omega}_{1,2}}{2} + \frac{\underline{\Omega}_{2,1}}{2} + \underline{\Omega}_{1,1,1}. \end{aligned} \quad (3.8)$$

The crucial result is then given by the following theorem [18].

Theorem 1 (a) *The combination $\underline{\Omega}_{a_1, \dots, a_d}$ with positive $\{a_i\}$ is parity-even iff*

$$(-1)^{a_1 + \dots + a_d} = (-1)^d. \quad (3.9)$$

(b) *If this condition is not satisfied, the expansion of $\underline{\Omega}_{a_1, \dots, a_d}$ is parity-odd, with the (trivial) exception of the leading constant term.*

(c) *The combination Ω_{a_1, \dots, a_d} with positive odd $\{a_i\}$ is parity-even.*

From this theorem we deduce the following

Theorem 2 (parity-invariance test) *A specific linear combination of harmonic sums is parity invariant iff it does not contain parity-odd terms when transformed from the S_a basis to the $\underline{\Omega}_a$ basis.*

To see how this test can be used let us consider an illustrative example, the two-loop anomalous dimension. One starts with the following ansatz of transcendentality three

$$\gamma_4 = a_1 S_3 + a_2 S_{1,2} + a_3 S_{2,1} + a_4 S_{1,1,1}, \quad (3.10)$$

with all sums evaluated at $M/2$. The corresponding P_4 -kernel, derived by inverting formula (3.1) and replacing the perturbative expansion (2.7), reads in a canonical basis

$$P_4 = \gamma_4 - \frac{1}{4}\gamma_2\gamma_2' \equiv (a_1 - 16)S_3 + (a_2 + 16)S_{1,2} + (a_3 + 16)S_{2,1} - 16\zeta_2 S_1 + a_4 S_{1,1,1}, \quad (3.11)$$

where we used the one-loop result (2.9). Writing (3.11) in terms of the $\underline{\Omega}$ basis one finds

$$P_4 = c_1 \underline{\Omega}_1 + c_3 \underline{\Omega}_3 + c_{1,2} \underline{\Omega}_{1,2} + c_{2,1} \underline{\Omega}_{2,1} + c_{1,1,1} \underline{\Omega}_{1,1,1} + \text{const}, \quad (3.12)$$

where the c_i are linear combinations of the coefficients a_i . The combinations $\underline{\Omega}_1$, $\underline{\Omega}_3$, $\underline{\Omega}_{1,1,1}$ are all reciprocity respecting, according to the above theorem. Imposing reciprocity on P_4 implies the vanishing of the coefficients of those $\underline{\Omega}$ with wrong parity, namely

$$c_{1,2} = a_2 + 16 + \frac{a_4}{2} = 0, \quad c_{2,1} = a_3 + 16 + \frac{a_4}{2} = 0. \quad (3.13)$$

This leads to the conditions $a_3 = a_2$ and $a_4 = -2(16 + a_2)$, that are indeed satisfied by the expression in (2.10). Thus, reciprocity has determined 2 of the 4 unknown coefficients in the initial ansatz for the anomalous dimension.³

4 Determination of γ_{10}^{ABA}

The strategy to derive the asymptotic part of the anomalous dimension at $n = 5$ loops incorporates a combined use of the maximum transcendentality principle, reciprocity and Bethe equations.

The starting point is to write γ_{10}^{ABA} as a linear combination of harmonic sums of transcendentality $\tau = 2n - 1 = 9$. For a given τ , basic combinatorics leads to the fact that there are $2^{\tau-1}$ linearly independent harmonic sums with positive indices. This means that there are in principle 256 terms which potentially contribute to the anomalous dimension.

From the numerical solution of the asymptotic Bethe equations it is possible to obtain a long list of rational values for $\gamma_{10}^{\text{ABA}}(M)$ for fixed values of M . The list-length is smaller than 256 due to rather hard computational limitations. However, these limitations can be overcome by means of parity-invariance.

To constrain the 256 unknown coefficients via reciprocity one has to impose parity in the sense of eq. (3.2) on the five-loop contribution P_{10} to the kernel P defined in eq. (3.1). This contribution can be derived from the anomalous dimension by simply inverting eq. (3.1) and taking into account the perturbative expansion eq. (2.7). Finally, we apply the previous parity-invariance test and obtain a large set of linear constraints on the unknown coefficients. The total number of constraints from Bethe equations and parity-invariance is now larger than 256 and we find an over-determined set of linear equations, which is solvable. The final result is given in table 1, in which terms multiplied by ζ_3 and ζ_5 are directly induced from the dressing factor. As is the case for lower-loop orders, only positive indices appear in the participating harmonic sums. The result that we have obtained by the above stated methods can be checked as follows:

1. *Scaling function (cusp anomaly)*. A consistency check of the formula presented in table 1 is given by its leading asymptotic behavior, namely

$$\gamma_{10}^{\text{ABA}}(M) \sim 32 \left(\frac{887}{14175} \pi^8 + \frac{4}{3} \pi^2 \zeta_3^2 + 40 \zeta_3 \zeta_5 \right) \log M, \quad \text{for } M \rightarrow \infty. \quad (4.1)$$

It coincides with the five-loop contribution in the weak coupling perturbative expansion of the integral equation obtained in [7], which is believed to describe the universal scaling function. Put differently, we confirm its universality [38, 39] for twist-three at five loops.

2. *Virtual scaling function*. A further confirmation can be found from the evaluation of the first finite-order correction to the asymptotic behavior (4.1). In general this quantity is twist-dependent and thus non-universal [40]. However, it has been shown

³The coefficient a_4 has only been kept to show the exact number of constraints coming from reciprocity. It could have been set to zero from the beginning because at large M the term $S_{1,1,1} \sim \log^3 M$ is not compatible with the universal leading logarithmic behavior (cusp anomaly).

$$\begin{aligned}
 \gamma_{10}^{\text{ABA}} = & 136S_9 + 368S_{1,8} + 2832S_{2,7} + 4272S_{3,6} + 848S_{4,5} - 3024S_{5,4} - 2736S_{6,3} - 1168S_{7,2} \\
 & - 496S_{8,1} - 5376S_{1,1,7} - 12352S_{1,2,6} - 8832S_{1,3,5} + 1600S_{1,4,4} + 3968S_{1,5,3} - 64S_{1,6,2} \\
 & - 1344S_{1,7,1} - 12352S_{2,1,6} - 13760S_{2,2,5} - 2112S_{2,3,4} + 4288S_{2,4,3} - 960S_{2,5,2} - 5440S_{2,6,1} \\
 & - 9088S_{3,1,5} - 2432S_{3,2,4} + 5120S_{3,3,3} + 2688S_{3,4,2} - 4160S_{3,5,1} + 1280S_{4,1,4} + 5824S_{4,2,3} \\
 & + 6400S_{4,3,2} + 2112S_{4,4,1} + 5120S_{5,1,3} + 6208S_{5,2,2} + 5312S_{5,3,1} + 3904S_{6,1,2} + 3904S_{6,2,1} \\
 & + 1728S_{7,1,1} + 21504S_{1,1,1,6} + 22784S_{1,1,2,5} + 5632S_{1,1,3,4} - 1280S_{1,1,4,3} + 6912S_{1,1,5,2} \\
 & + 11520S_{1,1,6,1} + 22784S_{1,2,1,5} + 9088S_{1,2,2,4} - 1024S_{1,2,3,3} + 6784S_{1,2,4,2} + 17152S_{1,2,5,1} \\
 & + 5504S_{1,3,1,4} - 3456S_{1,3,2,3} - 1536S_{1,3,3,2} + 7680S_{1,3,4,1} - 4480S_{1,4,1,3} - 6272S_{1,4,2,2} \\
 & - 3584S_{1,4,3,1} - 3840S_{1,5,1,2} - 3840S_{1,5,2,1} + 768S_{1,6,1,1} + 22784S_{2,1,1,5} + 9088S_{2,1,2,4} \\
 & - 1024S_{2,1,3,3} + 6784S_{2,1,4,2} + 17152S_{2,1,5,1} + 9088S_{2,2,1,4} - 2688S_{2,2,2,3} + 640S_{2,2,3,2} \\
 & + 13440S_{2,2,4,1} - 3456S_{2,3,1,3} - 7040S_{2,3,2,2} - 768S_{2,3,3,1} - 4480S_{2,4,1,2} - 4480S_{2,4,2,1} \\
 & + 2816S_{2,5,1,1} + 6272S_{3,1,1,4} - 2944S_{3,1,2,3} - 1536S_{3,1,3,2} + 7936S_{3,1,4,1} - 2944S_{3,2,1,3} \\
 & - 7296S_{3,2,2,2} - 768S_{3,2,3,1} - 6656S_{3,3,1,2} - 6656S_{3,3,2,1} - 1024S_{3,4,1,1} - 3968S_{4,1,1,3} \\
 & - 6528S_{4,1,2,2} - 3584S_{4,1,3,1} - 6528S_{4,2,1,2} - 6528S_{4,2,2,1} - 4864S_{4,3,1,1} - 5376S_{5,1,1,2} \\
 & - 5376S_{5,1,2,1} - 5376S_{5,2,1,1} - 4608S_{6,1,1,1} - 32768S_{1,1,1,1,5} - 10240S_{1,1,1,2,4} - 3072S_{1,1,1,3,3} \\
 & - 17920S_{1,1,1,4,2} - 30720S_{1,1,1,5,1} - 10240S_{1,1,2,1,4} - 8704S_{1,1,2,3,2} - 24064S_{1,1,2,4,1} \\
 & + 1024S_{1,1,3,1,3} + 2560S_{1,1,3,2,2} - 4096S_{1,1,3,3,1} - 512S_{1,1,4,1,2} - 512S_{1,1,4,2,1} - 10240S_{1,1,5,1,1} \\
 & - 10240S_{1,2,1,1,4} - 8704S_{1,2,1,3,2} - 24064S_{1,2,1,4,1} + 3072S_{1,2,2,2,2} - 6656S_{1,2,2,3,1} \\
 & + 512S_{1,2,3,1,2} + 512S_{1,2,3,2,1} - 10752S_{1,2,4,1,1} + 1024S_{1,3,1,1,3} + 3072S_{1,3,1,2,2} - 3584S_{1,3,1,3,1} \\
 & + 3072S_{1,3,2,1,2} + 3072S_{1,3,2,2,1} - 2560S_{1,3,3,1,1} + 3072S_{1,4,1,1,2} + 3072S_{1,4,1,2,1} + 3072S_{1,4,2,1,1} \\
 & + 3072S_{1,5,1,1,1} - 10240S_{2,1,1,1,4} - 8704S_{2,1,1,3,2} - 24064S_{2,1,1,4,1} + 3072S_{2,1,2,2,2} \\
 & - 6656S_{2,1,2,3,1} + 512S_{2,1,3,1,2} + 512S_{2,1,3,2,1} - 10752S_{2,1,4,1,1} + 3072S_{2,2,1,2,2} - 6656S_{2,2,1,3,1} \\
 & + 3072S_{2,2,2,1,2} + 3072S_{2,2,2,2,1} - 5632S_{2,2,3,1,1} + 3072S_{2,3,1,1,2} + 3072S_{2,3,1,2,1} + 3072S_{2,3,2,1,1} \\
 & + 3072S_{2,4,1,1,1} + 3072S_{3,1,1,2,2} - 4096S_{3,1,1,3,1} + 3072S_{3,1,2,1,2} + 3072S_{3,1,2,2,1} - 2560S_{3,1,3,1,1} \\
 & + 3072S_{3,2,1,1,2} + 3072S_{3,2,1,2,1} + 3072S_{3,2,2,1,1} + 4608S_{3,3,1,1,1} + 3072S_{4,1,1,1,2} + 3072S_{4,1,1,2,1} \\
 & + 3072S_{4,1,2,1,1} + 3072S_{4,2,1,1,1} + 3072S_{5,1,1,1,1} + 16384S_{1,1,1,1,3,2} + 32768S_{1,1,1,1,4,1} \\
 & + 8192S_{1,1,1,2,3,1} + 4096S_{1,1,1,3,1,2} + 4096S_{1,1,1,3,2,1} + 20480S_{1,1,1,4,1,1} + 8192S_{1,1,2,1,3,1} \\
 & + 12288S_{1,1,2,3,1,1} + 8192S_{1,2,1,1,3,1} + 12288S_{1,2,1,3,1,1} + 8192S_{2,1,1,1,3,1} + 12288S_{2,1,1,3,1,1} \\
 & - 16384S_{1,1,1,1,3,1,1} + \zeta_3 (896S_6 - 2304S_{1,5} - 1792S_{2,4} - 768S_{3,3} - 1792S_{4,2} - 2304S_{5,1} \\
 & + 2560S_{1,1,4} + 512S_{1,2,3} + 1536S_{1,3,2} + 3584S_{1,4,1} + 512S_{2,1,3} + 1536S_{2,3,1} + 512S_{3,1,2} \\
 & + 512S_{3,2,1} + 2560S_{4,1,1} - 2048S_{1,1,3,1} - 2048S_{1,3,1,1}) + 1280\zeta_5 (S_{1,3} + S_{3,1} - S_4)
 \end{aligned}$$

Table 1. The result for the five-loop asymptotic dimension $\gamma_{10}^{\text{ABA}}(\frac{M}{2})$, written in the canonical basis.

that this dependence is only linear and an all-loop integral equation can be written [24] (see also [25, 41]). The large M expansion of table 1 leads to the $\mathcal{O}(1/M^0)$ value

$$B_3^{(5)} = \frac{2048}{945}\pi^6\zeta_3 + 64\zeta_3^3 + \frac{8}{45}\pi^4\zeta_5 - \frac{440}{3}\pi^2\zeta_7 - 7448\zeta_9. \quad (4.2)$$

It coincides with the expression written explicitly in [25]. Further interesting observations on the other subleading terms in the asymptotic expansion of γ_{10}^{ABA} will be discussed in appendix B.

3. *BFKL-like poles.* An indirect indication of the correctness of the result emerges by looking at its analytical continuation to complex values of the spin. In particular, the structure of the expansion around $M = -2$ will be presented in section 6.
4. *Dressing self-consistency.* The dressing induced terms in γ_{10}^{ABA} are separately parity invariant. The dressing factor starts to contribute at four loops. It was observed that at this order terms of the anomalous dimensions of twist-two and -three operators coming from the dressing factor are reciprocity respecting *separately* [18, 34, 36]. The analysis of P_{10} in the case of γ_{10}^{ABA} confirms this feature. The five-loop term proportional to ζ_3 is reciprocity respecting if combined with the corresponding four-loop ζ_3 -term. The ζ_5 -term at five loops is reciprocity respecting separately (see formula B.16). This seems to indicate a perturbative pattern for the reciprocity of terms that are dressing-induced. Terms proportional to transcendental sums ζ_i , which newly appear at a given loop order should automatically be reciprocity respecting. Terms proportional to transcendental sums that are also present at lower-loop orders are invariant under (3.2) when combined altogether.
5. *Additional structural properties.* All coefficients of the harmonic sums are integers, likewise to the lower loop orders. Also, P_{10} turns out to be a combination of allowed parity-even combinations of type Ω , a condition being stronger than the general parity invariance.

5 The wrapping contribution

In this section we evaluate the leading wrapping correction to the asymptotic anomalous dimension of twist-three operators.

The Lüscher type formula for multi-particle states was conjectured in [13] and successfully applied to the Konishi-operator in [13] as well as twist-two operators of general spin in [17]. It consists of two parts. One describes the modification of the particle quantization condition due to the finite volume (which will not contribute at leading order), while the second comes from the propagation of virtual particles around the cylinder and is given by

$$\Delta E(L) = - \sum_{Q=1}^{\infty} \int_{-\infty}^{\infty} \frac{dq}{2\pi} \text{STr}_{a_1} [S_{a_1 a}^{a_2 a}(q, p_1) S_{a_2 a}^{a_3 a}(q, p_2) \dots S_{a_M a}^{a_1 a}(q, p_M)] e^{-\tilde{\epsilon}_{a_1}(q)L}. \quad (5.1)$$

This formula applies to a M -particle state of identical particles of type a , whose consecutive self-scatterings preserve the state and determine their momenta p_i via the ABA equations. The matrix $S_{ba}^{ca}(q, p)$ describes how a virtual particle of type b with momentum q scatters on a real particle of type a with momentum p . The exponential factor can be interpreted as the propagator of the virtual particle.

For twist-three operators the momenta of the particles are determined by the ABA equations in terms of the rapidities u by

$$u(p) = \frac{1}{2} \cot \frac{p}{2} \sqrt{1 + 16g^2 \sin^2 \frac{p}{2}}.$$

At one-loop the rapidities are given by roots of the Baxter- Q function $P_M(u)$ in (2.8). As this is an even polynomial of order M , we can repeat the derivation of [17], which leads to a result similar to the twist-two case. However, we have to take into account two differences. The first one is that the length is equal to $L = 3$, which renders the exponential part to be of the form

$$e^{-\tilde{\epsilon}_Q(q)L} = \frac{4^L g^{2L}}{(q^2 + Q^2)^L} \stackrel{L=3}{=} \frac{64g^6}{(q^2 + Q^2)^3}.$$

Additionally, the one-loop energy of twist-three operators differs from the the twist-two one. It is given by

$$\sum_{k=1}^M \frac{16}{1 + 4u_k^2} = 8S_1\left(\frac{M}{2}\right).$$

In the end, we can write the wrapping correction in a very elegant way as

$$\Delta\gamma = -64g^{10} S_1\left(\frac{M}{2}\right)^2 \sum_{Q=1}^{\infty} \int_{-\infty}^{\infty} \frac{dq}{2\pi} \frac{T_M(q, Q)^2}{R_M(q, Q)} \frac{64}{(q^2 + Q^2)^3}, \quad (5.2)$$

where R_M and T_M are functions given by the same expressions that are valid in the case of twist-two operators

$$\begin{aligned} R_M(q, Q) &= P_M\left(\frac{1}{2}(q - i(Q - 1))\right) P_M\left(\frac{1}{2}(q + i(Q - 1))\right) \\ &\quad \times P_M\left(\frac{1}{2}(q + i(Q + 1))\right) P_M\left(\frac{1}{2}(q - i(Q + 1))\right), \\ T_M(q, Q) &= \sum_{j=0}^{Q-1} \left[\frac{1}{2j - iq - Q} - \frac{1}{2(j + 1) - iq - Q} \right] P_M\left(\frac{1}{2}(q - i(Q - 1)) + ij\right). \end{aligned}$$

In order to obtain the wrapping contribution we calculated (5.2) for all even values of M up to $M = 40$. Assuming the maximal transcendentality principle, we expect the wrapping correction to have the following structure

$$\gamma^{\text{wrapping}}(M) = S_1\left(\frac{M}{2}\right)^2 (C_0(M)\zeta_7 + C_2(M)\zeta_5 + C_4(M)\zeta_3 + C_7(M)), \quad (5.3)$$

where the coefficients $C_n(M)$ have a degree of transcendentality n . We used the fact that S_1^2 is factored out in the Lüscher formula (5.2). Likewise to the case of the asymptotic Bethe Ansatz we are looking for coefficients that are linear combination of harmonic sums with positive indices. For a given degree of transcendentality n , there are 2^{n-1} independent sums. Thus, in order to obtain C_0 , C_2 and C_4 it is sufficient to know $\gamma^{\text{wrapping}}(M)$ to values of $M = 2$, $M = 4$ and $M = 16$, respectively, such that they can be determined from the results we computed. However, to unequivocally fix C_7 it is necessary to know $\gamma^{\text{wrapping}}(M)$ to values of the spin $M = 128$ which is far from our reach. Nevertheless, we can assume, as a natural refinement of the maximal transcendentality principle, that the coefficients of the

harmonic sums entering C_7 are integers. With this assumptions a result is easily found.⁴ The final result with all harmonic sums being of argument $M/2$ is given by

$$\begin{aligned} \gamma^{\text{wrapping}}(M) = & -64g^{10} S_1^2(35\zeta_7 - 40S_2\zeta_5 + (-8S_4 + 16S_{2,2})\zeta_3 \\ & + 2S_7 - 4S_{2,5} - 2S_{3,4} - 4S_{4,3} - 2S_{6,1} + 8S_{2,2,3} + 4S_{3,3,1}). \end{aligned} \quad (5.4)$$

For fixed values of M this result matches exactly numerical evaluations of the proposed Y -system [22].

Wrapping corrections, by their nature, should not modify the leading asymptotic behavior (4.1). The result (5.4) confirms this expectation, since the factor $S_1^2 \sim \log^2 M$ multiplies a linear combination of harmonic sums, which have a leading asymptotic behavior $\sim 1/M^2$. The first wrapping contribution to the asymptotic behavior therefore only enters at order $(\log^2 M/M^2)$ (see appendix B, formula (B.6)),

$$\gamma^{\text{wrapping}}(M) \sim -\left(768\zeta_3 - \frac{16\pi^4}{15}\right) \frac{\log^2 M}{M^2}, \quad \text{for } M \rightarrow \infty. \quad (5.5)$$

Thus, for large values of the spin wrapping corrections are of the same order as in the case of twist-two operators [18]. Further similarities with the asymptotic expansion of twist-two operators are discussed in appendix B.

In the previous section we stated that the asymptotic part given in table 1 is reciprocity invariant. Hence, for the complete anomalous dimension to be reciprocity invariant, (5.4) has to satisfy this property *separately*. Writing (5.4) in terms of $\underline{\Omega}$ and Ω

$$\gamma^{\text{wrapping}}(M) = -64g^{10} \Omega_1^2 \left(35\zeta_7 + 4\Omega_{3,3,1} + 8\underline{\Omega}_{2,2,3} + 24\underline{\zeta}_3 \underline{\Omega}_{2,2} - \Omega_7 \right), \quad (5.6)$$

one checks that this is indeed true, since according to the theorem in section 3.1 the appearing structures are all parity-invariant.

We conclude this section giving the prediction that our conjecture given in table 1 and (5.4), together with the formulas in (2.9)–(2.12), give for the five-loop anomalous dimension of the simplest twist-three operator with even spin ($M = 2$)

$$\gamma(2) = 8g^2 - 24g^4 + 136g^6 - (920 + 128\zeta_3)g^8 + 8(833 + 144\zeta_3 + 480\zeta_5 - 280\zeta_7)g^{10} + \mathcal{O}(g^{12}) \quad (5.7)$$

6 Analytic continuation

As already mentioned, no direct checks of the consistency of the multi-loop anomalous dimension (2.7) from its pole structure are possible.

However, it is worth to analyze the behavior of the anomalous dimension to four-loops (2.9)–(2.12) and the five-loop part given in table 1 and (5.4) at the singularity nearest

⁴As is usual in such kind of conjectures, there is a powerful numerical test that can be applied to any guesswork. Typically, one is able to compute spin dependent expressions like $C_7(M)$ up to a reasonable maximum value of M in exact (rational) form. On the other hand, numerical values can be obtained with a very high number of digits for quite larger values of M . Thus, given a conjectured expression obtained from data up to M_{max} one can always test it beyond that limit with a precision of several hundreds of digits. These kinds of tests are always passed by the expressions we derive in this paper.

to the origin, $M = -2$. Thus, we need the small ω expansion of general nested harmonic sums of the form

$$S_{a_1, \dots, a_d}(-1 + \omega), \quad a_i \in \mathbb{N}. \quad (6.1)$$

The analytic continuation we need can be obtained by observing that from the definition of harmonic sums it follows that

$$S_{a, \mathbf{b}}(-1 + \omega) = S_{a, \mathbf{b}}(\omega) - \frac{1}{\omega^a} S_{\mathbf{b}}(\omega). \quad (6.2)$$

This simple identity allows us to proceed by trivially expanding the *r.h.s* around $\omega = 0$. This is a straightforward task, once one makes use of the general formula for the derivatives of nested harmonic sums [18], and takes into account that $S_{\mathbf{a}}(0) = 0$.

The expansion of the n -th loop anomalous dimension γ_n has the general NNLO form

$$\gamma_n = a_n \omega^{1-2n} + b_n \zeta_2 \omega^{3-2n} + c_n \zeta_3 \omega^{4-2n} + \dots, \quad a_n, b_n, c_n \in \mathbb{Q}. \quad (6.3)$$

Up to four loops, the explicit formulas for the two highest terms (NLO) of the analytic continuation are given in [11].⁵ We recall them here for convenience, also adding the NNLO contribution

$$\gamma_2 = -\frac{8}{\omega} + 8\zeta_2\omega - 8\zeta_3\omega^2 + \dots, \quad \gamma_4 = -\frac{8}{\omega^3} + \frac{16\zeta_2}{\omega} + 16\zeta_3 + \dots, \quad (6.4)$$

$$\gamma_6 = -\frac{8}{\omega^5} + \frac{48\zeta_2}{\omega^3} + \frac{48\zeta_3}{\omega^2} + \dots, \quad \gamma_8 = -\frac{8}{\omega^7} + \frac{80\zeta_2}{\omega^5} + \frac{80\zeta_3}{\omega^4} + \dots. \quad (6.5)$$

In [11], an all-loop resummation at NLO was proposed⁶ and conjectured to be valid for the asymptotic γ^{ABA} part.

At five loops, we have found for the contributions of the γ^{ABA} and γ^{wrapping} parts, respectively, the expressions

$$\gamma_{10}^{\text{ABA}} = -\frac{136}{\omega^9} + \frac{496\zeta_2}{\omega^7} - \frac{784\zeta_3}{\omega^6} + \dots, \quad \gamma_{10}^{\text{wrapping}} = \frac{128}{\omega^9} - \frac{384\zeta_2}{\omega^7} - \frac{128\zeta_3}{\omega^6} + \dots. \quad (6.6)$$

The analytical continuation of the complete five-loop anomalous dimension is thus given by

$$\gamma_{10} = -\frac{8}{\omega^9} + \frac{112\zeta_2}{\omega^7} - \frac{912\zeta_3}{\omega^6} + \dots. \quad (6.7)$$

Interestingly enough, only the above formula for the *complete* anomalous dimension matches the proposed resummation *exactly*. The latter can therefore be rewritten as

$$\gamma_{\text{NLO}} = -8 \frac{g^2}{\omega} \left(\frac{1}{1-t} - \zeta_2 \frac{1+3t^2}{(1-t)^2} \omega^2 \right), \quad t = \frac{g^2}{\omega^2}, \quad (6.8)$$

with the equality valid in a perturbative sense. It is obviously tempting to extend such a conjecture to NNLO, trying to resum the poles that appear in (6.3) with ζ_3 as a coefficient. However, the five data-points available (one for each loop) hardly allow for a genuine resummation. In fact, the NNLO term in the above expression is likely to contain enough

⁵See eq. (5.12) there.

⁶See eq. (5.14) there.

more terms to be at least in a one-to-one correspondence with the available constraints. Nevertheless, we find intriguing that the following *simple* parameterization can be given, valid at five loops,

$$\gamma_{\text{NNLO}} = -8 \frac{g^2}{\omega} \left(\frac{1}{1-t} - \zeta_2 \frac{1+3t^2}{(1-t)^2} \omega^2 + \zeta_3 \frac{1-5t+3t^2+t^3+128t^4}{(1-t)^3} \omega^3 \right). \quad (6.9)$$

Acknowledgments

We benefited from discussion with Niklas Beisert, Yuri Dokshitzer, Romuald Janik, Gregory Korchemsky, Anatoly Kotikov, Giuseppe Marchesini, Domenico Seminara, Matthias Staudacher, Arkady A. Tseytlin and Pedro Vieira. The work of V. Forini is supported by the Alexander von Humboldt Foundation. The work of T. Lukowski is financed from Polish science funds during 2009-2011 as a research project. This research was supported in part by the National Science Foundation, under Grant No. PHY05-51164 (S.Z.).

A Harmonic sums

In this appendix we recall some useful formulas for harmonic sums *with positive indices*. The generalization to the case of arbitrary sign for the indices is treated in many references, for example [42] (see also appendix A of [18]).

The basic definition of nested harmonic sums S_{a_1, \dots, a_n} is recursive

$$S_a(N) = \sum_{n=1}^N \frac{1}{n^a}, \quad S_{a, \mathbf{b}}(N) = \sum_{n=1}^N \frac{1}{n^a} S_{\mathbf{b}}(n), \quad (A.1)$$

Given a particular sum $S_{\mathbf{a}} = S_{a_1, \dots, a_n}$ we define

$$\text{depth}(S_{\mathbf{a}}) = n, \quad (A.2)$$

$$\text{transcendentality}(S_{\mathbf{a}}) = \mathbf{a} \equiv a_1 + \dots + a_n. \quad (A.3)$$

For a product of S sums, we define transcendentality to be the sum of the transcendentalities of the factors.

Complementary harmonic sums are defined recursively by

$$\underline{S}_{\mathbf{a}} = S_{\mathbf{a}}, \quad (A.4)$$

$$\underline{S}_{\mathbf{a}} = S_{\mathbf{a}} - \sum_{k=1}^{\ell-1} S_{a_1, \dots, a_k} \underline{S}_{a_{k+1}, \dots, a_{\ell}}(\infty), \quad (A.5)$$

This definition is valid when the rightmost index of \mathbf{a} is not 1. Otherwise, the above recursive definition leads to a polynomial in the formal quantity $S_1(\infty)$. In this case our *definition* of $\underline{S}_{\mathbf{a}}$ prescribes to set $S_1(\infty) \rightarrow 0$ in the end.

B Analysis of the asymptotic structure of γ and P

Here we analyze the first few orders of the large M expansion of the twist-three anomalous dimension up to five-loops and its corresponding kernel P .⁷

The expansions of (2.9)–(2.12) to $\mathcal{O}(1/m^{-3})$ are given by

$$\gamma_2 = 8 \log \bar{m} + \frac{4}{m} - \frac{2}{3m^2} + \mathcal{O}\left(\frac{1}{m^4}\right), \quad (\text{B.1})$$

$$\begin{aligned} \gamma_4 = & -\frac{8}{3}\pi^2 \log \bar{m} - 8\zeta_3 + \frac{1}{m} \left[16 \log \bar{m} - \frac{4\pi^2}{3} \right] - \frac{1}{m^2} \left[8 \log \bar{m} - \frac{2\pi^2}{9} - 12 \right] \\ & + \frac{1}{m^3} \left[\frac{8}{3} \log \bar{m} - \frac{28}{3} \right] + \mathcal{O}\left(\frac{1}{m^4}\right), \end{aligned} \quad (\text{B.2})$$

$$\begin{aligned} \gamma_6 = & \frac{88}{45}\pi^4 \log \bar{m} - 8\zeta_5 + \frac{8}{3}\pi^2\zeta_3 - \frac{1}{m} \left[\frac{32}{3}\pi^2 \log \bar{m} + 16\zeta_3 - \frac{44\pi^4}{45} \right] \\ & - \frac{1}{m^2} \left[16 \log^2 \bar{m} - \left(32 + \frac{16\pi^2}{3} \right) \log \bar{m} - 8 + \frac{20\pi^2}{3} + \frac{22\pi^4}{135} - 8\zeta_3 \right] \\ & + \frac{1}{m^3} \left[16 \log^2 \bar{m} - \left(64 + \frac{16\pi^2}{9} \right) \log \bar{m} + 16 + \frac{44\pi^2}{9} - \frac{8\zeta_3}{3} \right] + \mathcal{O}\left(\frac{1}{m^4}\right), \end{aligned} \quad (\text{B.3})$$

$$\begin{aligned} \gamma_8 = & -\left(\frac{584\pi^6}{315} + 64\zeta_3^2 \right) \log \bar{m} - \frac{32}{15}\pi^4\zeta_3 + \frac{8}{3}\pi^2\zeta_5 + 440\zeta_7 + \frac{1}{m} \left[\frac{48\pi^4}{5} \log \bar{m} - \frac{292\pi^6}{315} + \frac{32}{3}\pi^2\zeta_3 \right. \\ & \left. - 32\zeta_3^2 - 16\zeta_5 \right] + \frac{1}{m^2} \left[16\pi^2 \log^2 \bar{m} - \left(64 + 32\pi^2 + \frac{24\pi^4}{5} - 64\zeta_3 \right) \log \bar{m} - 128 - \frac{8\pi^2}{3} \right. \\ & \left. + \frac{88\pi^4}{15} + \frac{146\pi^6}{945} - 32\zeta_3 - \frac{16}{3}\pi^2\zeta_3 + \frac{16\zeta_3^2}{3} + 8\zeta_5 \right] + \frac{1}{m^3} \left[\frac{64}{3} \log^3 \bar{m} - (96 + 16\pi^2) \log^2 \bar{m} \right. \\ & \left. + \left(96 + \frac{176\pi^2}{3} + \frac{8\pi^4}{5} - 64\zeta_3 \right) \log \bar{m} + 112 - \frac{56\pi^2}{3} - \frac{64\pi^4}{15} + 80\zeta_3 + \frac{16}{9}\pi^2\zeta_3 - \frac{8\zeta_5}{3} \right] \\ & + \mathcal{O}\left(\frac{1}{m^4}\right), \end{aligned} \quad (\text{B.4})$$

where $m = \frac{M}{2}$ and $\bar{m} = m \exp \gamma_E$.

At five loops, the large M expansion of table 1 and (5.4) leads to

$$\begin{aligned} \gamma_{10}^{\text{ABA}} = & \left(\frac{28384\pi^8}{14175} + \frac{128}{3}\pi^2\zeta_3^2 + 1280\zeta_3\zeta_5 \right) \log \bar{m} + \frac{2048}{945}\pi^6\zeta_3 + 64\zeta_3^3 + \frac{8}{45}\pi^4\zeta_5 - \frac{440}{3}\pi^2\zeta_7 \\ & - 7448\zeta_9 - \frac{1}{m} \left[\left(\frac{9472\pi^6}{945} + 256\zeta_3^2 \right) \log \bar{m} - \frac{14192\pi^8}{14175} + \frac{448}{45}\pi^4\zeta_3 - \frac{64}{3}\pi^2\zeta_3^2 - \frac{32}{3}\pi^2\zeta_5 \right. \\ & \left. - 640\zeta_3\zeta_5 - 880\zeta_7 \right] + \frac{1}{m^2} \left[\left(256 - \frac{272\pi^4}{15} - 128\zeta_3 \right) \log^2 \bar{m} + \left(1280 + \frac{128\pi^2}{3} \right. \right. \\ & \left. \left. + \frac{496\pi^4}{15} + \frac{4736\pi^6}{945} - 128\zeta_3 - \frac{160}{3}\pi^2\zeta_3 + 128\zeta_3^2 - 288\zeta_5 \right) \log \bar{m} + 1920 + \frac{128\pi^2}{3} \right] \end{aligned}$$

⁷In the case of higher twist $L > 2$, anomalous dimensions occupy a band [43]. In this paper we have considered the *minimal* anomalous dimension, see [44] for an asymptotic study of the full spectrum up to three loops.

$$\begin{aligned}
& + \frac{64\pi^4}{45} - \frac{128\pi^6}{21} - \frac{7096\pi^8}{42525} + 64\zeta_3 + 32\pi^2\zeta_3 + \frac{224}{45}\pi^4\zeta_3 - 208\zeta_3^2 - \frac{32}{9}\pi^2\zeta_3^2 - 32\zeta_5 \\
& - \frac{16}{3}\pi^2\zeta_5 - \frac{320}{3}\zeta_3\zeta_5 - 440\zeta_7 \Big] - \frac{1}{m^3} \left[\frac{256\pi^2}{9} \log^3 \bar{m} - \left(128\pi^2 \right. \right. \\
& + \frac{272\pi^4}{15} - 64\zeta_3 \Big) \log^2 \bar{m} + \left(768 + \frac{320\pi^2}{3} + \frac{2752\pi^4}{45} + \frac{4736\pi^6}{2835} - 320\zeta_3 - \frac{160}{3}\pi^2\zeta_3 \right. \\
& + \frac{128\zeta_3^2}{3} - 288\zeta_5 \Big) \log \bar{m} + 1536 + 32\pi^2 - \frac{904\pi^4}{45} - \frac{1792\pi^6}{405} + 160\zeta_3 \\
& \left. + \frac{208}{3}\pi^2\zeta_3 + \frac{224}{135}\pi^4\zeta_3 - \frac{496\zeta_3^2}{3} + 96\zeta_5 - \frac{16}{9}\pi^2\zeta_5 - \frac{440\zeta_7}{3} \right] + \mathcal{O}\left(\frac{1}{m^4}\right), \quad (\text{B.5})
\end{aligned}$$

$$\gamma^{\text{wrapping}} = - \left(768\zeta_3 - \frac{16\pi^4}{15} \right) \frac{\log^2 \bar{m}}{m^2} + \left(768\zeta_3 - \frac{16\pi^4}{15} \right) (\log^2 \bar{m} - \log \bar{m}) \frac{1}{m^3} + \mathcal{O}\left(\frac{1}{m^4}\right) \quad (\text{B.6})$$

As expected in the case of minimal anomalous dimension for operators of twist $L \leq 3$, logarithmic enhancements in the asymptotic expansions of γ are always positive in power.⁸

Notice that, when expressed in terms of the variable $M = 2m$, the maximal logarithmic terms $\log^p m/m^p$ in the expansions up to four loops, formulas (B.1)–(B.4), are compatible with a resummation of type

$$\gamma(M) = f(g) \log \left(M + \frac{1}{2}f(g) \log M + \dots \right) + \dots \quad (\text{B.7})$$

According to this, their coefficients are simply proportional to f^{m+1}

$$\gamma(M) \sim f \log M + \frac{f^2}{2} \frac{\log M}{M} - \frac{f^3}{8} \frac{\log^2 M}{M^2} + \dots \quad (\text{B.8})$$

where f is the universal scaling function, whose weak coupling expansion to five-loop order can be found in [7]. At five loops, the pattern (B.8) is broken by the term $\log^2 M/M^2$ in the expansion (B.5) above.⁹ Interestingly, it is precisely at this order in the large M expansion that wrapping corrections start to contribute. Explicitly, while on the basis of (B.8) one would expect at five loops a term of type

$$(c_{22})_5 \frac{\log^2 M}{M^2} \quad \text{with} \quad (c_{22})_5^{\text{naive}} = \left(-\frac{f^3}{8}\right)_5 = -\frac{1024}{15} \pi^4 = -6144 \zeta_4, \quad (\text{B.9})$$

reexpressing (B.5) and (B.6) in terms of M one finds

$$(c_{22})_5^{\text{ABA}} = 1024 - 512\zeta_3 - 6528\zeta_4 \quad \text{and} \quad (c_{22})_5^{\text{wrapping}} = -3072 \zeta_3 + 384 \zeta_4. \quad (\text{B.10})$$

⁸Terms with negative powers of the logarithm appear for twist $L \leq 3$, but for *non-minimal* anomalous dimensions [45]. For twist $L > 3$ terms $\sim \frac{1}{\log^p M}$ are present both for large L and M [46] and for finite twist, see a similar discussion in [47] and reference therein. For a general method to derive higher order terms in the $1/M$ expansion at fixed L see [48].

⁹Further maximal logarithmic terms as $\log^3 M/M^3, \log^4 M/M^4$ continue to obey the rule as dictated by further orders in (B.8). Coefficients of terms $\log^p M/M^p$ with $p > 4$ are absent in the expansion, as checked up to $1/M^{25}$. This is again consistent with (B.8), being such coefficients of the form $\frac{f^6}{160}, -\frac{f^7}{384}, \dots$, they would contribute starting at 6 loops.

The sum of these terms does clearly not reproduce (B.9). This is analogous to the case of twist two operators [18].¹⁰

It is interesting to notice that the structure, lost at higher orders in $1/m$, of the first terms in the expansion for γ^{wrapping} is also present in its twist-two analogue. The appearance of an overall coefficient multiplying the $1/m^2$ and $1/m^3$ terms was already noticed in formula (C.5) of [18]. Another analogy between the leading asymptotic behavior of twist-2 and twist-3 wrapping contributions is their negative sign and their common pattern $\sim c_n \zeta_n + c_{n+1} \zeta_{n+1}$, where n coincides with the twist.

Concerning the P -kernel, the logarithmic structure to four loops is remarkably *simpler* than the one of the corresponding anomalous dimension, as it is only *linear* in $\log M$ [36]. In particular, there are no maximally enhanced terms of the form $(\log M/M)^k$. As discussed in [32, 33], this feature of P translates into the chance of a resummation of type (B.7). This asymptotic structure changes at five loops.

The P function, derived by inverting formula (3.1), reads in terms of $m = \frac{M}{2}$ ($\partial \equiv \partial_m$) to five-loops

$$\mathcal{P}(m) = \sum_{k=1}^{\infty} \frac{1}{k!} \left(-\frac{1}{4}\partial\right)^{k-1} [\gamma(m)]^k = \gamma - \frac{1}{8}(\gamma^2)' + \frac{1}{96}(\gamma^3)'' - \frac{1}{1536}(\gamma^4)''' + \frac{1}{30720}(\gamma^5)'''' + \dots$$

Replacing γ by the perturbative expansion (2.7) we can formally write at five loops

$$P_{10} = \gamma_{10} - \frac{1}{4}(\gamma_4\gamma_6 + \gamma_2\gamma_8)' + \frac{1}{32}(\gamma_2\gamma_4^2 + \gamma_2^2\gamma_6)'' - \frac{1}{384}(\gamma_2^3\gamma_4)''' + \frac{1}{30720}(\gamma_2^5)'''' \quad (\text{B.11})$$

Expanded at large M , including the wrapping contribution, this becomes

$$\begin{aligned} P_{10} = & \left(\frac{28384\pi^8}{14175} + \frac{128}{3}\pi^2\zeta_3^2 + 1280\zeta_3\zeta_5 \right) \log \bar{m} + \frac{2048}{945}\pi^6\zeta_3 + 64\zeta_3^3 + \frac{8}{45}\pi^4\zeta_5 - \frac{440}{3}\pi^2\zeta_7 \\ & - 7448\zeta_9 + \frac{1}{m} \left[\frac{14192\pi^8}{14175} + \frac{64}{3}\pi^2\zeta_3^2 + 640\zeta_3\zeta_5 \right] + \frac{1}{m^2} \left[(256 - 896\zeta_3) \log^2 \bar{m} + \right. \\ & + \left(1280 + \frac{128\pi^2}{3} - \frac{16\pi^4}{15} - 128\zeta_3 - \frac{64}{3}\pi^2\zeta_3 - 320\zeta_5 \right) \log \bar{m} + 1920 + \frac{128\pi^2}{3} + \frac{64\pi^4}{45} + \\ & \left. - \frac{1024\pi^6}{945} - \frac{7096\pi^8}{42525} + 64\zeta_3 - 64\zeta_3^2 - \frac{32}{9}\pi^2\zeta_3^2 - \frac{320}{3}\zeta_3\zeta_5 \right] \\ & - \frac{1}{m^3} \left[(256 - 896\zeta_3) \log^2 \bar{m} + \left(1024 + \frac{128\pi^2}{3} - \frac{16\pi^4}{15} + 768\zeta_3 - \frac{64}{3}\pi^2\zeta_3 - 320\zeta_5 \right) \log \bar{m} \right. \\ & \left. + 1280 + \frac{64\pi^2}{3} + \frac{88\pi^4}{45} - \frac{1024\pi^6}{945} + 128\zeta_3 + \frac{32}{3}\pi^2\zeta_3 - 64\zeta_3^2 + 160\zeta_5 \right] + \mathcal{O}\left(\frac{1}{m^4}\right). \quad (\text{B.12}) \end{aligned}$$

The “simplicity” feature is lost, because at order $1/m^2$ a term $\log^2 m/m^2$ appears, which is responsible for the above formula (B.10).

We recall that the consequences (B.7) and (B.8) of the simplicity of the P function and the knowledge of f to presumably all loops [7] allow in principle an all-loop prediction for

¹⁰One difference is however that the degree of transcendentality of the asymptotic and wrapping contributions, that differs in (B.10), is the same in the twist-two case.

such maximal logarithmic terms, whose coefficients should be simply proportional to f^{m+1} . Indeed, such *inheritance* has been checked at strong coupling in [47] up to one-loop in the semiclassical sigma model expansion, as well as in [49] at the classical level. An independent strong coupling confirmation of (B.7) for twist-two operators has recently been given in [24]. To clarify if and how the difference in the simplicity of the P at weak and strong coupling works, further orders in the semiclassical sigma model expansion would be needed.

We conclude the appendix by reporting the separate contributions to P_{10} coming from the dressing factor, which obey the property described in point 4 of section 4. They read

$$\begin{aligned}
 P_{10}^{(\zeta_3)} \equiv \gamma_{10}^{(\zeta_3)} - \frac{1}{4}(\gamma_2 \gamma_8^{(\zeta_3)})' &= 896S_6 - 2304S_{1,5} - 1792S_{2,4} - 768S_{3,3} - 1792S_{4,2} - 2304S_{5,1} \\
 &+ 2560S_{1,1,4} + 512S_{1,2,3} + 1536S_{1,3,2} + 3584S_{1,4,1} + 512S_{2,1,3} + 1536S_{2,3,1} + 512S_{3,1,2} \\
 &+ 512S_{3,2,1} + 2560S_{4,1,1} - 2048S_{1,1,3,1} - 2048S_{1,3,1,1} + 512\zeta_2 S_1 S_3 - 512S_1 S_2 S_3 \\
 &+ 768\zeta_4 S_1^2 - 768S_1^2 S_4, \tag{B.13}
 \end{aligned}$$

$$P_{10}^{(\zeta_5)} \equiv \gamma_{10}^{(\zeta_5)} = 1280 S_1 S_3. \tag{B.14}$$

In the first line we included the dressing-induced contribution at four loops, which is proportional to ζ_3 . These contributions can be expressed in terms of parity invariant combinations (see Theorem (c) in section 3.1) as

$$P_{10}^{(\zeta_3)} = 256(8\Omega_1 \Omega_{1,1,3} - 4\Omega_1^2 \Omega_{1,3} + \Omega_{3,3} + S_1 \Omega_5 + 2\zeta_2 \Omega_1 \Omega_3 + 3\zeta_4 \Omega_1^2), \tag{B.15}$$

$$P_{10}^{(\zeta_5)} = 1280 \Omega_1 \Omega_3. \tag{B.16}$$

References

- [1] J.A. Minahan and K. Zarembo, *The Bethe-ansatz for $\mathcal{N} = 4$ super Yang-Mills*, *JHEP* **03** (2003) 013 [[hep-th/0212208](#)] [[SPIRES](#)];
 I. Bena, J. Polchinski and R. Roiban, *Hidden symmetries of the $AdS_5 \times S^5$ superstring*, *Phys. Rev. D* **69** (2004) 046002 [[hep-th/0305116](#)] [[SPIRES](#)];
 N. Beisert, C. Kristjansen and M. Staudacher, *The dilatation operator of $\mathcal{N} = 4$ super Yang-Mills theory*, *Nucl. Phys. B* **664** (2003) 131 [[hep-th/0303060](#)] [[SPIRES](#)].
- [2] N. Beisert and M. Staudacher, *Long-range $PSU(2, 2|4)$ Bethe ansätze for gauge theory and strings*, *Nucl. Phys. B* **727** (2005) 1 [[hep-th/0504190](#)] [[SPIRES](#)].
- [3] M. Staudacher, *The factorized S -matrix of CFT/AdS* , *JHEP* **05** (2005) 054 [[hep-th/0412188](#)] [[SPIRES](#)].
- [4] N. Beisert, *The $SU(2|2)$ dynamic S -matrix*, *Adv. Theor. Math. Phys.* **12** (2008) 945 [[hep-th/0511082](#)] [[SPIRES](#)].
- [5] R.A. Janik, *The $AdS_5 \times S^5$ superstring worldsheet S -matrix and crossing symmetry*, *Phys. Rev. D* **73** (2006) 086006 [[hep-th/0603038](#)] [[SPIRES](#)].
- [6] N. Beisert, R. Hernandez and E. Lopez, *A crossing-symmetric phase for $AdS_5 \times S^5$ strings*, *JHEP* **11** (2006) 070 [[hep-th/0609044](#)] [[SPIRES](#)].
- [7] N. Beisert, B. Eden and M. Staudacher, *Transcendentality and crossing*, *J. Stat. Mech.* (2007) P01021 [[hep-th/0610251](#)] [[SPIRES](#)].

- [8] Z. Bern, M. Czakon, L.J. Dixon, D.A. Kosower and V.A. Smirnov, *The four-loop planar amplitude and cusp anomalous dimension in maximally supersymmetric Yang-Mills theory*, *Phys. Rev. D* **75** (2007) 085010 [[hep-th/0610248](#)] [[SPIRES](#)];
F. Cachazo, M. Spradlin and A. Volovich, *Four-loop cusp anomalous dimension from obstructions*, *Phys. Rev. D* **75** (2007) 105011 [[hep-th/0612309](#)] [[SPIRES](#)].
- [9] A. Rej, M. Staudacher and S. Zieme, *Nesting and dressing*, *J. Stat. Mech.* (2007) P08006 [[hep-th/0702151](#)] [[SPIRES](#)];
K. Sakai and Y. Satoh, *Origin of dressing phase in $\mathcal{N} = 4$ super Yang-Mills*, *Phys. Lett. B* **661** (2008) 216 [[hep-th/0703177](#)] [[SPIRES](#)];
R.A. Janik and T. Lukowski, *From nesting to dressing*, *Phys. Rev. D* **78** (2008) 066018 [[arXiv:0804.4295](#)] [[SPIRES](#)].
- [10] J. Ambjørn, R.A. Janik and C. Kristjansen, *Wrapping interactions and a new source of corrections to the spin-chain/string duality*, *Nucl. Phys. B* **736** (2006) 288 [[hep-th/0510171](#)] [[SPIRES](#)].
- [11] A.V. Kotikov, L.N. Lipatov, A. Rej, M. Staudacher and V.N. Velizhanin, *Dressing and wrapping*, *J. Stat. Mech.* (2007) P10003 [[arXiv:0704.3586](#)] [[SPIRES](#)].
- [12] F. Fiamberti, A. Santambrogio, C. Sieg and D. Zanon, *Wrapping at four loops in $N = 4$ SYM*, *Phys. Lett. B* **666** (2008) 100 [[arXiv:0712.3522](#)] [[SPIRES](#)].
- [13] Z. Bajnok and R.A. Janik, *Four-loop perturbative Konishi from strings and finite size effects for multiparticle states*, *Nucl. Phys. B* **807** (2009) 625 [[arXiv:0807.0399](#)] [[SPIRES](#)].
- [14] V.N. Velizhanin, *The four-loop Konishi in $N = 4$ SYM*, [arXiv:0808.3832](#) [[SPIRES](#)].
- [15] M. Lüscher, *Volume dependence of the energy spectrum in massive quantum field theories. 1. Stable particle states*, *Commun. Math. Phys.* **104** (1986) 177 [[SPIRES](#)].
- [16] R.A. Janik and T. Lukowski, *Wrapping interactions at strong coupling — the giant magnon*, *Phys. Rev. D* **76** (2007) 126008 [[arXiv:0708.2208](#)] [[SPIRES](#)].
- [17] Z. Bajnok, R.A. Janik and T. Lukowski, *Four loop twist two, BFKL, wrapping and strings*, [arXiv:0811.4448](#) [[SPIRES](#)].
- [18] M. Beccaria and V. Forini, *Four loop reciprocity of twist two operators in $N = 4$ SYM*, [arXiv:0901.1256](#) [[SPIRES](#)].
- [19] V.N. Velizhanin, *Leading transcendentality contributions to the four-loop universal anomalous dimension in $N = 4$ SYM*, [arXiv:0811.0607](#) [[SPIRES](#)].
- [20] G. Arutyunov and S. Frolov, *On string S -matrix, bound states and TBA*, *JHEP* **12** (2007) 024 [[arXiv:0710.1568](#)] [[SPIRES](#)]; *String hypothesis for the $AdS_5 \times S^5$ mirror*, [arXiv:0901.1417](#) [[SPIRES](#)].
- [21] N. Gromov, V. Kazakov and P. Vieira, *Finite volume spectrum of 2D field theories from Hirota dynamics*, [arXiv:0812.5091](#) [[SPIRES](#)].
- [22] N. Gromov, V. Kazakov and P. Vieira, *Integrability for the full spectrum of planar AdS/CFT*, [arXiv:0901.3753](#) [[SPIRES](#)].
- [23] A.V. Kotikov and L.N. Lipatov, *DGLAP and BFKL evolution equations in the $N = 4$ supersymmetric gauge theory*, *Nucl. Phys. B* **661** (2003) 19 [Erratum *ibid.* **B 685** (2004) 405] [[hep-ph/0208220](#)] [[SPIRES](#)].
- [24] L. Freyhult and S. Zieme, *The virtual scaling function of AdS/CFT*, [arXiv:0901.2749](#) [[SPIRES](#)].

- [25] D. Fioravanti, P. Grinza and M. Rossi, *Beyond cusp anomalous dimension from integrability*, [arXiv:0901.3161](#) [[SPIRES](#)].
- [26] N. Beisert, V. Dippel and M. Staudacher, *A novel long range spin chain and planar $N = 4$ super Yang-Mills*, *JHEP* **07** (2004) 075 [[hep-th/0405001](#)] [[SPIRES](#)].
- [27] A.V. Kotikov, A. Rej and S. Zieme, *Analytic three-loop solutions for $N = 4$ SYM twist operators*, *Nucl. Phys. B* **813** (2009) 460 [[arXiv:0810.0691](#)] [[SPIRES](#)].
- [28] A.V. Belitsky, G.P. Korchemsky and D. Mueller, *Towards Baxter equation in supersymmetric Yang-Mills theories*, *Nucl. Phys. B* **768** (2007) 116 [[hep-th/0605291](#)] [[SPIRES](#)];
A.V. Belitsky, *Long-range SL(2) Baxter equation in $N = 4$ super-Yang-Mills theory*, *Phys. Lett. B* **643** (2006) 354 [[hep-th/0609068](#)] [[SPIRES](#)].
- [29] M. Beccaria, *Anomalous dimensions at twist-3 in the SL(2) sector of $N = 4$ SYM*, *JHEP* **06** (2007) 044 [[arXiv:0704.3570](#)] [[SPIRES](#)].
- [30] M. Beccaria and V. Forini, *Anomalous dimensions of finite size field strength operators in $N = 4$ SYM*, *JHEP* **11** (2007) 031 [[arXiv:0710.0217](#)] [[SPIRES](#)].
- [31] Y.L. Dokshitzer, G. Marchesini and G.P. Salam, *Revisiting parton evolution and the large- x limit*, *Phys. Lett. B* **634** (2006) 504 [[hep-ph/0511302](#)] [[SPIRES](#)].
- [32] Y.L. Dokshitzer and G. Marchesini, *$N = 4$ SUSY Yang-Mills: three loops made simple(r)*, *Phys. Lett. B* **646** (2007) 189 [[hep-th/0612248](#)] [[SPIRES](#)].
- [33] B. Basso and G.P. Korchemsky, *Anomalous dimensions of high-spin operators beyond the leading order*, *Nucl. Phys. B* **775** (2007) 1 [[hep-th/0612247](#)] [[SPIRES](#)].
- [34] M. Beccaria and V. Forini, *Reciprocity of gauge operators in $N = 4$ SYM*, *JHEP* **06** (2008) 077 [[arXiv:0803.3768](#)] [[SPIRES](#)].
- [35] V. Forini and M. Beccaria, *QCD-like properties for anomalous dimensions in $N = 4$ SYM*, talk at the conference *Nonlinear Physics. Theory and Experiment. V*, Gallipoli Italy June 12–21 2008, [[arXiv:0810.0101](#)] [[SPIRES](#)].
- [36] M. Beccaria, Y.L. Dokshitzer and G. Marchesini, *Twist 3 of the SL(2) sector of $N = 4$ SYM and reciprocity respecting evolution*, *Phys. Lett. B* **652** (2007) 194 [[arXiv:0705.2639](#)] [[SPIRES](#)].
- [37] M. Beccaria, *Three loop anomalous dimensions of twist-3 gauge operators in $N = 4$ SYM*, *JHEP* **09** (2007) 023 [[arXiv:0707.1574](#)] [[SPIRES](#)].
- [38] A.V. Belitsky, A.S. Gorsky and G.P. Korchemsky, *Gauge/string duality for QCD conformal operators*, *Nucl. Phys. B* **667** (2003) 3 [[hep-th/0304028](#)] [[SPIRES](#)].
- [39] B. Eden and M. Staudacher, *Integrability and transcendentality*, *J. Stat. Mech.* (2006) P11014 [[hep-th/0603157](#)] [[SPIRES](#)].
- [40] L. Freyhult, A. Rej and M. Staudacher, *A generalized scaling function for AdS/CFT*, *J. Stat. Mech.* (2008) P07015 [[arXiv:0712.2743](#)] [[SPIRES](#)].
- [41] D. Bombardelli, D. Fioravanti and M. Rossi, *Large spin corrections in $\mathcal{N} = 4$ SYM SL(2): still a linear integral equation*, *Nucl. Phys. B* **810** (2009) 460 [[arXiv:0802.0027](#)] [[SPIRES](#)].
- [42] J. Blumlein, *Algebraic relations between harmonic sums and associated quantities*, *Comput. Phys. Commun.* **159** (2004) 19 [[hep-ph/0311046](#)] [[SPIRES](#)].
- [43] A.V. Belitsky, V.M. Braun, A.S. Gorsky and G.P. Korchemsky, *Integrability in QCD and beyond*, *Int. J. Mod. Phys. A* **19** (2004) 4715 [[hep-th/0407232](#)] [[SPIRES](#)].

- [44] A.V. Belitsky, G.P. Korchemsky and R.S. Pasechnik, *Fine structure of anomalous dimensions in $N = 4$ super Yang-Mills theory*, *Nucl. Phys. B* **809** (2009) 244 [[arXiv:0806.3657](#)] [[SPIRES](#)].
- [45] V.M. Braun, S.E. Derkachov, G.P. Korchemsky and A.N. Manashov, *Baryon distribution amplitudes in QCD*, *Nucl. Phys. B* **553** (1999) 355 [[hep-ph/9902375](#)] [[SPIRES](#)];
S.E. Derkachov, G.P. Korchemsky and A.N. Manashov, *Evolution equations for quark gluon distributions in multi-color QCD and open spin chains*, *Nucl. Phys. B* **566** (2000) 203 [[hep-ph/9909539](#)] [[SPIRES](#)].
- [46] A.V. Belitsky, A.S. Gorsky and G.P. Korchemsky, *Logarithmic scaling in gauge/string correspondence*, *Nucl. Phys. B* **748** (2006) 24 [[hep-th/0601112](#)] [[SPIRES](#)].
- [47] M. Beccaria, V. Forini, A. Tirziu and A.A. Tseytlin, *Structure of large spin expansion of anomalous dimensions at strong coupling*, *Nucl. Phys. B* **812** (2009) 144 [[arXiv:0809.5234](#)] [[SPIRES](#)].
- [48] M. Beccaria and F. Catino, *Large spin expansion of the long-range Baxter equation in the $SL(2)$ sector of $N = 4$ SYM*, *JHEP* **01** (2008) 067 [[arXiv:0710.1991](#)] [[SPIRES](#)].
- [49] R. Ishizeki, M. Kruczenski, A. Tirziu and A.A. Tseytlin, *Spiky strings in $AdS_3 \times S^1$ and their AdS -pp-wave limits*, *Phys. Rev. D* **79** (2009) 026006 [[arXiv:0812.2431](#)] [[SPIRES](#)].