

Complex Langevin dynamics at finite chemical potential: mean field analysis in the relativistic Bose gas

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

JHEP05(2009)052

(<http://iopscience.iop.org/1126-6708/2009/05/052>)

[The Table of Contents](#) and [more related content](#) is available

Download details:

IP Address: 80.92.225.132

The article was downloaded on 03/04/2010 at 09:19

Please note that [terms and conditions apply](#).

Complex Langevin dynamics at finite chemical potential: mean field analysis in the relativistic Bose gas

Gert Aarts

*Department of Physics, Swansea University,
Singleton Park, Swansea SA2 8PP, U.K.*

E-mail: g.aarts@swan.ac.uk

ABSTRACT: Stochastic quantization can potentially be used to simulate theories with a complex action due to a nonzero chemical potential. We study complex Langevin dynamics in the relativistic Bose gas analytically, using a mean field approximation. We concentrate on the region with a Silver Blaze problem and discuss convergence, stability, fixed points, and the severeness of the sign problem. The real distribution satisfying the extended Fokker-Planck equation is constructed and its nonlocal form is explained. Finally, we compare the mean field results in finite volume with the numerical data presented in ref. [1].

KEYWORDS: Lattice Quantum Field Theory, Lattice QCD

ARXIV EPRINT: [0902.4686](https://arxiv.org/abs/0902.4686)

Contents

1	Introduction	1
2	Relativistic Bose gas and Langevin dynamics	2
3	Ignoring interactions	5
3.1	Standard results	5
3.2	Continuous Langevin dynamics	6
3.3	Discretized Langevin dynamics	8
4	Fokker-Planck equation	9
5	Mean field approximation	11
6	Conclusion	15
A	Dispersion relation	16

1 Introduction

Theories with a complex action are not easy to solve numerically, since approaches based on importance sampling break down. This is commonly referred to as the sign problem. An important theory in this class is QCD at finite baryon chemical potential, with a complex fermion determinant satisfying $[\det M(\mu)]^* = \det M(-\mu)$.¹ Several methods have been devised to circumvent the sign problem in QCD, mostly at small chemical potential and in the vicinity of the crossover between the confined and the deconfined phase [2–14]. For a detailed lattice QCD study of the sign problem at small chemical potential, see ref. [15]. Considerable insight in the QCD sign problem has also been obtained with Random Matrix Theory [16–21]. In some theories the sign problem can be eliminated altogether, using a reformulation in terms of different degrees of freedom [22–24].

Since stochastic quantization [25] does not rely on importance sampling, it can potentially be applied to theories with a complex action using complex Langevin dynamics [26, 27]. Studies in the 80’s, however, have given mixed results, see e.g. refs. [28, 29]. For an extensive review and more references, see ref. [30]. Recently the approach was reconsidered as a method to solve nonequilibrium quantum fields dynamics in Minkowski spacetime [31–33]. It was shown that instabilities, which plagued earlier studies, can be controlled by using small enough Langevin stepsizes. Moreover, insight in the convergence properties of the method can be obtained from features of classical flow diagrams. Other recent applications include PT symmetric theories [34] and unbounded actions [35]. In

¹In case of a complex chemical potential, this relation becomes $[\det M(\mu)]^* = \det M(-\mu^*)$.

ref. [36] we applied stochastic quantization to various theories with a nonzero chemical potential. In particular, we considered QCD with static quarks, in which the fermion determinant is approximated but the full gauge dynamics is preserved. First results on a 4^4 lattice are encouraging. The required extension from $SU(3)$ to $SL(3, \mathbb{C})$ is discussed in detail.

The sign problem in QCD at finite chemical potential does not arise because of the anticommuting nature of the quark fields. Also in bosonic theories with a nonzero chemical potential and an action that behaves under complex conjugation as $S^*(\mu) = S(-\mu)$, the sign problem appears. In ref. [1] we considered the relativistic Bose gas (a self interacting complex scalar field) in four dimensions in the presence of a chemical potential as one of the simplest examples of a relativistic field theory with a severe sign problem. Like QCD, this theory has a Silver Blaze problem [37]: at strictly zero temperature and small chemical potential, bulk physical observables are independent of the chemical potential, even though it enters explicitly in the microscopic dynamics. At larger chemical potential, the system enters a Bose condensed phase. The μ independence below onset and the formation of a state with nonzero density above onset is similar to what is expected to occur in QCD at zero temperature. It was demonstrated in ref. [1] that complex Langevin dynamics reproduces the expected physics, on lattices of size N^4 , with $N = 4, 6, 8, 10$. The sign problem was shown to be severe. However, no obstacles related to the sign problem, the Silver Blaze problem, or in taking the thermodynamic limit were encountered.

In this paper we complement the numerical study of ref. [1] with a detailed analytical study in the mean field approximation. We concentrate on the region with the Silver Blaze problem. The paper is organized as follows. In section 2 we remind the reader of the model and the corresponding complex Langevin equations. In order to prepare for the mean field analysis, we first discuss the case without interactions. In section 3 we summarize the exact results in the free field limit, using standard field theory. Subsequently the free Langevin equations are solved analytically, both for continuous and discretized dynamics. We discuss convergence and stability properties. In section 4 the stationary solution of the Fokker-Planck equation is given, again ignoring interactions, and shown to be in agreement with the solution of the Langevin equations in the limit of large Langevin time.² The criterium for convergence is derived from this distribution as well. We find that the real probability distribution is highly nonlocal and explain why. In section 5 interactions are included and the analysis is extended to the mean field approximation. We derive fixed points of the mean field Langevin equations at finite Langevin stepsize. Finally the mean field predictions in finite volume are compared with the nonperturbative results obtained by complex Langevin simulations [1]. The appendix contains a short remark about lattice dispersion relations.

2 Relativistic Bose gas and Langevin dynamics

We consider a self-interacting complex scalar field in the presence of a chemical potential μ , with the continuum action

$$S = \int d^4x \left[|\partial_\nu \phi|^2 + (m^2 - \mu^2) |\phi|^2 + \mu (\phi^* \partial_4 \phi - \partial_4 \phi^* \phi) + \lambda |\phi|^4 \right]. \quad (2.1)$$

²Recently, an interesting approach to study stationary solutions of complex Langevin dynamics was presented in ref. [38].

The euclidean action is complex and satisfies $S^*(\mu) = S(-\mu)$. We take $m^2 > 0$, so that at vanishing and small μ the theory is in its symmetric phase.

We study this theory on the lattice, with the action

$$S = \sum_x \left[(2d + m^2) \phi_x^* \phi_x + \lambda (\phi_x^* \phi_x)^2 - \sum_{\nu=1}^4 \left(\phi_x^* e^{-\mu \delta_{\nu,4}} \phi_{x+\hat{\nu}} + \phi_{x+\hat{\nu}}^* e^{\mu \delta_{\nu,4}} \phi_x \right) \right]. \quad (2.2)$$

As always, chemical potential is introduced as an imaginary constant vector potential in the temporal direction [39]. The number of euclidean dimensions is $d = 4$, the lattice spacing $a_{\text{lat}} \equiv 1$, and the lattice four-volume is $\Omega = N_s^3 N_\tau$, where N_s (N_τ) are the number of sites in a spatial (temporal) direction. We use periodic boundary conditions.

In order to formulate the complex Langevin equations for this theory, the complex field is first written in terms of two real fields ϕ_a ($a = 1, 2$) as $\phi = \frac{1}{\sqrt{2}}(\phi_1 + i\phi_2)$. The lattice action then reads

$$S = \sum_x \left[\frac{1}{2} (2d + m^2) \phi_{a,x}^2 + \frac{\lambda}{4} (\phi_{a,x}^2)^2 - \sum_{i=1}^3 \phi_{a,x} \phi_{a,x+\hat{i}} - \cosh \mu \phi_{a,x} \phi_{a,x+\hat{4}} + i \sinh \mu \varepsilon_{ab} \phi_{a,x} \phi_{b,x+\hat{4}} \right]. \quad (2.3)$$

We use the antisymmetric tensor ε_{ab} , with $\varepsilon_{12} = -\varepsilon_{21} = 1$, $\varepsilon_{11} = \varepsilon_{22} = 0$, and summation over repeated indices is implied throughout.

Since the Boltzmann weight e^{-S} in the partition function,

$$Z = \int D\phi_1 D\phi_2 e^{-S}, \quad (2.4)$$

is complex, the theory has a sign problem and one cannot rely on importance sampling. Writing the weight as $|e^{-S}| e^{i\varphi} = e^{-S_R} e^{-iS_I}$, one may consider the phase quenched theory

$$Z_{\text{pq}} = \int D\phi_1 D\phi_2 |e^{-S}| = \int D\phi_1 D\phi_2 e^{-S_R}, \quad (2.5)$$

where S_R is the real part of the action in eq. (2.3), i.e. the term proportional to $\sinh \mu$ is dropped. By analysing the average phase factor in the phase quenched theory, $\langle e^{i\varphi} \rangle_{\text{pq}}$, it was shown in ref. [1] that this theory has a severe sign problem: at nonzero chemical potential the average phase factor goes to zero exponentially fast in the thermodynamic limit.

We use stochastic quantization. The Langevin equations for the fields ϕ_a read

$$\frac{\partial}{\partial \theta} \phi_{a,x}(\theta) = -\frac{\delta S[\phi]}{\delta \phi_{a,x}(\theta)} + \eta_{a,x}(\theta), \quad (2.6)$$

where θ is the Langevin time. The noise η is Gaussian and normalized as

$$\langle \eta_{a,x}(\theta) \rangle = 0, \quad \langle \eta_{a,x}(\theta) \eta_{b,x'}(\theta') \rangle = 2\delta_{ab} \delta_{xx'} \delta(\theta - \theta'). \quad (2.7)$$

Since the force in eq. (2.6) is complex, the fields are complexified as

$$\phi_a \rightarrow \phi_a^{\text{R}} + i\phi_a^{\text{I}} \quad (a = 1, 2). \quad (2.8)$$

The complex Langevin equations we consider in this paper then read

$$\frac{\partial}{\partial \theta} \phi_{a,x}^{\text{R}}(\theta) = K_{a,x}^{\text{R}}(\theta) + \eta_{a,x}(\theta), \quad (2.9)$$

$$\frac{\partial}{\partial \theta} \phi_{a,x}^{\text{I}}(\theta) = K_{a,x}^{\text{I}}(\theta). \quad (2.10)$$

The noise is chosen to be real. The drift terms are defined as

$$K_{a,x}^{\text{R}} = -\text{Re} \left. \frac{\delta S}{\delta \phi_{a,x}} \right|_{\phi_a \rightarrow \phi_a^{\text{R}} + i\phi_a^{\text{I}}}, \quad (2.11)$$

$$K_{a,x}^{\text{I}} = -\text{Im} \left. \frac{\delta S}{\delta \phi_{a,x}} \right|_{\phi_a \rightarrow \phi_a^{\text{R}} + i\phi_a^{\text{I}}}, \quad (2.12)$$

and read explicitly

$$\begin{aligned} K_{a,x}^{\text{R}} = & - [2d + m^2 + \lambda (\phi_{b,x}^{\text{R}2} - \phi_{b,x}^{\text{I}2})] \phi_{a,x}^{\text{R}} + 2\lambda \phi_{b,x}^{\text{R}} \phi_{b,x}^{\text{I}} \phi_{a,x}^{\text{I}} + \sum_i (\phi_{a,x+i}^{\text{R}} + \phi_{a,x-i}^{\text{R}}) \\ & + \cosh \mu (\phi_{a,x+\hat{4}}^{\text{R}} + \phi_{a,x-\hat{4}}^{\text{R}}) + \sinh \mu \varepsilon_{ab} (\phi_{b,x+\hat{4}}^{\text{I}} - \phi_{b,x-\hat{4}}^{\text{I}}), \end{aligned} \quad (2.13)$$

$$\begin{aligned} K_{a,x}^{\text{I}} = & - [2d + m^2 + \lambda (\phi_{b,x}^{\text{R}2} - \phi_{b,x}^{\text{I}2})] \phi_{a,x}^{\text{I}} - 2\lambda \phi_{b,x}^{\text{R}} \phi_{b,x}^{\text{I}} \phi_{a,x}^{\text{R}} + \sum_i (\phi_{a,x+i}^{\text{I}} + \phi_{a,x-i}^{\text{I}}) \\ & + \cosh \mu (\phi_{a,x+\hat{4}}^{\text{I}} + \phi_{a,x-\hat{4}}^{\text{I}}) - \sinh \mu \varepsilon_{ab} (\phi_{b,x+\hat{4}}^{\text{R}} - \phi_{b,x-\hat{4}}^{\text{R}}). \end{aligned} \quad (2.14)$$

Observables are written in terms of the complexified fields (2.8) as well. We consider the square of the field modulus,

$$|\phi|^2 = \frac{1}{2} \phi_a^2 \rightarrow \frac{1}{2} (\phi_a^{\text{R}2} - \phi_a^{\text{I}2}) + i\phi_a^{\text{R}} \phi_a^{\text{I}}, \quad (2.15)$$

and the density $\langle n \rangle = (1/\Omega) \partial \ln Z / \partial \mu$, given by $n = (1/\Omega) \sum_x n_x$, with

$$\begin{aligned} n_x = & (\delta_{ab} \sinh \mu - i\varepsilon_{ab} \cosh \mu) \phi_{a,x} \phi_{b,x+\hat{4}} \\ & \rightarrow (\delta_{ab} \sinh \mu - i\varepsilon_{ab} \cosh \mu) \left(\phi_{a,x}^{\text{R}} \phi_{b,x+\hat{4}}^{\text{R}} - \phi_{a,x}^{\text{I}} \phi_{b,x+\hat{4}}^{\text{I}} + i \left[\phi_{a,x}^{\text{R}} \phi_{b,x+\hat{4}}^{\text{I}} + \phi_{a,x}^{\text{I}} \phi_{b,x+\hat{4}}^{\text{R}} \right] \right). \end{aligned} \quad (2.16)$$

After complexification all observables have a real and imaginary part.

Employing that the noise is random, we observe that the Langevin equations have the following symmetry,

$$\phi_{1,x}^{\text{R}} \rightarrow -\phi_{1,x}^{\text{R}}, \quad \phi_{2,x}^{\text{R}} \rightarrow \phi_{2,x}^{\text{R}}, \quad \phi_{1,x}^{\text{I}} \rightarrow \phi_{1,x}^{\text{I}}, \quad \phi_{2,x}^{\text{I}} \rightarrow -\phi_{2,x}^{\text{I}}, \quad (2.17)$$

for all x and similar with 1 and 2 interchanged. Under this transformation, the drift terms change as

$$K_{1,x}^{\text{R}} \rightarrow -K_{1,x}^{\text{R}}, \quad K_{2,x}^{\text{R}} \rightarrow K_{2,x}^{\text{R}}, \quad K_{1,x}^{\text{I}} \rightarrow K_{1,x}^{\text{I}}, \quad K_{2,x}^{\text{I}} \rightarrow -K_{2,x}^{\text{I}}. \quad (2.18)$$

Correlation functions odd under this transformation should vanish after noise averaging, which implies that

$$\langle \phi_{1,x}^{\text{R}} \phi_{2,y}^{\text{R}} \rangle = \langle \phi_{1,x}^{\text{I}} \phi_{2,y}^{\text{I}} \rangle = \langle \phi_{1,x}^{\text{R}} \phi_{1,y}^{\text{I}} \rangle = \langle \phi_{2,x}^{\text{R}} \phi_{2,y}^{\text{I}} \rangle = 0. \quad (2.19)$$

The nonzero combinations are

$$\langle \phi_{a,x}^R \phi_{b,y}^R \rangle \sim \delta_{ab}, \quad \langle \phi_{a,x}^I \phi_{b,y}^I \rangle \sim \delta_{ab}, \quad \langle \phi_{a,x}^R \phi_{b,y}^I \rangle \sim \varepsilon_{ab}. \quad (2.20)$$

Applying this to the expectation values of the observables in eqs. (2.15), (2.16), we find that they are purely real. This is indeed what was observed numerically in ref. [1].

3 Ignoring interactions

In order to set the stage for the mean field analysis, we first solve the Langevin dynamics without interactions ($\lambda = 0$), allowing for a detailed understanding of convergence and stability properties in the Silver Blaze regime.

3.1 Standard results

We start by summarizing the results obtained in the standard field theory approach (see e.g. ref. [40]). After going to momentum space, according to

$$\phi_{a,x} = \sum_p e^{ipx} \phi_{a,p}, \quad (3.1)$$

where $p_i = 2\pi n_i/N_s$, with $-N_s/2 < n_i \leq N_s/2$, and $p_4 = 2\pi n_4/N_\tau$, with $-N_\tau/2 < n_4 \leq N_\tau/2$, the action (2.3) reads

$$S = \sum_p \frac{1}{2} \phi_{a,-p} (\delta_{ab} A_p - \varepsilon_{ab} B_p) \phi_{b,p} = \sum_p \frac{1}{2} \phi_{a,-p} M_{ab,p} \phi_{b,p}, \quad (3.2)$$

where

$$M_p = \begin{pmatrix} A_p & -B_p \\ B_p & A_p \end{pmatrix}, \quad (3.3)$$

and³

$$A_p = m^2 + 4 \sum_{i=1}^3 \sin^2 \frac{p_i}{2} + 2(1 - \cosh \mu \cos p_4), \quad B_p = 2 \sinh \mu \sin p_4. \quad (3.4)$$

Note that $A_{-p} = A_p$, $B_{-p} = -B_p$, and that M_p is nonhermitian. The phase quenched theory is obtained by taking $B_p = 0$. Up to an irrelevant constant, the logarithm of the partition function is

$$\ln Z = -\frac{1}{2} \sum_p \ln \det M_p = -\frac{1}{2} \sum_p \ln(A_p^2 + B_p^2). \quad (3.5)$$

The observables we are interested in are given by

$$\langle |\phi|^2 \rangle = -\frac{1}{\Omega} \frac{\partial \ln Z}{\partial m^2} = \frac{1}{\Omega} \sum_p \frac{A_p}{A_p^2 + B_p^2}, \quad (3.6)$$

³In the formal continuum limit $A_p \rightarrow m^2 - \mu^2 + p_4^2 + \mathbf{p}^2$, $B_p \rightarrow 2\mu p_4$.

and

$$\langle n \rangle = \frac{1}{\Omega} \frac{\partial \ln Z}{\partial \mu} = -\frac{1}{\Omega} \sum_p \frac{A_p A'_p + B_p B'_p}{A_p^2 + B_p^2}, \quad (3.7)$$

where $A' = \partial A / \partial \mu = -2 \sinh \mu \cos p_4$, $B' = \partial B / \partial \mu = 2 \cosh \mu \sin p_4$.

As always, the severeness of the sign problem is estimated by the average phase factor in the phase quenched theory, given by the ratio of the partition functions of the full and phase quenched theories (2.4), (2.5),

$$\langle e^{i\varphi} \rangle_{\text{pq}} = \frac{Z}{Z_{\text{pq}}} = e^{-\Omega \Delta f}, \quad (3.8)$$

where Δf , the difference between the corresponding free energy densities, is given by

$$\Delta f = -\frac{1}{\Omega} \ln \frac{Z}{Z_{\text{pq}}} = \frac{1}{2\Omega} \sum_p \ln \frac{A_p^2 + B_p^2}{A_p^2}. \quad (3.9)$$

Note that this can be easily generalized to arbitrary powers of the phase factor in theories with nonvanishing phase factors.⁴

Finally, since the eigenvalues of M_p in the action (3.2) are $A_p \pm iB_p$, the theory without interactions exists provided that $A_p > 0$. This yields the standard stability criterium for a free Bose gas at finite chemical potential,

$$A_p > 0 \quad \Leftrightarrow \quad 4 \sinh^2 \frac{\mu}{2} < m^2, \quad (3.10)$$

corresponding to $\mu^2 < m^2$ in the formal continuum limit. We restrict the analysis below therefore to the case that $A_p > 0$; this is the Silver Blaze region.

3.2 Continuous Langevin dynamics

We now solve the complex Langevin equations to compare the outcome with the results given above. The Langevin equations (2.9), (2.10) read in momentum space

$$\frac{\partial}{\partial \theta} \phi_{a,p}^{\text{R}}(\theta) = K_{a,p}^{\text{R}}(\theta) + \eta_{a,p}(\theta), \quad (3.11)$$

$$\frac{\partial}{\partial \theta} \phi_{a,p}^{\text{I}}(\theta) = K_{a,p}^{\text{I}}(\theta), \quad (3.12)$$

where the noise is normalized as

$$\langle \eta_{a,-p}(\theta) \eta_{b,p'}(\theta') \rangle = 2\delta_{ab} \delta_{pp'} \delta(\theta - \theta'). \quad (3.13)$$

Ignoring interactions, we find for the drift terms

$$K_{a,p}^{\text{R}} = -A_p \phi_{a,p}^{\text{R}} + iB_p \varepsilon_{ab} \phi_{b,p}^{\text{I}}, \quad (3.14)$$

$$K_{a,p}^{\text{I}} = -A_p \phi_{a,p}^{\text{I}} - iB_p \varepsilon_{ab} \phi_{b,p}^{\text{R}}, \quad (3.15)$$

⁴Define the partition function $Z_\ell = \int D\phi_1 D\phi_2 |e^{-S}| e^{i\ell\varphi}$. Then $\langle e^{in\varphi} \rangle_\ell = Z_{n+\ell} / Z_\ell = \exp(-\Omega \Delta f_{n+\ell,\ell})$, with

$$\Delta f_{n+\ell,\ell} = -\frac{1}{\Omega} \ln \frac{Z_{n+\ell}}{Z_\ell} = \frac{1}{2\Omega} \sum_p \ln \frac{A_p^2 + (n+\ell)^2 B_p^2}{A_p^2 + \ell^2 B_p^2}.$$

where A_p and B_p are defined in eq. (3.4) above. In terms of

$$\mathcal{M}_p = \begin{pmatrix} A_p & 0 & 0 & -iB_p \\ 0 & A_p & iB_p & 0 \\ 0 & iB_p & A_p & 0 \\ -iB_p & 0 & 0 & A_p \end{pmatrix}, \quad \Phi_p = \begin{pmatrix} \phi_{1,p}^R \\ \phi_{1,p}^I \\ \phi_{2,p}^R \\ \phi_{2,p}^I \end{pmatrix}, \quad \Xi_p = \begin{pmatrix} \eta_{1,p} \\ 0 \\ \eta_{2,p} \\ 0 \end{pmatrix}, \quad (3.16)$$

the Langevin dynamics is written as

$$\frac{\partial}{\partial \theta} \Phi_p = -\mathcal{M}_p \Phi_p + \Xi_p. \quad (3.17)$$

The matrix \mathcal{M} can be diagonalized by an orthogonal transformation and has doubly degenerate eigenvalues $\lambda_p = A_p \pm iB_p$. The solution of the Langevin equations is

$$\begin{aligned} \phi_{a,p}^R(\theta) &= e^{-A_p \theta} \left[\cos(B_p \theta) \phi_{a,p}^R(0) + i \sin(B_p \theta) \varepsilon_{ab} \phi_{b,p}^I(0) \right] \\ &\quad + \int_0^\theta ds e^{-A_p(\theta-s)} \cos[B_p(\theta-s)] \eta_{a,p}(s), \end{aligned} \quad (3.18)$$

$$\begin{aligned} \phi_{a,p}^I(\theta) &= e^{-A_p \theta} \left[\cos(B_p \theta) \phi_{a,p}^I(0) - i \sin(B_p \theta) \varepsilon_{ab} \phi_{b,p}^R(0) \right] \\ &\quad - i \int_0^\theta ds e^{-A_p(\theta-s)} \sin[B_p(\theta-s)] \varepsilon_{ab} \eta_{b,p}(s), \end{aligned} \quad (3.19)$$

where $\phi_{a,p}^{R,I}(0)$ denote the initial conditions.

We are now in a position to discuss the convergence properties of the Langevin process in the limit of large Langevin time. First we note that there is independence of initial conditions provided that $A_p > 0$, i.e. in the region of interest here. Taking $\phi_{a,p}^{R,I}(0) = 0$, we find for the two-point functions, after using eq. (3.13) and performing the Langevin time integrals,

$$\begin{aligned} \langle \phi_{a,-p}^R(\theta) \phi_{b,p'}^R(\theta) \rangle &= \frac{1}{2A_p} \frac{\delta_{ab} \delta_{pp'}}{A_p^2 + B_p^2} \left(2A_p^2 + B_p^2 \right. \\ &\quad \left. - e^{-2A_p \theta} [A_p^2 + B_p^2 + A_p^2 \cos(2B_p \theta) - A_p B_p \sin(2B_p \theta)] \right), \\ \langle \phi_{a,-p}^I(\theta) \phi_{b,p'}^I(\theta) \rangle &= \frac{1}{2A_p} \frac{\delta_{ab} \delta_{pp'}}{A_p^2 + B_p^2} \left(B^2 \right. \\ &\quad \left. - e^{-2A_p \theta} [A_p^2 + B_p^2 - A_p^2 \cos(2B_p \theta) + A_p B_p \sin(2B_p \theta)] \right), \\ \langle \phi_{a,-p}^R(\theta) \phi_{b,p'}^I(\theta) \rangle &= \frac{i}{2} \frac{\varepsilon_{ab} \delta_{pp'}}{A_p^2 + B_p^2} \left(B_p - e^{-2A_p \theta} [B_p \cos(2B_p \theta) + A_p \sin(2B_p \theta)] \right). \end{aligned} \quad (3.20)$$

Most of the terms vanish in the limit that $\theta \rightarrow \infty$, again provided that $A_p > 0$. The surviving terms are

$$\begin{aligned} \lim_{\theta \rightarrow \infty} \langle \phi_{a,-p}^R(\theta) \phi_{b,p'}^R(\theta) \rangle &\equiv \langle \phi_{a,-p}^R \phi_{b,p'}^R \rangle = \delta_{ab} \delta_{pp'} \frac{1}{2A_p} \frac{2A_p^2 + B_p^2}{A_p^2 + B_p^2}, \\ \lim_{\theta \rightarrow \infty} \langle \phi_{a,-p}^I(\theta) \phi_{b,p'}^I(\theta) \rangle &\equiv \langle \phi_{a,-p}^I \phi_{b,p'}^I \rangle = \delta_{ab} \delta_{pp'} \frac{1}{2A_p} \frac{B_p^2}{A_p^2 + B_p^2}, \\ \lim_{\theta \rightarrow \infty} \langle \phi_{a,-p}^R(\theta) \phi_{b,p'}^I(\theta) \rangle &\equiv \langle \phi_{a,-p}^R \phi_{b,p'}^I \rangle = \varepsilon_{ab} \delta_{pp'} \frac{i}{2} \frac{B_p}{A_p^2 + B_p^2}. \end{aligned} \quad (3.21)$$

The structure of these two-point functions is in agreement with the symmetry (2.17), (2.20) discussed above.

For the observables we find the following. The square of the field modulus (2.15) is given by

$$\begin{aligned} \langle |\phi|^2 \rangle &= \frac{1}{2\Omega} \sum_p \langle \phi_{a,-p}^R \phi_{a,p}^R - \phi_{a,-p}^I \phi_{a,p}^I + 2i \phi_{a,-p}^R \phi_{a,p}^I \rangle \\ &= \frac{1}{\Omega} \sum_p \frac{A_p}{A_p^2 + B_p^2}, \end{aligned} \quad (3.22)$$

which agrees with eq. (3.6). After going to momentum space, the density (2.16) reads

$$\begin{aligned} \langle n \rangle &= \frac{1}{\Omega} \sum_p (\delta_{ab} \sinh \mu \cos p_4 + \varepsilon_{ab} \cosh \mu \sin p_4) \langle \phi_{a,-p}^R \phi_{b,p}^R - \phi_{a,-p}^I \phi_{b,p}^I + 2i \phi_{a,-p}^R \phi_{b,p}^I \rangle \\ &= \frac{2}{\Omega} \sum_p \left[\sinh \mu \cos p_4 \frac{A_p}{A_p^2 + B_p^2} - \cosh \mu \sin p_4 \frac{B_p}{A_p^2 + B_p^2} \right], \end{aligned} \quad (3.23)$$

which agrees with eq. (3.7). Note that all two-point functions in eq. (3.21) contribute to this answer.

We conclude therefore that the Langevin process is independent of initial conditions and converges to the correct result in the limit of infinite Langevin time, provided that $A_p > 0$, as required in the Silver Blaze region. Moreover, the complexification is essential, as exemplified by the observables above.

3.3 Discretized Langevin dynamics

We proceed by briefly considering the Langevin process after discretizing Langevin time as $\theta = n\epsilon$, where ϵ is the Langevin time step. The discretized Langevin equations are

$$\phi_{a,p}^R(n+1) = \phi_{a,p}^R(n) + \epsilon K_{a,p}^R(n) + \sqrt{\epsilon} \eta_{a,p}(n), \quad (3.24)$$

$$\phi_{a,p}^I(n+1) = \phi_{a,p}^I(n) + \epsilon K_{a,p}^I(n), \quad (3.25)$$

and the noise obeys $\langle \eta_{a,-p}(n) \eta_{b,p'}(n') \rangle = 2\delta_{nn'} \delta_{ab} \overline{\delta_{pp'}}$. In the notation of eq. (3.16), these equations are summarized as

$$\Phi_p(n+1) = (1 - \epsilon \mathcal{M}_p) \Phi_p(n) + \sqrt{\epsilon} \Xi_p(n), \quad (3.26)$$

and solved by

$$\Phi_p(n) = (1 - \epsilon \mathcal{M}_p)^n \Phi_p(0) + \sqrt{\epsilon} \sum_{i=0}^{n-1} (1 - \epsilon \mathcal{M}_p)^{n-1-i} \Xi_p(i), \quad (3.27)$$

where $\Phi_p(0)$ is the initial condition. Convergence is determined by the eigenvalues of \mathcal{M}_p . This yields the condition

$$|1 - \epsilon \lambda_p| < 1, \quad \lambda_p = A_p \pm i B_p, \quad (3.28)$$

resulting in the constraint

$$A_p - \frac{\epsilon}{2} (A_p^2 + B_p^2) > 0. \quad (3.29)$$

We find that the convergence criterium is modified by an explicit stepsize dependence. However, this restriction is not special for the complex Langevin process [30, 41]. Consider real Langevin dynamics at zero μ or in the phase quenched theory. In both cases $B_p = 0$ and the criterium reads

$$0 < \frac{\epsilon}{2} A_p < 1. \quad (3.30)$$

Since A_p is maximal at the edge of the Brillouin zone ($p = \pi$), this yields the modest constraint (for $\mu = 0$)

$$\epsilon < \frac{2}{4d + m^2}. \quad (3.31)$$

At nonzero chemical potential, this constraint is modified to

$$\epsilon < \frac{2}{4d + m^2 + 2(\cosh \mu - 1)}, \quad (3.32)$$

both in the full and the phase quenched theory. In the Silver Blaze region, where μ is bounded, this leads to only a slightly stronger bound on ϵ . However, since this bound is determined by ultraviolet modes at the scale of the lattice cutoff, it is likely that a similar constraint holds in the high-density phase as well. In the limit that $\mu \gg 1$, exponentially small stepsizes would eventually be required. It should be noted, however, that for such large chemical potentials lattice artefacts are severe.

The solution (3.27) can be used to study finite stepsize effects in two-point functions at infinite Langevin time. We come back to this below using a more elegant approach based on fixed points of the Langevin equations.

4 Fokker-Planck equation

In order to better understand the Langevin process, we now study properties of the associated distributions.

Consider first the Langevin process (2.6) and the distribution $P[\phi, \theta]$, defined via

$$\langle O[\phi, \theta] \rangle_\eta = \int D\phi P[\phi, \theta] O[\phi], \quad (4.1)$$

where the brackets on the l.h.s. denote noise averaging. The distribution satisfies the Fokker-Planck equation (in continuous Langevin time)

$$\frac{\partial P[\phi, \theta]}{\partial \theta} = \sum_x \frac{\delta}{\delta \phi_{a,x}(\theta)} \left(\frac{\delta}{\delta \phi_{a,x}(\theta)} + \frac{\delta S[\phi]}{\delta \phi_{a,x}(\theta)} \right) P[\phi, \theta]. \quad (4.2)$$

As always, the index $a = 1, 2$ is summed over. The stationary solution,

$$P[\phi] \sim e^{-S[\phi]}, \quad (4.3)$$

always exists. However, since the action is complex, this is not the probability distribution for the complex Langevin process.

More relevant for the complexified process (2.9), (2.10) we consider here, is the real distribution $\rho[\phi^{\text{R}}, \phi^{\text{I}}, \theta]$, defined via [26]

$$\langle O[\phi^{\text{R}} + i\phi^{\text{I}}, \theta] \rangle_{\eta} = \int D\phi^{\text{R}} D\phi^{\text{I}} \rho[\phi^{\text{R}}, \phi^{\text{I}}, \theta] O[\phi^{\text{R}} + i\phi^{\text{I}}]. \quad (4.4)$$

This distribution satisfies the extended Fokker-Planck equation

$$\frac{\partial \rho[\phi^{\text{R}}, \phi^{\text{I}}, \theta]}{\partial \theta} = \sum_x \left[\frac{\delta}{\delta \phi_{a,x}^{\text{R}}(\theta)} \left(\frac{\delta}{\delta \phi_{a,x}^{\text{R}}(\theta)} - K_{a,x}^{\text{R}}(\theta) \right) - \frac{\delta}{\delta \phi_{a,x}^{\text{I}}(\theta)} K_{a,x}^{\text{I}}(\theta) \right] \rho[\phi^{\text{R}}, \phi^{\text{I}}, \theta]. \quad (4.5)$$

If stochastic quantization is applicable for complex actions, the two expectation values (4.1) and (4.4) should be equal [26].

We focus on the stationary solution of eq. (4.5) and henceforth drop the θ dependence. Ignoring again interactions, the stationary solution should satisfy

$$\sum_p \left[\frac{\delta}{\delta \phi_{a,p}^{\text{R}}} \left(\frac{\delta}{\delta \phi_{a,-p}^{\text{R}}} - K_{a,p}^{\text{R}} \right) - \frac{\delta}{\delta \phi_{a,p}^{\text{I}}} K_{a,p}^{\text{I}} \right] \rho[\phi^{\text{R}}, \phi^{\text{I}}] = 0, \quad (4.6)$$

where the drift terms $K_{a,p}^{\text{R,I}}$ were given in eqs. (3.14), (3.15). Explicitly, this reads

$$\begin{aligned} \sum_p \left[\frac{\delta}{\delta \phi_{a,p}^{\text{R}}} \frac{\delta}{\delta \phi_{a,-p}^{\text{R}}} + (A_p \phi_{a,p}^{\text{R}} - iB_p \varepsilon_{ab} \phi_{b,p}^{\text{I}}) \frac{\delta}{\delta \phi_{a,p}^{\text{R}}} \right. \\ \left. + (A_p \phi_{a,p}^{\text{I}} + iB_p \varepsilon_{ab} \phi_{b,p}^{\text{R}}) \frac{\delta}{\delta \phi_{a,p}^{\text{I}}} + 2A_p \right] \rho[\phi^{\text{R}}, \phi^{\text{I}}] = 0. \end{aligned} \quad (4.7)$$

Based on the structure of the equation, the solution can be written as

$$\rho[\phi^{\text{R}}, \phi^{\text{I}}] = N \exp \left[- \sum_p (\alpha_p \phi_{a,-p}^{\text{R}} \phi_{a,p}^{\text{R}} + \beta_p \phi_{a,-p}^{\text{I}} \phi_{a,p}^{\text{I}} + 2i\varepsilon_{ab} \gamma_p \phi_{a,-p}^{\text{R}} \phi_{b,p}^{\text{I}}) \right], \quad (4.8)$$

where N is a normalization constant. Inserting this expression in eq. (4.7) yields the coefficients

$$\alpha_p = A_p, \quad \beta_p = \frac{A_p}{B_p^2} (2A_p^2 + B_p^2), \quad \gamma_p = \frac{A_p^2}{B_p}. \quad (4.9)$$

We have therefore found the stationary distribution corresponding to the complex Langevin process in the noninteracting case.⁵ Note that since $\gamma_{-p} = -\gamma_p$, the distribution is real in real space, as it should be.

We now verify that this stationary solution is indeed the distribution corresponding to the Langevin process in the limit of infinite Langevin time. Performing the Gaussian integrals, we find the partition function

$$Z = \prod_p \int d\phi_p^{\text{R}} d\phi_p^{\text{I}} \rho[\phi^{\text{R}}, \phi^{\text{I}}] = \mathcal{N} \prod_p \frac{1}{\alpha_p \beta_p - \gamma_p^2}, \quad (4.10)$$

⁵See refs. [42, 43] for other examples.

where \mathcal{N} is an irrelevant constant and

$$\alpha_p \beta_p - \gamma_p^2 = \frac{A_p^2}{B_p^2} (A_p^2 + B_p^2) > 0. \quad (4.11)$$

The two-point functions that follow from this distribution are

$$\begin{aligned} \langle \phi_{a,-p}^R \phi_{a,p}^R \rangle &= -\frac{\partial \ln Z}{\partial \alpha_p} = \frac{\beta_p}{\alpha_p \beta_p - \gamma_p^2} = \frac{1}{A_p} \frac{2A_p^2 + B_p^2}{A_p^2 + B_p^2}, \\ \langle \phi_{a,-p}^I \phi_{a,p}^I \rangle &= -\frac{\partial \ln Z}{\partial \beta_p} = \frac{\alpha_p}{\alpha_p \beta_p - \gamma_p^2} = \frac{1}{A_p} \frac{B_p^2}{A_p^2 + B_p^2}, \\ 2i\varepsilon_{ab} \langle \phi_{a,-p}^R \phi_{b,p}^I \rangle &= -\frac{\partial \ln Z}{\partial \gamma_p} = \frac{-2\gamma_p}{\alpha_p \beta_p - \gamma_p^2} = \frac{-2B_p}{A_p^2 + B_p^2}. \end{aligned} \quad (4.12)$$

These agree exactly with the results obtained by solving the Langevin equation, cf. eq. (3.21).

The theory with the probability distribution $\rho[\phi^R, \phi^I]$ exists provided that the eigenvalues of the quadratic form in eq. (4.8) are positive. We find the eigenvalues to be

$$\begin{aligned} \lambda_p &= \frac{1}{2} \left(\alpha_p + \beta_p \pm \sqrt{(\alpha_p - \beta_p)^2 + 4\gamma_p^2} \right) \\ &= \frac{A_p}{B_p^2} \sqrt{A_p^2 + B_p^2} \left(\sqrt{A_p^2 + B_p^2} \pm A_p \right). \end{aligned} \quad (4.13)$$

These are positive provided that $A_p > 0$. The criterium that determines the convergence of the Langevin dynamics also emerges in the stationary solution of the extended Fokker-Planck equation, as expected.

Let us discuss some more properties of the distribution (4.8). First we note that the distribution is highly nonlocal in real space and does not allow for e.g. a derivative expansion, due to the division by $B_p = 2 \sinh \mu \sin p_4$ in the coefficients (4.9). We find therefore that the complexity of the original local weight e^{-S} has been traded for the nonlocality of the real probability distribution. However, this nonlocal behaviour is expected: it follows from the Langevin equations that the modes with $p_4 = 0$ are purely real, i.e. $\phi_{a,(p_4=0,\mathbf{p})}^I = 0$. This is enforced in the probability distribution by the singular behaviour as $p_4 \rightarrow 0$. For the same reason the limit $\mu \rightarrow 0$ is singular, since there is no need to complexify the dynamics in this case and the distribution for the ϕ^I modes should reduce to a delta function, $\delta(\phi^I)$. These considerations fix the dependence on B_p .

In conclusion, we have found the stationary solution of the extended Fokker-Planck distribution. The real distribution is nonlocal and singular in the limit that $\mu, p_4 \rightarrow 0$.

5 Mean field approximation

We now return to the interacting theory, with discretized Langevin time $\theta = n\epsilon$, and consider the two-point functions

$$\begin{aligned} G_{ab,p}^{RR}(n) &= \langle \phi_{a,-p}^R(n) \phi_{b,p}^R(n) \rangle, \\ G_{ab,p}^{II}(n) &= \langle \phi_{a,-p}^I(n) \phi_{b,p}^I(n) \rangle, \\ G_{ab,p}^{RI}(n) &= \langle \phi_{a,-p}^R(n) \phi_{b,p}^I(n) \rangle. \end{aligned} \quad (5.1)$$

Using the Langevin equations (3.24), (3.25), we find that these correlation functions evolve according to

$$\begin{aligned}
G_{ab,p}^{\text{RR}}(n+1) &= G_{ab,p}^{\text{RR}}(n) + \epsilon \langle \phi_{a,-p}^{\text{R}}(n) K_{b,p}^{\text{R}}(n) + K_{a,-p}^{\text{R}}(n) \phi_{b,p}^{\text{R}}(n) \rangle \\
&\quad + \epsilon^2 \langle K_{a,-p}^{\text{R}}(n) K_{b,p}^{\text{R}}(n) \rangle + \epsilon \langle \eta_{a,-p} \eta_{b,p} \rangle, \\
G_{ab,p}^{\text{II}}(n+1) &= G_{ab,p}^{\text{II}}(n) + \epsilon \langle \phi_{a,-p}^{\text{I}}(n) K_{b,p}^{\text{I}}(n) + K_{a,-p}^{\text{I}}(n) \phi_{b,p}^{\text{I}}(n) \rangle \\
&\quad + \epsilon^2 \langle K_{a,-p}^{\text{I}}(n) K_{b,p}^{\text{I}}(n) \rangle, \\
G_{ab,p}^{\text{RI}}(n+1) &= G_{ab,p}^{\text{RI}}(n) + \epsilon \langle \phi_{a,-p}^{\text{R}}(n) K_{b,p}^{\text{I}}(n) + K_{a,-p}^{\text{R}}(n) \phi_{b,p}^{\text{I}}(n) \rangle \\
&\quad + \epsilon^2 \langle K_{a,-p}^{\text{R}}(n) K_{b,p}^{\text{I}}(n) \rangle.
\end{aligned} \tag{5.2}$$

Here we used that $\langle \eta_{a,-p}(n) \phi_{b,p}^{\text{R,I}}(n) \rangle = 0$, since the fields at time n do not depend on the noise at time n . The terms proportional to ϵ^2 are finite stepsize corrections. We then look for fixed points of the Langevin equations, and put

$$G_{ab,p}^{\text{RR}}(n+1) = G_{ab,p}^{\text{RR}}(n), \tag{5.3}$$

etc. This yields the fixed point equations

$$\begin{aligned}
\langle \phi_{a,-p}^{\text{R}} K_{b,p}^{\text{R}} + K_{a,-p}^{\text{R}} \phi_{b,p}^{\text{R}} \rangle + \epsilon \langle K_{a,-p}^{\text{R}} K_{b,p}^{\text{R}} \rangle &= -2\delta_{ab}, \\
\langle \phi_{a,-p}^{\text{I}} K_{b,p}^{\text{I}} + K_{a,-p}^{\text{I}} \phi_{b,p}^{\text{I}} \rangle + \epsilon \langle K_{a,-p}^{\text{I}} K_{b,p}^{\text{I}} \rangle &= 0, \\
\langle \phi_{a,-p}^{\text{R}} K_{b,p}^{\text{I}} + K_{a,-p}^{\text{R}} \phi_{b,p}^{\text{I}} \rangle + \epsilon \langle K_{a,-p}^{\text{R}} K_{b,p}^{\text{I}} \rangle &= 0.
\end{aligned} \tag{5.4}$$

To implement a mean field approximation and find a self-consistent set for the two-point functions (5.1), we factorize the interaction terms. Consider for example the term $\phi_{b,x}^{\text{R}} \phi_{b,x}^{\text{I}} \phi_{a,x}^{\text{R}}$ appearing in the drift term (2.14). We write

$$\phi_{b,x}^{\text{R}} \phi_{b,x}^{\text{I}} \phi_{a,x}^{\text{R}} \rightarrow \langle \phi_{b,x}^{\text{R}} \phi_{b,x}^{\text{I}} \rangle \phi_{a,x}^{\text{R}} + \langle \phi_{b,x}^{\text{R}} \phi_{a,x}^{\text{R}} \rangle \phi_{b,x}^{\text{I}} + \langle \phi_{b,x}^{\text{I}} \phi_{a,x}^{\text{R}} \rangle \phi_{b,x}^{\text{R}}. \tag{5.5}$$

Using the notation

$$\begin{aligned}
G_{ab}^{\text{RR}}(n) &= \langle \phi_{a,x}^{\text{R}}(n) \phi_{b,x}^{\text{R}}(n) \rangle = \frac{1}{\Omega} \sum_p G_{ab,p}^{\text{RR}}(n), \\
G_{ab}^{\text{II}}(n) &= \langle \phi_{a,x}^{\text{I}}(n) \phi_{b,x}^{\text{I}}(n) \rangle = \frac{1}{\Omega} \sum_p G_{ab,p}^{\text{II}}(n), \\
G_{ab}^{\text{RI}}(n) &= \langle \phi_{a,x}^{\text{R}}(n) \phi_{b,x}^{\text{I}}(n) \rangle = \frac{1}{\Omega} \sum_p G_{ab,p}^{\text{RI}}(n),
\end{aligned} \tag{5.6}$$

the drift terms in the mean field approximation read

$$\begin{aligned}
K_{a,p}^{\text{R}} &= - [A_p + \lambda (G_{bb}^{\text{RR}} - G_{bb}^{\text{II}})] \phi_{a,p}^{\text{R}} - 2\lambda (G_{ab}^{\text{RR}} - G_{ab}^{\text{II}}) \phi_{b,p}^{\text{R}} \\
&\quad + [iB_p \varepsilon_{ab} + 2\lambda (G_{ab}^{\text{RI}} + G_{ba}^{\text{RI}})] \phi_{b,p}^{\text{I}} + 2\lambda G_{bb}^{\text{RI}} \phi_{a,p}^{\text{I}},
\end{aligned} \tag{5.7}$$

$$\begin{aligned}
K_{a,p}^{\text{I}} &= - [A_p + \lambda (G_{bb}^{\text{RR}} - G_{bb}^{\text{II}})] \phi_{a,p}^{\text{I}} - 2\lambda (G_{ab}^{\text{RR}} - G_{ab}^{\text{II}}) \phi_{b,p}^{\text{I}} \\
&\quad - [iB_p \varepsilon_{ab} + 2\lambda (G_{ab}^{\text{RI}} + G_{ba}^{\text{RI}})] \phi_{b,p}^{\text{R}} - 2\lambda G_{bb}^{\text{RI}} \phi_{a,p}^{\text{R}}.
\end{aligned} \tag{5.8}$$

These can be further simplified by noting that both $G_{ab}^{\text{RR}}(n)$ and $G_{ab}^{\text{II}}(n)$ are proportional to δ_{ab} , for all Langevin times. We write therefore

$$\begin{aligned} G_{ab}^{\text{RR}}(n) &= \delta_{ab} G^{\text{RR}}(n), \\ G_{ab}^{\text{II}}(n) &= \delta_{ab} G^{\text{II}}(n), \\ G_{ab}^{\text{RI}}(n) &= \delta_{ab} G^{\text{RI}}(n) + \varepsilon_{ab} \bar{G}^{\text{RI}}(n), \end{aligned} \quad (5.9)$$

such that the drift terms reduce to

$$K_{a,p}^{\text{R}} = -\mathcal{A}_p \phi_{a,p}^{\text{R}} + i B_p \varepsilon_{ab} \phi_{b,p}^{\text{I}} + \mathcal{C} \phi_{a,p}^{\text{I}}, \quad (5.10)$$

$$K_{a,p}^{\text{I}} = -\mathcal{A}_p \phi_{a,p}^{\text{I}} - i B_p \varepsilon_{ab} \phi_{b,p}^{\text{R}} - \mathcal{C} \phi_{a,p}^{\text{R}}, \quad (5.11)$$

with

$$\mathcal{A}_p = A_p + 4\lambda (G^{\text{RR}} - G^{\text{II}}), \quad \mathcal{C} = 8\lambda G^{\text{RI}}. \quad (5.12)$$

Since $\mathcal{A}_p(n)$ and $\mathcal{C}(n)$ depend explicitly on the Langevin time, the time-dependent mean field Langevin equations cannot be solved analytically.⁶

We therefore look for fixed points. After substituting eqs. (5.10), (5.11) in the fixed point equations (5.4) and performing some algebra, we find that at the fixed point $\mathcal{C} = 0$ and that the two-point functions can be decomposed as

$$G_{ab,p}^{\text{RR}} = \delta_{ab} G_p^{\text{RR}}, \quad G_{ab,p}^{\text{II}} = \delta_{ab} G_p^{\text{II}}, \quad G_{ab,p}^{\text{RI}} = i \varepsilon_{ab} G_p^{\text{RI}}. \quad (5.13)$$

This is in agreement with the symmetry (2.17), (2.20). The three fixed point equations (5.4) then become

$$\begin{aligned} \mathcal{A}_p G_p^{\text{RR}} + B_p G_p^{\text{RI}} - \frac{\epsilon}{2} (\mathcal{A}_p^2 G_p^{\text{RR}} + B_p^2 G_p^{\text{II}} + 2\mathcal{A}_p B_p G_p^{\text{RI}}) &= 1, \\ \mathcal{A}_p G_p^{\text{II}} - B_p G_p^{\text{RI}} - \frac{\epsilon}{2} (\mathcal{A}_p^2 G_p^{\text{II}} + B_p^2 G_p^{\text{RR}} - 2\mathcal{A}_p B_p G_p^{\text{RI}}) &= 0, \\ \mathcal{A}_p G_p^{\text{RI}} - \frac{1}{2} B_p [G^{\text{RR}} - G^{\text{II}}] - \frac{\epsilon}{2} ([\mathcal{A}_p^2 - B_p^2] G_p^{\text{RI}} - \mathcal{A}_p B_p [G_p^{\text{RR}} - G_p^{\text{II}}]) &= 0. \end{aligned} \quad (5.14)$$

The solution is

$$\begin{aligned} G_p^{\text{RR}} + G_p^{\text{II}} &= \frac{1}{\mathcal{A}_p - \frac{1}{2}\epsilon (\mathcal{A}_p^2 + B_p^2)}, \\ G_p^{\text{RR}} - G_p^{\text{II}} &= \frac{1}{\mathcal{A}_p^2 + B_p^2} \frac{\mathcal{A}_p (1 - \frac{1}{2}\epsilon \mathcal{A}_p) + \frac{1}{2}\epsilon B_p^2}{(1 - \frac{1}{2}\epsilon \mathcal{A}_p)^2 + \frac{1}{4}\epsilon^2 B_p^2}, \\ G_p^{\text{RI}} &= \frac{1}{2} \frac{B_p}{\mathcal{A}_p^2 + B_p^2} \frac{1 - \epsilon \mathcal{A}_p}{(1 - \frac{1}{2}\epsilon \mathcal{A}_p)^2 + \frac{1}{4}\epsilon^2 B_p^2}. \end{aligned} \quad (5.15)$$

For vanishing Langevin stepsize this simplifies to

$$G_p^{\text{RR}} + G_p^{\text{II}} = \frac{1}{\mathcal{A}_p}, \quad G_p^{\text{RR}} - G_p^{\text{II}} = \frac{\mathcal{A}_p}{\mathcal{A}_p^2 + B_p^2}, \quad G_p^{\text{RI}} = \frac{1}{2} \frac{B_p}{\mathcal{A}_p^2 + B_p^2}, \quad (5.16)$$

⁶In fact, the problem is now very similar to that of nonequilibrium field dynamics using a self-consistent mean field approximation in the equal-time formalism [44].

while in the phase quenched theory ($B_p = 0$) where real Langevin dynamics is applicable, the solution reduces to

$$G_p^{\text{RR}} = \frac{1}{\mathcal{A}_p} \frac{1}{1 - \frac{1}{2}\epsilon\mathcal{A}_p}, \quad G_p^{\text{II}} = G_p^{\text{RI}} = 0. \quad (5.17)$$

We find finite stepsize corrections linear in ϵ , as expected [41]. Furthermore, we note that for large stepsize the denominator in the first line of eq. (5.15) can go negative. However, this occurs precisely when the stability criterium (3.29) is violated (after the replacement $A_p \rightarrow \mathcal{A}_p$) and is therefore excluded.

The expressions in eq. (5.16) agree precisely with the solutions (3.21) obtained by solving the Langevin equations without interactions, after making the mean field replacement $A_p \rightarrow \mathcal{A}_p$. This replacement corresponds to the standard mean field approximation in which the mass parameter receives a tadpole correction,

$$m^2 \rightarrow M^2 = m^2 + 4\lambda \langle |\phi|^2 \rangle, \quad (5.18)$$

or, in the notation of this section,

$$\begin{aligned} m^2 \rightarrow M^2 &= m^2 + 4\lambda (G^{\text{RR}} - G^{\text{II}}) \\ &= m^2 + \frac{4\lambda}{\Omega} \sum_p \frac{\mathcal{A}_p}{\mathcal{A}_p^2 + B_p^2}, \end{aligned} \quad (5.19)$$

with

$$\mathcal{A}_p = M^2 + 4 \sum_i \sin^2 \frac{p_i}{2} + 2(1 - \cosh \mu \cos p_4). \quad (5.20)$$

These equations define a self-consistent gap equation for M^2 . Given m and λ , the gap equation can be solved numerically after specifying the lattice size. For example, taking $m = \lambda = 1$, we find $M^2 = 1.47$ and $\langle |\phi|^2 \rangle = 0.119$ at $\mu = 0$ on a lattice of size 10^4 . The critical chemical potential then follows from $\mathcal{A}_0 = 0$ (or $M^2 = 4 \sinh^2 \frac{\mu}{2}$) and is found to be $\mu_c = 1.15$.⁷

We have solved the gap equation in the Silver Blaze region and used the outcome to compute $\langle |\phi|^2 \rangle$ and $\langle n \rangle$ in the mean field approximation as a function of chemical potential for different lattice sizes. The results are shown in figures 1 and 2 respectively, for $m = \lambda = 1$. The vertical dotted lines indicate the mean field estimate of the critical chemical potential. In the full theory (figures on the left) the expected μ independence emerges in the thermodynamic limit. In the phase quenched theory (figures on the right), observables depend on μ , since there is no Silver Blaze feature, see appendix A. Also shown in these plots are data points obtained from the numerical solution of the Langevin process, with stepsize $\epsilon = 5 \times 10^{-5}$ [1]. We observe surprisingly good agreement between the mean field and the nonperturbative results for all values of the chemical potential and all lattice sizes considered, indicating that the mean field approximation captures the most relevant interactions.

In figure 3 we show the average phase factor in the phase quenched theory $\langle e^{i\varphi} \rangle_{\text{pq}}$ (left) and the difference Δf between the free energy densities, given in eq. (3.9), again

⁷When $\lambda = 0$, μ_c follows from $\mathcal{A}_0 = 0$ (or $m^2 = 4 \sinh^2 \frac{\mu}{2}$), yielding $\mu_c^0 = 0.962$ for $m = 1$.




Figure 1. The lines represent the mean field results for $\langle|\phi|^2\rangle$ in the full (left) and phase quenched (right) theories for various lattice sizes, taking $m = \lambda = 1$. The vertical dotted line indicates the mean field estimate for the critical chemical potential. The data points are obtained with Langevin simulations [1].




Figure 2. As in figure 1 for the density $\langle n \rangle$.

after the replacement $A_p \rightarrow \mathcal{A}_p$. As already mentioned, the sign problem is severe in the thermodynamic limit: taking e.g. $\Delta f \sim 0.02$ and a lattice volume $\Omega = 10^4$, we find that the average phase factor $\langle e^{i\varphi} \rangle_{\text{pq}} = e^{-\Omega \Delta f}$ is indeed exponentially small.

To conclude this section, we note that it is straightforward to adapt the stationary solution of the extended Fokker-Planck equation, constructed in section 4, to the mean field approximation discussed here. Since in the mean field approximation only two-point functions appear, the mean field probability distribution remains of the form (4.8) with the simple replacement $A_p \rightarrow \mathcal{A}_p$ (or $m^2 \rightarrow M^2$). Existence of the Fokker-Planck distribution in the Silver Blaze region now requires $\mathcal{A}_p > 0$.

6 Conclusion

In order to further understand the applicability of complex Langevin dynamics for theories with a complex action due to finite chemical potential, we have studied the relativistic Bose gas in the Silver Blaze region analytically. Ignoring interactions, we have investigated




Figure 3. Left: as in figure 1 for the average phase factor in the phase quenched theory $\langle e^{i\varphi} \rangle_{\text{pq}}$. Right: difference Δf between the free energy densities of the full and the phase quenched theories in the mean field approximation.

convergence and stability, and constructed the stationary solution of the extended Fokker-Planck equation. We explained why this real probability distribution is nonlocal in real space. Subsequently, interactions were included on the mean field level and the fixed point of the mean field Langevin equations with finite stepsize was derived. We gave a comparison between the mean field predictions and the nonperturbative numerical data from ref. [1] in the Silver Blaze region. Surprisingly good agreement was found for all values of the chemical potential considered, including finite size effects, indicating that the mean field approximation captures the most important effects of the interactions. We have demonstrated analytically that the sign problem is severe for lattice volumes used in this study. From the combination of results obtained here and in ref. [1], it can be argued that complex Langevin dynamics in the Silver Blaze region is well understood in this theory.

One obvious next step is to extend the analysis to the high-density phase, which requires the introduction of the mean field $\langle \phi_{a,x}^R \rangle$ (note that $\langle \phi_{a,x}^I \rangle = 0$). Finally, it would be interesting to apply mean field approximations to other theories as well, especially in combination with numerical studies. In particular, this would be useful for QCD with static quarks [36].

Acknowledgments

I thank Kim Splittorff, Ion-Olimpiu Stamatescu and Simon Hands for their interest and discussion. This work is supported by an STFC Advanced Fellowship.

A Dispersion relation

The propagator corresponding to the action (3.2) is

$$G_{ab,p} = \frac{\delta_{ab}A_p + \varepsilon_{ab}B_p}{A_p^2 + B_p^2}. \tag{A.1}$$

Dispersion relations follow from the poles of the propagator, taking $p_4 = iE_{\mathbf{p}}$. We find

$$\cosh E_{\mathbf{p}}(\mu) = \left(1 + \frac{1}{2}\hat{\omega}_{\mathbf{p}}^2\right) \cosh \mu \pm \sqrt{1 + \frac{1}{4}\hat{\omega}_{\mathbf{p}}^2} \sinh \mu, \quad (\text{A.2})$$

where

$$\hat{\omega}_{\mathbf{p}}^2 = m^2 + 4 \sum_i \sin^2 \frac{p_i}{2}. \quad (\text{A.3})$$

This can be written as

$$\cosh E_{\mathbf{p}}(\mu) = \cosh [E_{\mathbf{p}}(0) \pm \mu], \quad (\text{A.4})$$

such that the (positive energy) solutions are

$$E_{\mathbf{p}}(\mu) = E_{\mathbf{p}}(0) \pm \mu, \quad (\text{A.5})$$

just as in the continuum theory. Lattice discretization effects only appear in the dispersion relation at zero chemical potential, $E_{\mathbf{p}}(0)$. The critical μ value is $\mu_c = E_{\mathbf{0}}(0) = 2\text{asinh}(m/2)$, so that one mode becomes exactly massless at the transition.

The phase quenched theory corresponds to $B_p = 0$. In the formal continuum limit, the phase quenched theory is a theory with a real action and mass parameter $m^2 - \mu^2$. The dispersion relation is

$$\cosh E_{\mathbf{p}}(\mu) = \frac{1}{\cosh \mu} \left(1 + \frac{1}{2}\hat{\omega}_{\mathbf{p}}^2\right), \quad (\text{A.6})$$

corresponding to $E_{\mathbf{p}}^2(\mu) = m^2 - \mu^2 + \mathbf{p}^2$ in the continuum limit, as anticipated.

These results are easily extended to the self-consistent mean field approximation, where the mass parameter receives a tadpole correction and is replaced by $M^2 = m^2 + 4\lambda \langle |\phi|^2 \rangle$.

References

- [1] G. Aarts, *Can stochastic quantization evade the sign problem? — the relativistic Bose gas at finite chemical potential*, *Phys. Rev. Lett.* **102** (2009) 131601 [[arXiv:0810.2089](#)] [[SPIRES](#)].
- [2] Z. Fodor and S.D. Katz, *A new method to study lattice QCD at finite temperature and chemical potential*, *Phys. Lett. B* **534** (2002) 87 [[hep-lat/0104001](#)] [[SPIRES](#)].
- [3] Z. Fodor and S.D. Katz, *Lattice determination of the critical point of QCD at finite T and μ* , *JHEP* **03** (2002) 014 [[hep-lat/0106002](#)] [[SPIRES](#)].
- [4] Z. Fodor, S.D. Katz and K.K. Szabo, *The QCD equation of state at nonzero densities: lattice result*, *Phys. Lett. B* **568** (2003) 73 [[hep-lat/0208078](#)] [[SPIRES](#)].
- [5] Z. Fodor and S.D. Katz, *Critical point of QCD at finite T and μ , lattice results for physical quark masses*, *JHEP* **04** (2004) 050 [[hep-lat/0402006](#)] [[SPIRES](#)].
- [6] C.R. Allton et al., *The QCD thermal phase transition in the presence of a small chemical potential*, *Phys. Rev. D* **66** (2002) 074507 [[hep-lat/0204010](#)] [[SPIRES](#)].
- [7] C.R. Allton et al., *The equation of state for two flavor QCD at non-zero chemical potential*, *Phys. Rev. D* **68** (2003) 014507 [[hep-lat/0305007](#)] [[SPIRES](#)].
- [8] C.R. Allton et al., *Thermodynamics of two flavor QCD to sixth order in quark chemical potential*, *Phys. Rev. D* **71** (2005) 054508 [[hep-lat/0501030](#)] [[SPIRES](#)].

- [9] R.V. Gavai and S. Gupta, *Pressure and non-linear susceptibilities in QCD at finite chemical potentials*, *Phys. Rev. D* **68** (2003) 034506 [[hep-lat/0303013](#)] [[SPIRES](#)].
- [10] P. de Forcrand and O. Philipsen, *The QCD phase diagram for small densities from imaginary chemical potential*, *Nucl. Phys. B* **642** (2002) 290 [[hep-lat/0205016](#)] [[SPIRES](#)].
- [11] P. de Forcrand and O. Philipsen, *The QCD phase diagram for three degenerate flavors and small baryon density*, *Nucl. Phys. B* **673** (2003) 170 [[hep-lat/0307020](#)] [[SPIRES](#)].
- [12] P. de Forcrand and O. Philipsen, *The chiral critical line of $N_f = 2 + 1$ QCD at zero and non-zero baryon density*, *JHEP* **01** (2007) 077 [[hep-lat/0607017](#)] [[SPIRES](#)].
- [13] M. D'Elia and M.-P. Lombardo, *Finite density QCD via imaginary chemical potential*, *Phys. Rev. D* **67** (2003) 014505 [[hep-lat/0209146](#)] [[SPIRES](#)].
- [14] Z. Fodor, S.D. Katz and C. Schmidt, *The density of states method at non-zero chemical potential*, *JHEP* **03** (2007) 121 [[hep-lat/0701022](#)] [[SPIRES](#)].
- [15] S. Ejiri, *Canonical partition function and finite density phase transition in lattice QCD*, *Phys. Rev. D* **78** (2008) 074507 [[arXiv:0804.3227](#)] [[SPIRES](#)].
- [16] J.C. Osborn, *Universal results from an alternate random matrix model for QCD with a baryon chemical potential*, *Phys. Rev. Lett.* **93** (2004) 222001 [[hep-th/0403131](#)] [[SPIRES](#)].
- [17] G. Akemann, J.C. Osborn, K. Splittorff and J.J.M. Verbaarschot, *Unquenched QCD Dirac operator spectra at nonzero baryon chemical potential*, *Nucl. Phys. B* **712** (2005) 287 [[hep-th/0411030](#)] [[SPIRES](#)].
- [18] J.C. Osborn, K. Splittorff and J.J.M. Verbaarschot, *Chiral symmetry breaking and the Dirac spectrum at nonzero chemical potential*, *Phys. Rev. Lett.* **94** (2005) 202001 [[hep-th/0501210](#)] [[SPIRES](#)].
- [19] K. Splittorff and J.J.M. Verbaarschot, *Phase of the fermion determinant at nonzero chemical potential*, *Phys. Rev. Lett.* **98** (2007) 031601 [[hep-lat/0609076](#)] [[SPIRES](#)].
- [20] J. Han and M.A. Stephanov, *A random matrix study of the QCD sign problem*, *Phys. Rev. D* **78** (2008) 054507 [[arXiv:0805.1939](#)] [[SPIRES](#)].
- [21] J. Bloch and T. Wettig, *Random matrix analysis of the QCD sign problem for general topology*, *JHEP* **03** (2009) 100 [[arXiv:0812.0324](#)] [[SPIRES](#)].
- [22] S. Chandrasekharan and U.-J. Wiese, *Meron-cluster solution of a fermion sign problem*, *Phys. Rev. Lett.* **83** (1999) 3116 [[cond-mat/9902128](#)] [[SPIRES](#)].
- [23] M.G. Endres, *Method for simulating $O(N)$ lattice models at finite density*, *Phys. Rev. D* **75** (2007) 065012 [[hep-lat/0610029](#)] [[SPIRES](#)].
- [24] S. Chandrasekharan, *A new computational approach to lattice quantum field theories*, PoS(LATTICE 2008)003 [[arXiv:0810.2419](#)] [[SPIRES](#)].
- [25] G. Parisi and Y.s. Wu, *Perturbation theory without gauge fixing*, *Sci. Sin.* **24** (1981) 483 [[SPIRES](#)].
- [26] G. Parisi, *On complex probabilities*, *Phys. Lett. B* **131** (1983) 393 [[SPIRES](#)].
- [27] J.R. Klauder and W.P. Petersen, *Numerical integration of multiplicative noise stochastic differential equations*, PRINT-85-0297 (BTL) [[SPIRES](#)]; *Spectrum of certain non-self-adjoint operators and solutions of Langevin equations with complex drift*, *J. Stat. Phys.* **39** (1985) 53.
- [28] F. Karsch and H.W. Wyld, *Complex Langevin simulation of the SU(3) spin model with nonzero chemical potential*, *Phys. Rev. Lett.* **55** (1985) 2242 [[SPIRES](#)].

- [29] J. Ambjørn, M. Flensburg and C. Peterson, *The complex Langevin equation and Monte Carlo simulations of actions with static charges*, *Nucl. Phys. B* **275** (1986) 375 [SPIRES].
- [30] P.H. Damgaard and H. Huffel, *Stochastic quantization*, *Phys. Rept.* **152** (1987) 227 [SPIRES].
- [31] J. Berges and I.O. Stamatescu, *Simulating nonequilibrium quantum fields with stochastic quantization techniques*, *Phys. Rev. Lett.* **95** (2005) 202003 [hep-lat/0508030] [SPIRES].
- [32] J. Berges, S. Borsányi, D. Sexty and I.O. Stamatescu, *Lattice simulations of real-time quantum fields*, *Phys. Rev. D* **75** (2007) 045007 [hep-lat/0609058] [SPIRES].
- [33] J. Berges and D. Sexty, *Real-time gauge theory simulations from stochastic quantization with optimized updating*, *Nucl. Phys. B* **799** (2008) 306 [arXiv:0708.0779] [SPIRES].
- [34] C.W. Bernard and V.M. Savage, *Numerical simulations of PT-symmetric quantum field theories*, *Phys. Rev. D* **64** (2001) 085010 [hep-lat/0106009] [SPIRES].
- [35] C. Pehlevan and G. Guralnik, *Complex Langevin equations and Schwinger-Dyson equations*, *Nucl. Phys. B* **811** (2009) 519 [arXiv:0710.3756] [SPIRES].
- [36] G. Aarts and I.-O. Stamatescu, *Stochastic quantization at finite chemical potential*, *JHEP* **09** (2008) 018 [arXiv:0807.1597] [SPIRES].
- [37] T.D. Cohen, *Functional integrals for QCD at nonzero chemical potential and zero density*, *Phys. Rev. Lett.* **91** (2003) 222001 [hep-ph/0307089] [SPIRES].
- [38] G. Guralnik and C. Pehlevan, *Effective potential for complex Langevin equations*, [arXiv:0902.1503] [SPIRES].
- [39] P. Hasenfratz and F. Karsch, *Chemical potential on the lattice*, *Phys. Lett. B* **125** (1983) 308 [SPIRES].
- [40] J.I. Kapusta, *Finite-temperature field theory*, Cambridge University Press, Cambridge U.K. (1994).
- [41] G.G. Batrouni et al., *Langevin simulations of lattice field theories*, *Phys. Rev. D* **32** (1985) 2736 [SPIRES].
- [42] J. Ambjørn and S.K. Yang, *Numerical problems in applying the Langevin equation to complex effective actions*, *Phys. Lett. B* **165** (1985) 140 [SPIRES].
- [43] H. Nakazato and Y. Yamanaka, *Minkowski stochastic quantization*, *Phys. Rev. D* **34** (1986) 492 [SPIRES].
- [44] G. Aarts, G.F. Bonini and C. Wetterich, *Exact and truncated dynamics in nonequilibrium field theory*, *Phys. Rev. D* **63** (2001) 025012 [hep-ph/0007357] [SPIRES].