

Constraints on the nonuniversal Z couplings from $B \rightarrow \pi K$, πK^* and ρK decays

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

JHEP05(2009)056

(<http://iopscience.iop.org/1126-6708/2009/05/056>)

[The Table of Contents](#) and [more related content](#) is available

Download details:

IP Address: 80.92.225.132

The article was downloaded on 03/04/2010 at 09:19

Please note that [terms and conditions apply](#).

Constraints on the nonuniversal Z' couplings from $B \rightarrow \pi K, \pi K^*$ and ρK Decays

Qin Chang,^{a,b} Xin-Qiang Li^{c,1} and Ya-Dong Yang^{a,d,2}

^a*Institute of Particle Physics, Huazhong Normal University,
Wuhan, Hubei 430079, P. R. China*

^b*Department of Physics, Henan Normal University,
Xinxiang, Henan 453007, P. R. China*

^c*Institut für Theoretische Physik E, RWTH Aachen University,
D-52056 Aachen, Germany*

^d*Key Laboratory of Quark & Lepton Physics, Ministry of Education,
P. R. China*

*E-mail: changqin@iopp.cnu.edu.cn, xinqiang@physik.rwth-aachen.de,
yangyd@iopp.cnu.edu.cn*

ABSTRACT: Motivated by the large difference between the direct CP asymmetries $A_{CP}(B^- \rightarrow \pi^0 K^-)$ and $A_{CP}(\bar{B}^0 \rightarrow \pi^+ K^-)$, we combine the up-to-date experimental information on $B \rightarrow \pi K, \pi K^*$ and ρK decays to pursue possible solutions with the nonuniversal Z' model. Detailed analyses of the relative impacts of different types of couplings are presented in four specific cases. Numerically, we find that the new coupling parameters, ξ^{LL} and ξ^{LR} with a common nontrivial new weak phase $\phi_L \sim -86^\circ$, which are relevant to the Z' contributions to the electroweak penguin sector ΔC_9 and ΔC_7 , are crucial to the observed “ πK puzzle”. Furthermore, they are found to be definitely unequal and opposite in sign. We also find that $A_{CP}(B^- \rightarrow \rho^0 K^-)$ can put a strong constraint on the new Z' couplings, which implies the Z' contributions to the coefficient of QCD penguins operator O_3 involving the parameter ζ^{LL} required.

KEYWORDS: Rare Decays, Beyond Standard Model, B-Physics, CP violation

ARXIV EPRINT: [0903.0275](https://arxiv.org/abs/0903.0275)

¹Alexander-von-Humboldt Fellow

²Corresponding author

Contents

1	Introduction	1
2	The SM results with two schemes for the end-point divergence	3
3	Solution to the “πK puzzle” with nonuniversal Z' model	5
3.1	Formalism of the family nonuniversal Z' model	5
3.2	Numerical analyses and discussions	7
4	Conclusions	14
A	Decay amplitudes in the SM with QCDF	15
B	The hard-spectator and annihilation corrections with the infrared finite gluon propagator	15
C	Theoretical input parameters	16
C.1	CKM matrix elements	16
C.2	Quark masses and lifetimes	16
C.3	The decay constants and form factors	17
C.4	The LCDAs of mesons and light-cone projector operators.	17

1 Introduction

During the past several years, the observed discrepancies between the experimental measurements and the theoretical predications within the Standard Model (SM) for several observables in $B \rightarrow \pi K$ decays, the so-called “ πK puzzle” [1], have attracted much attention. Extensive investigations both within the SM [2–7], as well as with various specific New Physics (NP) scenarios [8–10], have been performed.

Averaging the recent experimental data from BABAR [11], Belle [12], CLEO [13] and CDF [14], the Heavy Flavor Averaging Group (HFAG) gives the following up-to-date results [15]

$$\begin{aligned}
 A_{CP}(B^- \rightarrow K^- \pi^0) &= 0.050 \pm 0.025, \\
 A_{CP}(\bar{B}^0 \rightarrow K^- \pi^+) &= -0.097 \pm 0.012,
 \end{aligned}
 \tag{1.1}$$

from which the difference between direct CP violations in the charged and the neutral modes

$$\Delta A \equiv A_{CP}(B^- \rightarrow K^- \pi^0) - A_{CP}(\bar{B}^0 \rightarrow K^- \pi^+) = 0.147 \pm 0.028
 \tag{1.2}$$

is now established at about 5σ level.

Theoretically, it is generally expected that within the SM, these two CP asymmetries $A_{CP}(\bar{B}_d^0 \rightarrow \pi^+ K^-)$ and $A_{CP}(B_u^- \rightarrow \pi^0 K^-)$ should be approximately equal. For example, based on the QCD factorization approach (QCDF) [16], the recent theoretical predictions with two different schemes for the end-point divergence are

$$\begin{cases} A_{CP}(B_u^- \rightarrow \pi^0 K^-) = -3.6\%, \\ A_{CP}(\bar{B}_d^0 \rightarrow \pi^+ K^-) = -4.1\%, \end{cases} \text{Scheme I (Scenario S4) [3],} \quad (1.3)$$

$$\begin{cases} A_{CP}(B_u^- \rightarrow \pi^0 K^-) = -10.8\%, \\ A_{CP}(\bar{B}_d^0 \rightarrow \pi^+ K^-) = -12.4\%, \end{cases} \text{Scheme II } (m_g = 0.5 \text{ MeV}) [8]. \quad (1.4)$$

Here, the Scheme I is the way to parameterize the end-point divergence appearing in hard-spectator and annihilation corrections, by complex parameters $X_{A,H} = \int_0^1 dy/y = \ln(m_b/\Lambda)(1 + \rho_{A,H}e^{i\phi_{A,H}})$, with $\rho_{A,H} \leq 1$ and unrestricted $\phi_{A,H}$ [3]. The Scheme II, as an alternative to the first one, is the way to quote the infrared finite gluon propagator to regulate the divergence. It is interesting to note that an infrared finite behavior of gluon propagator are not only obtained by solving the well-known Schwinger-Dyson equation [17–19], but also supported by recent Lattice QCD simulations [20]. However, both of these two schemes suffer the mismatch of ΔA given by eq. (1.2). Furthermore, within the framework of perturbative QCD approach (pQCD) [21], and the soft-collinear effective theory (SCET) [22], the theoretical predictions read

$$\begin{cases} A_{CP}(B_u^- \rightarrow \pi^0 K^-)_{PQCD} = (-1_{-5}^{+3})\%, \\ A_{CP}(\bar{B}_d^0 \rightarrow \pi^+ K^-)_{PQCD} = (-9_{-8}^{+6})\%, \end{cases} \text{pQCD [5],} \quad (1.5)$$

$$\begin{cases} A_{CP}(B_u^- \rightarrow \pi^0 K^-)_{SCET} = (-11 \pm 9 \pm 11 \pm 2)\%, \\ A_{CP}(\bar{B}_d^0 \rightarrow \pi^+ K^-)_{SCET} = (-6 \pm 5 \pm 6 \pm 2)\%. \end{cases} \text{SCET [6].} \quad (1.6)$$

Obviously, the present theoretical estimations within the SM are not consistent with the established ΔA . The mismatch might be due to our current limited understanding of the strong dynamics involved in hadronic B decays, but equally also to possible NP effects [23, 24].

In some well-motivated extensions of the SM, additional $U(1)'$ gauge symmetries and associated Z' gauge boson could arise. Searching for the extra Z' boson is an important mission in the experimental programs of Tevatron [25] and LHC [26]. Performing the constraints on the new Z' couplings through low-energy physics, on the other hand, is very important for the direct searches and understanding its phenomenology. Theoretically, the flavor changing neutral current (FCNC) is forbidden at tree level in the SM. One of the simple extensions is the family nonuniversal Z' model, which could be naturally derived in certain string constructions [27], E_6 models [28] and so on. It is interesting to note that the nonuniversal Z' couplings could lead to FCNC and new CP-violating effect [29], which possibly provide a solution to the afore mentioned “ πK puzzle”. With some simplifications of the nonuniversal Z' model and neglecting the color-suppressed electroweak (EW)

penguins and the annihilation amplitudes, ref. [9] gets four possible solutions

$$\begin{aligned}
 A_L : \{\xi^{LL}, \phi_L\} &= \{0.0055, 110^\circ\}, & B_L : \{\xi^{LL}, \phi_L\} &= \{0.0098, -97^\circ\}, \\
 & & & \text{with } \xi^{LR} = 0; \\
 A_{LR} : \{\xi^{LL} = \xi^{LR}, \phi_L\} &= \{0.0104, -70^\circ\}, & B_{LR} : \{\xi^{LL} = \xi^{LR}, \phi_L\} &= \{0.0186, 83^\circ\}. \quad (1.7)
 \end{aligned}$$

However, the corresponding prediction $A_{CP}(B_u^- \rightarrow \pi^0 K^-) = -0.03 \pm 0.01$ [9] of solution A_L and A_{LR} in eq. (1.7) is obviously inconsistent with the up-to-date experimental data 0.050 ± 0.025 . Moreover, the annihilation amplitudes, which could generate some strong-interaction phases, are important for predicting CP violations.

Based on the above observations, in this paper we shall adopt the QCDF approach and reevaluate the effects of the nonuniversal Z' model on these decay modes with the updated experimental data. Furthermore, since the $B \rightarrow \pi K^*$ and ρK decays also involve the same quark level $b \rightarrow s\bar{q}q$ ($q = u, d$) transition, it is necessary to take into account these decay modes.

In section 2, we provide a quick survey of $B \rightarrow \pi K$, πK^* and ρK decays in the SM within the QCDF formalism; our numerical results, with two different schemes for the end-point divergence, are also presented. In section 3, after reviewing the nonuniversal Z' model briefly, we present our analyses and numerical results in detail. Section 4 contains our conclusions. Appendix A recapitulates the decay amplitudes for the twelve decay modes within the SM [3]. Appendix B contains the formulas for hard-spectator and annihilation amplitudes with the infrared finite gluon propagator [8]. All the theoretical input parameters are summarized in appendix C.

2 The SM results with two schemes for the end-point divergence

In the SM, the effective Hamiltonian responsible for $b \rightarrow s$ transitions is given as [31]

$$\begin{aligned}
 \mathcal{H}_{\text{eff}} &= \frac{G_F}{\sqrt{2}} \left[V_{ub}V_{us}^* (C_1 O_1^u + C_2 O_2^u) + V_{cb}V_{cs}^* (C_1 O_1^c + C_2 O_2^c) \right. \\
 &\quad \left. - V_{tb}V_{ts}^* \left(\sum_{i=3}^{10} C_i O_i + C_{7\gamma} O_{7\gamma} + C_{8g} O_{8g} \right) \right] + \text{h.c.}, \quad (2.1)
 \end{aligned}$$

where $V_{qb}V_{qs}^*$ ($q = u, c$ and t) are products of the Cabibbo-Kobayashi-Maskawa (CKM) matrix elements [30], C_i the Wilson coefficients, and O_i the relevant four-quark operators whose explicit forms could be found, for example, in refs. [2, 31].

In recent years, the QCDF approach has been employed extensively to study the hadronic B-meson decays. The $B \rightarrow \pi K$, πK^* and ρK decays have been studied comprehensively within the SM in refs. [2–4, 32], and the relevant decay amplitudes within this formalism are shown in appendix A. It is also noted that the framework contains estimates of the hard-spectator and annihilation corrections. Even though they are power-suppressed, their strength and associated strong-interaction phase are numerically important to evaluate the branching ratio and the CP asymmetry. However, unfortunately, the end-point

singularities appear in twist-3 spectator and annihilation amplitudes. So, how to regulate the end-point divergence becomes important and necessary within this formalism. Here we shall adopt the following two schemes:

Scheme I: Parametrization. As the most popular way, the end-point divergent integrals are treated as signs of infrared sensitive contributions and phenomenologically parameterized by [2, 3]

$$\int_0^1 \frac{dy}{y} \rightarrow X_A = (1 + \rho_A e^{i\phi_A}) \ln \frac{m_B}{\Lambda_h}, \quad \int_0^1 dy \frac{\ln y}{y} \rightarrow -\frac{1}{2}(X_A)^2, \quad (2.2)$$

with $\Lambda_h = 0.5 \text{ GeV}$, $\rho_A \leq 1$ and ϕ_A unrestricted. X_H is treated in the same manner. The different choices of ρ_A and ϕ_A correspond to different scenarios as discussed in ref. [3], and S4 is mentioned as the most favorable one. It presents the moderate value of nonuniversal annihilation phase $\phi_A = -55^\circ$ (PP), -20° (PV) and -70° (VP). Conservatively, in our calculations we quote $\pm 5^\circ$ as their theoretical uncertainties. Taking $\rho_A = 1$ and $X_{A,H}$ universal for all decay processes belonging to the same modes (PP, PV or VP), we present our numerical results of branching ratios and direct CP asymmetries for $B \rightarrow \pi K$, πK^* and ρK decays in the third column of tables 2 and 3, respectively.

As is known, the mixing-induced CP asymmetry A_{CP}^{mix} is well suited for testing the SM and searching for new physics effects. For example, the investigation of mixing-induced CP asymmetries in penguin dominated $\bar{B}^0 \rightarrow \pi^0 K_S^0$ and $\bar{B}^0 \rightarrow \rho^0 K_S^0$ decay modes has attracted much attention recently [33–36]. After neglecting the $K_0 - \bar{K}_0$ mixing effect, the mixing-induced asymmetry could be written as

$$A_{CP}^{\text{mix}}(\bar{B}^0 \rightarrow f) = \frac{2 \text{Im}\lambda_f}{1 + |\lambda_f|^2}, \quad (f = \pi^0 K_S^0, \rho^0 K_S^0), \quad (2.3)$$

with $\lambda_f = -\exp\left\{i \arg \left[\frac{V_{td} V_{tb}^*}{V_{td}^* V_{tb}} \right]\right\} \bar{A}_f / A_f$ in our phase convention. Our numerical predictions are listed in table 4, which agree with the measurements within large experimental errors.

Scheme II: Infrared finite dynamical gluon propagator. In our previous paper [8], we have thoroughly studied the end-point divergence with an infrared finite dynamical gluon propagator. It is interesting to note that recent theoretical and phenomenological studies are now accumulating supports for a softer infrared behavior of the gluon propagator [19, 37, 38]. Furthermore, the infrared finite dynamical gluon propagator, which is shown to be not divergent as fast as $\frac{1}{q^2}$, has been successfully applied to the hadronic B-meson decays [39, 40]. In our evaluations, we shall quote the gluon propagator derived by Cornwall (in Minkowski space) [17]

$$D(q^2) = \frac{1}{q^2 - M_g^2(q^2) + i\epsilon}, \quad (2.4)$$

where q is the gluon momentum. The corresponding strong coupling constant reads

$$\alpha_s(q^2) = \frac{4\pi}{\beta_0 \ln\left(\frac{q^2 + 4M_g^2(q^2)}{\Lambda_{QCD}^2}\right)}, \quad (2.5)$$

where $\beta_0 = 11 - \frac{2}{3}n_f$ is the first coefficient of the beta function, with n_f being the number of active quark flavors. The dynamical gluon mass square $M_g^2(q^2)$ is obtained as [17]

$$M_g^2(q^2) = m_g^2 \left[\frac{\ln\left(\frac{q^2 + 4m_g^2}{\Lambda_{QCD}^2}\right)}{\ln\left(\frac{4m_g^2}{\Lambda_{QCD}^2}\right)} \right]^{-\frac{12}{11}}, \quad (2.6)$$

where m_g is the effective gluon mass and $\Lambda_{QCD} = 225$ MeV. In ref. [8], we present our suggestion, $m_g = 0.50 \pm 0.05$ GeV, which is a reasonable choice so that most of the observables (except for $A_{CP}(B \rightarrow \pi^0 K^-)$) are in good agreement with the experimental data. In this way, we find that the hard-spectator scattering contributions are real, and the annihilation contributions are complex with a large imaginary part [8]. Our numerical predictions for branching ratios, direct CP asymmetries and mixing-induced CP asymmetries are listed in the fourth column of tables 2, 3 and 4, respectively.

Although numerically these two schemes have some differences, both of their predictions are consistent with most of the experimental data within errors. However, as expected in the SM, we again find that $A_{CP}(B_u^- \rightarrow \pi^0 K^-) = -0.041 \pm 0.008$ (-0.100 ± 0.008), are very close to $A_{CP}(\bar{B}_d^0 \rightarrow \pi^+ K^-) = -0.077 \pm 0.009$ (-0.116 ± 0.008) in the first (second) scheme. So, it is still hard to accommodate the measured large difference ΔA in the SM within the QCDF formalism, irrespective of adopting which scheme. In the following, we pursue possible solutions to this problem with a family nonuniversal Z' model [29].

3 Solution to the “ πK puzzle” with nonuniversal Z' model

3.1 Formalism of the family nonuniversal Z' model

A possible heavy Z' boson is predicted in many extensions of the SM, such as grand unified theories, superstring theories, and theories with large extra dimensions. The simplest way to extend the SM gauge structure is to include a new U(1) gauge group. A family nonuniversal Z' model can lead to FCNC processes even at tree level due to the non-diagonal chiral coupling matrix. The formalism of the model has been detailed in ref. [29]. The relevant studies in the context of B physics have also been extensively performed in refs. [9, 42, 43, 45].

After neglecting the $Z-Z'$ mixing with small mixing angle $\theta \sim \mathcal{O}(10^{-3})$ [44], and taking all the fields being the physical eigenstates, the Z' part of the neutral-current Lagrangian can be written as [29]

$$\mathcal{L}' = -g' J'_\mu Z'^\mu, \quad (3.1)$$

where g' is the gauge coupling constant of extra $U'(1)$ group at the EW M_W scale. The Z' chiral current is

$$J'_\mu = \bar{\psi}_i \gamma_\mu \left[(B_q^L)_{ij} P_L + (B_q^R)_{ij} P_R \right] \psi_j, \quad (3.2)$$

where ψ is the mass eigenstate of chiral fields and $P_{L,R} = (1 \mp \gamma_5)/2$. The effective chiral Z' coupling matrices are given as

$$B_q^X = V_{qX} \epsilon_{qX} V_{qX}^\dagger, \quad (q = u, d; X = L, R). \quad (3.3)$$

With the assumption of flavor-diagonal right-handed couplings, the Z' part of the effective Hamiltonian for $b \rightarrow s\bar{q}q$ ($q = u, d$) transitions can be written as [9]

$$\mathcal{H}_{eff}^{Z'} = \frac{2G_F}{\sqrt{2}} \left(\frac{g' M_Z}{g_1 M_{Z'}} \right)^2 B_{sb}^L (\bar{s}b)_{V-A} \sum_q (B_{qq}^L (\bar{q}q)_{V-A} + B_{qq}^R (\bar{q}q)_{V+A}) + h.c., \quad (3.4)$$

where $g_1 = e/(\sin \theta_W \cos \theta_W)$ and $M_{Z'}$ the new gauge boson mass. It is noted that the forms of the above operators already exist in the SM. As a result, eq. (3.4) can be modified as

$$\mathcal{H}_{eff}^{Z'} = -\frac{G_F}{\sqrt{2}} V_{ts}^* V_{tb} \sum_q (\Delta C_3 O_3^q + \Delta C_5 O_5^q + \Delta C_7 O_7^q + \Delta C_9 O_9^q) + h.c., \quad (3.5)$$

where O_i^q ($i = 3, 5, 7, 9$) are the effective operators in the SM, and ΔC_i the modifications to the corresponding SM Wilson coefficients caused by Z' boson, which are expressed as

$$\begin{aligned} \Delta C_{3,5} &= -\frac{2}{3V_{ts}^* V_{tb}} \left(\frac{g' M_Z}{g_1 M_{Z'}} \right)^2 B_{sb}^L (B_{uu}^{L,R} + 2B_{dd}^{L,R}), \\ \Delta C_{9,7} &= -\frac{4}{3V_{ts}^* V_{tb}} \left(\frac{g' M_Z}{g_1 M_{Z'}} \right)^2 B_{sb}^L (B_{uu}^{L,R} - B_{dd}^{L,R}), \end{aligned} \quad (3.6)$$

in terms of the model parameters at the M_W scale.

Generally, the diagonal elements of the effective coupling matrices $B_{qq}^{L,R}$ are real as a result of the hermiticity of the effective Hamiltonian. However, the off-diagonal ones of B_{sb}^L can contain a new weak phase ϕ_L . Then, conveniently we can represent ΔC_i as¹

$$\begin{aligned} \Delta C_{3,5} &= 2 \frac{|V_{ts}^* V_{tb}|}{V_{ts}^* V_{tb}} \zeta^{LL,LR} e^{i\phi_L}, \\ \Delta C_{9,7} &= 4 \frac{|V_{ts}^* V_{tb}|}{V_{ts}^* V_{tb}} \xi^{LL,LR} e^{i\phi_L}, \end{aligned} \quad (3.7)$$

where the real NP parameters $\zeta^{LL,LR}$, $\xi^{LL,LR}$ and ϕ_L are defined, respectively, as

$$\begin{aligned} \zeta^{LL,LR} &= -\frac{1}{3} \left(\frac{g' M_Z}{g_1 M_{Z'}} \right)^2 \left| \frac{B_{sb}^L}{V_{ts}^* V_{tb}} \right| (B_{uu}^{L,R} + 2B_{dd}^{L,R}), \\ \xi^{LL,LR} &= -\frac{1}{3} \left(\frac{g' M_Z}{g_1 M_{Z'}} \right)^2 \left| \frac{B_{sb}^L}{V_{ts}^* V_{tb}} \right| (B_{uu}^{L,R} - B_{dd}^{L,R}), \\ \phi_L &= \text{Arg} [B_{sb}^L]. \end{aligned} \quad (3.8)$$

It is noted that the other SM Wilson coefficients may also receive contributions from the Z' boson through renormalization group (RG) evolution. With our assumption that no significant RG running effect between M_Z' and M_W scales, the RG evolution of the modified Wilson coefficients is exactly the same as the ones in the SM [31, 41]. For simplicity, we define

$$\begin{aligned} X' &= \zeta^{LL} e^{i\phi_L}, & Y' &= \zeta^{LR} e^{i\phi_L}, \\ X &= \xi^{LL} e^{i\phi_L}, & Y &= \xi^{LR} e^{i\phi_L}. \end{aligned} \quad (3.9)$$

¹ For comparison, we take the same phase convention as ref. [9].

Wilson coefficients	$\mu = m_b$		$\mu_h = \sqrt{\Lambda_h m_b}$	
	C_i^{SM}	$\Delta C_i^{Z'}$	C_i^{SM}	$\Delta C_i^{Z'}$
C_1	1.075	-0.006X	1.166	-0.008X
C_2	-0.170	-0.009X	-0.336	-0.014X
C_3	0.013	$0.05X - 0.01Y - 2.20X' - 0.05Y'$	0.025	$0.11X - 0.02Y - 2.37X' - 0.12Y'$
C_4	-0.033	$-0.13X + 0.01Y + 0.55X' + 0.02Y'$	-0.057	$-0.24X + 0.02Y + 0.92X' + 0.09Y'$
C_5	0.008	$0.03X + 0.01Y - 0.06X' - 1.83Y'$	0.011	$0.03X + 0.02Y - 0.10X' + 0.09Y'$
C_6	-0.038	$-0.15X + 0.01Y + 0.1X' - 0.6Y'$	-0.076	$-0.32X + 0.04Y + 0.16X' - 1.26Y'$
C_7/α_{em}	-0.015	$4.18X - 473Y + 0.25X' + 1.27Y'$	-0.034	$5.7X - 459Y + 0.4X' + 1.7Y'$
C_8/α_{em}	0.045	$1.18X - 166Y + 0.01X' + 0.56Y'$	0.089	$3.2X - 355Y + 0.2X' + 1.5Y'$
C_9/α_{em}	-1.119	$-561X + 4.52Y - 0.8X' + 0.4Y'$	-1.228	$-611X + 6.7Y - 1.2X' + 0.6Y'$
C_{10}/α_{em}	0.190	$118X - 0.5Y + 0.2X' - 0.05Y'$	0.356	$207X - 1.4Y + 0.5X' - 0.1Y'$
$C_{7\gamma}$	-0.297	—	0.360	—
C_{8g}	-0.143	—	-0.168	—

Table 1. The Wilson coefficients C_i within the SM and with the contribution from Z' boson included in NDR scheme at the scale $\mu = m_b$ and $\mu_h = \sqrt{\Lambda_h m_b}$.

Decay Mode	Exp. data	SM		Z' model			
		Scheme I	Scheme II	Case I	Case II	Case III	Case IV
$B_u^- \rightarrow \pi^- \bar{K}^0$	23.1 ± 1.0	19.0 ± 2.5	23.4 ± 3.9	23.3 ± 0.7	23.3 ± 0.6	23.2 ± 0.6	23.3 ± 0.7
$B_u^- \rightarrow \pi^0 K^-$	12.9 ± 0.6	10.5 ± 1.3	12.7 ± 2.0	12.5 ± 0.6	12.6 ± 0.6	12.5 ± 0.5	12.6 ± 0.6
$\bar{B}_d^0 \rightarrow \pi^+ K^-$	19.4 ± 0.6	16.2 ± 2.2	20.1 ± 3.4	19.9 ± 0.5	19.8 ± 0.5	19.9 ± 0.5	20.0 ± 0.5
$\bar{B}_d^0 \rightarrow \pi^0 \bar{K}^0$	9.8 ± 0.6	7.3 ± 1.1	9.3 ± 1.7	9.4 ± 0.6	9.5 ± 0.6	9.1 ± 0.4	9.1 ± 0.4
$B_u^- \rightarrow \pi^- \bar{K}^{*0}$	10.0 ± 0.8	11.7 ± 1.2	10.3 ± 3.3	8.4 ± 1.0	8.5 ± 0.9	8.7 ± 0.6	8.6 ± 0.7
$B_u^- \rightarrow \pi^0 K^{*-}$	6.9 ± 2.3	7.0 ± 0.7	6.0 ± 1.8	4.7 ± 0.6	4.7 ± 0.5	4.9 ± 0.3	4.8 ± 0.3
$\bar{B}_d^0 \rightarrow \pi^+ K^{*-}$	10.3 ± 1.1	9.9 ± 1.1	9.2 ± 2.8	7.5 ± 1.0	7.7 ± 0.9	8.0 ± 0.6	8.0 ± 0.6
$\bar{B}_d^0 \rightarrow \pi^0 \bar{K}^{*0}$	2.4 ± 0.7	4.1 ± 0.5	3.9 ± 1.3	3.6 ± 0.5	3.7 ± 0.4	3.5 ± 0.4	3.5 ± 0.4
$B_u^- \rightarrow \rho^- \bar{K}^0$	$8.0^{+1.5}_{-1.4}$	5.2 ± 0.9	10.6 ± 3.7	9.6 ± 1.4	9.7 ± 1.3	10.6 ± 1.3	10.7 ± 1.5
$B_u^- \rightarrow \rho^0 K^-$	$3.81^{+0.48}_{-0.46}$	2.5 ± 0.4	5.4 ± 1.6	4.22 ± 0.62	4.47 ± 0.63	4.7 ± 0.6	4.8 ± 0.7
$\bar{B}_d^0 \rightarrow \rho^+ K^-$	$8.6^{+0.9}_{-1.1}$	6.3 ± 1.0	13.0 ± 3.8	10.8 ± 1.4	10.9 ± 1.4	11.9 ± 1.4	12.5 ± 1.8
$\bar{B}_d^0 \rightarrow \rho^0 \bar{K}^0$	$5.4^{+0.9}_{-1.0}$	3.7 ± 0.5	7.3 ± 2.1	7.1 ± 0.9	7.4 ± 0.9	6.8 ± 0.9	6.9 ± 1.0

Table 2. The CP -averaged branching ratios (in units of 10^{-6}) of $B \rightarrow \pi K$, πK^* and ρK decays in the SM with two end-point divergence regulation schemes, and in the nonuniversal Z' model with four different cases.

The numerical results of Wilson coefficients in the naive dimensional regularization (NDR) scheme at the scale $\mu = m_b$ ($\mu_h = \sqrt{\Lambda_h m_b}$) are listed in table 1. The values at the scale μ_h , with $m_b = 4.79$ GeV and $\Lambda_h = 500$ MeV, should be used in the calculation of hard-spectator and weak annihilation contributions.

3.2 Numerical analyses and discussions

With the theoretical formulas and the input parameters summarized in appendix A, B and C, we now present our numerical analyses and discussions. Our analyses are divided into the following four cases with different simplifications for our attention, namely,

Decay Mode	Exp. data	SM		Z' model			
		Scheme I	Scheme II	Case I	Case II	Case III	Case IV
$B_u^- \rightarrow \pi^- \bar{K}^0$	0.9 ± 2.5	0.4 ± 0.1	0.04 ± 0.07	-1.6 ± 0.3	-2.7 ± 0.9	5.2 ± 0.5	5.1 ± 0.6
$B_u^- \rightarrow \pi^0 K^-$	5.0 ± 2.5	-4.1 ± 0.8	-10.0 ± 0.8	2.4 ± 1.6	2.3 ± 1.5	0.9 ± 0.7	1.2 ± 0.9
$\bar{B}_d^0 \rightarrow \pi^+ K^-$	$-9.8_{-1.1}^{+1.2}$	-7.7 ± 0.9	-11.6 ± 0.3	-11.7 ± 0.3	-11.0 ± 0.7	-10.5 ± 1.1	-10.5 ± 1.2
$\bar{B}_d^0 \rightarrow \pi^0 \bar{K}^0$	-1 ± 10	-1.5 ± 0.3	0.7 ± 0.3	-17 ± 2	-18 ± 2	-6 ± 2	-6 ± 2
$B_u^- \rightarrow \pi^- \bar{K}^{*0}$	$-2_{-6.1}^{+6.7}$	0.6 ± 0.1	0.09 ± 0.15	-2.1 ± 0.4	-3.3 ± 0.5	-0.6 ± 2.4	-3.0 ± 6.7
$B_u^- \rightarrow \pi^0 K^{*-}$	4 ± 29	-6 ± 2	-37 ± 9	6.8 ± 7.1	9.1 ± 7.2	-17 ± 4	-18 ± 6
$\bar{B}_d^0 \rightarrow \pi^+ K^{*-}$	-25 ± 11	-13 ± 2	-43 ± 10	-48 ± 3	-46 ± 3	-49 ± 3	-50 ± 5
$\bar{B}_d^0 \rightarrow \pi^0 \bar{K}^{*0}$	-15 ± 12	-4 ± 1	4 ± 2	-58 ± 9	-62 ± 9	-34 ± 7	-36 ± 11
$B_u^- \rightarrow \rho^- \bar{K}^0$	-12 ± 17	0.4 ± 0.1	0.5 ± 0.2	1.5 ± 0.1	-0.15 ± 0.7	5.3 ± 1.1	6.5 ± 4.5
$B_u^- \rightarrow \rho^0 K^-$	$41.9_{-10.4}^{+8.1}$	57.3 ± 5.8	42.3 ± 9.5	-36 ± 10	-46 ± 12	27 ± 4	27 ± 5
$\bar{B}_d^0 \rightarrow \rho^+ K^-$	15 ± 6	36 ± 4	29 ± 6	31 ± 3	33 ± 3	25 ± 2	25 ± 2
$\bar{B}_d^0 \rightarrow \rho^0 \bar{K}^0$	1 ± 20	-2.1 ± 1.3	-2.4 ± 1.4	45 ± 5	50 ± 5	8 ± 3	9 ± 4

Table 3. The direct CP asymmetries (in unit of 10^{-2}) of $B \rightarrow \pi K$, πK^* and ρK decays. The other captions are the same as table. 2.

Decay Mode	Exp. data	SM		Z' model			
		Scheme I	Scheme II	Case I	Case II	Case III	Case IV
$\bar{B}_d^0 \rightarrow \pi^0 K_S^0$	57 ± 17	77 ± 2	77 ± 2	46 ± 6	44 ± 6	61 ± 3	62 ± 5
$\bar{B}_d^0 \rightarrow \rho^0 K_S^0$	63_{-21}^{+17}	60 ± 2	66 ± 2	87 ± 2	84 ± 3	85 ± 3	86 ± 9

Table 4. The mixing-induced CP asymmetries (in unit of 10^{-2}) of $\bar{B}^0 \rightarrow \pi^0 K_S^0$ and $\rho^0 K_S^0$ decays. The other captions are the same as table. 2.

- Case I: With the simplifications $B_{uu}^{L,R} \simeq -2B_{dd}^{L,R}$ (i.e., $\zeta^{LL,LR} = 0$) and $\xi^{LR} = 0$,
- Case II: With the simplifications $B_{uu}^{L,R} \simeq -2B_{dd}^{L,R}$ only (i.e., $\zeta^{LL,LR} = 0$),
- Case III: Taking $B_{uu}^R \simeq -2B_{dd}^R$ (i.e., $\zeta^{LR} \simeq 0$), and leaving ζ^{LL} and $\xi^{LL,LR}$ arbitrary,
- Case IV: Without any simplifications for $B_{uu}^{L,R}$ and $B_{dd}^{L,R}$, i.e., arbitrary values for $\zeta^{LL,LR}$ and $\xi^{LL,LR}$ are allowed.

Our fitting is performed with the experimental data varying randomly within their 2σ error-bars, while the theoretical uncertainties are obtained by varying the input parameters within the regions specified in appendix C. In addition, we quote the Scheme II (taking $m_g = 0.5\text{GeV}$) to regulate the appearing end-point divergences.

With the assumption $B_{uu}^{L,R} \simeq -2B_{dd}^{L,R}$ and neglecting the color-suppressed EW penguins and the annihilation amplitudes, four possible solutions eq. (1.7) to the “ πK puzzle” are obtained in ref. [9]. It is still worth to recheck these solutions with the updated experiment data and taken into account the neglected corrections. Furthermore, the possible solutions may also suffer strong constraints from $B \rightarrow \pi K^*$ and ρK decays, since they are also mediated by the same quark level $b \rightarrow s\bar{q}q$ transitions.




Figure 1. The dependence of $A_{CP}(B \rightarrow \pi K)$ on the new weak phase ϕ_L for the values of ξ^{LL} , ξ^{LR} , ζ^{LL} , and ζ^{LR} as marked by the legends.

Case I: With the simplifications $B_{uu}^{L,R} \simeq -2B_{dd}^{L,R}$ (i.e., $\zeta^{LL,LR} = 0$) and $\xi^{LR} = 0$. In this case, assuming $B_{uu}^{L,R} \simeq -2B_{dd}^{L,R}$ as in ref. [9], the NP effect primarily manifests itself in the EW penguin sector and the Z' contribution to the Wilson coefficients eq. (3.6) can be simplified as

$$\begin{aligned} \Delta C_{3,5} &= 0, \\ \Delta C_{9,7} &= 4 \frac{|V_{ts}^* V_{tb}|}{V_{ts}^* V_{tb}} \xi^{LL,LR} e^{i\phi_L}, \quad \text{with } \xi^{LL,LR} = \left(\frac{g' M_Z}{g_1 M_{Z'}} \right)^2 \left| \frac{B_{sb}^L}{V_{ts}^* V_{tb}} \right| B_{dd}^{L,R}. \end{aligned} \quad (3.10)$$

As shown in figure 1 (a), taking $\xi^{LL} = 0.004$ and $\xi^{LR} = 0$, we find that $A_{CP}(B^- \rightarrow \pi^0 K^-)$ is enhanced to be consistent with the experimental data when $\phi_L \sim -90^\circ$. Moreover, $A_{CP}(B^- \rightarrow \pi^- K^0)$ and $A_{CP}(B^0 \rightarrow \pi^+ K^-)$, which agree roughly with the experimental data in the SM, are not sensitive to the parameter ξ^{LL} . So, a possible solution to the observed “ πK puzzle” eq. (1.2) in Case I is naively favored.

Taking $\mathcal{B}(B \rightarrow \pi K)$ and $A_{CP}(B \rightarrow \pi K)$ as constraints on ξ^{LL} and ϕ_L , the allowed region for these two parameters are shown in figure 3 and the corresponding numerical results are listed in table. 5, i.e., $\xi^{LL} = (3.96 \pm 0.70) \times 10^{-3}$ and $\phi_L = -88^\circ \pm 7^\circ$. Our result confirms that the solution B_L in eq. (1.7) is helpful to resolve the “ πK puzzle” (note that a bit of difference might be due to the fact that the annihilation corrections are not




Figure 2. The dependence of $A_{CP}(B \rightarrow \rho K)$ on the new weak phase ϕ_L .




Figure 3. The allowed regions for the parameters ξ^{LL} and ϕ_L in Case I.

included in ref. [9]). However, the solution A_L is excluded by the updated experimental data $A_{CP}(B^- \rightarrow \pi^0 K^-) = 0.050 \pm 0.025$ as indicated in figure 1 (a).

With $\xi^{LL} = (3.96 \pm 0.70) \times 10^{-3}$ and $\phi_L = -88^\circ \pm 7^\circ$ as input parameters, we present our predictions for $\mathcal{B}(B \rightarrow \pi K^*, \rho K)$, $A_{CP}(B \rightarrow \pi K^*, \rho K)$ and $A_{CP}^{\text{mix}}(B^0 \rightarrow \pi^0 K_S, \rho^0 K_S)$




Figure 4. The allowed regions for the parameters $\xi^{LL,LR}$ and ϕ_L in Case II.

in the fifth column of tables 2, 3 and 4, respectively. We can see that most of them are consistent with the experimental data within 2σ . Especially, the predicted $A_{CP}^{\text{mix}}(B^0 \rightarrow \pi^0 K_S) = 0.46 \pm 0.06$ is very close to the measurement 0.57 ± 0.17 [15]. However, the prediction for $A_{CP}(B^- \rightarrow \rho^0 K^-) = -0.36 \pm 0.10$ presents a large discrepancy (larger than 6σ errors) with the current experiment data $0.419_{-0.104}^{+0.081}$ [15], which is also shown in figure 2 (a). This fact implies that $A_{CP}(B^- \rightarrow \rho^0 K^-)$ can provide a strong constraint on the Z' couplings, at least in Case I, and some more general Z' models might be required to explain all of these measurements.

Case II: With the simplification $B_{uu}^{L,R} \simeq -2B_{dd}^{L,R}$ only (i. e. $\zeta^{LL,LR} = 0$). It is interesting to note that, as shown in figure 1 (b), a region of minus ξ^{LR} with $\phi_L \sim -90^\circ$ can bridge the discrepancy of $A_{CP}(B^- \rightarrow \pi^0 K^-)$ between theoretical predictions and experimental data. Moreover, it is also possible to moderate the problem of $A_{CP}(B^- \rightarrow \rho^0 K^-)$ induced by ξ^{LL} as shown in figure 2 (b). So, in Case II we give up the simplification $\xi^{LR} = 0$ and pursue possible solutions to these discrepancies.

Taking $\mathcal{B}(B \rightarrow \pi K)$ and $A_{CP}(B \rightarrow \pi K)$ as constraints, we present the allowed regions for ξ^{LL} , ξ^{LR} and ϕ_L in figure 4. Unfortunately, we find that the required region of minus ξ^{LR} with $\phi_L \sim -90^\circ$ is excluded by $A_{CP}(B^0 \rightarrow \pi^+ K^-)$, because it will induce a large negative $A_{CP}(B^0 \rightarrow \pi^+ K^-)$ as shown in figure 1 (b). In addition, as shown in figure 1 (b), the region of plus ξ^{LR} with $\phi_L \sim -90^\circ$ is helpless to resolve the “ πK puzzle”. The Z' effects are therefore still dominated by large ξ^{LL} , and the problem of $A_{CP}(B^- \rightarrow \rho^0 K^-)$ induced by ξ^{LL} still exist.

In fact, with ξ^{LL} and ξ^{LR} having the same sign, the corresponding Z' contributions counteract with each other in the $B \rightarrow \pi^0 K^-$ decay as shown in figures 1 (a) and (b). It is also easily understood from the expression for the effective coefficient $\alpha_{3,EW}^p(PP) = a_9^p(PP) - a_7^p(PP)$ [3], which involves the leading-order Z' contribution in this case. Thus, we conclude that any attempt to explain the $B \rightarrow \pi K$ anomaly in the non-universal Z' model with the assumption $\xi^{LL} = \xi^{LR} = \xi$, as made in ref. [45], is frangible and excluded in our case.

Parameters	Case I	Case II	Case III	Case IV
$\xi^{LL}(\times 10^{-3})$	3.96 ± 0.70	4.32 ± 0.75	1.52 ± 0.24	1.65 ± 0.35
$\xi^{LR}(\times 10^{-3})$	—	0.21 ± 0.15	-0.53 ± 0.13	-0.54 ± 0.15
$\zeta^{LL}(\times 10^{-3})$	—	—	-11.8 ± 3.1	-14.6 ± 7.1
$\zeta^{LR}(\times 10^{-3})$	—	—	—	1.04 ± 2.70
ϕ^L	$-88^\circ \pm 7^\circ$	$-88^\circ \pm 7^\circ$	$-86^\circ \pm 14^\circ$	$-85^\circ \pm 16^\circ$

Table 5. The numerical results for the parameters $\xi^{LL,LR}$, $\zeta^{LL,LR}$ and ϕ_L in the four different cases. The dashes mean that the corresponding parameters are neglected in each case.




Figure 5. The allowed regions for the parameters $\xi^{LL,LR}$, ζ^{LL} , and ϕ_L in Case III.

In a word, although the Z' contributions with a positive ξ^{LL} or a negative ξ^{LR} and $\phi_L \sim -90^\circ$ are helpful to bridge the discrepancy of $A_{CP}(B^- \rightarrow \pi^0 K^-)$, they would induce the unmatched $A_{CP}(B^- \rightarrow \rho^0 K^-)$ and $A_{CP}(B^0 \rightarrow \pi^+ K^-)$, respectively. Thus, with both $\mathcal{B}(B \rightarrow \pi K)$ and $A_{CP}(B \rightarrow \pi K, \rho K)$ as constraints, our results indicate that all of the parameter spaces in Case I and Case II are excluded with the assumption $B_{uu}^{L,R} \simeq -2B_{dd}^{L,R}$. As an alternative, in the following, we proceed to pursue possible solutions to these observations by considering the Z' contributions to the QCD penguins $\Delta C_{3,5}$.




Figure 6. The allowed regions for the parameters $\xi^{LL,LR}$, $\zeta^{LL,LR}$ and ϕ_L in Case IV.

Case III: Taking $B_{uu}^R \simeq -2B_{dd}^R$ (i.e., $\zeta^{LR} \simeq 0$), and leaving ζ^{LL} and $\xi^{LL,LR}$ arbitrary. As shown in figure 1 (c), we find that the variation trends of $A_{CP}(B^0 \rightarrow \pi^+ K^-)$ and $A_{CP}(B^- \rightarrow \pi^0 K^-)$ are always the same, indicating that the Z' contributions in this case could not give a solution to the observed “ πK puzzle” directly, as well as the unmatched $A_{CP}(B^- \rightarrow \rho^0 K^-)$ induced by ξ^{LL} . However, it is interesting to note that, with $\phi_L \sim -90^\circ$, both $A_{CP}(B^0 \rightarrow \pi^+ K^-)$ and $A_{CP}(B^- \rightarrow \pi^0 K^-)$ could be enhanced simultaneously, which may relax the constraints on ξ^{LR} . As mentioned in Case II, a negative ξ^{LR} is favored by the “ πK puzzle” and can moderate the problem of $A_{CP}(B^- \rightarrow \rho^0 K^-)$ induced by ξ^{LL} . So, the parameter ζ^{LL} may play an important role.

With $\mathcal{B}(B \rightarrow \pi K)$, $A_{CP}(B \rightarrow \pi K)$ and $A_{CP}(B \rightarrow \rho K)$ as constraints, the allowed regions for ξ^{LL} , ξ^{LR} , ζ^{LL} and ϕ_L are shown in figures 5. We find that none of ξ^{LL} , ξ^{LR} and ζ^{LL} could be neglected. Especially, the ζ^{LL} part moderates the contradictions caused by ξ^{LL} and ξ^{LR} . Furthermore, it is interesting to note that our predictions for $\mathcal{B}(B \rightarrow \pi K^*, \rho K)$, $A_{CP}(B \rightarrow \pi K^*)$ and $A_{CP}^{\text{mix}}(B^0 \rightarrow \pi^0 K_S, \rho^0 K_S)$, listed in tables 2, 3 and 4, respectively, are all consistent with the experimental data within 2σ .

Case IV: Without any simplification of $B_{uu}^{L,R}$ and $B_{dd}^{L,R}$, i.e., arbitrary values of $\zeta^{LL,LR}$ and $\xi^{LL,LR}$ are allowed. More generally, we give up any assumptions of the couplings $B_{uu}^{L,R}$ and $B_{dd}^{L,R}$. Then, there are five arbitrary NP parameters. As in Case III,

we take $\mathcal{B}(B \rightarrow \pi K)$, $A_{CP}(B \rightarrow \pi K)$ and $A_{CP}(B \rightarrow \rho K)$ as constraints and present the predictions for the other observables.

The allowed regions for $\xi^{LL,LR}$, $\zeta^{LL,LR}$ and ϕ_L are shown in figure 6, while the numerical results are listed in the last column of table 5. We find that, similar to Case III, the values of $\xi^{LL,LR}$ are definitely nonzero. The values of ζ^{LL} is a little larger than the one in Case III, due to the interference effect caused by the parameter ζ^{LR} . Our predictions for $\mathcal{B}(B \rightarrow \pi K^*, \rho K)$, $A_{CP}(B \rightarrow \pi K^*)$ and $A_{CP}^{\text{mix}}(B^0 \rightarrow \pi^0 K_S, \rho^0 K_S)$, listed in tables 2, 3 and 4, respectively, are consistent with the experimental data within 2σ .

4 Conclusions

Motivated by the recent observed large difference ΔA between $\mathcal{A}_{CP}(B^\mp \rightarrow \pi^0 K^\mp)$ and $A_{CP}(B^0 \rightarrow K^\pm \pi^\mp)$, we have investigated the effect of family non-universal Z' model and pursued possible solutions to the observed “ πK puzzle”. Moreover, we have also taken into account the constraints from the $B \rightarrow \pi K^*$, ρK decays, which also involve the same quark level $b \rightarrow s\bar{q}q$ ($q = u, d$) transitions. Our main conclusions are summarized as:

- The Z' contributions to the coefficients of operators O_7 and O_9 (ξ^{LL} and ξ^{LR}) with $\phi_L \sim -86^\circ$ are crucial to bridge the discrepancy of $A_{CP}(B^- \rightarrow \pi^0 K^-)$ between theoretical prediction and experimental data. However, they are definitely unequal and opposite in sign.
- The Z' contributions to the coefficients of QCD penguins operator O_3 related to ζ_{LL} are required to moderate the contradiction of $A_{CP}(B^- \rightarrow \rho^0 K^-)$ and $A_{CP}(B^0 \rightarrow \pi^+ K^-)$ to their experimental values induced by ξ^{LL} and ξ^{LR} , respectively, even though they are helpless to resolve the observed “ πK puzzle”. On the other hand, the Z' contributions to $C_5(\zeta^{LR})$ are inessential.
- For all of the four cases, a new weak phase associated with the chiral Z' couplings, with a value about -86° , is always required for the “ πK puzzle”.

Combing the up-to-date experimental measurements of $B \rightarrow \pi K$, πK^* and ρK decays, the family non-universal Z' model is found to be helpful to resolve the observed “ πK puzzle”. It is also reminded that more refined measurements of the mix-induced CP asymmetries in the $B^0 \rightarrow \pi^0 K_S$ and $\rho^0 K_S$ decays are required to confirm or refute the NP signals. In the following years, the precision of measurements for these observables is expected to be much improved, which will then shrink and reveal the Z' parameter spaces.

Note added: When the paper is finished, we are aware of the interesting paper by Barger et al. [56]. Although our topics are very similar, we have taken into account of not only the CP asymmetries but also the branching ratios of the correlated decay modes to constrain Z' couplings. Moreover, our approaches for the hadronic dynamics are different.

Acknowledgments

The work is supported by National Science Foundation under contract Nos.10675039 and 10735080. X. Q. Li acknowledges support from the Alexander-von-Humboldt Foundation.

A Decay amplitudes in the SM with QCDF

The decay amplitudes for $B \rightarrow \pi K$ decays are recapitulated from ref. [3]

$$\mathcal{A}_{B^- \rightarrow \pi^- \bar{K}}^{\text{SM}} = \sum_{p=u,c} V_{pb} V_{ps}^* A_{\pi \bar{K}} \left[\delta_{pu} \beta_2 + \alpha_4^p - \frac{1}{2} \alpha_{4,\text{EW}}^p + \beta_3^p + \beta_{3,\text{EW}}^p \right], \quad (\text{A.1})$$

$$\begin{aligned} \sqrt{2} \mathcal{A}_{B^- \rightarrow \pi^0 K^-}^{\text{SM}} &= \sum_{p=u,c} V_{pb} V_{ps}^* \left\{ A_{\pi^0 K^-} \left[\delta_{pu} (\alpha_1 + \beta_2) + \alpha_4^p + \alpha_{4,\text{EW}}^p + \beta_3^p + \beta_{3,\text{EW}}^p \right] \right. \\ &\quad \left. + A_{K^- \pi^0} \left[\delta_{pu} \alpha_2 + \frac{3}{2} \alpha_{3,\text{EW}}^p \right] \right\}, \end{aligned} \quad (\text{A.2})$$

$$\mathcal{A}_{\bar{B}^0 \rightarrow \pi^+ K^-}^{\text{SM}} = \sum_{p=u,c} V_{pb} V_{ps}^* A_{\pi^+ K^-} \left[\delta_{pu} \alpha_1 + \alpha_4^p + \alpha_{4,\text{EW}}^p + \beta_3^p - \frac{1}{2} \beta_{3,\text{EW}}^p \right], \quad (\text{A.3})$$

$$\begin{aligned} \sqrt{2} \mathcal{A}_{\bar{B}^0 \rightarrow \pi^0 \bar{K}^0}^{\text{SM}} &= \sum_{p=u,c} V_{pb} V_{ps}^* \left\{ A_{\pi^0 \bar{K}^0} \left[-\alpha_4^p + \frac{1}{2} \alpha_{4,\text{EW}}^p - \beta_3^p + \frac{1}{2} \beta_{3,\text{EW}}^p \right] \right. \\ &\quad \left. + A_{\bar{K}^0 \pi^0} \left[\delta_{pu} \alpha_2 + \frac{3}{2} \alpha_{3,\text{EW}}^p \right] \right\}, \end{aligned} \quad (\text{A.4})$$

where the explicit expressions for the coefficients $\alpha_i^p \equiv \alpha_i^p(M_1 M_2)$ and $\beta_i^p \equiv \beta_i^p(M_1 M_2)$ can also be found in ref. [3]. Note that expressions of the hard-spectator terms H_i appearing in α_i^p and the weak annihilation ones appearing in β_i^p should be replaced by our recalculated ones listed in appendix B. The decay amplitudes of $B \rightarrow \pi K^*$ and $B \rightarrow \rho K$ decays could be obtained from the above results by replacing $(\pi K) \rightarrow (\pi K^*)$ and $(\pi K) \rightarrow (\rho K)$, respectively.

B The hard-spectator and annihilation corrections with the infrared finite gluon propagator

With the infrared finite gluon propagator to cure the end-point divergences, the hard-spectator corrections in $B \rightarrow PP$ and PV decays can be expressed as [8]

$$\begin{aligned} H_i(M_1 M_2) &= \frac{B_{M_1 M_2}}{A_{M_1 M_2}} \int_0^1 dx dy d\xi \frac{\alpha_s(q^2)}{\xi} \Phi_{B1}(\xi) \Phi_{M_2}(x) \\ &\quad \times \left[\frac{\Phi_{M_1}(y)}{\bar{x}(\bar{y} + \omega^2(q^2)/\xi)} + r_\chi^{M_1} \frac{\phi_{m_1}(y)}{x(\bar{y} + \omega^2(q^2)/\xi)} \right], \end{aligned} \quad (\text{B.1})$$

for the insertion of operators $Q_{i=1-4,9,10}$,

$$\begin{aligned} H_i(M_1 M_2) &= -\frac{B_{M_1 M_2}}{A_{M_1 M_2}} \int_0^1 dx dy d\xi \frac{\alpha_s(q^2)}{\xi} \Phi_{B1}(\xi) \Phi_{M_2}(x) \\ &\quad \times \left[\frac{\Phi_{M_1}(y)}{x(\bar{y} + \omega^2(q^2)/\xi)} + r_\chi^{M_1} \frac{\phi_{m_1}(y)}{\bar{x}(\bar{y} + \omega^2(q^2)/\xi)} \right], \end{aligned} \quad (\text{B.2})$$

for $Q_{i=5,7}$, and $H_i(M_1M_2) = 0$ for $Q_{i=6,8}$. When both M_1 and M_2 are pseudoscalars, the final building blocks for annihilation contributions can be expressed as [8]

$$A_1^i = \pi \int_0^1 dx dy \alpha_s(q^2) \left\{ \left[\frac{\bar{x}}{(\bar{x}y - \omega^2(q^2) + i\epsilon)(1 - x\bar{y})} + \frac{1}{(\bar{x}y - \omega^2(q^2) + i\epsilon)\bar{x}} \right] \Phi_{M_1}(y) \Phi_{M_2}(x) + \frac{2}{\bar{x}y - \omega^2(q^2) + i\epsilon} r_\chi^{M_1} r_\chi^{M_2} \phi_{m_1}(y) \phi_{m_2}(x) \right\}, \quad (\text{B.3})$$

$$A_2^i = \pi \int_0^1 dx dy \alpha_s(q^2) \left\{ \left[\frac{y}{(\bar{x}y - \omega^2(q^2) + i\epsilon)(1 - x\bar{y})} + \frac{1}{(\bar{x}y - \omega^2(q^2) + i\epsilon)y} \right] \Phi_{M_1}(y) \Phi_{M_2}(x) + \frac{2}{\bar{x}y - \omega^2(q^2) + i\epsilon} r_\chi^{M_1} r_\chi^{M_2} \phi_{m_1}(y) \phi_{m_2}(x) \right\}, \quad (\text{B.4})$$

$$A_3^i = \pi \int_0^1 dx dy \alpha_s(q^2) \left\{ \frac{2\bar{y}}{(\bar{x}y - \omega^2(q^2) + i\epsilon)(1 - x\bar{y})} r_\chi^{M_1} \phi_{m_1}(y) \Phi_{M_2}(x) - \frac{2x}{(\bar{x}y - \omega^2(q^2) + i\epsilon)(1 - x\bar{y})} r_\chi^{M_2} \phi_{m_2}(x) \Phi_{M_1}(y) \right\}, \quad (\text{B.5})$$

$$A_1^f = A_2^f = 0, \quad (\text{B.6})$$

$$A_3^f = \pi \int_0^1 dx dy \alpha_s(q^2) \left\{ \frac{2(1 + \bar{x})}{(\bar{x}y - \omega^2(q^2) + i\epsilon)\bar{x}} r_\chi^{M_1} \phi_{m_1}(y) \Phi_{M_2}(x) + \frac{2(1 + y)}{(\bar{x}y - \omega^2(q^2) + i\epsilon)y} r_\chi^{M_2} \phi_{m_2}(x) \Phi_{M_1}(y) \right\}. \quad (\text{B.7})$$

When M_1 is a vector meson and M_2 a pseudoscalar, the sign of the second term in A_1^i , the first term in A_2^i , and the second terms in A_3^i and A_3^f need to be changed. When M_2 is a vector meson and M_1 a pseudoscalar, one only has to change the overall sign of A_2^i .

C Theoretical input parameters

C.1 CKM matrix elements

For the CKM matrix elements, we adopt the Wolfenstein parameterization [46] and choose the four parameters A , λ , ρ and η as [47]

$$A = 0.798_{-0.017}^{+0.023}, \quad \lambda = 0.22521_{-0.00082}^{+0.00083}, \quad \bar{\rho} = 0.141_{-0.021}^{+0.035}, \quad \bar{\eta} = 0.340 \pm 0.016, \quad (\text{C.1})$$

with $\bar{\rho} = \rho(1 - \frac{\lambda^2}{2})$ and $\bar{\eta} = \eta(1 - \frac{\lambda^2}{2})$.

C.2 Quark masses and lifetimes

As for the quark masses, there are two different classes appearing in our calculation. One type is the current quark mass which appears in the factor r_χ^M through the equation of

motion for quarks. This type of quark masses is scale dependent and denoted by \overline{m}_q . Here we take

$$\begin{aligned}\overline{m}_s(\mu)/\overline{m}_q(\mu) &= 27.4 \pm 0.4 \quad [48], \quad \overline{m}_s(2 \text{ GeV}) = 87 \pm 6 \text{ MeV} \quad [48], \\ \overline{m}_b(\overline{m}_b) &= 4.20^{+0.17}_{-0.07} \text{ GeV} \quad [49],\end{aligned}\tag{C.2}$$

where $\overline{m}_q(\mu) = (\overline{m}_u + \overline{m}_d)(\mu)/2$, and the difference between u and d quark is not distinguished.

The other one is the pole quark mass appearing in the evaluation of penguin loop corrections, and denoted by m_q . In this paper, we take [49]

$$m_u = m_d = m_s = 0, \quad m_c = 1.61^{+0.08}_{-0.12} \text{ GeV}, \quad m_b = 4.79^{+0.19}_{-0.08} \text{ GeV}.\tag{C.3}$$

As for the B-meson lifetimes, we take [49] $\tau_{B_u} = 1.638 \text{ ps}$ and $\tau_{B_d} = 1.530 \text{ ps}$, respectively.

C.3 The decay constants and form factors

In this paper, we take the heavy-to-light transition form factors [51]

$$\begin{aligned}F_0^{B \rightarrow \pi}(0) &= 0.258 \pm 0.031, & F_0^{B \rightarrow K}(0) &= 0.331 \pm 0.041, & V^{B \rightarrow K^*}(0) &= 0.411 \pm 0.033, \\ A_0^{B \rightarrow K^*}(0) &= 0.374 \pm 0.034, & A_1^{B \rightarrow K^*}(0) &= 0.292 \pm 0.028, & V^{B \rightarrow \rho}(0) &= 0.323 \pm 0.030, \\ A_0^{B \rightarrow \rho}(0) &= 0.303 \pm 0.029, & A_1^{B \rightarrow \rho}(0) &= 0.242 \pm 0.023.\end{aligned}\tag{C.4}$$

and the decay constants

$$\begin{aligned}f_B &= (216 \pm 22) \text{ MeV} \quad [50], & f_\pi &= (130.4 \pm 0.2) \text{ MeV} \quad [49], \\ f_K &= (155.5 \pm 0.8) \text{ MeV} \quad [49], & f_{K^*} &= (217 \pm 5) \text{ MeV} \quad [51], \\ f_\rho &= (209 \pm 2) \text{ MeV} \quad [51].\end{aligned}\tag{C.5}$$

C.4 The LCDAs of mesons and light-cone projector operators.

The light-cone projector operators of light mesons in momentum space read [3, 52]

$$\begin{aligned}M_{\alpha\beta}^P &= \frac{if_P}{4} \left[\not{p} \gamma_5 \Phi_P(x) - \mu_P \gamma_5 \frac{k_2 k_1}{k_2 \cdot k_1} \Phi_P(x) \right]_{\alpha\beta}, \\ \left(M_{\parallel}^V \right)_{\alpha\beta} &= -\frac{if_V}{4} \left[\not{p} \Phi_V(x) - \frac{m_V f_V^\perp}{f_V} \frac{k_2 k_1}{k_2 \cdot k_1} \Phi_V(x) \right]_{\alpha\beta},\end{aligned}\tag{C.6}$$

where $f_{P,V}$ are the decay constants, and $\mu_P = m_b r_\chi^P / 2$, with the chirally-enhanced factor r_χ^P defined as

$$r_\chi^\pi(\mu) = \frac{2m_\pi^2}{m_b(\mu)2m_q(\mu)}, \quad r_\chi^K(\mu) = \frac{2m_K^2}{m_b(\mu)(m_q + m_s)(\mu)},\tag{C.7}$$

where the quark masses are all running masses defined in the $\overline{\text{MS}}$ scheme. For the LCDAs of mesons, we use their asymptotic forms [53, 54]

$$\Phi_{P,V}(x) = 6x(1-x), \quad \phi_p(x) = 1, \quad \phi_v(x) = 3(2x-1).\tag{C.8}$$

As for the B-meson wave function, we take the form [55]

$$\Phi_B(\xi) = N_B \xi(1 - \xi) \exp \left[- \left(\frac{M_B}{M_B - m_b} \right)^2 (\xi - \xi_B)^2 \right], \quad (\text{C.9})$$

where $\xi_B \equiv 1 - m_b/M_B$, and N_B is the normalization constant to insure that $\int_0^1 d\xi \Phi_B(\xi) = 1$.

References

- [1] A.J. Buras, R. Fleischer, S. Recksiegel and F. Schwab, *B* → ππ, *new physics in B* → π*K* and implications for rare *K* and *B* decays, *Phys. Rev. Lett.* **92** (2004) 101804 [[hep-ph/0312259](#)] [[SPIRES](#)]; *Anatomy of prominent B and K decays and signatures of CP-violating new physics in the electroweak penguin sector*, *Nucl. Phys.* **B 697** (2004) 133 [[hep-ph/0402112](#)] [[SPIRES](#)].
- [2] M. Beneke, G. Buchalla, M. Neubert and C.T. Sachrajda, *QCD factorization in B* → π*K*, ππ decays and extraction of Wolfenstein parameters, *Nucl. Phys.* **B 606** (2001) 245 [[hep-ph/0104110](#)] [[SPIRES](#)];
M. Beneke and M. Neubert, *Flavor-singlet B decay amplitudes in QCD factorization*, *Nucl. Phys.* **B 651** (2003) 225 [[hep-ph/0210085](#)] [[SPIRES](#)].
- [3] M. Beneke and M. Neubert, *QCD factorization for B* → *PP* and *B* → *PV* decays, *Nucl. Phys.* **B 675** (2003) 333 [[hep-ph/0308039](#)] [[SPIRES](#)].
- [4] T. Muta, A. Sugamoto, M.-Z. Yang and Y.-D. Yang, *B* → ππ, *K*π decays in QCD improved factorization approach, *Phys. Rev.* **D 62** (2000) 094020 [[hep-ph/0006022](#)] [[SPIRES](#)];
M.-Z. Yang and Y.-D. Yang, *B* → *PV* decays in QCD improved factorization approach, *Phys. Rev.* **D 62** (2000) 114019 [[hep-ph/0007038](#)] [[SPIRES](#)].
- [5] H.-N. Li, S. Mishima and A.I. Sanda, *Resolution to the B* → π*K* puzzle, *Phys. Rev.* **D 72** (2005) 114005 [[hep-ph/0508041](#)] [[SPIRES](#)].
- [6] A.R. Williamson and J. Zupan, *Two body B decays with isosinglet final states in SCET*, *Phys. Rev.* **D 74** (2006) 014003 [*Erratum ibid.* **D 74** (03901) 2006] [[hep-ph/0601214](#)] [[SPIRES](#)].
- [7] M. Gronau and J.L. Rosner, *Rate and CP-asymmetry sum rules in B* → *K*π, *Phys. Rev.* **D 74** (2006) 057503 [[hep-ph/0608040](#)] [[SPIRES](#)];
X.-q. Li and Y.-d. Yang, *Reexamining charmless B* → *PV* decays in QCD factorization approach, *Phys. Rev.* **D 73** (2006) 114027 [[hep-ph/0602224](#)] [[SPIRES](#)]; *Revisiting B* → ππ, π*K* decays in QCD factorization approach, *Phys. Rev.* **D 72** (2005) 074007 [[hep-ph/0508079](#)] [[SPIRES](#)];
J. Chay, H.-N. Li and S. Mishima, *Possible complex annihilation and B* → *K*π direct CP asymmetry, *Phys. Rev.* **D 78** (2008) 034037 [[arXiv:0711.2953](#)] [[SPIRES](#)];
C.W. Bauer, I.Z. Rothstein and I.W. Stewart, *SCET analysis of B* → *K*π, *B* → *K* \bar{K} and *B* → ππ decays, *Phys. Rev.* **D 74** (2006) 034010 [[hep-ph/0510241](#)] [[SPIRES](#)];
C.S. Kim, S. Oh and C. Yu, *A critical study of the B* → *K*π puzzle, *Phys. Rev.* **D 72** (2005) 074005 [[hep-ph/0505060](#)] [[SPIRES](#)].
- [8] Q. Chang, X.-Q. Li and Y.-D. Yang, *Revisiting B* → π*K*, π*K** and ρ*K* decays: direct CP-violation and implication for New Physics, *JHEP* **09** (2008) 038 [[arXiv:0807.4295](#)] [[SPIRES](#)].

- [9] V. Barger, C.-W. Chiang, P. Langacker and H.-S. Lee, *Solution to the $B \rightarrow \pi K$ puzzle in a flavor-changing Z' model*, *Phys. Lett. B* **598** (2004) 218 [[hep-ph/0406126](#)] [[SPIRES](#)].
- [10] W.-S. Hou, M. Nagashima and A. Soddu, *Difference in B^+ and B^0 direct CP asymmetry as effect of a fourth generation*, *Phys. Rev. Lett.* **95** (2005) 141601 [[hep-ph/0503072](#)] [[SPIRES](#)];
 S. Khalil, *CP asymmetries and branching ratios of $B \rightarrow K\pi$ in supersymmetric models*, *Phys. Rev. D* **72** (2005) 035007 [[hep-ph/0505151](#)] [[SPIRES](#)];
 R. Arnowitt et al., *The $B \rightarrow K\pi$ puzzle and supersymmetric models*, *Phys. Lett. B* **633** (2006) 748 [[hep-ph/0509233](#)] [[SPIRES](#)];
 Y.D. Yang, R.M. Wang and G.R. Lu, *The puzzles in $B \rightarrow \pi\pi$ and πK decays: Possible implications for R-parity violating supersymmetry*, *Phys. Rev. D* **73** (2006) 015003 [[hep-ph/0509273](#)] [[SPIRES](#)];
 S. Baek, *New physics in $B \rightarrow \pi\pi$ and $B \rightarrow \pi K$ decays*, *JHEP* **07** (2006) 025 [[hep-ph/0605094](#)] [[SPIRES](#)];
 W.S.Hou, M. Nagashima, G. Raz and A. Soddu, *Four generation CP violation in $B \rightarrow \phi K^0, \pi^0 K^0, \eta - \text{prime} K^0$ and hadronic uncertainties*, *JHEP* **09** (2006) 012 [[hep-ph/0603097](#)] [[SPIRES](#)];
 C.S. Kim, S. Oh, C. Sharma, R. Sinha and Y.W. Yoon, *Generalized analysis on $B \rightarrow K^* \rho$ within and beyond the Standard Model: Can it help understand the $B \rightarrow K\pi$ puzzle?*, *Phys. Rev. D* **76** (2007) 074019 [[arXiv:0706.1150](#)] [[SPIRES](#)];
 C.S. Kim, S. Oh and Y.W. Yoon, *Analytic resolution of puzzle in $B \rightarrow K\pi$ decays*, *Phys. Lett. B* **665** (2008) 231 [[arXiv:0707.2967](#)] [[SPIRES](#)];
 M. Ciuchini, E. Franco, G. Martinelli, M. Pierini and L. Silvestrini, *Searching for New Physics with $B \rightarrow K\pi$ decays*, *Phys. Lett. B* **674** (2009) 197 [[arXiv:0811.0341](#)] [[SPIRES](#)];
 N. Mahajan, *CP asymmetry in $B^+ \rightarrow K^+ \pi^0$ and New Physics*, [[arXiv:0812.0230](#)] [[SPIRES](#)].
- [11] BABAR collaboration, B. Aubert et al., *Observation of CP violation in $B^0 \rightarrow K^+ \pi^-$ and $B^0 \rightarrow \pi^+ \pi^-$* , *Phys. Rev. Lett.* **99** (2007) 021603 [[hep-ex/0703016](#)] [[SPIRES](#)].
- [12] BELLE collaboration, S.W. Lin et al., *Difference in direct charge-parity violation between charged and neutral B meson decays*, *Nature* **452** (2008) 332.
- [13] CLEO collaboration, S. Chen et al., *Measurement of charge asymmetries in charmless hadronic in b meson decays*, *Phys. Rev. Lett.* **85** (2000) 525 [[hep-ex/0001009](#)] [[SPIRES](#)].
- [14] CDF collaboration, M. Morello, *Branching fractions and direct CP asymmetries of charmless decay modes at the Tevatron*, *Nucl. Phys. Proc. Suppl.* **170** (2007) 39 [[hep-ex/0612018](#)] [[SPIRES](#)].
- [15] HEAVY FLAVOR AVERAGING GROUP, E. Barberio et al., *Averages of b-hadron properties at the end of 2006*, [[arXiv:0704.3575](#)] [[SPIRES](#)] and online update at <http://www.slac.stanford.edu/xorg/hfag>.
- [16] M. Beneke, G. Buchalla, M. Neubert and C.T. Sachrajda, *QCD factorization for $B \rightarrow \pi\pi$ decays: Strong phases and CP violation in the heavy quark limit*, *Phys. Rev. Lett.* **83** (1999) 1914 [[hep-ph/9905312](#)] [[SPIRES](#)]; *QCD factorization for exclusive, nonleptonic B meson decays: General arguments and the case of heavy light final states*, *Nucl. Phys. B* **591** (2000) 313 [[hep-ph/0006124](#)] [[SPIRES](#)].
- [17] J.M. Cornwall, *Dynamical mass generation in continuum QCD*, *Phys. Rev. D* **26** (1982) 1453 [[SPIRES](#)];
 J. Papavassiliou and J.M. Cornwall, *Coupled fermion gap and vertex equations for chiral symmetry breakdown in QCD*, *Phys. Rev. D* **44** (1991) 1285 [[SPIRES](#)].

- [18] A.C. Aguilar, A.A. Natale and P.S. Rodrigues da Silva, *Relating a gluon mass scale to an infrared fixed point in pure gauge QCD*, *Phys. Rev. Lett.* **90** (2003) 152001 [[hep-ph/0212105](#)] [[SPIRES](#)];
A.C. Aguilar and A.A. Natale, *A dynamical gluon mass solution in a coupled system of the Schwinger-Dyson equations*, *JHEP* **08** (2004) 057 [[hep-ph/0408254](#)] [[SPIRES](#)].
- [19] L. Von Smekal, A. Hauck and R. Alkofer, *The infrared behavior of gluon and ghost propagators in Landau gauge QCD*, *Phys. Rev. Lett.* **79** (1997) 3591 [[hep-ph/9705242](#)] [[SPIRES](#)];
R. Alkofer and L. Von Smekal, *The infrared behavior of QCD Green's functions - Confinement, dynamical symmetry breaking, and hadrons as relativistic bound states*, *Phys. Rep.* **353** (2001) 281 [[hep-ph/0007355](#)] [[SPIRES](#)];
C.S. Fisher and R. Alkofer, *Non-perturbative propagators, running coupling and dynamical quark mass of Landau gauge QCD*, *Phys. Rev. D* **67** (2003) 094020 [[hep-ph/0301094](#)] [[SPIRES](#)];
R. Alkofer, W. Detmold, C.S. Fischer and P. Maris, *Analytic structure of the gluon and quark propagators in Landau gauge QCD*, *Nucl. Phys. Proc. Suppl.* **141** (2005) 122 [[hep-ph/0309078](#)] [[SPIRES](#)].
- [20] I.L. Bogolubsky, E.M. Ilgenfritz, M. Muller-Preussker and A. Sternbeck, *The Landau gauge gluon and ghost propagators in 4DSU(3) gluodynamics in large lattice volumes*, *PoS(LATTICE 2007)290* [[arXiv:0710.1968](#)] [[SPIRES](#)];
A. Cucchieri and T. Mendes, *What's up with IR gluon and ghost propagators in Landau gauge? A puzzling answer from huge lattices*, *PoS(LATTICE 2007)297* [[arXiv:0710.0412](#)] [[SPIRES](#)];
P.O. Bowman et al., *Scaling behavior and positivity violation of the gluon propagator in full QCD*, *Phys. Rev. D* **76** (2007) 094505 [[hep-lat/0703022](#)] [[SPIRES](#)].
- [21] Y.-Y. Keum, H.-N. Li and A.I. Sanda, *Fat penguins and imaginary penguins in perturbative QCD*, *Phys. Lett. B* **504** (2001) 6 [[hep-ph/0004004](#)] [[SPIRES](#)]; *Penguin enhancement and $B \rightarrow K\pi$ decays in perturbative QCD*, *Phys. Rev. D* **63** (2001) 054008 [[hep-ph/0004173](#)] [[SPIRES](#)].
- [22] C.W. Bauer, S. Fleming and M.E. Luke, *Summing Sudakov logarithms in $B \rightarrow X_s \gamma$ in effective field theory*, *Phys. Rev. D* **63** (2000) 014006 [[hep-ph/0005275](#)] [[SPIRES](#)];
C.W. Bauer, S. Fleming, D. Pirjol and I.W. Stewart, *An effective field theory for collinear and soft gluons: heavy to light decays*, *Phys. Rev. D* **63** (2001) 114020 [[hep-ph/0011336](#)] [[SPIRES](#)];
C.W. Bauer and I.W. Stewart, *Invariant operators in collinear effective theory*, *Phys. Lett. B* **516** (2001) 134 [[hep-ph/0107001](#)] [[SPIRES](#)];
C.W. Bauer, D. Pirjol and I.W. Stewart, *Soft-collinear factorization in effective field theory*, *Phys. Rev. D* **65** (2002) 054022 [[hep-ph/0109045](#)] [[SPIRES](#)].
- [23] M.E. Peskin, *Particle physics: song of the electroweak penguin*, *Nature* **452** (2008) 293 [[SPIRES](#)].
- [24] T. Feldmann, M. Jung and T. Mannel, *Is there a non-Standard-Model contribution in non-leptonic $b \rightarrow s$ decays?*, *JHEP* **08** (2008) 066 [[arXiv:0803.3729](#)] [[SPIRES](#)].
- [25] M.S. Carena, A. Daleo, B.A. Dobrescu and T.M.P. Tait, *Z' gauge bosons at the Tevatron*, *Phys. Rev. D* **70** (2004) 093009 [[hep-ph/0408098](#)] [[SPIRES](#)].
- [26] T.G. Rizzo, *Z' phenomenology and the LHC*, [hep-ph/0610104](#) [[SPIRES](#)]; *Z' physics at the LHC and the LHeC*, [arXiv:0808.1906](#) [[SPIRES](#)].

- [27] G. Buchalla, G. Burdman, C.T. Hill and D. Kominis, *GIM violation and new dynamics of the third generation*, *Phys. Rev. D* **53** (1996) 5185 [[hep-ph/9510376](#)] [[SPIRES](#)];
 G. Burdman, K.D. Lane and T. Rador, *\bar{B} B mixing constrains topcolor-assisted technicolor*, *Phys. Lett. B* **514** (2001) 41 [[hep-ph/0012073](#)] [[SPIRES](#)];
 A. Martin and K. Lane, *CP violation and flavor mixing in technicolor models*, *Phys. Rev. D* **71** (2005) 015011 [[hep-ph/0404107](#)] [[SPIRES](#)].
- [28] E. Nardi, *Z' , new fermions and flavor changing processes. Constraints on E_6 models from $\mu \rightarrow eee$* , *Phys. Rev. D* **48** (1993) 1240 [[hep-ph/9209223](#)] [[SPIRES](#)];
 J. Bernabeu, E. Nardi and D. Tommasini, *$\mu - e$ conversion in nuclei and Z' physics*, *Nucl. Phys. B* **409** (1993) 69 [[hep-ph/9306251](#)] [[SPIRES](#)];
 V.D. Barger, M.S. Berger and R.J.N. Phillips, *Quark singlets: implications and constraints*, *Phys. Rev. D* **52** (1995) 1663 [[hep-ph/9503204](#)] [[SPIRES](#)];
 M.B. Popovic and E.H. Simmons, *Weak singlet fermions: models and constraints*, *Phys. Rev. D* **62** (2000) 035002 [[hep-ph/0001302](#)] [[SPIRES](#)];
 T.G. Rizzo, *Gauge kinetic mixing and leptophobic Z' in E_6 and $SO(10)$* , *Phys. Rev. D* **59** (1999) 015020 [[hep-ph/9806397](#)] [[SPIRES](#)].
- [29] P. Langacker and M. Plümacher, *Flavor changing effects in theories with a heavy Z' boson with family nonuniversal couplings*, *Phys. Rev. D* **62** (2000) 013006 [[hep-ph/0001204](#)] [[SPIRES](#)].
- [30] N. Cabibbo, *Unitary symmetry and leptonic decays*, *Phys. Rev. Lett.* **10** (1963) 531 [[SPIRES](#)];
 M. Kobayashi and T. Maskawa, *CP violation in the renormalizable theory of weak interaction*, *Prog. Theor. Phys.* **49** (1973) 652 [[SPIRES](#)].
- [31] G. Buchalla, A.J. Buras and M.E. Lautenbacher, *Weak decays beyond leading logarithms*, *Rev. Mod. Phys.* **68** (1996) 1125 [[hep-ph/9512380](#)] [[SPIRES](#)].
- [32] D.-s. Du, J.-f. Sun, D.-s. Yang and G.-h. Zhu, *Charmless two-body B decays: a global analysis with QCD factorization*, *Phys. Rev. D* **67** (2003) 014023 [[hep-ph/0209233](#)] [[SPIRES](#)];
 D.-s. Du, H.-J. Gong, J.-f. Sun, D.-s. Yang and G.-h. Zhu, *Phenomenological analysis of charmless decays $B \rightarrow PV$ with QCD factorization*, *Phys. Rev. D* **65** (2002) 094025 [[Erratum ibid. D66](#) (2002) 079904] [[hep-ph/0201253](#)] [[SPIRES](#)];
 D.-s. Du, D.-s. Yang and G.-h. Zhu, *QCD factorization for $B \rightarrow PP$* , *Phys. Rev. D* **64** (2001) 014036 [[hep-ph/0103211](#)] [[SPIRES](#)]; *Infrared divergence and twist-3 distribution amplitudes in QCD factorization for $B \rightarrow PP$* , *Phys. Lett. B* **509** (2001) 263 [[hep-ph/0102077](#)] [[SPIRES](#)]; *Analysis of the decays $B \rightarrow \pi\pi$ and πK in QCD factorization in the heavy quark limit*, *Phys. Lett. B* **488** (2000) 46 [[hep-ph/0005006](#)] [[SPIRES](#)].
- [33] M. Beneke, *Corrections to $\sin(2\beta)$ from CP asymmetries in $B^0 \rightarrow (\pi^0, \rho^0, \eta, \eta', \omega, \Phi)K_S$ decays*, *Phys. Lett. B* **620** (2005) 143 [[hep-ph/0505075](#)] [[SPIRES](#)].
- [34] G. Buchalla, G. Hiller, Y. Nir and G. Raz, *The pattern of CP asymmetries in $b \rightarrow s$ transitions*, *JHEP* **09** (2005) 074 [[hep-ph/0503151](#)] [[SPIRES](#)].
- [35] R. Fleischer, S. Jager, D. Pirjol and J. Zupan, *Benchmarks for the New-Physics search through CP-violation in $B^0 \rightarrow \pi^0 K_S$* , *Phys. Rev. D* **78** (2008) 111501 [[arXiv:0806.2900](#)] [[SPIRES](#)].
- [36] M. Gronau and J.L. Rosner, *Implications for CP asymmetries of improved data on $B \rightarrow K^0 \pi^0$* , *Phys. Lett. B* **666** (2008) 467 [[arXiv:0807.3080](#)] [[SPIRES](#)].

- [37] D. Zwanziger, *Non-perturbative Faddeev-Popov formula and infrared limit of QCD*, *Phys. Rev. D* **69** (2004) 016002 [[hep-ph/0303028](#)] [[SPIRES](#)];
 D.M. Howe and C.J. Maxwell, *All-orders infra-red freezing of $R_{e^+e^-}$ in perturbative QCD*, *Phys. Lett. B* **541** (2002) 129 [[hep-ph/0204036](#)] [[SPIRES](#)]; *All-orders infrared freezing of observables in perturbative QCD*, *Phys. Rev. D* **70** (2004) 014002 [[hep-ph/0303163](#)] [[SPIRES](#)];
 S. Furui and H. Nakajima, *Infrared features of lattice Landau gauge QCD and the Gribov copy problem*, *AIP Conf. Proc.* **717** (2004) 685 [[hep-lat/0309166](#)] [[SPIRES](#)].
- [38] S. Brodsky, S. Menke, C. Merino and J. Rathsmann, *On the behavior of the effective QCD coupling $\alpha_{\text{tau}}(s)$ at low scales*, *Phys. Rev. D* **67** (2003) 055008 [[hep-ph/0212078](#)] [[SPIRES](#)];
 A.C. Mattingly and P.M. Stevenson, *Optimization of $R_{e^+e^-}$ and "Freezing" of the QCD couplant at low energies*, *Phys. Rev. D* **49** (1994) 437 [[hep-ph/9307266](#)] [[SPIRES](#)];
 M. Baldicchi and G.M. Prosperi, *Infrared behavior of the running coupling constant and bound states in QCD*, *Phys. Rev. D* **66** (2002) 074008 [[hep-ph/0202172](#)] [[SPIRES](#)].
- [39] F. Su, Y.D. Yang, G.R. Lu and H.J. Hao, *Revisiting the annihilation decay $\bar{B}_s \rightarrow \pi^+\pi^-$* , *Eur. Phys. J. C* **44** (2005) 243 [[hep-ph/0507326](#)] [[SPIRES](#)];
 F. Su, Y.L. Wu, Y.D. Yang and C. Zhuang, *Large strong phases and CP violation in the annihilation processes $\bar{B}^0 \rightarrow K^+K^-$, $K^{*\pm}K^\mp$, $K^{*+}K^*$* , *Eur. Phys. J. C* **48** (2006) 401 [[hep-ph/0604082](#)] [[SPIRES](#)]; *QCD Approach to $B \rightarrow D\pi$ decays and CP violation*, *Commun. Theor. Phys.* **49** (2008) 716 [[arXiv:0705.1575](#)] [[SPIRES](#)].
- [40] A.A. Natale and C.M. Zanetti, *Non-leptonic annihilation decays of B mesons with a natural infrared cutoff*, [arXiv:0803.0154](#) [[SPIRES](#)].
- [41] A.J. Buras, P. Gambino, and U.A. Haisch, *Electroweak penguin contributions to non-Leptonic $\Delta F = 1$ decays at NNLO*, *Nucl. Phys. B* **570** (2000) 117 [[hep-ph/9911250](#)] [[SPIRES](#)].
- [42] V. Barger, C.W. Chiang, P. Langacker and H.S. Lee, *Z' mediated flavor changing neutral currents in B meson decays*, *Phys. Lett. B* **580** (2004) 186 [[hep-ph/0310073](#)] [[SPIRES](#)];
 V. Barger, C.W. Chiang, Jing Jiang and P. Langacker, *$B_s - \bar{B}_s$ mixing in Z' models with flavor-changing neutral currents*, *Phys. Lett. B* **596** (2004) 229 [[hep-ph/0405108](#)] [[SPIRES](#)].
- [43] K. Cheung, C.W. Chiang, N.G. Deshpande and J. Jiang, *Constraints on flavor-changing Z' models by B_s mixing, Z' production, and $B_s \rightarrow \mu^+\mu^-$* , *Phys. Lett. B* **652** (2007) 285 [[hep-ph/0604223](#)] [[SPIRES](#)];
 J.H. Jeon, C.S. Kim, J. Lee and C. Yu, *Exclusive $B \rightarrow M\nu\bar{\nu}$ ($M = \pi, K, \rho, K^*$) decays and leptophobic Z' model*, *Phys. Lett. B* **636** (2006) 270 [[hep-ph/0602156](#)] [[SPIRES](#)];
 C.H. Chen and H. Hatanaka, *Nonuniversal Z' couplings in B decays*, *Phys. Rev. D* **73** (2006) 075003 [[hep-ph/0602140](#)] [[SPIRES](#)].
- [44] DELPHI collaboration, P. Abreu et al., *A Study of radiative muon pair events at Z^0 energies and limits on an additional Z' gauge boson*, *Z. Phys. C* **65** (1995) 603 [[SPIRES](#)].
- [45] R. Mohanta and A.K. Giri, *Explaining $B \rightarrow K\pi$ anomaly with non-universal Z' boson*, [arXiv:0812.1842](#) [[SPIRES](#)].
- [46] L. Wolfenstein, *Parametrization of the Kobayashi-Maskawa matrix*, *Phys. Rev. Lett.* **51** (1983) 1945 [[SPIRES](#)].
- [47] CKMFITTER GROUP, J. Charles et al., *CP violation and the CKM matrix: Assessing the impact of the asymmetric B factories*, *Eur. Phys. J. C* **41** (2005) 1 [[hep-ph/0406184](#)] [[SPIRES](#)] updated results and plots available at: <http://ckmfitter.in2p3.fr>.

- [48] HPQCD Collaboration, Q. Mason et al., *High-precision determination of the light-quark masses from realistic lattice QCD*, *Phys. Rev. D* **73** (2006) 114501 [[hep-ph/0511160](#)] [[SPIRES](#)].
- [49] PARTICLE DATA GROUP, C. Amsler et al., *Review of particle physics*, *Phys. Lett. B* **667** (2008) 1 [[SPIRES](#)].
- [50] HPQCD collaboration, A. Gray et al., *The B meson decay constant from unquenched lattice QCD*, *Phys. Rev. Lett.* **95** (2005) 212001 [[hep-lat/0507015](#)] [[SPIRES](#)].
- [51] P. Ball and R. Zwicky, *New results on $B \rightarrow \pi, K, \eta$ decay formfactors from light-cone sum rules*, *Phys. Rev. D* **71** (2005) 014015 [[hep-ph/0406232](#)] [[SPIRES](#)]; *$B_{d,s} \rightarrow \rho, \omega, K^*, \phi$ decay form factors from light-cone sum rules revisited*, *Phys. Rev.* **71** (2005) 014029 [[hep-ph/0412079](#)] [[SPIRES](#)]; *SU(3) breaking of leading-twist K and K^* distribution amplitudes: A reprise*, *Phys. Lett. B* **633** (2006) 289 [[hep-ph/0510338](#)] [[SPIRES](#)].
- [52] B.V. Geshkenbein and M.V. Terentev, *The phenomenological formula for the pion electromagnetic form-factor*, *Yad. Fiz.* **40** (1984) 758 [*Sov. J. Nucl. Phys.* **40** (1984) 487] [[SPIRES](#)].
- [53] M. Beneke and Th. Feldmann, *Symmetry breaking corrections to heavy to light B meson form-factors at large recoil*, *Nucl. Phys. B* **592** (2001) 3 [[hep-ph/0008255](#)] [[SPIRES](#)].
- [54] A. Ali et al., *A comparative study of the decays $B \rightarrow (K, K^*) \ell^+ \ell^-$ in standard model and supersymmetric theories*, *Phys. Rev. D* **61** (2000) 074024 [[hep-ph/9910221](#)] [[SPIRES](#)]; P. Ball and V.M. Braun, *Exclusive semileptonic and rare B meson decays in QCD*, *Phys. Rev. D* **58** (1998) 094016 [[hep-ph/9805422](#)] [[SPIRES](#)]; P. Ball et al., *Higher twist distribution amplitudes of vector mesons in QCD: Formalism and twist - three distributions*, *Nucl. Phys. B* **529** (1998) 323 [[hep-ph/9802299](#)] [[SPIRES](#)].
- [55] G. Eilam, M. Ladisa and Y.D. Yang, *Study of $B^0 \rightarrow J/\Psi D^{(*)}$ and $\eta_c D^{(*)}$ in perturbative QCD*, *Phys. Rev. D* **65** (2002) 037504 [[hep-ph/0107043](#)] [[SPIRES](#)].
- [56] V. Barger et al., *Family non-universal $U(1)'$ gauge symmetries and $b \rightarrow s$ transitions*, [arXiv:0902.4507](#) [[SPIRES](#)].