

$H \rightarrow \gamma\gamma$ beyond the Standard Model

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

JHEP06(2009)054

(<http://iopscience.iop.org/1126-6708/2009/06/054>)

[The Table of Contents](#) and [more related content](#) is available

Download details:

IP Address: 80.92.225.132

The article was downloaded on 03/04/2010 at 09:14

Please note that [terms and conditions apply](#).

$H \rightarrow \gamma\gamma$ beyond the Standard Model

Giacomo Cacciapaglia, Aldo Deandrea and Jérémie Llodra-Perez

*Université de Lyon,
Caserne Sergent Blandan, 37, rue du Repos, 69365 Lyon Cedex 07, France*

*Université Lyon 1,
43 Boulevard du 11 Novembre 1918, 69622 Villeurbanne Cedex, France*

*CNRS/IN2P3, UMR5822, Institut de Physique Nucléaire de Lyon,
F-69622 Villeurbanne Cedex, France*

*E-mail: g.cacciapaglia@ipnl.in2p3.fr, a.deandrea@ipnl.in2p3.fr,
jllodra@ipnl.in2p3.fr*

ABSTRACT: We consider the $H \rightarrow \gamma\gamma$ decay process and the gluon fusion production of a light Higgs, and provide a general framework for testing models of new physics beyond the Standard Model. We apply our parametrisation to typical models extending the Standard Model in 4 and 5 dimensions, and show how the parametrisation can be used to discriminate between different scenarios of new physics at the Large Hadron Collider and at future Linear Colliders.

KEYWORDS: Higgs Physics, Beyond Standard Model

ARXIV EPRINT: [0901.0927](https://arxiv.org/abs/0901.0927)

Contents

1	Introduction	1
2	Definitions and notations	2
2.1	Observables at the LHC and linear colliders	5
3	Survey of models of new physics in 4 dimensions	6
3.1	A 4 th generation	6
3.2	Supersymmetry	7
3.3	Little Higgs models	9
3.4	Extended scalar sector: colour octet	13
3.5	Lee-Wick Standard Model	13
4	Survey of models of new physics in extra dimensions	16
4.1	Gauge bosons in a flat extra dimension	17
4.1.1	Gauge Higgs	18
4.1.2	Brane Higgs	19
4.1.3	Bulk Higgs	20
4.2	Gauge bosons in a warped extra dimension	20
4.2.1	Gauge Higgs	21
4.2.2	Brane Higgs	22
4.3	Bulk fermions in a flat extra dimension	23
4.3.1	Bulk fermions	23
4.3.2	Bulk fermion in UED models	25
4.3.3	Odd bulk masses: fermions in models of flavour	26
4.4	Fermions in a warped extra dimension	28
5	Numerical results	31
6	Conclusions	34
A	Higgs couplings in extended Higgs sectors	35
A.1	Multiple Higgs	35
A.2	Higgs mixing	36
A.3	Charged Higgs couplings	36
B	Gauge bosons in 5D	38
B.1	Gauge Higgs unification models	39
B.2	Brane Higgs models	41

C Fermionic fields	43
C.1 Gauge Higgs unification models	44
C.2 Brane Yukawas	44

1 Introduction

The decay of the Higgs in two photons is one of the most important discovery channels at the Large Hadron Collider (LHC), and it is certainly the golden mode at low masses, where the decay channels into heavy gauge bosons are closed. Detailed studies, including detector simulations, in the Standard Model (SM) and in its supersymmetric extensions are available [1]. This mode is also a powerful probe of the electroweak symmetry breaking sector of the theory, because it is a loop-induced process, therefore it is sensitive to any particle with a large coupling to the Higgs. In the SM it depends primarily on the couplings of the Higgs boson with heavy quarks (the top) and gauge bosons (the W), whose masses are tightly related to the electroweak scale. In any extension of the SM, particles that do couple strongly to the Higgs, and therefore play a role in the breaking of the electroweak symmetry, will also contribute to this loop and modify the SM prediction. For instance, new particles at the TeV scale are required to soften the divergences that appear in the corrections to the Higgs mass generated by top and W - Z loops. Many models in fact predict the existence of partners of the top and W : stops and gauginos in supersymmetry, heavy W 's and tops in extra dimensional models and Little Higgs models, and so on. Studying this channel will therefore give an indirect access to the mechanism underlying the electroweak symmetry breaking. At the LHC, we also need to take into account the Higgs production mechanism.

In the SM there are four main production mechanisms: gluon fusion ($gg \rightarrow H$), weak vector boson fusion, weak boson associated production (WH , ZH) and top associated production ($t\bar{t}H$). Gluon fusion dominates the inclusive production at LHC energies and it is roughly an order of magnitude larger than vector boson fusion and other processes.

While some of the production channels may have additional leptons, jets or missing energy in their final state, in the photon channel it will be difficult, at least at low luminosity, to take advantage of these different signatures. We shall therefore consider mainly the inclusive $H \rightarrow \gamma\gamma$ process. The interest of performing exclusive studies like the production via vector boson fusion, will be also discussed as it allows to better discriminate the kind of new physics that can be tested in the $H \rightarrow \gamma\gamma$ mode, especially when large integrated luminosity is available [2]. The main production process $gg \rightarrow H$ is a loop induced process like the decay $H \rightarrow \gamma\gamma$, and it is sensitive to the same particles and physics.

In this paper we study the photon channel with the purpose of performing a model independent analysis, allowing to determine the possibility and the limits for discriminating various scenarios of new physics. In the following we shall propose a model independent parametrisation of these loop processes in order to test the possibility of discrimination of various models of new physics. We shall provide a general and simple formalism to

easily calculate the contribution of the new heavy states given their spectrum. We will assume that the new physics only affects those two processes, and corrections to the other production and decay channels are ignored. In the SM it is well known that the contribution of heavy particles to $H \rightarrow \gamma\gamma$ and $H \rightarrow gg$ processes does not decouple for particle masses much larger than the Higgs boson one. The reason is that these SM masses are uniquely generated by the coupling to the Higgs boson and the mass dependence of their coupling cancels the mass dependence in the loop integral. In general extensions of the SM this is not necessarily the case, as the masses may receive other contributions. The effect on the decay can therefore be sensitive to the mass scale of the new physics. Studying this channel in detail can give some hints about the model of new physics, and this information will be complementary to the direct discovery of new states at the LHC. Finally, the precise determination of the Higgs branching ratios at future Linear Collider will be an even more powerful discrimination tool, even when the new particles are well beyond the direct production threshold at the Linear Collider.

In the next section we settle our notation and define our parametrisation of the loop induced processes $H \rightarrow \gamma\gamma$ and $H \rightarrow gg$. In the sections 3 and 4 we consider various scenarios of new physics in 4 and 5 dimensions, in section 5 we discuss numerical results in various models and how the parametrisation we propose can provide a hint to what kind of new physics can be deduced from data both at the LHC and at Linear Colliders. Finally we give our conclusions, and we leave details on the calculation to the appendices.

2 Definitions and notations

In order to establish our notations, we will briefly review the decay of the Higgs in photons and gluons (the decay width in gluons is directly related to the gluon-fusion production cross section at hadronic colliders). The decay widths can be written as:

$$\Gamma_{\gamma\gamma} = \frac{G_F \alpha^2 m_H^3}{128 \sqrt{2} \pi^3} \left| A_W(\tau_W) + \sum_{\text{fermions}} N_{c,f} Q_f^2 A_F(\tau_f) + \sum_{\text{NP}} N_{c, \text{NP}} Q_{\text{NP}}^2 A_{\text{NP}}(\tau_{\text{NP}}) \right|^2, \quad (2.1)$$

$$\Gamma_{gg} = \frac{G_F \alpha_s^2 m_H^3}{16 \sqrt{2} \pi^3} \left| \frac{1}{2} \sum_{\text{quarks}} A_F(\tau_f) + \sum_{\text{NP}} C(r_{\text{NP}}) A_{\text{NP}}(\tau_{\text{NP}}) \right|^2, \quad (2.2)$$

where $\tau_x = \frac{m_H^2}{4m_x^2}$, $N_{c,x}$ is the number of colour states in the colour representation (3 for quarks, 1 for leptons), the constant $C(r)$ is an SU(3) colour factor (defined as $\text{Tr}[t_r^a t_r^b] = C(r) \delta^{ab}$ where t_r^a are the SU(3) generators in the representation r ; it is equal to 1/2 for the quarks and 3 for an adjoint), Q_x is the electric charge of the particle in the loop, and the functions $A(\tau)$ depend on the spin and couplings to the Higgs of the particle running in the loop. Note that G_F here is a numerical normalisation of the widths, defined in terms of the SM Higgs VEV v_{SM} ($\sqrt{2}G_F = 1/v_{\text{SM}}^2$), and not the physical Fermi constant, which may receive corrections from the New Physics.

In the SM, all masses are proportional to the Higgs vacuum expectation value (VEV) v_{SM} , therefore the couplings to the Higgs can be written as

$$y_{h\bar{f}f}^{\text{SM}} = \frac{m_f}{v_{\text{SM}}} \quad \text{for fermions,} \quad (2.3)$$

$$g_{h\phi\phi}^{\text{SM}} = 2\frac{m_\phi^2}{v_{\text{SM}}} \quad \text{for bosons.} \quad (2.4)$$

Under this assumption, the amplitudes are given by (F stands for spin-1/2 fermions, W for vector bosons and S for scalar bosons) [3]

$$A_F(\tau) = \frac{2}{\tau^2} (\tau + (\tau - 1)f(\tau)) , \quad (2.5)$$

$$A_W(\tau) = -\frac{1}{\tau^2} (2\tau^2 + 3\tau + 3(2\tau - 1)f(\tau)) , \quad (2.6)$$

$$A_S(\tau) = -\frac{1}{\tau^2} (\tau - f(\tau)) ; \quad (2.7)$$

where

$$f(\tau) = \begin{cases} \arcsin^2 \sqrt{\tau} & \tau \leq 1 \\ -\frac{1}{4} \left[\log \frac{1+\sqrt{1-\tau^{-1}}}{1-\sqrt{1-\tau^{-1}}} - i\pi \right]^2 & \tau > 1 \end{cases} . \quad (2.8)$$

For our study we are particularly interested in the limit of such functions for large mass of the particle in the loop with respect to the Higgs mass, $\tau \ll 1$:

$$A_F(0) = \frac{4}{3}, \quad A_W(0) = -7, \quad A_S(0) = \frac{1}{3} . \quad (2.9)$$

Note that the particle in the loop does not decouple for large mass because the (SM) coupling to the Higgs is also proportional to the mass of the particle. As we are interested in Higgs masses below the W threshold and above the LEP limit (where the $\gamma\gamma$ signal is non negligible), the light Higgs approximation is useful for the top and the new physics. For the W , this approximation is not valid, and the function $A_W(\tau_W)$ ranges from -8 for $m_H = 115$ GeV to -9.7 for $m_H = 150$ GeV.

However, the mass of new particles in most models is not proportional to the Higgs VEV v , but receives only a small correction from the electroweak symmetry breaking. Therefore, the amplitude for new physics is given by the same formulae as above up to a factor taking into account the different coupling to the Higgs (which is in general not proportional to the mass). The coupling to the Higgs for a fermion (boson) can be written in general as

$$y_{h\bar{f}f} = \frac{\partial m_f(v)}{\partial v}, \quad g_{h\phi\phi} = \frac{\partial m_\phi^2(v)}{\partial v} . \quad (2.10)$$

Therefore we can write the A function for the new physics contribution for fermions (bosons) as

$$A_{\text{NP}}^F = \frac{y_{h\bar{f}f}^{\text{NP}}}{y_{h\bar{f}f}^{\text{SM}}} A_F \quad \text{for fermions,} \quad (2.11)$$

$$A_{\text{NP}}^{W,S} = \frac{g_{h\phi\phi}^{\text{NP}}}{g_{h\phi\phi}^{\text{SM}}} A_{W,S} \quad \text{for bosons;} \quad (2.12)$$

which can be written without loss of generality, as

$$A_{\text{NP}} = \frac{v_{\text{SM}}}{m_{\text{NP}}} \frac{\partial m_{\text{NP}}}{\partial v} A_{F,W,S}. \quad (2.13)$$

As the mass can be a generic function of v , this formula allows to treat a wide range of physical situations beyond the Standard Model, as long as the particle mass is at least partially generated by the Higgs VEV (those formulae are valid for a SM Higgs sector; when the Higgs sector is extended, and for scalars which do mix with the Higgs doublet, more general formulae apply: see appendix A for details). When the mass of the new physics is not proportional to the Higgs VEV, A_{NP} will decouple for large masses. Examples of such cases will be discussed in detail in sections 3, 4. Note also that in general $v \neq v_{\text{SM}}$, however, as it will be clear in the following, this difference only introduces higher order corrections in an expansion for large new physics scale.

The new physics can be parametrised by two independent parameters describing the contribution of the new particles to the two decay widths, however using the actual amplitude is not a convenient way of treating the new contributions. Here we propose to normalise the new contribution to the top one. The main reason is that the top gives the main contribution to the amplitudes in the SM, and any new physics, which addresses the problem of the Higgs mass naturalness, will have a tight relation with the top. Moreover, as it will soon be clear, those two parameters can give some intuitive information about what kind of new physics runs into the loop. The widths can be rewritten as

$$\Gamma_{\gamma\gamma} = \frac{G_F \alpha^2 m_H^3}{128 \sqrt{2} \pi^3} \left| A_W(\tau_W) + 3 \left(\frac{2}{3} \right)^2 A_t(\tau_t) [1 + \kappa_{\gamma\gamma}] + \dots \right|^2, \quad (2.14)$$

$$\Gamma_{gg} = \frac{G_F \alpha_s^2 m_H^3}{16 \sqrt{2} \pi^3} \left| \frac{1}{2} A_t(\tau_t) [1 + \kappa_{gg}] + \dots \right|^2, \quad (2.15)$$

where the dots stand for the negligible contribution of the light quarks and leptons, and the coefficients κ can be written as:

$$\kappa_{\gamma\gamma} = \sum_{\text{NP}} \frac{3}{4} N_{c,\text{NP}} Q_{\text{NP}}^2 \frac{v_{\text{SM}}}{m_{\text{NP}}} \frac{\partial m_{\text{NP}}}{\partial v} \frac{A_{F,W,S}(m_{\text{NP}})}{A_t}, \quad (2.16)$$

$$\kappa_{gg} = \sum_{\text{NP}} 2C(r_{\text{NP}}) \frac{v_{\text{SM}}}{m_{\text{NP}}} \frac{\partial m_{\text{NP}}}{\partial v} \frac{A_{F,W,S}(m_{\text{NP}})}{A_t}, \quad (2.17)$$

where the ratio of A functions depends on the spin and masses of the new particles (and top). In the light Higgs approximation, however, the ratio only depends on the spin of the new particle:

$$\frac{A_{\text{NP}}}{A_t} = \begin{cases} 1 & \text{for fermions} \\ -\frac{21}{4} & \text{for vectors} \\ \frac{1}{4} & \text{for scalars} \end{cases} \quad (2.18)$$

An interesting feature of this parameterisation is that a particle with the same quantum numbers of the top will give $\kappa_{\gamma\gamma} = \kappa_{gg}$, and a single particle will give a contribution to

the two coefficients with the same sign. In this way, if the experimental data allow to point to a specific quadrant in the $\kappa_{\gamma\gamma}-\kappa_{gg}$ parameter space, we can have a hint of the underlying new physics model. This will be illustrated in various examples in the following sections. Note also that positive κ 's enhance the top contribution, therefore inducing an enhancement in the gluon channel but a suppression in the photon one, where there is a numerical cancellation between the dominant W contribution and the top one.

The presence of new physics often modifies the tree level relation between the mass of the SM particles and the Higgs VEV. This modification of the SM contribution can also be cast in the κ parameters. For the top it will read:

$$\kappa_{\gamma\gamma}(top) = \kappa_{gg}(top) = \frac{v_{SM}}{m_t} \frac{\partial m_t}{\partial v} - 1. \quad (2.19)$$

For the W :

$$\kappa_{\gamma\gamma}(W) = \frac{3}{4} \left(\frac{v_{SM}}{m_W} \frac{\partial m_W}{\partial v} - 1 \right) \frac{A_W(\tau_W)}{A_F(\tau_{top})}, \quad (2.20)$$

$$\kappa_{gg}(W) = 0. \quad (2.21)$$

Here, the difference between the VEVs does introduce relevant corrections and they must be taken into account.

Note that the modification of the SM couplings will also affect the other production channels, and the branching ratio in heavy gauge bosons. Those effects will however have a minor impact on our analysis, and their inclusion will be necessary in a later model-dependent analysis, after (and if) a model is preferred by data

2.1 Observables at the LHC and linear colliders

The LHC will measure the inclusive $\gamma\gamma$ Higgs decays and the new physics will modify both the total production cross section and the branching fraction in photons. For large masses, close to the W threshold, the decay in two heavy gauge bosons (one is virtual) becomes relevant and will also yield a relatively early measurement. At large luminosities, one may also measure the $\gamma\gamma$ decays in a specific production channel, for instance the vector boson fusion one that can be isolated using two forward jet tagging: in this case one may probe directly the branching ratios.

In the Higgs mass range of interest, between 115 and 150 GeV, the main production channel is gluon fusion with a SM cross section of 40 – 25 pb, followed by vector boson fusion (5 – 4 pb) and by other channels ($WH, ZH, \bar{t}tH$) which sum up to 4 – 2 pb. Here we will assume that the new physics significantly contributes only to the loop in the gluon fusion channel, while the other cross sections are unaffected. The total production cross section normalised with the SM one, that we denote as $\bar{\sigma}$, can be written as:

$$\bar{\sigma}(H) = \left(\frac{\sigma_{gg}^{NP} + \sigma_{VBF}^{SM} + \sigma_{VH,\bar{t}tH}^{SM}}{\sigma_{gg}^{SM} + \sigma_{VBF}^{SM} + \sigma_{VH,\bar{t}tH}^{SM}} \right) \simeq \left(\frac{(1 + \kappa_{gg})^2 \sigma_{gg}^{SM} + \sigma_{VBF}^{SM} + \sigma_{VH,\bar{t}tH}^{SM}}{\sigma_{gg}^{SM} + \sigma_{VBF}^{SM} + \sigma_{VH,\bar{t}tH}^{SM}} \right). \quad (2.22)$$

In the SM the Higgs branching fraction in photons amounts to $2 \cdot 10^{-3}$. In presence of new physics, the branching fraction will also be sensitive to the gluon loop via the

total width, as the gluon channel is significant: it amounts to 7% of the total for $m_H = 115$ GeV, decreasing to 3% for $m_H = 150$ GeV. Also in this case, we define a branching ratio normalised to the SM value, \overline{BR}

$$\begin{aligned} \overline{BR}(H \rightarrow \gamma\gamma) &= \frac{\Gamma_{\gamma\gamma}^{\text{NP}}}{\Gamma_{\gamma\gamma}^{\text{SM}}} \frac{\Gamma_{\text{tot}}^{\text{SM}}}{\Gamma_{gg}^{\text{NP}} + \Gamma_{\gamma\gamma}^{\text{NP}} + \Gamma_{\text{others}}^{\text{SM}}} \\ &\simeq \left(1 + \frac{\kappa_{\gamma\gamma}}{\frac{9}{16}A_W(\tau_W) + 1} \right)^2 \frac{\Gamma_{\text{tot}}^{\text{SM}}}{(1 + \kappa_{gg})^2\Gamma_{gg}^{\text{SM}} + (\Gamma_{\text{tot}}^{\text{SM}} - \Gamma_{gg}^{\text{SM}})}. \end{aligned} \quad (2.23)$$

The branching ratio in heavy vectors will depend on κ_{gg} via the total width of the Higgs, therefore the normalised \overline{BR} is

$$\overline{BR}(H \rightarrow VV^*) = \frac{\Gamma_{\text{tot}}^{\text{SM}}}{\Gamma_{gg}^{\text{NP}} + \Gamma_{\gamma\gamma}^{\text{NP}} + \Gamma_{\text{others}}^{\text{SM}}} \simeq \frac{\Gamma_{\text{tot}}^{\text{SM}}}{(1 + \kappa_{gg})^2\Gamma_{gg}^{\text{SM}} + (\Gamma_{\text{tot}}^{\text{SM}} - \Gamma_{gg}^{\text{SM}})}. \quad (2.24)$$

For completeness, the normalised gluon branching fraction can be written as

$$\overline{BR}(H \rightarrow gg) = \frac{\Gamma_{gg}^{\text{NP}}}{\Gamma_{gg}^{\text{SM}}} \frac{\Gamma_{\text{tot}}^{\text{SM}}}{\Gamma_{gg}^{\text{NP}} + \Gamma_{\gamma\gamma}^{\text{NP}} + \Gamma_{\text{others}}^{\text{SM}}} \simeq \frac{(1 + \kappa_{gg})^2\Gamma_{\text{tot}}^{\text{SM}}}{(1 + \kappa_{gg})^2\Gamma_{gg}^{\text{SM}} + (\Gamma_{\text{tot}}^{\text{SM}} - \Gamma_{gg}^{\text{SM}})}. \quad (2.25)$$

The branching ratios will be measured with an accuracy of few % at a TeV e^+e^- Linear Collider.

3 Survey of models of new physics in 4 dimensions

In this section we will summarise the values of the two parameters $\kappa_{\gamma\gamma}$ and κ_{gg} in a variety of models of new physics. It is not intended to be a complete survey, but rather a collection of examples of the usefulness of our proposed parametrisation, and of the impact of new physics on the Higgs search. Here, we will briefly discuss a fourth generation, supersymmetry, Little Higgs models, a scalar colour octet and the Lee-Wick SM. As the new particles and mass scales are often heavier than the top, we will use the light Higgs approximation to derive some simple analytical formulae.

3.1 A 4th generation

As for SM fermions, the masses of a chiral fourth generation are proportional to the Higgs VEV, and they cannot be arbitrarily large due to the perturbativity of the Yukawa couplings, naively $m_4 < 4\pi v \sim 2$ TeV. It has been shown that the impact of a relatively light 4th generation on the electroweak precision tests can be minimised if the spectrum follows a specific pattern [4]: in particular if the splitting between the up and down type quarks is about 50 GeV (and similarly for the leptons). For masses of a few hundred GeV, this is not a severe fine tuning. Finally, let us remind that direct bounds on such new particles are of the order of 190 GeV (for a fourth generation bottom type quark in $p\bar{p}$ collisions [5]) and 100 GeV for a charged lepton.

In the light Higgs approximation, the mass dependence disappears: κ_{gg} simply counts the number of new colour triplet quarks, $\kappa_{gg} = 2$, while $\kappa_{\gamma\gamma}$ depends on the charges

$$\kappa_{\gamma\gamma} = \frac{3}{4} \left[3 \left(\frac{2}{3} \right)^2 + 3 \left(-\frac{1}{3} \right)^2 + 1 \right] = 2. \quad (3.1)$$

Due to an accident in the charges, therefore, a complete extra generation contributes like two tops. Another accident is that the width in photons is largely suppressed, while the gluon one is enhanced by almost the same amount: overall, the inclusive $\gamma\gamma$ signal will be similar to the SM one [4] (for a light Higgs).

3.2 Supersymmetry

The supersymmetric contributions to the $h \rightarrow \gamma\gamma$ and $h \rightarrow gg$ amplitudes are well studied in supersymmetric extensions of the Standard Model (see for example [6] for few sample benchmark scenarios). Here we will focus on the common scenario where the heavier Higgses are above the WW threshold, so that the $\gamma\gamma$ decay mode is only relevant for the light Higgs h . However, the parametrisation we propose in this paper cannot be used in general for supersymmetric models. In fact, due to the presence of two Higgses which develop a VEV, the tree level couplings of the SM particles to the Higgs are modified at order $\mathcal{O}(1)$ compared to the SM case. If we define $\tan \beta = v_u/v_d$ the ratio of the two VEVs, and α the mixing angle in the neutral Higgs sector [7], the couplings of W , top (up-type fermions) and bottom (down-type fermions) compared to the SM values are corrected by the following factors:

$$\frac{g_{W+W-h}}{g_{SM}} = \sin(\beta - \alpha), \quad \frac{g_{tth}}{g_{SM}} = \frac{\cos \alpha}{\sin \beta}, \quad \frac{g_{bbh}}{g_{SM}} = -\frac{\sin \alpha}{\cos \beta}. \quad (3.2)$$

Those corrections can be large, even for heavy susy masses. In the large $\tan \beta$ case, which is preferred by the top Yukawa perturbativity and experimental constraints, the bottom (and tau) Yukawas are enhanced by a large factor $\sim \tan \beta$: the Higgs width increases and the branching ratio in photons can be easily suppressed by orders of magnitudes, making this channel unobservable. In order to keep the $\gamma\gamma$ channel alive, one needs to compensate the large $\tan \beta$ with a small mixing angle in the Higgs sector: $\alpha \sim \pm(\pi/2 - \beta)$. This requirements means that we are close to the decoupling limit in which the behaviour of the MSSM Higgs sector is Standard Model like. Indeed a limit $m_A \gg m_Z$ (where m_A is the mass of the pseudoscalar Higgs) implies that only the Standard Model like Higgs boson stays light while all the other scalars are heavy and that $\alpha \rightarrow (-\pi/2 + \beta)$ (for more details concerning this decoupling limit see [8]).

In order to safely use our formalism, we need to make sure that the corrections to the bottom Yukawa (and couplings to the W) are negligible. In the left panel of figure 1 we plotted the region in the α - β parameter space where both the W and bottom couplings deviate by less than 5% from the SM value (up to the overall sign). Note that the region delimited by the solid (red) lines in the left part of the left panel in figure 1 is precisely the one around the line of points where $\alpha = (-\pi/2 + \beta)$ which corresponds to the decoupling limit discussed above. We also superimposed the region where corrections to the top




Figure 1. Left panel: in solid red the region where the couplings of the W and bottom are both within 5% from the SM values (up to the sign), in dashed blue the same for the top couplings. Right panel: lower bound on the heavy Higgs masses as a function of $\tan\beta$ requiring deviations of 1% (solid) or 5% (dashed).

Yukawa are smaller than 5%. There is a tiny region where a fine tuning between the two angles allow for our formalism to be used. Note that a larger mixing angle in the Higgs sector will soon enhance the bottom Yukawa and kill the $\gamma\gamma$ signal, so the region is not a negligible part of the parameter space where the signal is observable. On the right panel we used a three level relation between the masses in the Higgs sector and α

$$\frac{\tan 2\alpha}{\tan 2\beta} = \frac{m_A^2 + m_Z^2}{m_A^2 - m_Z^2}, \quad (3.3)$$

where m_A is the mass of the pseudoscalar (which also sets the mass scale of the other heavy Higgses), to set a lower bound on the heavy Higgs masses. Therefore, we expect that for masses above 1 TeV, the corrections to the bottom (and tau) Yukawa can be safely neglected. In this region, corrections to the W and top couplings are small too.

For the purpose of illustrating our parameterisation, we will focus on some approximate expressions that arise in a simple scenario: the MSSM golden region [9]. This scenario is motivated by naturalness in the Higgs mass, minimal fine tuning and precision tests. The main features are large soft masses for the gauginos and for the light generations, and large mixing in the stop sector induced by a large soft trilinear term. A general analysis of the $\gamma\gamma$ channel can be found in ref. [10]. As a numerical example we will consider a variation of the benchmark point in ref. [9]: here, $\tan\beta = 10$ and all the soft masses except the stop and Higgs ones are at 1 TeV, $\mu = 250$ GeV and the soft trilinear term for the stops A_t is at 1 TeV to induce a large mixing in the stop sector and reduce the fine tuning in the Higgs potential. In this benchmark point, the light Higgs is at 129 GeV. Charginos and neutralinos (mostly higgsinos) are at 250 GeV (set by μ), while the stops are at 400 and

700 GeV. All other masses are above a TeV, and we will neglect the contribution of those sparticles. The only difference is that, in order to avoid the bottom Yukawa problem, we will push the heavy Higgses above 1 TeV: to do that it is enough to increase the H_d soft mass above the TeV scale. This will not introduce a severe fine tuning, as the contribution of this mass to the Higgs VEV is also suppressed by the large $\tan\beta$ [7]. In this scenario, only the stops contribute to $h \rightarrow gg$ and $h \rightarrow \gamma\gamma$. We neglect the contribution of charginos because they are mostly higgsinos: the coupling to the Higgs is suppressed by the large gaugino masses. For the stops, assuming that the soft masses for left and right handed scalars are equal (~ 550 GeV at the benchmark point), the contribution to the κ parameters can be expressed as

$$\kappa_{gg}(stops) = \kappa_{\gamma\gamma}(stops) \simeq \frac{m_2^2 + m_1^2}{4m_1^2m_2^2}m_t^2 - \frac{(m_2^2 - m_1^2)^2}{16m_1^2m_2^2} \sim -0.02, \quad (3.4)$$

where $m_{1,2}$ are the masses of the two eigenstates, and the second term is proportional to the soft trilinear term: $m_2^2 - m_1^2 \simeq 2|A_t|m_t$.

Those formulae are presented here for illustration purpose only, and we will use exact one loop expressions for the numerical analysis, including the contribution of charginos. A more general analysis of the region in the MSSM parameter space is beyond the scope of this paper, and it is postponed to a following publication.

3.3 Little Higgs models

In Little Higgs models [11], the gauge symmetries of the SM are a subgroup of a larger global symmetry. The breaking of such symmetry at a higher scale f produces light pseudo-Goldstone bosons, which we want to identify with the Higgs boson. The symmetry structure removes the divergences from the Higgs mass at one loop: the reason being that one loop is not sensitive to the explicit breaking of such global symmetry (while higher loops are). This is however enough to solve the little hierarchy problem, because the scale of new physics required beyond the Little Higgs mechanism is pushed above 10 TeV. In general new gauge bosons are introduced in order to eat up unwanted Goldstone bosons, and they also generate the loops that cancel the divergences from the SM gauge bosons. Similarly, new fermionic states, cousins of the top, required by the global symmetry, will cancel the divergences of the top loop.

We will first derive some very general formulae, and then apply them to explicit examples. In models with only one extra massive gauge boson, W' , the cancellation works thanks to the different sign between the couplings of the W and W' [12]:

$$g_{hW'W'} = -g_{hWW}. \quad (3.5)$$

This is a consequence of the fact that $m_W^2 + m_{W'}^2$ does not depend on the Higgs VEV, but it is fixed by the scale f at which the global symmetry is broken. The coupling of the W with the Higgs is also modified

$$g_{hWW} = g_{hWW}^{\text{SM}} (1 - \delta_W) = \frac{2m_W^2}{v_{\text{SM}}} (1 - \delta_W), \quad (3.6)$$

where δ_W contains corrections in v/f . After recalling that $g_{hWW} = \frac{\partial m_W^2}{\partial v}$, the contribution of the W and W' to the κ parameters is (in the light Higgs approximation):

$$\kappa_{\gamma\gamma}(W') = \frac{63}{16} \left(\frac{m_W}{m_{W'}} \right)^2 (1 - \delta_W), \quad (3.7)$$

$$\kappa_{\gamma\gamma}(W) = -\frac{9}{16} A_W(\tau_W) \delta_W; \quad (3.8)$$

while $\kappa_{gg} = 0$.

The precise value of δ_W depends on the symmetry structure of the model: in the Simplest Little Higgs (SLH) model [13], which is based on an SU(3) gauge symmetry,

$$m_W = gf \sin \frac{v}{2f}, \quad (3.9)$$

$$m_{W'} = gf \cos \frac{v}{2f}; \quad (3.10)$$

so that

$$\delta_W = 1 - \cos \frac{v}{2f} = 1 - \frac{m_{W'}}{\sqrt{m_{W'}^2 + m_W^2}} \simeq \frac{1}{2} \left(\frac{m_W}{m_{W'}} \right)^2 + \dots \quad (3.11)$$

At leading order in $m_W/m_{W'} \sim v/f$:

$$\kappa_{\gamma\gamma} \simeq \frac{9}{16} \left(7 - \frac{1}{2} A_W(\tau_W) \right) \left(\frac{m_W}{m_{W'}} \right)^2 \sim (6.2 \div 6.7) \cdot \left(\frac{m_W}{m_{W'}} \right)^2 \quad (3.12)$$

where we have varied the Higgs mass between 115 and 150 GeV.

The top sector is more complicated because doubling of fields is usually required in order to generate a realistic spectrum for the light states. For instance, the simplest way to introduce the top is to embed the SM left-handed doublet in a triplet of SU(3) that couples via the two Higgses to two right-handed singlets. In this case we need to double the right-handed tops in order to give mass both to the top and to its heavy partner T . The symmetry structure of the model implies that $m_t^2 + m_T^2$ does not depend on the SM Higgs VEV, therefore the following relation holds between the couplings to the Higgs ($g_{hff} = \frac{\partial m_f}{\partial v}$):

$$g_{hTT} = -\frac{m_t}{m_T} g_{htt}. \quad (3.13)$$

As in the gauge sector, the coupling of the SM top also receives deviations from the usual Yukawa coupling, which we can parametrise as

$$g_{htt} = \frac{m_t}{v_{\text{SM}}} (1 - \delta_t). \quad (3.14)$$

In terms of this parameterisation, the contribution to $\kappa_{\gamma\gamma}$ and κ_{gg} of the top and T are, in the light Higgs approximation:

$$\kappa_{\gamma\gamma}(top) = \kappa_{gg}(top) = -\delta_t, \quad (3.15)$$

$$\kappa_{\gamma\gamma}(T) = \kappa_{gg}(T) = -\frac{m_t^2}{m_T^2} (1 - \delta_t). \quad (3.16)$$

In the SLH model, the masses can be written as

$$m_{t,T}^2 = \lambda_T^2 f^2 \left(1 \mp \sqrt{1 - \frac{\lambda_t^2}{\lambda_T^2} \sin^2 \frac{v}{f}} \right), \quad (3.17)$$

where $\lambda_{t,T}$ are related to the Yukawa couplings to the two Higgses $\lambda_{1,2}$:

$$\lambda_T = \sqrt{\frac{\lambda_1^2 + \lambda_2^2}{2}}, \quad \lambda_t = \frac{\lambda_1 \lambda_2}{\lambda_T}. \quad (3.18)$$

Therefore

$$\delta_t = 1 - \frac{m_T^2}{m_T^2 - m_t^2} \frac{m_{W'}^2 - m_W^2}{m_{W'} \sqrt{m_{W'}^2 + m_W^2}} \simeq -\frac{m_t^2}{m_T^2} + \frac{3}{2} \frac{m_W^2}{m_{W'}^2} + \dots \quad (3.19)$$

and, at leading order,

$$\kappa_{\gamma\gamma}(top + T) = \kappa_{gg}(top + T) \simeq -\frac{3}{2} \frac{m_W^2}{m_{W'}^2} + \dots \quad (3.20)$$

Note that at leading order the result is independent on the heavy top mass, but only depends on the heavy gauge boson W' .

The total contribution is therefore:

$$\kappa_{gg}(SLH) \simeq -\frac{3}{2} \frac{m_W^2}{m_{W'}^2} \sim -0.002 \cdot \left(\frac{2\text{TeV}}{m_{W'}} \right)^2, \quad (3.21)$$

$$\kappa_{\gamma\gamma}(SLH) \simeq \left(\frac{141}{32} - \frac{9}{32} (7 + A_W) \right) \frac{m_W^2}{m_{W'}^2} \sim 0.007 \cdot \left(\frac{2\text{TeV}}{m_{W'}} \right)^2; \quad (3.22)$$

the Higgs mass dependence in A_W is very mild due to the small coefficient. In the numerical values we have chosen a W' mass of 2 TeV, which is roughly the one required by electroweak precision measurements [14]. Note however that the implementation of a T parity [15] would reduce the bound by almost an order of magnitude.

Another simple model using the Little Higgs mechanism was proposed in ref. [16] and dubbed Littlest Higgs. Here a global $SU(5)$ is spontaneously broken down to $SO(5)$, and a subgroup $SU(2)^2 \times U(1)^2$ is gauged. The mechanism acts thanks to the presence of two copies of the SM gauge group, which are broken to the diagonal by the spontaneous breaking of $SU(5)$. The Higgs again is a pseudo-Goldstone boson of the global symmetry breaking. The model, together with a heavy W (W_H) and top (T) also contains a heavy charged scalar Φ from a triplet of $SU(2)$ that develops a VEV (v'). The model therefore contains more parameters than the SLH, and its contribution to $H \rightarrow \gamma\gamma$ and $H \rightarrow gg$ has been computed in ref. [17]. Here we will simply translate those results in our notation: the contribution of the W and heavy gauge states is (expressed in terms of the masses at

leading order in the Higgs VEV v):

$$\kappa_{\gamma\gamma}(W) \simeq \frac{9}{16} \left(\frac{m_W^2}{m_{W_H}^2} - \frac{5-x^2}{8} \frac{v^2}{f^2} \right) A_W(\tau_W), \quad (3.23)$$

$$\kappa_{\gamma\gamma}(W_H) \simeq \frac{63}{16} \frac{m_W^2}{m_{W_H}^2}, \quad (3.24)$$

$$\kappa_{\gamma\gamma}(\Phi) \simeq -\frac{4-3x^2}{64} \frac{v^2}{f^2}; \quad (3.25)$$

$$\kappa_{gg}(W, W_H, \Phi) = 0. \quad (3.26)$$

Here x is proportional to the ratio between the triplet and doublet VEVs ($0 \leq x < 1$). Note that in the limit where the Higgs is much lighter than the W -threshold, the contributions proportional to the W mass tend to cancel, while the ones proportional to v^2 do not. The top and heavy top contributions are

$$\kappa_{\gamma\gamma}(top) = \kappa_{gg}(top) \simeq \frac{m_t^2}{m_T^2} - \frac{7-4x+x^2}{8} \frac{v^2}{f^2}, \quad (3.27)$$

$$\kappa_{\gamma\gamma}(T) = \kappa_{gg}(T) \simeq -\frac{m_t^2}{m_T^2}. \quad (3.28)$$

Note that as in the SLH, the contribution proportional to the top mass cancels out. Therefore, the main corrections in this model are proportional to the Higgs VEV and suppressed by the global symmetry breaking f :

$$\kappa_{gg}(LH) \simeq -\frac{7-4x+x^2}{8} \frac{v^2}{f^2}, \quad (3.29)$$

$$\begin{aligned} \kappa_{\gamma\gamma}(LH) \simeq & \left(\frac{195}{64} + x - \frac{73}{64}x^2 \right) \frac{1}{2} \frac{v^2}{f^2} + \\ & + \frac{9}{16} (7 + A_W) \left(\frac{m_W^2}{m_{W_H}^2} - \frac{5-x^2}{8} \frac{v^2}{f^2} \right). \end{aligned} \quad (3.30)$$

Note that the second term in $\kappa_{\gamma\gamma}$, which depends on the W_H mass, is negligible due to a small coefficient, therefore the result only depends on f and the triplet VEV x .

The bound from precision measurements on the scale f is around 5 TeV [14], which correspond roughly to masses of order 2 TeV. To give a numerical example, if x is negligible

$$\kappa_{gg}(LH) \simeq -\frac{7}{8} \frac{v^2}{f^2} \sim -0.002 \cdot \left(\frac{5 \text{ TeV}}{f} \right)^2, \quad (3.31)$$

$$\kappa_{\gamma\gamma}(LH) \simeq \frac{195}{128} \frac{v^2}{f^2} \sim 0.0036 \cdot \left(\frac{5 \text{ TeV}}{f} \right)^2. \quad (3.32)$$

Not that when a T parity is implemented on this model [15], the bound on f is lowered to 500 GeV [18], therefore the contribution to the κ parameters is 100 times bigger.

3.4 Extended scalar sector: colour octet

The scalar sector is experimentally the least tested part of the Standard Model and may be more complicated than the minimal content of the SM. It has been shown [19] that in order to avoid tree level flavour changing neutral currents, the extra scalar should be either a copy of the SM one (leading to the two Higgs model) or a colour octet with the same weak quantum numbers as the SM Higgs. Here we will focus on the latter possibility [19]. The most general potential contains 3 terms that are bilinear in both the Higgs H and the colour octet S :

$$\mathcal{L} = \lambda_1 H^{\dagger i} H_i S^{\dagger j} S_j + \lambda_2 H^{\dagger i} H_j S^{\dagger j} S_i + \left[\lambda_3 H^{\dagger i} H^{\dagger j} S_i S_j + h.c. \right] + \dots \quad (3.33)$$

where i and j are SU(2) indices and we have left implicit the colour contractions. Note that imposing custodial symmetry would require $\lambda_2 = 2\lambda_3$. After the Higgs develops a VEV $\langle H \rangle = v/\sqrt{2}$, the spectrum contains one charged and two neutral scalar octets with masses

$$m_{S^\pm}^2 = m_S^2 + \lambda_1 \frac{v^2}{4} = m_S^2 (1 + X_1), \quad (3.34)$$

$$m_{S_{1,2}^0}^2 = m_S^2 + (\lambda_1 + \lambda_2 \pm 2\lambda_3) \frac{v^2}{4} = m_S^2 (1 + X_1 + X_2 \pm 2X_3); \quad (3.35)$$

where $X_i = \lambda_i v^2/4$. At loop level, the octet will contribute to the electroweak precision tests [19]: the corrections can be encoded in the S and ρ parameters, and for small $v \ll m_S$:

$$S \simeq \frac{2}{3\pi} X_2, \quad (3.36)$$

$$\Delta\rho \simeq \frac{\sin^2 \theta_W m_W^2}{96\alpha\pi^3 m_S^2} (\lambda_2^2 - 4\lambda_3^3). \quad (3.37)$$

The corrections to the ρ parameter can be minimised by imposing (approximate) custodial symmetry, while S will give a direct constraint on X_2 . Note that X_1 is not strongly constrained.

Using the formalism developed in the previous section we can compute the contribution of the scalar octet to the κ parameters (for $v \ll m_S$):

$$\kappa_{\gamma\gamma}(S) \simeq \frac{3}{2} \frac{\lambda_1 v^2}{4m_{S^\pm}^2} \sim \frac{3}{2} X_1, \quad (3.38)$$

$$\begin{aligned} \kappa_{gg}(S) &\simeq \frac{C(8)}{2} \left(\frac{\lambda_1 v^2}{4m_{S^\pm}^2} + \frac{1}{2} \frac{(\lambda_1 + \lambda_2 + 2\lambda_3)v^2}{4m_{S_1^0}^2} + \frac{1}{2} \frac{(\lambda_1 + \lambda_2 - 2\lambda_3)v^2}{4m_{S_2^0}^2} \right) \\ &\sim \frac{3}{2} (2X_1 + X_2); \end{aligned} \quad (3.39)$$

where $C(8) = 3$. As a numerical example, we will use $\lambda_1 = 4$, $\lambda_2 = 1$ and $m_S = 750$ GeV. In this case, $X_1 \sim 1/9$ and $X_2 \sim 1/36$, therefore $\kappa_{\gamma\gamma} \sim 0.17$ and $\kappa_{gg} \sim 0.37$.

3.5 Lee-Wick Standard Model

Lee and Wick (LW) proposed a modification of the particle propagators in QED by means of higher derivative terms in order to improve the ultraviolet convergence of the theory and

make loop corrections finite. This modification of the propagator can also be parametrised by the presence of a new degree of freedom with large and negative kinetic term, so that the corrected propagator looks like a Pauli-Villars regularised one, where the Pauli-Villars cutoff scale is replaced by the mass scale of such new degree of freedom. This idea has recently been extended to the full SM [20]: in this case the loops are not finite, however the softening of the divergences is enough to address the hierarchy problem in the Higgs mass.

Notwithstanding the theoretical issues arisen by this formulation, the contribution of the LW degrees of freedom to the $H \rightarrow gg$ and $H \rightarrow \gamma\gamma$ amplitudes has been computed [21]: here we will sketch the calculation, making use of the general formulae given in section 2, and give some simple results in the large LW mass approximation.

In this model, to each SM particle, a new LW degree of freedom is associated (2 for each chiral fermion). For more details of the construction we refer the reader to the refs. [20, 21]. The Higgs VEV will generate a mixing between the standard and LW particles, which has been studied in detail in [21]: in the following we will review just the results needed to complete our calculation.

In the Higgs sector, the SM Higgs h and the LW scalar \tilde{h} mix via the Higgs VEV: the mixing can be described by a symplectic rotation

$$\begin{pmatrix} \cosh \theta & \sinh \theta \\ \sinh \theta & \cosh \theta \end{pmatrix}, \quad \text{with} \quad \tanh 2\theta = -2 \frac{m_{\tilde{h}}^2 \tilde{m}_h^2}{m_h^4 + \tilde{m}_h^4}. \quad (3.40)$$

A very similar mixing takes place in the gauge sector, between the W and the LW \tilde{W} . The two mass eigenstates are

$$m_W^2 = \frac{1}{2} \left(M_2^2 - \sqrt{M_2^4 - g^2 v^2 M_2^2} \right), \quad (3.41)$$

$$\tilde{m}_W^2 = \frac{1}{2} \left(M_2^2 + \sqrt{M_2^4 - g^2 v^2 M_2^2} \right); \quad (3.42)$$

where M_2 is the mass of the LW partner of the SU(2) gauge bosons. Note that there is no trilinear coupling between the W and the LW Higgs \tilde{h} , therefore (using the formulae in appendix A):

$$\begin{aligned} \frac{v_{\text{SM}}}{m_W} \frac{\partial m_W}{\partial v} &\rightarrow \cosh \theta \frac{gv}{2m_W} \frac{M_2^2}{\sqrt{M_2^4 - g^2 v^2 M_2^2}} \\ &= \frac{\tilde{m}_h^2}{\sqrt{m_h^4 + \tilde{m}_h^4}} \frac{\tilde{m}_W \sqrt{\tilde{m}_W^2 + m_W^2}}{\tilde{m}_W^2 - m_W^2} \simeq 1 + \frac{3}{2} \frac{m_W^2}{\tilde{m}_W^2}. \end{aligned} \quad (3.43)$$

For the \tilde{W} , the coupling to the Higgs is given by $-\frac{\partial \tilde{m}_W^2}{\partial v}$: the minus sign comes from the negative sign of the kinetic term. This can be proved by an explicit calculation, and it is true for all the LW fields (for fermions, the coupling to the Higgs is $-\frac{\partial \tilde{m}_f}{\partial v}$). However, another minus sign comes from the loops: compared to the SM ones, propagators and couplings to the gauge bosons (photons and gluons) have a minus sign from the negative

kinetic term of the LW fields. All in all, a minus sign from the loop compensates the minus sign in the Higgs coupling and we can safely use the formulae in section 2:

$$\frac{v_{\text{SM}}}{\tilde{m}_W} \frac{\partial \tilde{m}_W}{\partial v} \rightarrow -\cosh \theta \frac{g v m_W}{2 \tilde{m}_W^2} \frac{M_2^2}{\sqrt{M_2^4 - g^2 v^2 M_2^2}} \simeq -\frac{m_W^2}{\tilde{m}_W^2}. \quad (3.44)$$

Putting the two results together, and expanding for $m \ll \tilde{m}$:

$$\kappa_{\gamma\gamma}(W + \tilde{W}) \simeq \frac{27}{32} \frac{m_W^2}{\tilde{m}_W^2} (7 + A_W(\tau_W)) - \frac{63}{32} \frac{m_W^2}{\tilde{m}_W^2}. \quad (3.45)$$

This result is numerically small,¹ and the contribution of the W and its LW partner partially cancel each other in the light Higgs limit.

The spectrum also contains a LW charged scalar: in fact the LW Higgs does not develop a VEV and its charged component \tilde{h}^+ is not eaten up. The mass of such scalar is simply given by the Higgs LW mass $\tilde{m}_{h^\pm}^2 = M_H^2$. Nevertheless, as it happens in the SM with the Goldstone boson in the Higgs, the Lagrangian contains a coupling between \tilde{h}^+ and the Higgs field which can be calculated explicitly and enters the formulae in section 2 as (we are using here the same notation as in ref.s [20, 21])

$$\begin{aligned} \frac{v_{\text{SM}}}{\tilde{m}_{h^\pm}} \frac{\partial \tilde{m}_{h^\pm}}{\partial v} &\rightarrow -\frac{v_{\text{SM}}}{2 \tilde{m}_{h^\pm}^2} (\cosh \theta - \sinh \theta) \frac{\lambda v}{2} \\ &= -\frac{1}{2} \frac{\sqrt{\tilde{m}_W^2 + m_W^2}}{\tilde{m}_W} \frac{m_h^2 + \tilde{m}_h^2}{\sqrt{m_h^4 + \tilde{m}_h^4}} \frac{m_h^2 \tilde{m}_h^2}{\tilde{m}_{h^\pm}^2 (m_h^2 + \tilde{m}_h^2)} \simeq -\frac{1}{2} \frac{m_h^2}{\tilde{m}_{h^\pm}^2}, \end{aligned} \quad (3.46)$$

where an extra minus sign comes from the propagators in the loop. Therefore

$$\kappa_{\gamma\gamma}(\tilde{h}^\pm) \simeq -\frac{3}{32} \frac{m_h^2}{\tilde{m}_{h^\pm}^2}. \quad (3.47)$$

This result is also different from the result in ref. [21], where the contribution of the charged LW Higgs vanishes at this order.²

The top sector is more complicated because for each chiral SM fermion one needs to add a massive Dirac fermion (with negative kinetic term). The Yukawa couplings, however, have a simple form: in particular they have the same structure as the SM Yukawas, and they are functions of the field combination $H - \tilde{H} = 1/\sqrt{2}(v + h - \tilde{h} + \dots)$: the presence of a LW Higgs will only manifest itself in the fact that the couplings to the standard Higgs are

¹The authors of ref. [21] find that the contribution of the W is proportional to the SM amplitude by a factor $s_{(A+\tilde{A})^2} = \frac{\tilde{m}_W^2 + m_W^2}{\tilde{m}_W^2 - m_W^2}$. In fact, the coupling of the W with the Higgs can be written as $\cosh \theta s_{(A+\tilde{A})^2} \frac{g^2 v}{2}$; however $\frac{g v}{2} = \frac{m_W m_W}{\sqrt{\tilde{m}_W^2 + m_W^2}} \neq m_W$.

²In ref. [21], the authors include the contribution of the charged LW Higgs using the amplitude of the SM Goldstone boson in the Feynman gauge rescaled by the ratio $\frac{m_W^2}{\tilde{m}_{h^\pm}^2}$. However the coupling of a Goldstone boson (which is the same as \tilde{h}^+) is not proportional to its mass. See appendix A.3 for more details.

proportional to $\cosh \theta - \sinh \theta$. The spectrum can be calculated as a series for large LW top mass M_t (assuming the same mass for the LW partners of the left- and right-handed tops):

$$m_t = M_t \epsilon (1 + \epsilon^2 + \dots), \quad (3.48)$$

$$\tilde{m}_{t1,2} = M_t \left(1 \mp \frac{1}{2} \epsilon - \frac{3}{8} \epsilon^2 + \dots \right); \quad (3.49)$$

where $\epsilon = \frac{y_t v}{\sqrt{2} M_t}$. The contribution of the top (and partners) is therefore:

$$\begin{aligned} \kappa_{gg}(\text{tops}) = \kappa_{\gamma\gamma}(\text{tops}) = & \left((\cosh \theta - \sinh \theta) \frac{2m_W}{gv} \frac{\epsilon}{m_t} \frac{\partial m_t}{\partial \epsilon} - 1 \right) + \\ & + (\cosh \theta - \sinh \theta) \left(\frac{\epsilon}{\tilde{m}_{t1}} \frac{\partial \tilde{m}_{t1}}{\partial \epsilon} + \frac{\epsilon}{\tilde{m}_{t2}} \frac{\partial \tilde{m}_{t2}}{\partial \epsilon} \right) \simeq \frac{m_h^2}{\tilde{m}_h^2} + \frac{1}{2} \frac{m_W^2}{\tilde{m}_W^2} + \dots \end{aligned} \quad (3.50)$$

The ϵ dependence cancels out between the top and LW tops contributions, at the end the result only depends on the LW Higgs mass.

In total (at this order $\tilde{m}_{h\pm} = \tilde{m}_h$):

$$\kappa_{gg} \simeq \frac{m_h^2}{\tilde{m}_h^2} + \frac{1}{2} \frac{m_W^2}{\tilde{m}_W^2} \sim 0.015 \cdot \left(\frac{m_h}{120 \text{ GeV}} \frac{1 \text{ TeV}}{\tilde{m}_h} \right)^2, \quad (3.51)$$

$$\kappa_{\gamma\gamma} \simeq \frac{29}{32} \frac{m_h^2}{\tilde{m}_h^2} - \frac{47}{32} \frac{m_W^2}{\tilde{m}_W^2} + \frac{9}{16} \frac{m_W^2}{\tilde{m}_W^2} (7 + A_W(\tau_W)) \sim 0.017 \cdot \left(\frac{m_h}{120 \text{ GeV}} \frac{1 \text{ TeV}}{\tilde{m}_h} \right)^2, \quad (3.52)$$

where in the numerical example we neglected the contribution proportional to the W mass because of the higher bound on \tilde{m}_W from electroweak precision tests ($\tilde{m}_W \gg 3 \text{ TeV}$) [22].

4 Survey of models of new physics in extra dimensions

In this section we will focus on models of new physics in one extra dimension, in particular on the different ways one can employ the Higgs mechanism in this context. Most models can be divided in 3 main categories: bulk Higgs (BH), brane Higgs (bH) and Gauge Higgs (GH). In the first case, the Higgs is just a 5D scalar field in the bulk, which picks up a VEV due to a potential, which may be localised on one brane. In this class of models we find Universal Extra Dimensions [23–25] and gaugephobic Higgs models [26] in warped space, as an example. In brane Higgs models, the Higgs is a 4 dimensional field localised on one brane or end-point of the compact space: the advantage of these models is that there is no tower of massive scalars and, if the brane where the Higgs is localised plays a special role like the TeV brane in warped space, the model may address the little hierarchy problem. A model of this kind was proposed by Randall-Sundrum [27]. Finally, a new possibility allowed only in extra dimensional models is that the Higgs is part of a gauge group [28]: in fact the 5th component of a bulk gauge vector is a scalar from the 4D point of view. The interactions and potential of such particle are however constrained by 5D Lorentz and gauge invariance: in particular, the Higgs potential (including its mass) is finite and insensitive to the physics at the cutoff. The limit of this mechanism is that the model is only valid below an effective scale of few TeV (few Kaluza-Klein modes). In this

class we can find examples both in flat [29–31] and warped space [32–34]. It is interesting to note that the sign of the Higgs couplings, which are also related to the cancellation of the UV sensitivity of the Higgs mass, determines the sign of the contribution to the loop decays. For instance, in models of Gauge-Higgs unification we expect a reduction of the gluon coupling due to negative interference, contrary to what happens in models of Universal Extra dimensions (see for instance [31]).

Extra dimensional models are by nature non normalisable: from the 4D point of view, they are an effective description of the physics below a cutoff scale where some of the bulk interactions become strong. Such scale lies typically above a few tens of Kaluza-Klein (KK) modes. In general, if the symmetries allow so, we can add a tree level higher order operator which describes the coupling between the Higgs and the massless gauge bosons: in this way, the decay widths would be non-calculable. Adding such operator is actually necessary in order to act as a counter-term to the divergences that will arise at loop level. However, the loops we are interested in are effectively a box diagram if one considers a VEV insertion in the loop, therefore the one loop calculation turns out to be finite in all 5 dimension models. The counter-term will only be required at higher loops, and we will take the finite one loop result as a good approximation. In some cases, like in the Barbieri-Hall-Nomura model [23], the operator is actually forbidden by an extra symmetry (supersymmetry in this case). Models of Gauge Higgs are special: the Higgs interactions are constrained not only by gauge symmetry, but by 5D Lorentz invariance as well. This is enough to forbid a tree level potential for the Higgs, and also tree level contributions to the decay widths. Therefore, in this models the Higgs mass is really protected by symmetry and our calculation can be trusted as UV insensitive [35].

As we want to keep the discussion here as model independent as possible, we will express the spectra as a function of a dimensionless parameter α that is proportional to the Higgs VEV. Its precise definition depends on the specific model, and it will be specified case by case. In general, we will write:

$$\frac{v_{\text{SM}}}{m_{\text{NP}}} \frac{\partial m_{\text{NP}}}{\partial v} = \frac{v_{\text{SM}}}{v} \cdot \frac{\alpha}{m_{\text{NP}}} \frac{\partial m_{\text{NP}}}{\partial \alpha}. \quad (4.1)$$

The factor $v_{\text{SM}}/v = 1 - \delta_v$ contains eventual deviations in the numerical value of the Higgs VEV, and its effect is only relevant for the W and top contributions. We will present some general results on two different geometries: a flat extra dimension compactified on an interval (which is equivalent to an orbifold) and a warped extra dimension.

4.1 Gauge bosons in a flat extra dimension

In the flat case, the metric is an extended Minkowski, where the extra coordinate y lies on an interval $[0, \pi L]$. The notation is such that typically the mass of the first Kaluza-Klein state is $m_{KK} = 1/L$: this will be our reference mass scale in the following. Note that this is the only mass scale introduced by the extra space structure. This scale should be much larger than the W mass due to direct and indirect constraints: the electroweak precision tests usually push it above ~ 2 TeV (see for example [29, 36]). It is possible to relax this bound by adding symmetries: as a typical number in this scenario we will use

$m_{KK} \sim 500 \text{ GeV}$. This is the case, for example, in Universal Extra dimensions due to a Kaluza-Klein parity or the BHN model.

4.1.1 Gauge Higgs

One of the peculiarities of this models is the presence of a tower of charged vectors, H_μ^+ , associated with the charged component of the Higgs. They will necessarily mix with the W_μ via the Higgs VEV. Here we will focus on the simplest example, a $SU(3)$ gauge symmetry in the bulk, broken to $SU(2) \times U(1)$ at both endpoints. The value of the Higgs VEV can be expressed in terms of a dimensionless parameter α , which is indeed proportional to the field expectation value. We postpone all details of the calculation of the spectrum and the precise definition of α in the appendix B.1. The spectrum for the W and its KK tower is simply given, in terms of α , as

$$m_n^2 = \frac{(n + \alpha)^2}{L^2}, \quad n = 0, \pm 1, \pm 2 \dots \quad (4.2)$$

where $n = 0$ corresponds to the W mass:

$$m_W = \frac{\alpha}{L}. \quad (4.3)$$

For the purpose of this section, this can be considered as the definition of $\alpha = m_W L = m_W/m_{KK}$: it is typically a small number because we want the mass of the first KK mode to be much larger than the W mass in order to avoid direct and indirect bounds. The W mass is proportional to the Higgs VEV, so that its contribution to the loop is equal to the SM one and one finds $\delta_v(\text{GHflat}) = 0$. The KK tower contribution to the κ 's is proportional to

$$\sum_n \frac{\alpha}{m_n} \frac{\partial m_n}{\partial \alpha} = \alpha \sum_{n=1}^{\infty} \left(\frac{1}{n + \alpha} - \frac{1}{n - \alpha} \right) = \pi \alpha \cot \pi \alpha - 1 = -\frac{\pi^2 \alpha^2}{3} + \mathcal{O}(\alpha^4). \quad (4.4)$$

We can use the definition of α to express the result in terms of the W mass and the mass of the first KK mode $m_{KK} = 1/L$:

$$\begin{aligned} \kappa_{\gamma\gamma}(W_{KK}) &= -\frac{63}{16} \left(\pi \frac{m_W}{m_{KK}} \cot \left(\pi \frac{m_W}{m_{KK}} \right) - 1 \right) \\ &\simeq \frac{63}{16} \frac{\pi^2}{3} \frac{m_W^2}{m_{KK}^2} \sim 0.021 \cdot \left(\frac{2 \text{ TeV}}{m_{KK}} \right)^2; \end{aligned} \quad (4.5)$$

and $\kappa_{gg} = 0$. Note that the contribution has an opposite sign compared to the W .

Models with a gauge group larger than $SU(3)$ may also contain gauge bosons with different boundary conditions on the two endpoints. Those fields consist only of a tower of massive vector bosons, and they do not give rise to any massless vector or scalar modes. Also, they cannot mix with the W due to the flatness of the Higgs profile, therefore their presence will not affect the previous result. If they do couple to the Higgs, their spectrum is given by

$$m_n^2 = \frac{(n + 1/2 + c\alpha)^2}{L^2}, \quad n = 0, \pm 1, \pm 2 \dots \quad (4.6)$$

where c is a coefficient determined by gauge group factors. Their contribution to $\kappa_{\gamma\gamma}$ is proportional to the sum of all modes. The sum can be rewritten in terms of a sum from zero to infinity

$$c\alpha \sum_{n=0}^{\infty} \left(\frac{1}{n + 1/2 + c\alpha} - \frac{1}{n + 1/2 - c\alpha} \right) = -\pi c\alpha \tan \pi c\alpha = -\pi^2 c^2 \alpha^2 + \mathcal{O}(\alpha^4); \quad (4.7)$$

where therefore (Q_X being the charge of the extra gauge boson)

$$\kappa_{\gamma\gamma}(X) \simeq -\frac{63}{16} \pi^2 Q_X^2 c^2 \frac{m_W^2}{m_{KK}^2} \sim -0.063 Q_X^2 c^2 \cdot \left(\frac{2 \text{TeV}}{m_{KK}} \right)^2. \quad (4.8)$$

Note that it has an opposite sign compared to the W tower contribution, and that it tends to be larger by a factor of 3.

4.1.2 Brane Higgs

Let us first consider a bulk $SU(2) \times U(1)$ gauge symmetry, so that there is a single W tower. The spectrum is determined by the zeros of the equation (for more details, see the appendix B.2)

$$f(m, \alpha) = \pi L m \tan \pi L m - \pi^2 \alpha^2 = 0, \quad (4.9)$$

where α is again a dimensionless quantity proportional to the Higgs VEV. The spectrum can be computed in an expansion for small α :

$$m_W^2 L^2 = \alpha^2 \left(1 - \frac{\pi^2}{3} \alpha^2 + \mathcal{O}(\alpha^4) \right), \quad (4.10)$$

$$m_n^2 L^2 = n^2 + 2\alpha^2 + \mathcal{O}(\alpha^4). \quad (4.11)$$

In first approximation, $\alpha \sim m_W L = m_W/m_{KK}$: however higher order corrections in α will modify the couplings of the W to the Higgs, and they must be taken into account. The VEV is modified compared to the SM one, and in appendix B.2 we calculated

$$\delta_v(\text{bHflat}) \simeq \frac{\pi^2}{6} \alpha^2. \quad (4.12)$$

Even though the spectrum cannot be calculated analytically, in the appendix we showed that

$$\begin{aligned} 1 - \frac{\alpha}{m_W} \frac{\partial m_W}{\partial \alpha} &= \sum_{n=1}^{\infty} \frac{\alpha}{m_n} \frac{\partial m_n}{\partial \alpha} = 1 - \frac{2 \sin(2\pi L m_W)}{2\pi L m_W + \sin(2\pi L m_W)} \\ &= \frac{\pi^2}{3} m_W^2 L^2 + \mathcal{O}(m_W^4 L^4). \end{aligned} \quad (4.13)$$

The contribution to κ can be therefore written as:

$$\begin{aligned} \kappa_{\gamma\gamma}(W) &= -\frac{9}{16} (7 + A_W(\tau_W)) \left(1 - \frac{2 \sin(2\pi L m_W)}{2\pi L m_W + \sin(2\pi L m_W)} \right) (1 - \delta_v) \\ &\quad - \frac{9}{16} \delta_v A_W(\tau_W). \end{aligned} \quad (4.14)$$

Expanding for small L :

$$\begin{aligned} \kappa_{\gamma\gamma}(W) &\simeq -\frac{9}{32}\pi^2 (7 + A_W(\tau_W)) (m_W L)^2 + \frac{21}{32}\pi^2 (m_W L)^2 \\ &\sim (0.015 \div 0.022) \cdot \left(\frac{2 \text{ TeV}}{m_{KK}}\right)^2 \end{aligned} \quad (4.15)$$

where we have varied the H mass from 115 to 150 GeV. Note that the contribution of the KK tower is the same as in the Gauge Higgs case (up to a sign): however, the total contribution is suppressed by a partial cancellation between the KK tower and the W .

We can also consider the case of a bulk custodial symmetry, which will contain a W_R gauge boson which mixes with the W . In order to make it massive, one can impose Dirichlet boundary conditions on $y = 0$, the opposite brane to where the Higgs is localised. The spectrum is very similar, the only difference is to replace $L \rightarrow 2L$ in all the equations. However, this effect is compensated by the fact that the lightest KK mode from the W_R tower has mass $1/(2L)$ instead of $1/L$, therefore the result is the same as a function of the lowest KK mass.

4.1.3 Bulk Higgs

When the Higgs is in the bulk, it will generate a bulk mass for the gauge bosons. The VEV therefore will shift the spectrum

$$m_n^2 = \frac{n^2 + \alpha^2}{L^2}. \quad (4.16)$$

In this case, the W mass ($n = 0$) is proportional to the Higgs VEV (α), so that no corrections will come from the ordinary W ($\delta_v(\text{BHflat}) = 0$). We postpone details of the precise definition of α in the appendix B.1. The contribution of the tower is proportional to the sum

$$\sum_{n=1}^{\infty} \frac{m_W^2 L^2}{n^2 + m_W^2 L^2} = \frac{\pi m_W L \coth \pi m_W L - 1}{2} = \frac{\pi^2}{6} m_W^2 L^2 + \mathcal{O}(m_W^4). \quad (4.17)$$

The contribution to $\kappa_{\gamma\gamma}$ is therefore:

$$\kappa_{\gamma\gamma}(W) \simeq -\frac{63}{16} \frac{\pi^2}{6} m_W^2 L^2 \sim -0.01 \left(\frac{2 \text{ TeV}}{m_{KK}}\right)^2. \quad (4.18)$$

Note that the sign is different from the previous two cases, so that this contribution tends to sum up with the ordinary W one.

4.2 Gauge bosons in a warped extra dimension

A warped extra dimension is characterised by a non-trivial metric that, in the covariant coordinates that we will be using in this paper, can be written as

$$ds^2 = \left(\frac{R}{z}\right)^2 (\eta_{\mu\nu} dx^\mu dx^\nu - dz^2), \quad (4.19)$$

where R is the curvature of the space. Moving along the extra coordinate z , the unit length in 4D is rescaled, so that the natural energy scale of the model depends on the position along the warped extra coordinate. Now, z spans over an interval, but the two endpoints have a very different meaning: the brane at small $z = \epsilon$ is called the UltraViolet (UV) brane, and its typical scale represents the ultimate UV cutoff of the theory, $1/\epsilon = \Lambda$. One can imagine that this scale is very large, say the Planck scale M_{Pl} . On the other hand, at large $z = R'$ one places an InfraRed (IR) brane: its energy scale is directly related to the mass of the KK modes, so that $m_{KK} \sim 1/R'$ of order TeV. The proportionality factor depends on the particle we are considering: for a gauge boson like the W , the first KK mode is at $2.4 m_{KK}$.

The large splitting between the UV and IR scale, beyond explaining the weakness of gravitational interactions, also introduces a gap between the W mass and the KK mass scale

$$m_{KK} \simeq m_W \sqrt{\log \Lambda R'} \sim 6 m_W \tag{4.20}$$

for $\Lambda = M_{Pl}$ and $R' = 1 \text{ TeV}^{-1}$. This feature makes those models much more attractive than the flat cases, because the Higgs VEV can be closer to the IR scale. Finally, indirect bounds will usually require $m_{KK} \geq 1 \text{ TeV}$, which corresponds to a W' above 2 TeV [32, 37], similar to the flat case.

Here we will show some features of those models, and use a numerical evaluation of the κ 's in generic models. We focus on Gauge Higgs and IR brane Higgs models, as generic bulk Higgs models are much more complicated to deal with, both analytically and numerically [26].

4.2.1 Gauge Higgs

The spectrum of gauge bosons is determined by a complicated equation involving Bessel functions of order 1 and 0 (more details in the appendix B.1). If we expand for large UV scale, we can get a very good approximate spectrum which depends only logarithmically on Λ . For the W , assuming that it is much lighter than the KK mass, we can expand for small $m_W R' \ll 1$:

$$m_W^2 R'^2 \simeq \frac{2}{\log \Lambda R'} \sin^2 \pi \alpha \simeq \frac{2\pi^2}{\log \Lambda R'} \alpha^2, \tag{4.21}$$

where we have neglected higher order corrections in the log. The first KK mode will be given by the zeros of Bessel functions and one finds $m_{W'} \sim 2.4/R'$. Note also that this expression fixes α as a function of the KK scale R' . Contrary to the flat case, the W mass is not linear in the Higgs VEV, so that there will be corrections coming from the deviations from the SM coupling to the Higgs. We found (see appendix B.1):

$$\delta_v(\text{GHwarped}) \simeq 1 - \frac{\sin \pi \alpha}{\pi \alpha}, \tag{4.22}$$

and the total correction is therefore proportional to

$$(1 - \delta_v) \frac{\alpha}{m_W} \frac{\partial m_W}{\partial \alpha} - 1 \simeq \cos \pi \alpha - 1 \simeq -\frac{m_W^2 R'^2}{4} \log \Lambda R' \sim -0.055 \left(\frac{1/R'}{1 \text{ TeV}} \right)^2. \tag{4.23}$$

Note that one can use the same trick that we used in the flat brane Higgs (see appendix B.2) to calculate this quantity exactly: however, for our purpose, this approximate result is more than sufficient, and we can check that sub-leading terms will be suppressed by a log compared to this result:

$$\frac{\alpha}{m_W} \frac{\partial m_W}{\partial \alpha} - 1 = -\frac{m_W^2 R'^2}{6} \log \Lambda R' \left(1 - \frac{9}{4} \frac{1}{\log \Lambda R'} + \frac{3}{2} \frac{1}{\log^2 \Lambda R'} \right) + \mathcal{O}(m_W^4 R'^4). \quad (4.24)$$

The contribution to $\kappa_{\gamma\gamma}$ is

$$\kappa_{\gamma\gamma}(W) \sim \frac{9}{16} (-0.055) A_W(\tau_W) \sim (0.25 \div 0.30) \cdot \left(\frac{1/R'}{1\text{TeV}} \right)^2 \quad (4.25)$$

where $m_H = 115 \div 150$ GeV.

One can also numerically compute the KK tower contribution to the κ 's and find (for a W tower):

$$\kappa_{\gamma\gamma}(W_{KK}) \sim 0.009 \cdot \left(\frac{1\text{TeV}}{1/R'} \right)^2 \quad (4.26)$$

for $\Lambda = M_{Pl}$. This contribution is much smaller than the contribution of the W .

4.2.2 Brane Higgs

Expanding for large Λ and small $m_W R' \ll 1$, the mass of the W is (for more details, see the appendix B.2):

$$m_W^2 R'^2 = \frac{\alpha^2}{(1 + \alpha^2/2) \log \Lambda R'} + \dots \quad (4.27)$$

and

$$\delta_v(\text{bHwarped}) \simeq \frac{\alpha^2}{4} \simeq \frac{m_W^2 R'^2}{4} \log \Lambda R'. \quad (4.28)$$

Similarly to the Gauge Higgs case, the coupling of the W to the Higgs will receive corrections, and the contribution to the κ will be proportional to

$$\frac{\alpha}{m_W} \frac{\partial m_W}{\partial \alpha} - 1 = -\frac{m_W^2 R'^2}{2} \log \Lambda R' \left(1 - 2 \frac{1}{\log \Lambda R'} + \frac{1}{\log^2 \Lambda R'} \right) + \mathcal{O}(m_W^4 R'^4). \quad (4.29)$$

We also numerically computed the contribution of the KK tower, and, as in the flat case, the following relation holds:

$$\sum_{n=1}^{\infty} \frac{\alpha}{m_n} \frac{\partial m_n}{\partial \alpha} = 1 - \frac{\alpha}{m_W} \frac{\partial m_W}{\partial \alpha}. \quad (4.30)$$

Numerically:

$$\begin{aligned} \kappa_{\gamma\gamma}(W) &\sim -\frac{9}{16} (0.12) (7 + A_W(\tau_W)) - \frac{9}{16} (0.06) A_W(\tau_W) \\ &\sim (0.34 \div 0.51) \cdot \left(\frac{1/R'}{1\text{TeV}} \right)^2 \end{aligned} \quad (4.31)$$

where $m_H = 115 \div 150$ GeV.

4.3 Bulk fermions in a flat extra dimension

Bulk fermions are easier to analyse because the basic structure is common to all kind of Higgs models: we always need two bulk fermions, a doublet and a singlet of SU(2), that couple via the Higgs (either in the bulk or on the brane). In Gauge Higgs, those fields are in the same representation of the extended bulk gauge symmetry, while in other models they can be independent fields. Both in Gauge Higgs and in the brane Higgs case, the Higgs appears in the boundary conditions, which have the same form in the two cases: the reason is that one can use a gauge transformation to remove the Gauge Higgs VEV from the bulk equations of motion, as explained in more detail in the appendix C. The only difference is that the boundary conditions depend differently on the Higgs VEV. In the following we will use the notation of the Gauge Higgs models, where the Higgs VEV enters via trigonometric functions of a dimensionless parameter β . Note that the β parameter is different in general from the one for the gauge bosons α , due to either gauge group factors or Yukawa couplings. In Gauge Higgs, both β and α are proportional to the Higgs VEV. In the brane Higgs case, we can also define a fictitious β parameter that is related to the actual brane Higgs VEV V (see the appendix C for more details) as

$$\tan \pi\beta = yV, \tag{4.32}$$

where y is an effective Yukawa coupling. The spectrum will be the same in the two cases, as a function of β , however the couplings to the Higgs are different. The results in the brane Higgs case are equal to the Gauge Higgs case, up to a correction factor

$$\frac{V}{\beta} \frac{\partial\beta}{\partial V} = \frac{\sin \pi\beta \cos \pi\beta}{\pi\beta}. \tag{4.33}$$

This factor takes into account the non linear relation between β and the brane Higgs VEV V . Like in the gauge case, we will use eq. (4.1) (with α replaced by β), and take into account the contribution of δ_v on the top one. For simplicity, we will leave this effect implicit through this section.

In the Bulk Higgs case, the spectrum is different: the calculation is more complicated in the case of a generic Higgs profile, and we will only study the case of a constant Higgs VEV in a flat extra dimension, which is relevant for the UED model.

4.3.1 Bulk fermions

The first spectrum we will consider is the following:

$$m_n^2 = M^2 + \frac{(n + \beta)^2}{L^2}, \quad n = 0, \pm 1, \pm 2 \dots \tag{4.34}$$

This spectrum can arise in many scenarios: in Gauge Higgs models, one can generate the masses of light fermions by using two copies of bulk fermions with opposite boundary conditions, and connected by a bulk mass term M [38]. The light fermions are localised degrees of freedom that can mix with the massive bulk fields via localised mass terms: the electroweak symmetry breaking is mediated to the localised fields by the massive ones, like

in the Froggatt-Nielsen model. In this case, the smallness of the light mass can be achieved either by a small mixing, or by a large bulk mass M .

For $M = 0$

$$m_n^2 = \frac{(n + \beta)^2}{L^2}, \quad n = 0, \pm 1, \pm 2 \dots \quad (4.35)$$

and we have a light mode with mass

$$m_f = \frac{\beta}{L}. \quad (4.36)$$

In Gauge Higgs models, this can be identified with the top, whose mass is of order the electroweak scale: a $m_{top} \neq m_W$ can be obtained by using a large representation for the field containing the top [29], or by an explicit breaking of Lorentz invariance [30]. As we will shortly see, warped geometry automatically solves this problem. In brane Higgs models, all fermion masses can be generated in this way, because the relation between β and the Higgs VEV depends on Yukawa couplings and each field will feel a different effective β parameter. Finally, this spectrum will be generated also by a Bulk Higgs with a flat profile, where β is proportional to the Higgs VEV. For more detail about the spectra, see the appendix C.

The κ 's will be proportional to the sum

$$\begin{aligned} \sum_n \frac{\beta}{m_n} \frac{\partial m_n}{\partial \beta} &= \frac{\beta^2}{(ML)^2 + \beta^2} + \beta \sum_{n=1}^{\infty} \frac{n + \beta}{(ML)^2 + (n + \beta)^2} - \frac{n - \beta}{(ML)^2 + (n - \beta)^2} = \\ &= \frac{\pi \beta \sin(2\pi\beta)}{\cosh(2\pi ML) - \cos(2\pi\beta)}, \end{aligned} \quad (4.37)$$

with the proportionality coefficient determined by the quantum numbers of the 5D field (charge and colour), and a correction factor in the brane Higgs case. For large ML , this contribution is exponentially suppressed: the mass of the localised fermions is also suppressed by $\exp(-\pi ML)$, therefore we find

$$\kappa \sim \frac{m_f^2}{m_{KK}^2}. \quad (4.38)$$

In this class of models, one can safely neglect the contribution of the light fermion towers.

For the top in Gauge Higgs, or in the case of brane Higgs and Bulk Higgs models ($M = 0$), $\beta = m_f L$ and

$$\sum_n \frac{\beta}{m_n} \frac{\partial m_n}{\partial \beta} = \pi \beta \cot \pi \beta. \quad (4.39)$$

The contribution of the top tower is:

$$\kappa_{\gamma\gamma} = \kappa_{gg} = \left(\pi \frac{m_t}{m_{KK}} \cot \left(\pi \frac{m_t}{m_{KK}} \right) - 1 \right) \simeq -\frac{\pi^2}{3} \frac{m_t^2}{m_{KK}^2} \sim -0.025 \left(\frac{2 \text{ TeV}}{m_{KK}} \right)^2, \quad (4.40)$$

where we have subtracted the top contribution. In the brane Higgs case:

$$\begin{aligned} \kappa_{\gamma\gamma} = \kappa_{gg} &= \left((1 - \delta_v) \cos^2 \left(\pi \frac{m_t}{m_{KK}} \right) - 1 \right) \\ &\simeq -\pi^2 \frac{m_t^2}{m_{KK}^2} - \frac{\pi^2}{6} \frac{m_W^2}{m_{KK}^2} \sim -0.076 \left(\frac{2 \text{ TeV}}{m_{KK}} \right)^2. \end{aligned} \quad (4.41)$$

The contribution from other light fermion towers are negligible, as they will be proportional to the light fermion mass squared.

For completeness, let us also report the contribution from a tower of states with twisted boundary conditions, which may be present in models with large bulk representation and have a spectrum

$$m_n^2 = M^2 + \frac{(n + 1/2 + \beta)^2}{L^2}, \quad n = 0, \pm 1, \pm 2 \dots \quad (4.42)$$

In this case:

$$\begin{aligned} \sum_n \frac{\beta}{m_n} \frac{\partial m_n}{\partial \beta} &= \beta \sum_{n=0}^{\infty} \frac{n + 1/2 + \beta}{(ML)^2 + (n + 1/2 + \beta)^2} - \frac{n + 1/2 - \beta}{(ML)^2 + (n + 1/2 - \beta)^2} = \\ &= -\frac{\pi \beta \sin(2\pi\beta)}{\cosh(2\pi ML) + \cos(2\pi\beta)}. \end{aligned} \quad (4.43)$$

This contribution is also suppressed for large bulk masses. In the $M \rightarrow 0$ limit we get:

$$\sum_n \frac{\beta}{m_n} \frac{\partial m_n}{\partial \beta} = -\pi \beta \tan \pi \beta \simeq -\pi^2 \beta^2. \quad (4.44)$$

The contribution to the κ 's is

$$\kappa_{gg}(\text{twisted}) = -\left\{ \begin{array}{l} \pi \beta \tan \pi \beta \\ (1 - \delta_v) \sin^2 \pi \beta \end{array} \right\} \sim -0.075 \frac{m_f^2}{m_t^2} \left(\frac{2 \text{TeV}}{m_{KK}} \right)^2, \quad (4.45)$$

$$\kappa_{\gamma\gamma}(\text{twisted}) = \frac{9}{4} Q_f^2 \kappa_{gg}, \quad (4.46)$$

for a colour-triplet with $m_f L = \beta$. The two results in the brackets correspond to GH/BH (up) and bH (down), and they give the same contribution in the small β limit.

4.3.2 Bulk fermion in UED models

In this case the spectrum is (similarly to the gauge case)

$$m_n^2 = \frac{n^2 + \beta^2}{L^2}, \quad n = 0, 1, 2 \dots \quad (4.47)$$

with β proportional to the Higgs VEV via the bulk Yukawa coupling: for any SM fermion, $\beta = m_f L$, therefore only the top quark is relevant. The contribution to the amplitude is the same as for the gauge bosons, so that the top KK tower gives

$$\kappa_{\gamma\gamma}(\text{top}) = \kappa_{gg}(\text{top}) \simeq \frac{\pi^2}{6} m_t^2 L^2 \sim 0.01 \left(\frac{2 \text{TeV}}{m_{KK}} \right)^2. \quad (4.48)$$

Note that this contribution has an opposite sign compared to the GH/bH cases, and, in $\kappa_{\gamma\gamma}$, it tends to cancel the contribution of the W tower.

4.3.3 Odd bulk masses: fermions in models of flavour

Lorentz invariance in 5D allows to write down a mass term for a single 5D fermion \tilde{M} : this mass term is however forbidden in orbifold models, because the two components of the 5D fermions have opposite parities (unless the mass has an odd profile). A model defined on an interval is less constrained as it allows for the presence of such masses. Those odd masses have a very important phenomenological feature [39]: the zero mode of the 5D fermion is chiral, therefore \tilde{M} cannot give it a mass! Its effect is to exponentially localise the zero mode, and therefore modify the overlap with other fields, in particular with the Higgs (either bulk or localised). This feature can be used in a variety of models to generate the hierarchies in the fermion masses using order 1 Yukawas and bulk masses for all fermion fields! This is an alternative mechanism to generate light fermions in GH models, where the Yukawa couplings are equal due to gauge invariance, but it can also be used in bH and BH models. There is however a crucial difference between the two: in GH models the bulk masses are the same for the two SM fields that couple to the Higgs, because they come from the same bulk multiplet, while in bH/BH models they can be different. As we will see, this has dramatic consequences for the Higgs phenomenology.

Here we will focus on the GH and bH cases: the Higgs VEV enters via a dimensionless parameter β , and we will define

$$m_f = \beta/L. \tag{4.49}$$

As a reference, we will assume $m_f = m_{top}$, but this may not be the case in all models. The equation determining the spectrum is more complicated than in the previous case, therefore we will limit ourselves to an expansion for small β . In the GH case, the odd masses are the same for the two bulk fields, \tilde{M} . Expanding for $mL \ll \tilde{M}L$, we can calculate the mass of the light mode, which would be identified with the SM fermion:

$$m_l^2 = \frac{2\tilde{M}^2 \sin^2 \pi\beta}{\cosh(2\pi\tilde{M}L) - \cos(2\pi\beta)} \simeq \frac{2\tilde{M}^2 L^2 \pi^2}{\sinh^2 \pi\tilde{M}L} m_f^2. \tag{4.50}$$

It is clear from this formula that the light mode mass is suppressed by $\exp(-\pi\tilde{M}L)$ compared to the Higgs VEV β . Therefore, a $\tilde{M}L \sim \mathcal{O}(1)$ can explain the lightness of the fermions in the SM. The spectrum of the heavy modes, $m_n > \tilde{M}$, is more complicated:

$$m_n^2 L^2 = \tilde{M}^2 L^2 + n^2 \pm \frac{2n^2}{\sqrt{n^2 + \tilde{M}^2 L^2}} \beta + \frac{n^4 + 3\tilde{M}^2 L^2 n^2}{(n^2 + \tilde{M}^2 L^2)^2} \beta^2 + \mathcal{O}(\beta^3). \tag{4.51}$$

The couplings of each mode to the Higgs are large, however like in the previous case the modes have a different sign in the coupling. The sum, therefore, gives at leading order in β :

$$\begin{aligned} \sum_n \frac{\beta}{m_n} \frac{\partial m_n}{\partial \beta} &= -2\beta^2 \sum_{n=1}^{\infty} \frac{n^4 - 3\tilde{M}^2 L^2 n^2}{(n^2 + \tilde{M}^2 L^2)^2} \\ &= -\frac{\pi^2 \beta^2}{\sinh^2 \pi\tilde{M}L} \left(\frac{\pi\tilde{M}L}{\tanh \pi\tilde{M}L} - 1 \right) \simeq -\frac{m_l^2}{2\tilde{M}^2} \left(\frac{\pi\tilde{M}L}{\tanh \pi\tilde{M}L} - 1 \right). \end{aligned} \tag{4.52}$$

We find again that the tower of light modes does not contribute significantly to the widths: the only contribution will come from the top tower ($\tilde{M} \rightarrow 0$) which is approximate by the result in the previous subsection. Note again that this exponential suppression comes from a non trivial cancellation between modes.

In models with brane Yukawa couplings, the fields containing the SU(2) doublet and singlet are not necessarily the same, so they can have different bulk masses, \tilde{M}_L and \tilde{M}_R . In this case, the spectra of the two bulk fermions are different, the two KK towers are not degenerate in the $\beta \rightarrow 0$ limit and there are no cancellations between modes. As we will see, the spectra are degenerate if $\tilde{M}_R = -\tilde{M}_L$, however, even in this case, the cancellation between modes that we observe in the GH case does not occur. In conclusion, in models of this kind, the contribution of the KK tower of light modes can be large, as it is proportional to the 5D Yukawa coupling and not to the effective light-mode Yukawa (light fermion mass). As an example, we can study the latter case $\tilde{M}_L = -\tilde{M}_R = \tilde{M}$, where some simple analytical results can be obtained. The zero mode mass is, at leading order,

$$m_l \simeq 4\tilde{M}e^{-2\pi\tilde{M}L} \sin \pi\beta, \tag{4.53}$$

suppressed by $\exp(-2\pi\tilde{M}L)$. As before, we can compute the approximate KK spectrum for small β

$$m_n^2 L^2 = \tilde{M}^2 L^2 + n^2 \pm \frac{2n^2}{\sqrt{n^2 + \tilde{M}^2 L^2}} \beta + \frac{(1 - 2\pi\tilde{M}L)n^4 + (3 - 2\pi\tilde{M}L)\tilde{M}^2 L^2 n^2}{(n^2 + \tilde{M}^2 L^2)^2} \beta^2 + \mathcal{O}(\beta^3), \tag{4.54}$$

which differs to the $\tilde{M}_L = \tilde{M}_R$ case only at order β^2 , and the sum over the massive modes (the SM fermion is not included)

$$\begin{aligned} \sum_n \frac{\beta}{m_n} \frac{\partial m_n}{\partial \beta} &= -2\beta^2 \sum_{n=1}^{\infty} \frac{(1 + 2\pi\tilde{M}L)n^4 - (3 - 2\pi\tilde{M}L)\tilde{M}^2 L^2 n^2}{(n^2 + \tilde{M}^2 L^2)^2} = \\ &= -\frac{\pi^2 \beta^2}{4 \sinh^3 \pi\tilde{M}L} \left(\cosh(3\pi\tilde{M}L) + (4\pi\tilde{M}L - 1) \cosh(\pi\tilde{M}L) \right. \\ &\quad \left. - 4(\pi\tilde{M}L + 1) \sinh(\pi\tilde{M}L) \right) \\ &\simeq -\pi^2 \beta^2 \sim -0.075 \left(\frac{2\text{TeV}}{m_{KK}} \right)^2 \left(\frac{m_f}{m_{top}} \right)^2. \end{aligned} \tag{4.55}$$

The result is not very sensitive to the precise value of the bulk masses, even in the case of $\tilde{M}_L \neq \tilde{M}_R$ (we checked this numerically). Moreover, corrections from the non-linear relation between β and the Higgs VEV (the same multiplicative factor as in the previous section) will only affect this result at higher orders in β , while the light mode is negligible. The contribution of the top tower will be the same as in the massless case and, at leading order, it also gives $-\pi^2 \beta^2$. For a model with this flavour structure, contributions of the light fermion and top towers are:

$$\kappa_{\gamma\gamma} = \kappa_{gg} \simeq 6(-\pi^2 \beta^2) - \frac{\pi^2 \alpha^2}{6} \sim -0.45 \left(\frac{2\text{TeV}}{m_{KK}} \right)^2, \tag{4.56}$$

where the factor of 6 takes into account 3 complete SM generations, and we assumed that all the Yukawa couplings are of the same order (as the top one).

4.4 Fermions in a warped extra dimension

The localisation mechanism in warped extra dimension is much more effective than in the flat case: the reason is that the localisation is exponential with the large number $\Lambda R'$. The geometry itself generates two hierarchical mass scales: the UV cutoff Λ on the UV brane and the KK scale $1/R'$ on the IR brane. Here we will use the usual notation to call c the odd bulk mass in units of the curvature, $c = \tilde{M}R$. A left-handed (right-handed) zero mode is localised on the UV brane for $c > 1/2$ ($c < -1/2$) and IR brane for $c < 1/2$ ($c > -1/2$) [40].

GH models are characterised by the same odd mass c for the two fields that couple to the Higgs, because they are part of the same bulk multiplet. Like in the gauge boson case, we can expand for large UV cutoff, however in the fermionic case the expansion is more complicate and depends on the value of the bulk mass c . For $-1/2 < c < 1/2$ (when both zero modes are localised on the IR brane), the mass of the light mode is

$$m_f R' \simeq \sqrt{1 - 4c^2} \pi \beta \left(1 - \frac{3 + 4c^2}{9 - 4c^2} \pi^2 \beta^2 + \mathcal{O}(\beta^4) \right). \quad (4.57)$$

The mass is not suppressed compared to the Higgs VEV β ; notice also that the log suppression between the W mass and β is not present here, therefore one can fit the top mass without using a large representation (therefore, $\beta = \alpha$ is acceptable)! The mass does not depend linearly on β , thus the coupling with the Higgs receives corrections compared to the SM value, that will contribute to the κ 's. For a fermion with the same quantum numbers of the top (and in the light-Higgs approximation):

$$\kappa_{\gamma\gamma}(t) = \kappa_{gg}(t) = (1 - \delta_v) \frac{\beta}{m_f} \frac{\partial m_f}{\partial \beta} - 1 \simeq -\frac{32}{3} \frac{c^2}{(9 - 4c^2)(1 - 4c^2)} m_f^2 R' - \delta_v. \quad (4.58)$$

We also calculated this contribution exactly, and verified that this approximation is good for $|c| < 0.4$ at a few percent level. The κ 's vanish for $c \rightarrow 0$: in fact, in this limit the Bessel functions reduce to sines and cosines and we recover the flat case result where the light fermion mass is linear in the Higgs VEV. The coupling of the Higgs to the KK modes is also large. In summary, for $-1/2 < c < 1/2$, the light mass is un-suppressed compared to the Higgs VEV and the contribution of the tower to the κ 's is sizable. Numerically we found

$$\kappa_{\gamma\gamma}(t_{KK}) = \kappa_{gg}(t_{KK}) \sim -\frac{1}{3} m_f^2 R'^2 \quad (4.59)$$

for a fermion tower with the same quantum numbers of the top. This contribution will sum with the one coming from the top; notice that it is very similar to the flat case result (factor of 1/3). For the top quark in GH (with $1/R = 1 \text{ TeV}$, and $\beta = \alpha$ fixed by the W mass), we need $c \sim 0.43$: numerically

$$\kappa_{\gamma\gamma}(top) = \kappa_{gg}(top) \sim (-0.029 - 0.018) + (-0.011) = -0.06 \cdot \left(\frac{1 \text{ TeV}}{1/R} \right)^2. \quad (4.60)$$




Figure 2. Contribution to κ from a bulk fermion in GHU as a function of the bulk mass c , $1/R = 1 \text{ TeV}$, and $m_H = 130 \text{ GeV}$. The dotted line represents the contribution of the light mode (deviation from the SM one), the dashed line is for the KK tower and the thick one corresponds to the sum of both.

Note finally that this result can also be generalised to the brane-Higgs case with equal masses, taking into account the correction mentioned in the previous section, which is the same independently on the geometry.

For $c > 1/2$ and $c < -1/2$ the two zero modes are localised on different endpoints, and the light mass is suppressed:

$$m_f R' \simeq \left(\frac{1}{\Lambda R'} \right)^{|c| - \frac{1}{2}} \sqrt{4c^2 - 1} \sin \pi \beta. \tag{4.61}$$

In this case the coupling of the KK modes to the Higgs is also suppressed by:

$$\left(\frac{1}{2\Lambda R'} \right)^{2|c| - 1} \sim \frac{m_f^2}{m_{KK}^2}. \tag{4.62}$$

The contribution of a light fermion KK tower is negligible,³ however, contrary to the flat case, there is no cancellation involved and each KK mode coupling is suppressed by the light fermion mass. In figure 2 we computed numerically the contribution of a bulk fermion as a function of the bulk mass c ($\beta = \alpha$ fits the W mass and $1/R' = 1 \text{ TeV}$). The contribution of the light fermion is the deviation from a SM fermion of the same mass: it vanishes for $c = 0$, grows towards $c = 1/2$ reaching the value calculated for the W , and then goes down due to the decrease in the SM amplitude for a light fermion. The contribution of the KK modes, on the other hand, decreases for large c . The total contribution is almost constant for $c < 1/2$, then reaches a peak when the light fermion is at the Higgs decay threshold (in the plot, $m_H = 130 \text{ GeV}$), and then goes rapidly to zero.

It is straightforward to understand the suppression if we analyse in detail the structure of the wave functions: let us consider first the doublet, which contains a left-handed zero

³This result agrees with ref. [34].

mode. The wave functions, after applying the UV boundary conditions, are

$$\chi_L = A_L z^{5/2} \left[J_{-c+\frac{1}{2}}(p/\Lambda) J_{c+\frac{1}{2}}(pz) + J_{c-\frac{1}{2}}(p/\Lambda) J_{-c-\frac{1}{2}}(pz) \right] \quad (4.63)$$

$$\psi_L = A_L z^{5/2} \left[J_{-c+\frac{1}{2}}(p/\Lambda) J_{c-\frac{1}{2}}(pz) - J_{c-\frac{1}{2}}(p/\Lambda) J_{-c+\frac{1}{2}}(pz) \right] \quad (4.64)$$

where χ (ψ) is the left-handed (right-handed) wave function component. The expansion for small p/Λ depends on the value of c ($J_\nu(x) \sim x^\nu$ for small x):

$$\chi_L \simeq \begin{cases} J_{c+\frac{1}{2}}(pz) & \text{for } c > 1/2 \\ J_{-c-\frac{1}{2}}(pz) & \text{for } c < 1/2 \end{cases} \quad \psi_L \simeq \begin{cases} J_{c-\frac{1}{2}}(pz) & \text{for } c > 1/2 \\ J_{-c+\frac{1}{2}}(pz) & \text{for } c < 1/2 \end{cases} \quad (4.65)$$

plus corrections suppressed by $(\frac{p}{\Lambda})^{|2c-1|}$. For the singlet field, that contains the right-handed zero mode

$$\chi_R \simeq \begin{cases} J_{c+\frac{1}{2}}(pz) & \text{for } c > -1/2 \\ J_{-c-\frac{1}{2}}(pz) & \text{for } c < -1/2 \end{cases} \quad \psi_R \simeq \begin{cases} J_{c-\frac{1}{2}}(pz) & \text{for } c > -1/2 \\ J_{-c+\frac{1}{2}}(pz) & \text{for } c < -1/2 \end{cases} \quad (4.66)$$

plus corrections suppressed by $(\frac{p}{\Lambda})^{|2c+1|}$. The IR boundary conditions are:

$$\psi_L \cos \pi\beta + i\psi_R \sin \pi\beta = 0, \quad (4.67)$$

$$\chi_R \cos \pi\beta + i\chi_L \sin \pi\beta = 0. \quad (4.68)$$

It is clear that if the wave functions are proportional to each other, $\psi_L \propto \psi_R$ and $\chi_L \propto \chi_R$, the β dependence drops out from the equations: this is indeed the case at leading order for $c > 1/2$ and $c < -1/2$. In this case the Higgs VEV will affect the spectrum only via a suppressed contribution.

We can apply the same discussion to the generic brane localised Higgs: in this case, there are two different bulk masses c_L and c_R . Unless $c_L = c_R$, the wave functions are different and cannot be proportional to each other, therefore the coupling of the Higgs will be sizable even though the zero mode mass is suppressed due to the localisation of its wave functions. As in the flat case, the towers of light modes will give a large contribution to the κ 's. It can be calculated numerically and we found that for $c_L > 1/2$ and $c_R < -1/2$ it can be approximated by

$$\kappa_{\gamma\gamma} = \kappa_{gg} \simeq -\pi^2 \beta^2 \sim -0.12 \cdot \left(\frac{1 \text{ TeV}}{1/R'} \right)^2 \quad (4.69)$$

for a fermion tower with the same quantum numbers of the top, $\Lambda = M_{Pl}$ and using $\beta = \alpha$ that fits the W mass. Like in the flat case, the contribution of the KK towers of the light fermions is very large. To conclude, we can quote the number for a realistic quark and lepton spectrum. We use $c_L^{top} = 0.37$, $c_R^{top} = 0$, $c_R^{bot} = -0.55$:

$$\kappa_{\gamma\gamma}(\text{fermions}) \simeq \kappa_{gg}(\text{fermions}) \sim -0.71 \cdot \left(\frac{1 \text{ TeV}}{1/R'} \right)^2. \quad (4.70)$$

5 Numerical results

In this section we present exact numerical results for the models we considered in the previous two sections: in all cases, the analytic formulae are a very good approximation. We considered the following models:

- [♦] a fourth generation (the result is independent on the masses and Yukawa couplings);
- [♣] supersymmetry in the MSSM golden region: we only included the contribution of the stops with the spectrum given by the benchmark point in [9]. In this case the result is very sensitive to the parameters in the superpotential and in the susy breaking terms, therefore the general MSSM will cover a region of the parameter space;
- [▲] Simplest Little Higgs, the result scales with the W' mass (in the plots, $m_{W'} = 2 \text{ TeV}$);
- [*] Littlest Higgs, the result scales with the symmetry breaking scale f and has a mild dependence on the triplet VEV x (we set $x = 0$): for a model with T-parity we use $f = 500 \text{ GeV}$, without T parity $f = 5 \text{ TeV}$;
- [■] colour octet model, the result depends on 2 free parameters: for illustration we use in the plots $X_1 = 1/9$ and $X_2 = 1/36$ (see section 3.4);
- [▶] Lee-Wick Standard Model, the result scales with the LW Higgs mass: in the plots we set it to 1 TeV for illustration;
- [⊗] Universal Extra Dimension model [25], where only the top and W resonances contribute and the result scales with the size of the extra dimension: here we set $m_{KK} = 500 \text{ GeV}$ close to the experimental bound;
- [★] the model of Gauge Higgs unification in flat space in ref. [30], where only the W and top towers contribute ($\beta = m_t L$), with the first W resonance at 2 TeV;
- [●] the Minimal Composite Higgs [32] (Gauge Higgs unification in warped space) with the IR brane at $1/R' = 1 \text{ TeV}$: only W and top towers contribute significantly. The point only depends on the overall scale of the KK masses, as the other parameters are fixed by the W and top masses;
- [▼] a flat (W' at 2 TeV) and [♠] warped ($1/R'$ at 1 TeV) version of brane Higgs models, in both cases the hierarchy in the fermionic spectrum is explained by the localisation, and all light fermion towers contribute. Notwithstanding the many parameters in the fermion sector, the result only depends on the overall scale of the KK masses.

In the numerical results, the value of the mass of the new particles is at or around the lower bound given by precision electroweak tests; for larger masses, the contribution scales like the inverse squared mass (with the exception of the fourth generation). Note that in




Figure 3. $\kappa_{\gamma\gamma}$ and κ_{gg} at the LHC for a light Higgs ($m_H = 120$ GeV). The two solid lines correspond to the SM values of the inclusive $\gamma\gamma$ channel (**A**), and the vector boson fusion production channel (**B**). On the left panel, the dashed lines are spaced by 0.5, while the dotted ones by 0.1. On the right, we zoomed near the SM point.

many cases, the result only depends on one mass scale, and is insensitive to other free parameters present in the model: for example, in extra dimensional models with flavour, the final result does not depend on the precise localisation pattern of the bulk fields. Therefore, changing the parameters of the model can only move the point towards the origin by increasing such mass scale (except for supersymmetry and the colour octet model, where a wide region of the parameter space may be covered). The models are displayed in figures 3 and 4: different classes of models point in different quadrants of the parameter space. Therefore, if we could measure experimentally the two parameters, depending on the accuracy of the measurements, we may be able to distinguish between models and have an hint of what kind of mechanism lies behind the breaking of the electroweak symmetry. The direct discovery of the new particles would then be a confirmation of the model. The complementarity between the two measurements is crucial, because this indirect probe is sensitive to the quantum numbers and couplings to the Higgs of the new particles. This information is hardly accessible at the LHC, except in some special cases: most of the models analysed here predict new states above 1 TeV, at which mass scale one can only probe states produced by strong interactions, and their couplings to the Higgs and weak bosons will generally not play any significant role. It is crucial to understand the reach and discrimination power of the LHC in this parameter space.

The LHC will surely be able to measure the inclusive cross section $\sigma(pp \rightarrow H \rightarrow \gamma\gamma)$, as this is one of the golden channels for the discovery of a light Higgs. For an integrated luminosity of 10 fb^{-1} we can expect a 10% accuracy with respect to the Standard Model one [41]. We plotted the inclusive cross section normalised by the SM value in the $\kappa_{\gamma\gamma}-\kappa_{gg}$




Figure 4. $\kappa_{\gamma\gamma}$ and κ_{gg} at the LHC for a Higgs near the WW threshold ($m_H = 150$ GeV). The two solid lines correspond to the SM values of the inclusive $\gamma\gamma$ channel (**A**), and the inclusive V^*V channel ($V = W, Z$) (**B**). On the left panel, the dashed lines are spaced by 0.5, while the dotted ones by 0.1. On the right, we zoomed near the SM point.

parameter space for a light Higgs ($m_H = 120$ GeV) in figure 3 and for a Higgs near the VV -threshold ($m_H = 150$ GeV) in figure 4: many models lie very far from such line, and a 10% measurement would allow to probe new physics masses up to few TeV in some cases. Note that many of the models we studied predict a reduction of the inclusive signal: the measurement of an enhancement at the LHC may be a sign of unexpected new physics. Note also that some very different models can give the same prediction, like the fourth generation case where a suppression in the $\gamma\gamma$ decay is accidentally compensated by an enhancement in the gluon fusion cross section. Therefore, we need to measure another observable at the LHC in order to distinguish such models. For the light Higgs case, in figure 3 we plotted the vector boson fusion channel, which is sensitive to the $\gamma\gamma$ branching fraction directly. This channel is orthogonal to the inclusive one, and therefore offers the best discrimination power. Experimentally, this channel is very promising even at low luminosity, for the observation of the Higgs Boson in the $H \rightarrow \gamma\gamma$ decay mode [42]. However a detailed study of this channel, as required for the precise determination of the κ parameters demands a high luminosity [43]. A precise study requires a detailed simulation and will not be given here. For a heavier Higgs, in figure 4, the decay in massive gauge bosons $H \rightarrow V^*V$ (with one virtual) becomes relevant and offers another discovery channel. This channel, sensitive to the total cross section, will allow for a discrimination for Higgs masses near the WW threshold.

The Linear Collider will have a much better chance to discriminate between models than the LHC. In fact, an experiment at a linear collider will be able to measure directly




Figure 5. $\kappa_{\gamma\gamma}$ and κ_{gg} at the ILC ($m_H = 120$ GeV). The two solid lines correspond to the SM values of the $\gamma\gamma$ (**A**) and gluon (**B**) branching ratios. On the left panel, the dashed lines are spaced by 0.5, while the dotted ones by 0.1. On the right, we zoomed near the SM point.

the branching fractions into gluons and photons. After 100 fb^{-1} of data, in the photon channel an accuracy of 5–7 % is expected (reduced to 2–3 % with the $\gamma\gamma$ collider option), while the gluon channel offers a 2 % accuracy (assuming SM values) [44]. We compared the models with the ILC measurements in figure 5.

6 Conclusions

The decay in a pair of photons is the golden channel for the discovery of the Higgs boson at the LHC for an intermediate mass, below the WW threshold, where the dominant decay mode would be $b\bar{b}$. This decay occurs via a loop diagram, where the heaviest particles in the SM (W and top) contribute the most. Furthermore, the production cross section at the LHC is dominated by a similar loop diagram that mediates the coupling of the Higgs to a pair of gluons. This situation offers a precious handle on new physics: in fact, new particles that may be present at the TeV scale will also contribute to those loops, therefore modifying the SM predictions for the Higgs production and decay rates.

One of the main motivations to expect new physics at the TeV scale is the naturalness of the Higgs mass (electroweak scale): the new particles, partners of the gauge bosons and of the top, will cancel or soften the divergences in the loop corrections to the Higgs mass. If this is the case, the new particles will have a significant coupling to the Higgs and therefore contribute significantly to the loop couplings of the Higgs. The LHC will be able to discover such new particles, with masses up to few TeV for particles with strong interactions and 1 TeV for weakly interacting ones. However, little information on the couplings will be directly accessible: the discovery of new states will not tell us if they play any role in the Higgs physics. Measuring deviations in the $H \rightarrow \gamma\gamma$ and $H \rightarrow gg$ couplings

at later times will give us an important hint to understand the nature of the new states and of the underlying model of electroweak symmetry breaking.

In this paper, we studied the contribution of new physics to the $H \rightarrow \gamma\gamma$ and $H \rightarrow gg$ decay widths (the latter is proportional to the production cross section). We propose a convenient parameterisation of the new contributions, by introducing two independent parameters $\kappa_{\gamma\gamma}$ and κ_{gg} . Such a simple parameterisation neglects contributions to the tree level processes, such as production channels other than gluon fusion and decays, that are generically present in models of new physics. This parameterisation is especially useful in models where such effects are small. They could be taken into account in a later model-dependent analysis once a specific model or class of models is preferred by data. On more general grounds, more parameters can be introduced and the analysis extended in a similar fashion: for instance, in supersymmetry, a parameter describing the variation of the total width of the Higgs due to the bottom Yukawa coupling can be used. We avoided doing so as many models do have small corrections and in order to keep the parameterisation as simple as possible.

Simple new physics scenarios give rise to simple correlations in this parameter space: for instance, a top partner will have $\kappa_{\gamma\gamma} = \kappa_{gg}$, while a single new particle will generate same-sign κ 's. In order to illustrate the power of a model independent measurement at the LHC (and at future Linear Colliders) we compiled a necessarily incomplete survey of models of new physics both in 4 and 5 dimensions. Our results show that there are classes of models pointing in different quadrants of the parameter space, and that the deviations from the SM predictions can be as large as 50%. Moreover, in most cases those results do not depend on the details of the model and they are sensitive to just one mass scale of the new physics. Therefore, a cross section measurement at the LHC will allow to discriminate models even with new particle masses at the TeV scale. At the Linear Collider, the few percent level measurement of the Higgs branching ratios will allow an even better discrimination. Note also that most of the models in our survey populate the $\kappa_{\gamma\gamma} < \kappa_{gg}$ region, where we generically expect a suppression of the inclusive cross section. In this parameterisation it would be easy to discover hints of unconventional or unexpected new physics, independently on direct and/or indirect signals in other channels.

Acknowledgments

We thank Suzanne Gascon and Nicolas Chanon for useful discussions and comments. This work is supported in part by the ANR project SUSYPHENO (ANR-06-JCJC-0038).

A Higgs couplings in extended Higgs sectors

A.1 Multiple Higgs

The Higgs sector may contain multiple scalar fields which develop a VEV, like for instance in supersymmetry where two Higgs doublets are required in order to allow up and down type Yukawa interactions. Let us imagine that there are n such Higgs multiplets ϕ_i , such

that

$$\phi_i = \frac{1}{\sqrt{2}}(v_i + c_i h + \dots), \tag{A.1}$$

where h is the lightest mass eigenstate (that we would identify with the SM Higgs), and dots represent the other (heavier) scalar mass eigenstates. The VEVs are all non zero $v_i \neq 0$. In this case, in the formulae in section 2, one needs to replace:

$$\frac{v_{\text{SM}}}{m} \frac{\partial m}{\partial v} \rightarrow \frac{v_{\text{SM}}}{m} \sum_i \frac{\partial m}{\partial v_i} c_i. \tag{A.2}$$

For example, in the case of supersymmetry, there are two Higgs doublets, $H_{u,d}$ with $H_u = 1/\sqrt{2}(v_u + h \cos \alpha + \sin \alpha H)$ and $H_d = 1/\sqrt{2}(v_d - h \sin \alpha + \cos \alpha H)$, and $v_{\text{SM}}^2 = v_u^2 + v_d^2$ ($\tan \beta = v_u/v_d$). The W mass is given by $m_W^2 = g^2/4(v_u^2 + v_d^2)$, so that:

$$\frac{v_{\text{SM}}}{m_W} \frac{\partial m_W}{\partial v} \rightarrow \frac{v_{\text{SM}}}{m_W} \left(\frac{m_W}{v_{\text{SM}}} \sin \beta \cos \alpha - \frac{m_W}{v_{\text{SM}}} \cos \beta \sin \alpha \right) = \sin(\beta - \alpha). \tag{A.3}$$

For the top, $m_t = y v_u$:

$$\frac{v_{\text{SM}}}{m_t} \frac{\partial m_t}{\partial v} \rightarrow \frac{v_{\text{SM}}}{m_t} y \cos \alpha = \frac{\cos \alpha}{\sin \beta}. \tag{A.4}$$

A.2 Higgs mixing

Another interesting case is when the Higgs mixes with additional scalars that do not develop a VEV. This situation may be realised in multiple Higgs models, or in the Lee-Wick SM. We will call S_j those inert scalars, which contain the light Higgs field h :

$$S_j = s_j h + \dots \tag{A.5}$$

As before, we are assuming that the other mass eigenstates are heavier than the h , which we want to identify with the SM Higgs. The scalars S_j may couple to a particle p with coupling

$$g_{S_j} S_j \bar{p} p, \tag{A.6}$$

which will contribute to the coupling of p to the Higgs h via the mixing. In this case, one can use the formulae in section 2 with

$$\frac{v_{\text{SM}}}{m} \frac{\partial m}{\partial v} \rightarrow \frac{v_{\text{SM}}}{m} \left(\sum_i \frac{\partial m}{\partial v_i} c_i + \sum_j g_{S_j} s_j \right). \tag{A.7}$$

A.3 Charged Higgs couplings

Another situation where the coupling to the Higgs does not come via the v -dependence of the mass, is when the particle in question does couple with the Higgs potential. In fact, the Higgs potential implicitly contains the VEV, and this fact may lead to cancellations in the particle mass. One may calculate the mass of the particle as a function of the Higgs field VEV $\langle H \rangle$ and v (which are numerically equal), derive in $\langle H \rangle$ and then impose $\langle H \rangle = v$.

In most cases it is easier to compute the coupling directly from the Higgs potential: here we will summarise three cases that are useful for the calculations in this paper.

The most trivial example is the charged Goldstone boson in the SM, which is eaten by the W in the Unitary gauge. In Feynman gauge, the charged component of the Higgs doublet remains in the spectrum and its mass is $m_{\phi^\pm} = m_W$. This may lead to the wrong conclusion that its couplings to the Higgs are the same as the W . The Higgs potential can be written as

$$V(H) = \frac{\lambda}{4} \left(H^\dagger H - \frac{v^2}{2} \right)^2 ; \tag{A.8}$$

after expanding the Higgs field as $H = \begin{pmatrix} \phi^+ \\ \frac{v+h+i\phi}{\sqrt{2}} \end{pmatrix}$, it does not generate any mass for the Goldstone bosons ϕ , because of a cancellation between the mass term $-\lambda v^2/4$ and a contribution from the quartic coupling (the mass is given by the gauge fixing term). However, the quartic coupling does generate a trilinear coupling with the Higgs h :

$$\frac{\lambda v}{2} \phi^+ \phi^- h = \frac{m_h^2}{v} \phi^+ \phi^- h . \tag{A.9}$$

The coupling to the Higgs is therefore proportional to the Higgs mass. The amplitude generated by the Goldstone boson can be computed starting from the amplitude of a standard scalar

$$A_{\phi^\pm}(\tau_W) = \frac{v}{2m_W^2} \frac{m_h^2}{v} A_S(\tau_W) = 2\tau_W A_S(\tau_W) , \tag{A.10}$$

where $\tau_W = \frac{m_h^2}{4m_W^2}$.

A similar situation happens in the Lee-Wick SM: together with the standard Higgs field H , there exist a LW scalar \tilde{H} with negative kinetic term. The potential is:

$$V_{LW}(H, \tilde{H}) = V(H - \tilde{H}) - M_H^2 \tilde{H}^\dagger \tilde{H} . \tag{A.11}$$

Only the standard Higgs develops a VEV, while the LW Higgs does not thanks to its large LW mass M_H . The charged component of the Higgs is eaten by the massive W ; the charged component of the LW field \tilde{h}^+ is a physical degree of freedom with mass given simply by the LW mass: the v dependence cancels out like for the Goldstone bosons. Nevertheless, a trilinear coupling $\tilde{h}^+ \tilde{h}^- h$ is present with coefficient proportional to $\lambda v/2$ (the proportionality coefficient depends on the mixing in the neutral sector, and it is discussed in section 3). The amplitude for the LW field can be written as:

$$A_{\tilde{h}^\pm}(\tilde{\tau}_{h^\pm}) = \frac{v_{SM}}{2\tilde{m}_{h^\pm}^2} \frac{\lambda v}{2} A_S(\tilde{\tau}_h) = \frac{\lambda v v_{SM}}{2m_h^2} 2\tilde{\tau}_h A_S(\tilde{\tau}_h) ; \tag{A.12}$$

where $\tilde{\tau}_h = \frac{m_h^2}{4\tilde{m}_{h^\pm}^2}$. This formula is different from the one used in ref. [21].

Finally, let us discuss the case of the charged Higgs in the MSSM: the model contains two Higgs doublets $H_{u,d}$ with opposite hypercharge to generate up- and down-type Yukawas. The potential contains two quartic couplings [7]:

$$\begin{aligned}
 V_{MSSM}(H_u, H_d) &= \frac{g^2 + g'^2}{8} (|H_u|^2 - |H_d|^2)^2 + \frac{g^2}{2} |H_u H_d^\dagger|^2 + \dots \\
 &= \frac{g^2 + g'^2}{8} (|H_u^0|^2 - |H_d^0|^2 + H_u^- H_u^+ - H_d^+ H_d^-)^2 \\
 &\quad + \frac{g^2}{2} |H_u^+(H_d^0)^* + H_u^0 H_d^+|^2 + \dots
 \end{aligned}
 \tag{A.13}$$

where the dots stand for quadratic terms. The two neutral components develop a VEV: $\sqrt{2} \langle H_u^0 \rangle = v_u = v_{\text{SM}} \sin \beta$ and $\sqrt{2} \langle H_d^0 \rangle = v_d = v_{\text{SM}} \cos \beta$. However, only one combination actually acquires a VEV: we can define $H_1 = \sin \beta H_u - \cos \beta H_d^\dagger$ and $H_2 = \cos \beta H_u + \sin \beta H_d^\dagger$ such that $\sqrt{2} \langle H_1 \rangle = v_{\text{SM}}$ and $\sqrt{2} \langle H_2 \rangle = 0$. The charged component of H_1 is eaten by the W , while the charged component of H_2 is the physical charged Higgs: $H_u^+ = \cos \beta H^+$ and $H_d^+ = \sin \beta H^+$. Plugging those solutions in the potential, and expanding around the VEV $\sqrt{2} \langle H_u^0 \rangle = v_u + \cos \alpha h + \sin \alpha H$ and $\sqrt{2} \langle H_d^0 \rangle = v_d - \sin \alpha h + \cos \alpha H$, we find that

$$m_{H^\pm}^2 = m_A^2 + m_W^2, \tag{A.14}$$

$$g_{H^+ H^- h} = \frac{2m_W^2}{v_{\text{SM}}} \sin(\beta - \alpha) + \frac{m_Z^2}{v_{\text{SM}}} \cos(2\beta) \sin(\beta + \alpha), \tag{A.15}$$

where m_A is a mass term independent on the VEVs. The coupling to the light Higgs has a term proportional to the W mass square, coming from the second term in the potential (this is what we would obtain from the mass formula), and a term proportional to the Z mass square, from the first quartic term in the potential: the latter cancels out in the mass formula but does contribute to the Higgs couplings.

B Gauge bosons in 5D

In this appendix, we propose a more detailed description of the models that we consider in section 4, we sketch how to extract the spectra of masses for Gauge bosons for different choices of geometry and compactification of the fifth dimension. We first derive general results in a generic metric with the extra coordinate $y \in [y_1, y_2]$ and

$$ds^2 = w(y)^2 (dx_\mu dx^\mu - dy^2), \tag{B.1}$$

and then we discuss the limits of flat ($w = 1$, $y_1 = 0$ and $y_2 = \pi L$) and warped AdS ($w = R/y$, $y_1 = R$ and $y_2 = R'$) metrics. The action of a pure gauge theory in one extra dimension, after fixing the R_ξ gauge, is given by:

$$\mathcal{S} = \int d^4x \int_{y_1}^{y_2} dy w \left\{ -\frac{1}{4} F_{\mu\nu}^a F^{\mu\nu a} - \frac{1}{2} F_{\mu 5}^a F^{\mu 5 a} - \frac{1}{2\xi} \left[\partial_\mu A^{\mu a} - \xi \frac{1}{w} \partial_5 (w A_5^a) \right]^2 \right\}, \tag{B.2}$$

where $F_{MN}^a = \partial_M A_N^a - \partial_N A_M^a + g_5 f^{abc} A_M^b A_N^c$ and g_5 is the 5D gauge coupling. In the Unitary gauge ($\xi \rightarrow \infty$), the massive modes of the fifth component A_5 are removed and they become the longitudinal polarisation of the massive KK vectors. In the following, we will discuss two models: one where the Higgs is part of the gauge field, namely a zero mode of the A_5 , and another where the Higgs is a brane localised field.

B.1 Gauge Higgs unification models

A zero mode for the A_5 component of the gauge field is a physical scalar in the spectrum because it is not eaten up in the Unitary gauge. However, it is a special scalar because its potential is constrained by Lorentz and gauge invariance: in 5D no potential is allowed at tree level, therefore it is generated at loop level and it is finite. This property makes the A_5 an ideal candidate to play the role of the Higgs boson. In order to obtain a zero mode, we need to enlarge the SM gauge group such that a doublet of $SU(2)$ is part of the gauge fields, and break the gauge directions of this doublet on both end points by imposing Dirichlet boundary conditions on the vectors (and therefore Neumann boundary conditions on the A_5 component).

For simplicity, we work on the minimal model where the gauge symmetry is enlarged to $SU(3)$, broken to the electroweak $SU(2) \times U(1)$ at the boundaries. The bulk fields can be written as:

$$A_\mu = \begin{bmatrix} \mathbf{W}_\mu^{(3)+1/\sqrt{3}\mathbf{B}_\mu^{(8)}} & \mathbf{W}_\mu^+ & D_\mu^+ \\ \mathbf{W}_\mu^- & -\mathbf{W}_\mu^{(3)+1/\sqrt{3}\mathbf{B}_\mu^{(8)}} & D_\mu^0 \\ D_\mu^- & D_\mu^{0\dagger} & 2/\sqrt{3}\mathbf{B}_\mu^{(8)} \end{bmatrix} \quad \text{and} \quad A_5 = \begin{bmatrix} 0 & 0 & \mathbf{H}^+ \\ 0 & 0 & \mathbf{H}^0 \\ \mathbf{H}^- & \mathbf{H}^{0\dagger} & 0 \end{bmatrix} \quad (\text{B.3})$$

where W and B are towers with a zero mode, D are massive gauge bosons and H is the Higgs field (only the zero mode). We assume that the radiative potential will generate a VEV for the Higgs

$$\langle H^0 \rangle = \frac{V}{\sqrt{2}} \frac{1}{w(y)}, \quad (\text{B.4})$$

where V is a constant and the y dependence is encoded in the metric factor w . The presence of this VEV will affect the bulk equation of motions for all fields: however, being H part of gauge fields, we can use an $SU(3)$ gauge transformation to remove the VEV from the bulk equation of motions, and cast it into the boundary conditions [45]. For the gauge bosons, we can define:

$$\begin{aligned} \tilde{A}_M &= \Omega(y) A_M \Omega^\dagger(y) - \frac{i}{g_5} \Omega(y) \partial_M \Omega^\dagger(y) \\ \text{so that} \quad \langle \tilde{A}_5 \rangle &= \Omega(y) \langle A_5 \rangle \Omega^\dagger(y) - \frac{i}{g_5} \Omega(y) \partial_y \Omega^\dagger(y) = 0. \end{aligned} \quad (\text{B.5})$$

The gauge transformation that does this job can be written as:

$$\Omega(y) = \exp \left[i g_5 v / 2 \int_{y_1}^y dy' \frac{1}{w(y')} \lambda_7 \right], \quad (\text{B.6})$$

where λ_7 is the generator of $SU(3)$ aligned with H^0 . Note that we fixed the gauge transformation such that it only affects one brane: in fact $\Omega(y_1) = 1$, and

$$\Omega(y_2) = \exp [i\pi\alpha\lambda_7] = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \pi\alpha & i \sin \pi\alpha \\ 0 & i \sin \pi\alpha & \cos \pi\alpha \end{pmatrix}, \quad (\text{B.7})$$

where

$$\alpha = \frac{g_5 V}{2} \int_{y_1}^{y_2} \frac{dy}{\pi} \frac{1}{w(y)} \quad (\text{B.8})$$

is a dimensionless parameter proportional to the Higgs VEV V . The equations of motion of the new fields do not depend on the Higgs VEV, however the boundary conditions on one end will be affected. For example, for the charged gauge bosons, the gauge transformation will mix W^+ and D^+ , which have respectively Neumann and Dirichlet boundary conditions on both endpoints: in the new basis

$$\begin{cases} D_\mu^+(y_1) = \tilde{D}_\mu^+(y_1) = 0 \\ \partial_5 W_\mu^+(y_1) = \partial_5 \tilde{W}_\mu^+(y_1) = 0 \end{cases} \quad \text{and} \quad \begin{cases} D_\mu^+(y_2) = \cos \pi\alpha \tilde{D}_\mu^+ + i \sin \pi\alpha \tilde{D}_\mu^+ = 0 \\ \partial_5 W_\mu^+(y_2) = \cos \pi\alpha \partial_5 \tilde{W}_\mu^+ + i \sin \pi\alpha \partial_5 \tilde{D}_\mu^+ = 0 \end{cases}$$

Eq. (B.8) can be used to calculate δ_v , however we need to first identify the 4-dimensional VEV v . The physical Higgs field has, with a good approximation, the same profile as the 5D VEV V , therefore the couplings of the Higgs can be calculated by replacing $V \rightarrow V + h/N$, where N is the normalisation of the Higgs wave function:

$$N^2 = \int_{y_1}^{y_2} dy w(y) \cdot \frac{1}{w(y)^2}, \quad (\text{B.9})$$

and $v = VN$. Therefore, α can be written in terms of the 4-dimensional VEV v as:

$$\alpha = \frac{g_5 v N}{2\pi} = \frac{g_4 v}{2\pi} N \sqrt{\mathcal{V}}, \quad \text{where} \quad \mathcal{V} = \int_{y_1}^{y_2} dy w(y) \quad (\text{B.10})$$

is the volume of the extra dimension, and g_4 is the 4-dimensional gauge coupling (equal to the SM one up to electroweak precision corrections). Therefore:

$$\frac{v_{\text{SM}}}{v} = 1 - \delta_v = \frac{g_4 v_{\text{SM}}}{2\pi\alpha} N \sqrt{\mathcal{V}} = \frac{m_W N \sqrt{\mathcal{V}}}{\pi\alpha}. \quad (\text{B.11})$$

The specific form of the spectrum depends on the metric: in the flat case

$$\alpha = \frac{g_5 V}{2} \int_0^{\pi L} \frac{dy}{\pi} = \frac{g_5 L}{2} V, \quad (\text{B.12})$$

and

$$\begin{Bmatrix} \tilde{W}(x, y) \\ \tilde{D}(x, y) \end{Bmatrix} = \sum_n \begin{Bmatrix} (A_n \cos m_n y + B_n \sin m_n y) \\ (C_n \cos m_n y + D_n \sin m_n y) \end{Bmatrix} W_n(x). \quad (\text{B.13})$$

Applying the boundary conditions to such wave functions, we obtain that the spectrum is determined by the solutions of the following equation:

$$\sin \pi (m_n L \pm \alpha) = 0. \quad (\text{B.14})$$

Given $m_W L = \alpha$, and $N^2 = \mathcal{V} = \pi L$:

$$\delta_v = 1 - \frac{\pi m_W L}{\pi \alpha} = 0. \quad (\text{B.15})$$

The warped case is more complicated because the solutions of the equations of motion are Bessel functions of the first and second kind of order 1:

$$\alpha = \frac{g_5 V}{2} \int_{1/\Lambda}^{R'} \frac{dy}{\pi} \frac{y}{R} = \frac{g_5 R'^2}{4\pi R} V \left(1 - \frac{1}{(\Lambda R')^2} \right), \quad (\text{B.16})$$

and

$$\left\{ \begin{array}{l} \tilde{W}(x, y) \\ \tilde{D}(x, y) \end{array} \right\} = \sum_n \left\{ \begin{array}{l} y(A_n J_1(m_n y) + B_n Y_1(m_n y)) \\ y(C_n J_1(m_n y) + D_n Y_1(m_n y)) \end{array} \right\} W_n(x). \quad (\text{B.17})$$

Finally

$$1 - \delta_v = \frac{m_W R'}{\pi \alpha} \sqrt{\frac{\log \Lambda R'}{2}} \simeq \frac{\sin \pi \alpha}{\pi \alpha}, \quad (\text{B.18})$$

where we used that $N \simeq R'/\sqrt{2R}$, $\mathcal{V} = R \log \Lambda R'$ and

$$m_W R' \simeq \sqrt{\frac{2}{\log \lambda R'}} \sin \pi \alpha. \quad (\text{B.19})$$

B.2 Brane Higgs models

In these models, the Higgs boson is a 4D field which couples with the 5D gauge bulk field only on a boundary, so that the Higgs VEV only enters in the boundary conditions. We will first focus on the case where the bulk gauge symmetry is the same as in the SM, without extra fields that mix with the W : the action in the bulk is the same as in (B.2) and the 5D field can be KK decomposed as we have done before. The boundary conditions on the two endpoints can be written as (here we assume the Higgs localised on y_2 , but the results do not depend on this choice)

$$\left\{ \begin{array}{l} \partial_5 W_\mu^+(y_1) = 0 \\ \partial_5 W_\mu^+(y_2) - \frac{g_5^2 v^2}{4w(y_2)} W_\mu^+(y_2) = 0 \end{array} \right. \quad (\text{B.20})$$

where v is the Higgs VEV and g_5 the 5D gauge coupling. If we decompose the 5D fields as usual

$$W^+(y, x) = \sum_n f_n(m_n y) W_n(x)^+, \quad (\text{B.21})$$

the second boundary condition determines the spectrum as the solutions of the equation

$$m_n \frac{f'(m_n y_2)}{f(m_n y_2)} - \frac{g_5^2 v^2}{4w(y_2)} = 0. \quad (\text{B.22})$$

The precise form of the function f depends on the geometry after imposing the boundary condition on the other endpoint. The Higgs VEV can be written in terms of the SM one as:

$$\frac{v_{\text{SM}}}{v} = 1 - \delta_v = \frac{g_5 v_{\text{SM}}}{2\sqrt{w(y_2)}} \sqrt{\frac{f(m_W y_2)}{m_W f'(m_W y_2)}} = \sqrt{\frac{m_W \mathcal{V}}{w(y_2)}} \sqrt{\frac{f(m_W y_2)}{f'(m_W y_2)}}. \quad (\text{B.23})$$

In the flat case

$$f(m_n y) = \cos(m_n y) \Rightarrow \pi L m_n \tan \pi L m_n - \pi^2 \alpha^2 = 0, \quad (\text{B.24})$$

where we have defined for convenience

$$\alpha = \sqrt{\frac{L}{\pi}} \frac{g_5 V}{2}. \quad (\text{B.25})$$

Eq. (B.23) gives

$$\delta_v = 1 - \sqrt{m_W \pi L \cot(m_W \pi L)} \simeq \frac{\pi^2}{6} (m_W L)^2. \quad (\text{B.26})$$

In the warped case

$$f(m_n y) = y(Y_0(m_n R)J_1(m_n y) - J_0(m_n R)Y_1(m_n y)), \quad (\text{B.27})$$

and

$$\alpha = \frac{g_5 v R'}{2\sqrt{R}}. \quad (\text{B.28})$$

Expanding eq. (B.23) we obtain

$$\delta_v \simeq \frac{\alpha^2}{4}. \quad (\text{B.29})$$

Note finally that the simple form of eq. (B.22) allows us to calculate the couplings of the n -th mode to the Higgs as a function of the mass, even though the mass cannot be explicitly calculated: in fact, taking the total derivative with respect to V and eliminating V by using eq. (B.22), we obtain

$$\frac{v}{m_n} \frac{\partial m_n}{\partial v} = \frac{2 \frac{f'}{f}}{\frac{f'}{f} + m_n y_2 \left(\frac{f''}{f} - \left(\frac{f'}{f} \right)^2 \right)}. \quad (\text{B.30})$$

By studying this expression numerically or in an expansion for small α , we found that the sum rule

$$\sum_n \frac{v}{m_n} \frac{\partial m_n}{\partial v} = 1, \quad (\text{B.31})$$

where we are summing over all the mass eigenstates, is respected both in the flat and warped case.

C Fermionic fields

Here we are considering the minimal 5D bulk action for a fermionic field Ψ :

$$\mathcal{S} = \int d^4x \int_{y_1}^{y_2} dy w(y)^4 \left[\frac{i}{2} (\bar{\Psi} \Gamma^M \partial_M \Psi - \partial_M \bar{\Psi} \Gamma^M \Psi) - w(y) \tilde{M} \bar{\Psi} \Psi \right] \quad (\text{C.1})$$

where Γ^M with $M = 1 \dots 5$ are the five Dirac 4×4 matrices for 5D representation of the Clifford Algebra and \tilde{M} is the odd bulk mass of the 5D fermion. We remind that in 5D, the irreducible Lorentz representation for Ψ is a Dirac spinor, which is not chiral. For convenience we can use the Weyl spinor notation $\Psi = \begin{pmatrix} \chi \\ \bar{\psi} \end{pmatrix}$; thus, the equations of motion for the fermionic bulk fields are given by:

$$\begin{cases} -i\bar{\sigma}^\mu \partial_\mu \chi - \partial_5 \bar{\psi} + \left(w\tilde{M} - 2\frac{w'}{w} \right) \bar{\psi} = 0 \\ -i\sigma^\mu \partial_\mu \bar{\psi} + \partial_5 \chi + \left(w\tilde{M} + 2\frac{w'}{w} \right) \chi = 0 \end{cases} \quad (\text{C.2})$$

The next step consists in using the KK decomposition of the 5D spinors to extract the evolution along the fifth dimension. The components χ and ψ are defined by:

$$\chi(x, y) = \sum_n g_n(y) \chi_n(x) \quad \text{and} \quad \bar{\psi}(x, y) = \sum_n f_n(y) \bar{\psi}_n(x), \quad (\text{C.3})$$

where $\chi_n(x)$ and $\bar{\psi}_n(x)$ are the two 4D-components of the Dirac field with the mass m_n and satisfying usual 4D Dirac equations

$$\begin{cases} -i\bar{\sigma}^\mu \partial_\mu \chi^{(n)} + m_n \bar{\psi}^{(n)} = 0 \\ -i\sigma^\mu \partial_\mu \bar{\psi}^{(n)} + m_n \chi^{(n)} = 0 \end{cases} \quad (\text{C.4})$$

The wave functions therefore will satisfy the following equations

$$\text{flat case:} \implies \begin{cases} g'_n + \tilde{M} g_n - m_n f_n = 0 \\ f'_n - \tilde{M} f_n + m_n g_n = 0 \end{cases} \quad (\text{C.5})$$

$$\text{AdS case:} \implies \begin{cases} g'_n + \frac{c-2}{y} g_n - m_n f_n = 0 \\ f'_n - \frac{c+2}{y} f_n + m_n g_n = 0 \end{cases} \quad (\text{C.6})$$

where we have defined $c = \tilde{M}R$ in the warped case. The solutions of those equations in the flat case will be combinations of $\sin(\sqrt{m_n^2 - \tilde{M}^2}y)$ and $\cos(\sqrt{m_n^2 - \tilde{M}^2}y)$ (which become hyperbolic for the massless/light mode). In the warped case we have Bessel functions $J_{1/2 \pm c}(m_n y)$ and $J_{-1/2 \pm c}(m_n y)$ [46].

Finally we have to consider models with chiral SM fermions. This is achieved by taking boundary conditions for the Dirac fields which allow light chiral zero modes:

$$\left. \begin{array}{l} \text{Left-handed} \\ \text{fermion} \end{array} \right\} \rightarrow \psi \Big|_{y_1, y_2} = 0 \quad \left\| \quad \begin{array}{l} \text{Right-handed} \\ \text{fermion} \end{array} \right\} \rightarrow \chi \Big|_{y_1, y_2} = 0 \quad (\text{C.7})$$

To complete the description of fermions and to relate it to SM phenomenology, we need to introduce two bulk fields, a singlet Ψ_R with a right-handed zero mode and a doublet of $SU(2)$ Ψ_L with a left-handed zero mode, and their couplings with the Higgs boson. From here, we need to specify some properties of the 5D models.

C.1 Gauge Higgs unification models

In this case, the singlet Ψ_R and the doublet Ψ_L are embedded in the same bulk field, a representation of the larger bulk gauge symmetry. Consequently the odd bulk mass \tilde{M} is the same for the doublet and the singlet components. The interaction with the Higgs boson appears in the covariant derivative of Ψ in the kinetic term. This additional term in the action is given by $-ig_5 \bar{\Psi} \Gamma^5 A_5 \Psi$ in the bulk: the bulk Yukawa coupling is therefore proportional to the gauge coupling g_5 and the proportionality factor depends on the specific representation of Ψ . This term will modify the bulk equations of motion: however, as in the gauge boson case, we can use a gauge transformation to remove the Higgs VEV, and recast its effects on one of the boundary conditions.

Here we will focus on the SU(3) case described in the text for simplicity. The gauge transformed fields on the y_2 brane are

$$\tilde{\Psi}(y_2) = \Omega_f(y_2) \Psi(y_2) \quad \text{where} \quad \Omega_f(y_2) = \exp \left[i \pi \alpha \tilde{\lambda}_7 \right], \quad (\text{C.8})$$

where $\tilde{\lambda}_7$ is the SU(3) generator in the representation of Ψ . The matrix Ω_f will mix the singlet and the component of the doublet which picks up a mass (for simplicity we will denote it with Ψ_L). The mixing angle however, is not α in general: in fact it will depend on the representation of the bulk field, and the proportionality factor can be calculated by explicitly computing the generator $\tilde{\lambda}_7$ for the bulk fermion representation. In general, we will define a new parameter β to describe the mixing. Note that in the case of a bulk fundamental, Ω is the same as the one used for the gauge bosons in the previous section, therefore $\beta(\mathbf{3}) = \alpha$. The new boundary conditions for the transformed fields are

$$\begin{cases} \psi_L(y_2) = \cos \pi \beta \tilde{\psi}_L(y_2) - i \sin \pi \beta \tilde{\psi}_R(y_2) = 0 \\ \chi_R(y_2) = \cos \pi \beta \tilde{\chi}_R(y_2) - i \sin \pi \beta \tilde{\chi}_L(y_2) = 0 \end{cases} \quad (\text{C.9})$$

This boundary conditions will determine the spectrum: for instance, the spectrum m_n in the flat case is given by the solutions of

$$-\cos 2\pi L \sqrt{-\tilde{M}^2 + m_n^2} + \cos 2\pi \beta + 2 \frac{\tilde{M}^2}{m_n^2} \sin^2 \pi \beta = 0. \quad (\text{C.10})$$

C.2 Brane Yukawas

Fermionic masses can also be generated by Yukawa couplings localised on an endpoint of the extra dimension: this is possible both in the bulk Higgs model and in the localised Higgs case. Like in the Gauge Higgs case, the Higgs VEV only enters in the boundary conditions. However, boundary conditions for fermions are more tricky than for bosons, due to the fact that the equations of motion are first order differential equations: therefore how the VEV enters the boundary conditions depends crucially on the localisation mechanism for the Higgs field or for the Yukawa couplings (see ref. [46]). Here we will consider the simplest possibility: that the boundary conditions are linear in the Higgs VEV:

$$\begin{cases} \psi_L - yvL\psi_R = 0 \\ \chi_R + yvL\chi_L = 0 \end{cases} \quad (\text{C.11})$$

Those boundary conditions are the same as in the gauge Higgs case if we identify $\tan \pi\beta = yvL$ (and removing the i with a phase redefinition of the fields). The only difference is that β is not proportional to the Higgs VEV, therefore additional corrections to the couplings will arise. Another novelty is that the singlet and doublet fields are part of different bulk fields, therefore they can have different bulk masses \tilde{M}_L and \tilde{M}_R .

References

- [1] M. Pieri, S. Bhattacharya, I. Fisk, J. Letts, V. Litvin and J.G. Branson, *Inclusive search for the Higgs boson in the $H \rightarrow \gamma\gamma$ channel*, CERN-CMS-NOTE-2006-112 [SPIRES]; ATLAS collaboration, L. Carminati, *Search for a Standard Model Higgs boson in the $H \rightarrow \gamma\gamma$ channel with the ATLAS detector*, *Acta Phys. Polon.* **B 38** (2007) 747 [SPIRES].
- [2] M. Dührssen et al., *Extracting Higgs boson couplings from LHC data*, *Phys. Rev.* **D 70** (2004) 113009 [hep-ph/0406323] [SPIRES].
- [3] M. Spira, A. Djouadi, D. Graudenz and P.M. Zerwas, *Higgs boson production at the LHC*, *Nucl. Phys.* **B 453** (1995) 17 [hep-ph/9504378] [SPIRES].
- [4] G.D. Kribs, T. Plehn, M. Spannowsky and T.M.P. Tait, *Four generations and Higgs physics*, *Phys. Rev.* **D 76** (2007) 075016 [arXiv:0706.3718] [SPIRES].
- [5] PARTICLE DATA GROUP collaboration, C. Amsler et al., *Review of particle physics*, *Phys. Lett.* **B 667** (2008) 1 [SPIRES].
- [6] M.S. Carena, S. Heinemeyer, C.E.M. Wagner and G. Weiglein, *Suggestions for benchmark scenarios for MSSM Higgs boson searches at hadron colliders*, *Eur. Phys. J.* **C 26** (2003) 601 [hep-ph/0202167] [SPIRES].
- [7] S.P. Martin, *A supersymmetry primer*, hep-ph/9709356 [SPIRES].
- [8] A. Dobado, M.J. Herrero and S. Penaranda, *The Higgs sector of the MSSM in the decoupling limit*, *Eur. Phys. J.* **C 17** (2000) 487 [hep-ph/0002134] [SPIRES].
- [9] M. Perelstein and C. Spethmann, *A collider signature of the supersymmetric golden region*, *JHEP* **04** (2007) 070 [hep-ph/0702038] [SPIRES].
- [10] I. Low and S. Shalgar, *Implications of the Higgs discovery in the MSSM golden region*, *JHEP* **04** (2009) 091 [arXiv:0901.0266] [SPIRES].
- [11] N. Arkani-Hamed, A.G. Cohen and H. Georgi, *Electroweak symmetry breaking from dimensional deconstruction*, *Phys. Lett.* **B 513** (2001) 232 [hep-ph/0105239] [SPIRES].
- [12] M. Schmaltz and D. Tucker-Smith, *Little Higgs review*, *Ann. Rev. Nucl. Part. Sci.* **55** (2005) 229 [hep-ph/0502182] [SPIRES].
- [13] M. Schmaltz, *The simplest little Higgs*, *JHEP* **08** (2004) 056 [hep-ph/0407143] [SPIRES].
- [14] G. Marandella, C. Schappacher and A. Strumia, *Little-Higgs corrections to precision data after LEP2*, *Phys. Rev.* **D 72** (2005) 035014 [hep-ph/0502096] [SPIRES].
- [15] I. Low, *T parity and the littlest Higgs*, *JHEP* **10** (2004) 067 [hep-ph/0409025] [SPIRES].
- [16] N. Arkani-Hamed, A.G. Cohen, E. Katz and A.E. Nelson, *The littlest Higgs*, *JHEP* **07** (2002) 034 [hep-ph/0206021] [SPIRES].

- [17] T. Han, H.E. Logan, B. McElrath and L.-T. Wang, *Loop induced decays of the little Higgs: $h \rightarrow gg, \gamma\gamma$* , *Phys. Lett. B* **563** (2003) 191 [Erratum *ibid.* **B 603** (2004) 257] [[hep-ph/0302188](#)] [[SPIRES](#)].
- [18] J. Hubisz, P. Meade, A. Noble and M. Perelstein, *Electroweak precision constraints on the lightest Higgs model with T parity*, *JHEP* **01** (2006) 135 [[hep-ph/0506042](#)] [[SPIRES](#)].
- [19] A.V. Manohar and M.B. Wise, *Flavor changing neutral currents, an extended scalar sector and the Higgs production rate at the LHC*, *Phys. Rev. D* **74** (2006) 035009 [[hep-ph/0606172](#)] [[SPIRES](#)].
- [20] B. Grinstein, D. O'Connell and M.B. Wise, *The Lee-Wick standard model*, *Phys. Rev. D* **77** (2008) 025012 [[arXiv:0704.1845](#)] [[SPIRES](#)].
- [21] F. Krauss, T.E.J. Underwood and R. Zwicky, *The Process $gg \rightarrow h_0 \rightarrow \gamma\gamma$ in the Lee-Wick Standard Model*, *Phys. Rev. D* **77** (2008) 015012 [[arXiv:0709.4054](#)] [[SPIRES](#)].
- [22] T.E.J. Underwood and R. Zwicky, *Electroweak precision data and the Lee-Wick Standard Model*, *Phys. Rev. D* **79** (2009) 035016 [[arXiv:0805.3296](#)] [[SPIRES](#)].
- [23] R. Barbieri, L.J. Hall and Y. Nomura, *A constrained Standard Model from a compact extra dimension*, *Phys. Rev. D* **63** (2001) 105007 [[hep-ph/0011311](#)] [[SPIRES](#)].
- [24] G. Cacciapaglia, M. Cirelli and G. Cristadoro, *Gluon fusion production of the Higgs boson in a calculable model with one extra dimension*, *Phys. Lett. B* **531** (2002) 105 [[hep-ph/0111287](#)] [[SPIRES](#)].
- [25] T. Appelquist, H.-C. Cheng and B.A. Dobrescu, *Bounds on universal extra dimensions*, *Phys. Rev. D* **64** (2001) 035002 [[hep-ph/0012100](#)] [[SPIRES](#)].
- [26] G. Cacciapaglia, C. Csáki, G. Marandella and J. Terning, *The gaugephobic Higgs*, *JHEP* **02** (2007) 036 [[hep-ph/0611358](#)] [[SPIRES](#)].
- [27] L. Randall and R. Sundrum, *A large mass hierarchy from a small extra dimension*, *Phys. Rev. Lett.* **83** (1999) 3370 [[hep-ph/9905221](#)] [[SPIRES](#)];
C. Csáki, J. Erlich and J. Terning, *The effective Lagrangian in the Randall-Sundrum model and electroweak physics*, *Phys. Rev. D* **66** (2002) 064021 [[hep-ph/0203034](#)] [[SPIRES](#)].
- [28] Y. Hosotani, *Dynamical mass generation by compact extra dimensions*, *Phys. Lett. B* **126** (1983) 309 [[SPIRES](#)].
- [29] G. Cacciapaglia, C. Csáki and S.C. Park, *Fully radiative electroweak symmetry breaking*, *JHEP* **03** (2006) 099 [[hep-ph/0510366](#)] [[SPIRES](#)].
- [30] G. Panico, M. Serone and A. Wulzer, *A model of electroweak symmetry breaking from a fifth dimension*, *Nucl. Phys. B* **739** (2006) 186 [[hep-ph/0510373](#)] [[SPIRES](#)].
- [31] N. Maru and N. Okada, *Gauge-Higgs unification at LHC*, *Phys. Rev. D* **77** (2008) 055010 [[arXiv:0711.2589](#)] [[SPIRES](#)].
- [32] K. Agashe, R. Contino and A. Pomarol, *The minimal composite Higgs model*, *Nucl. Phys. B* **719** (2005) 165 [[hep-ph/0412089](#)] [[SPIRES](#)];
K. Agashe and R. Contino, *The minimal composite Higgs model and electroweak precision tests*, *Nucl. Phys. B* **742** (2006) 59 [[hep-ph/0510164](#)] [[SPIRES](#)].
- [33] A. Djouadi and G. Moreau, *Higgs production at the LHC in warped extra-dimensional models*, *Phys. Lett. B* **660** (2008) 67 [[arXiv:0707.3800](#)] [[SPIRES](#)].

- [34] A. Falkowski, *Pseudo-Goldstone Higgs production via gluon fusion*, *Phys. Rev. D* **77** (2008) 055018 [[arXiv:0711.0828](#)] [[SPIRES](#)].
- [35] For an explicit calculation see N. Maru, *Finite gluon fusion amplitude in the gauge-Higgs unification*, *Mod. Phys. Lett. A* **23** (2008) 2737 [[arXiv:0803.0380](#)] [[SPIRES](#)].
- [36] G. Panico, M. Serone and A. Wulzer, *Electroweak symmetry breaking and precision tests with a fifth dimension*, *Nucl. Phys. B* **762** (2007) 189 [[hep-ph/0605292](#)] [[SPIRES](#)].
- [37] K. Agashe, A. Delgado, M.J. May and R. Sundrum, *RS1, custodial isospin and precision tests*, *JHEP* **08** (2003) 050 [[hep-ph/0308036](#)] [[SPIRES](#)].
- [38] C.A. Scrucca, M. Serone, L. Silvestrini and A. Wulzer, *Gauge-Higgs unification in orbifold models*, *JHEP* **02** (2004) 049 [[hep-th/0312267](#)] [[SPIRES](#)].
- [39] N. Arkani-Hamed and M. Schmaltz, *Hierarchies without symmetries from extra dimensions*, *Phys. Rev. D* **61** (2000) 033005 [[hep-ph/9903417](#)] [[SPIRES](#)].
- [40] Y. Grossman and M. Neubert, *Neutrino masses and mixings in non-factorizable geometry*, *Phys. Lett. B* **474** (2000) 361 [[hep-ph/9912408](#)] [[SPIRES](#)];
T. Gherghetta and A. Pomarol, *Bulk fields and supersymmetry in a slice of AdS*, *Nucl. Phys. B* **586** (2000) 141 [[hep-ph/0003129](#)] [[SPIRES](#)].
- [41] CMS collaboration, G.L. Bayatian et al., *CMS technical design report, volume II: physics performance*, *J. Phys. G* **34** (2007) 995 [[SPIRES](#)].
- [42] D.L. Rainwater and D. Zeppenfeld, *Searching for $H \rightarrow \gamma\gamma$ in weak boson fusion at the LHC*, *JHEP* **12** (1997) 005 [[hep-ph/9712271](#)] [[SPIRES](#)].
- [43] CMS collaboration, K. Mazumdar, *Search for light Higgs boson at LHC via production through weak boson fusion*, [hep-ex/0308070](#) [[SPIRES](#)];
CMS collaboration, M. Dubinin et al., *Vector boson fusion production with $H \rightarrow \gamma\gamma$* , CERN-CMS-NOTE-2006-097 [[SPIRES](#)].
- [44] For more information and recent updates see the talks at the INFN Workshop *LC08: e^+e^- Physics at the TeV scale*, Laboratori Nazionali di Frascati Italy September 22–25 2008, <http://www.lnf.infn.it/conference/lc08/>.
- [45] N. Haba, M. Harada, Y. Hosotani and Y. Kawamura, *Dynamical rearrangement of gauge symmetry on the orbifold S^1/Z_2* , *Nucl. Phys. B* **657** (2003) 169 [*Erratum ibid.* **B 669** (2003) 381] [[hep-ph/0212035](#)] [[SPIRES](#)].
- [46] C. Csaki, C. Grojean, J. Hubisz, Y. Shirman and J. Terning, *Fermions on an interval: quark and lepton masses without Higgs*, *Phys. Rev. D* **70** (2004) 015012 [[hep-ph/0310355](#)] [[SPIRES](#)].