

Finite temperature Casimir effect for massive scalar field in spacetime with extra dimensions

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

JHEP06(2009)076

(<http://iopscience.iop.org/1126-6708/2009/06/076>)

[The Table of Contents](#) and [more related content](#) is available

Download details:

IP Address: 80.92.225.132

The article was downloaded on 03/04/2010 at 09:13

Please note that [terms and conditions apply](#).

Finite temperature Casimir effect for massive scalar field in spacetime with extra dimensions

L.P. Teo

*Faculty of Information Technology, Multimedia University, Jalan Multimedia,
63100, Cyberjaya, Selangor, Malaysia*

E-mail: lptheo@mmu.edu.my

ABSTRACT: We compute the finite temperature Casimir energy for massive scalar field with general curvature coupling subject to Dirichlet or Neumann boundary conditions on the walls of a closed cylinder with arbitrary cross section, located in a background spacetime of the form $M^{d_1+1} \times \mathcal{N}^n$, where M^{d_1+1} is the $(d_1 + 1)$ -dimensional Minkowski spacetime and \mathcal{N}^n is an n -dimensional internal manifold. The Casimir force acting on a piston moving freely inside the closed cylinder is derived and it is shown that it is independent of the regularization procedure. By letting one of the chambers of the cylinder divided by the piston to be infinitely long, we obtain the Casimir force acting on two parallel plates embedded in the cylinder. It is shown that if both the plates assume Dirichlet or Neumann boundary conditions, the strength of the Casimir force is reduced by the increase in mass. Under certain conditions, the passage from massless to massive will change the nature of the force from long range to short range. Other properties of the Casimir force such as its sign, its behavior at low and high temperature, and its behavior at small and large plate separations, are found to be similar to the massless case. Explicit exact formulas and asymptotic behaviors of the Casimir force at different limits are derived. The Casimir force when one plate assumes Dirichlet boundary condition and one plate assumes Neumann boundary condition is also derived and shown to be repulsive.

KEYWORDS: Thermal Field Theory, Field Theories in Higher Dimensions

ARXIV EPRINT: [0903.3765](https://arxiv.org/abs/0903.3765)

Contents

1	Introduction	1
2	Basic formalism	4
3	The Casimir energy inside a cylinder	6
3.1	Zero temperature Casimir energy	6
3.2	Finite temperature Casimir energy	10
3.3	Special case	12
4	The Casimir force acting on a piston or two parallel plates	14
4.1	The Casimir force acting on a piston embedded in a closed cylinder or on two parallel plates embedded in an infinitely long cylinder	14
4.2	The Casimir force density acting on a pair of infinite parallel plates	22
4.3	Special case	23
5	Conclusion	29
A	High temperature asymptotic behavior of $\tilde{\Sigma}_1$	30
B	The asymptotic behavior of the Casimir force $F_{\text{Cas}}^\infty(a)$ in different limits	30

1 Introduction

In the endeavor to solve some fundamental problems in physics, such as the unification of fundamental forces and the dark energy and cosmological constant problem, it has been proposed that we should consider higher dimensional spacetime. For example, string theory [1] predicts that we live in spacetime of ten or eleven dimensions, where the extra six or seven space dimensions are curled up to a tiny invisible compact manifold. Therefore, there is a strong motivation to study physics in spacetimes with extra dimensions. Since Casimir effect is an important quantum effect, the influence of extra dimensions on Casimir effect becomes an important issue. The Casimir effect in spacetime with extra dimensions was considered in the works [2–17]. The role of Casimir effect in stabilizing extra dimensions was discussed in [18–24]. The possible role of Casimir effect as cosmological constant responsible for the observed dark energy was considered in [10, 25–30]. In [4, 5], the Casimir effect for massless scalar field in Kaluza-Klein spacetime of the form $M^{3+1} \times \mathcal{N}^n$, where M^{3+1} is the $(3+1)$ -dimensional Minkowski spacetime and \mathcal{N}^n is an n -dimensional internal manifold, was considered in the piston setting. In [14, 15], we extended the results of [4, 5] and considered the finite temperature correction to the Casimir effect. In the work [16], the




Figure 1. A movable piston inside a closed cylinder divides the cylinder into two chambers.

Casimir effect due to massive scalar field with general curvature coupling constant subject to Robin boundary conditions in spacetime of the form $M^{d_1+1} \times \mathcal{N}^n$, where M^{d_1+1} is the $(d_1 + 1)$ -dimensional Minkowski spacetime, was considered. In this paper, we generalize the work of [16] by taking into account the temperature correction. However, we restrict ourselves to only consider either Dirichlet or Neumann boundary conditions, or combinations of these two conditions. We study in detail the influences of the mass and the temperature corrections to the Casimir effect.

In the studies of Casimir effects, one of the most popular geometric configurations is the infinite parallel plate configuration since this configuration is mathematically tractable for obtaining explicit exact results. In this article, we consider a more general setup, where the parallel plates has finite instead of infinite cross section. To avoid the divergence on the boundary of the plates, we consider the case where the two parallel plates is embedded orthogonally in an infinitely long cylinder of arbitrary finite cross section. In the three dimensional case, such a setup has been considered for example in [31]. In practice, one should consider the parallel plates being embedded in a finite cylinder and take the limit where one end of the cylinder is moved to infinite distance away. Since we are only concerned with the Casimir force acting on one of the plates, one can treat the other plate as fixed. This setup has become fashionable nowadays, and is known as the piston setup [32]. In this setup, a piston is moving freely inside a closed cylinder and dividing it into two chambers (see figure 1).

The Casimir energy of this system is the sum of the Casimir energy of the left chamber, the Casimir energy of the right chamber and the Casimir energy of the region exterior to the cylinder. Except for some special cases such as when the cylinder is infinitely long and the cross section of the cylinder is a disc or a ball, the Casimir energy of the region exterior to the cylinder is not easy to be explicitly calculated. However, the Casimir energy outside the cylinder does not have any effect on the piston inside the cylinder. The Casimir force acting on the piston is the result of the interplay between the zero modes of the field in the left and the right chambers. Since our main concern is the effect of the Casimir force acting on the piston, we will not calculate the Casimir energy outside the cylinder. We would only calculate the Casimir energies of the left and right chambers. For this, it is enough to consider the Casimir energy inside a finite closed cylinder, with arbitrary cross section.

In section 2, we give the basic formalisms needed for computing the Casimir energy inside a closed cylinder due to a massive scalar field confined in a cylinder of arbitrary cross section in the background spacetime of the form $M^{d_1+1} \times \mathcal{N}^n$. In section 3, we compute the Casimir energy using exponential cut-off method. Since a field should not have quantum fluctuations in the infinite mass limit [33], we impose the condition that the Casimir energy should vanish when the mass approaches infinity. We show that this condition allows us to renormalize the Casimir energy and explicit formula of the renormalized Casimir energy is given. The result can also be interpreted as the Casimir energy in a cylindrical universe. In section 4, we consider the Casimir force in the piston setting. It is observed that the regularization and renormalization procedures do not affect the Casimir force that acts on the piston. More precisely, the Casimir force acting on the piston is the same whether we compute it using the cut-off dependent Casimir energy or the renormalized Casimir energy. From the results for piston, we deduce the formula for the Casimir force acting on a pair of parallel plates embedded orthogonally in an infinitely long cylinder. By taking the limit where the cross section of the cylinder is infinitely large, we obtain the finite temperature Casimir force density acting on a pair of infinite parallel plates in a $(d_1 + 1)$ -dimensional macroscopic Minkowski spacetime, with the presence of an n -dimensional internal manifold.

An advantage of our approach is that we obtain a formula for the Casimir force acting on a pair of parallel plates embedded in an infinitely long cylinder as a series over elementary functions, which enables us to derive some properties of the Casimir force easily. It is shown that the Casimir force is attractive if both the plates assume Dirichlet boundary conditions or both plates assume Neumann boundary conditions. Moreover, the magnitude of the Casimir force is always a decreasing function of the plate separation a and the mass m . The latter supports the assumption that quantum fluctuations vanish in the infinite mass limit. In the case where the surrounding cylinder assumes Neumann boundary conditions, we show that taking the massless limit will change the nature of the Casimir force from short range to long range. For the influence of the internal manifold, it is shown that the Casimir force is enhanced in the presence of extra dimensions. A stronger result shows that the Casimir force is an increasing function of the size of the internal manifold. By passing to the limit of infinite parallel plates, all these properties are preserved, although some of them is not obvious from the formulas for the case of infinite parallel plates.

Besides the properties of the Casimir force, we derive explicit formulas for the asymptotic behaviors of the Casimir force in different limits, such as low and high temperature, small plate separation, small mass and large cross section. It is shown that when the plate separation is small, the Casimir force is dominated by terms that are independent of mass. This shows that the effect of mass is less significant if the plate separation is small. On the other hand, the leading order term of the Casimir force is linear in temperature in the high temperature regime. In the case that the size of the internal manifold is comparable to the plate separations, the behavior of the Casimir force is quite complicated and it depends strongly on the geometry of the internal manifold.

Although we assume that the macroscopic spacetime is Minkowskian, the results of this paper can be easily generalized to the case where the macroscopic spacetime is also curved. By setting the size of the internal manifold to be zero, one can obtain the corresponding

results for spacetime without extra dimensions.

Throughout this paper, we use the units where $\hbar = c = k_B = 1$ except for the figures.

2 Basic formalism

We consider a background $(d + 1)$ -dimensional spacetime of the form $M^{d_1+1} \times \mathcal{N}^n$, where M^{d_1+1} is the $(d_1 + 1)$ -dimensional Minkowski spacetime, \mathcal{N}^n is an n -dimensional internal space, which is assumed to be a compact connected manifold without boundary and $d = d_1 + n$. Let the spacetime metric be given by

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = \eta_{\alpha\beta} dx^\alpha dx^\beta - G_{ab} dy^a dy^b,$$

$$\mu, \nu = 0, 1, \dots, d; \quad \alpha, \beta = 0, 1, \dots, d_1; \quad a, b = 1, \dots, n,$$

where $\eta_{\alpha\beta} = \text{diag}(1, -1, \dots, -1)$, $y^a = x^{d_1+a}$ for $a = 1, \dots, n$ and $G_{ab} dy^a dy^b$ is a Riemannian metric on \mathcal{N}^n . In this article, we are interested in studying the finite temperature Casimir force acting on a pair of parallel plates due to a massive scalar field $\varphi(x)$ satisfying the equation of motion

$$\left(\frac{1}{\sqrt{|g|}} \partial_\mu \sqrt{|g|} g^{\mu\nu} \partial_\nu + m^2 + \xi \mathfrak{R} \right) \varphi(x) = 0, \tag{2.1}$$

where \mathfrak{R} is the scalar curvature of the background spacetime and ξ is a coupling constant — $\xi = 0$ corresponds to minimal coupling and $\xi = (d - 1)/4d$ corresponds to conformal coupling. According to [4], a correct approach to this problem is the piston setup (see figure 1), where we first compute the Casimir force acting on a piston moving freely inside a closely cylinder and then take the limit where one end of the cylinder is moved to infinity. Notice that for the piston system, the Casimir energy is the sum of the Casimir energy of the left chamber, the Casimir energy of the right chamber and the Casimir energy of the region outside the cylinder [32], i.e.,

$$E_{\text{Cas}}^{\text{piston}} = E_{\text{Cas}}^{\text{left}} + E_{\text{Cas}}^{\text{right}} + E_{\text{Cas}}^{\text{out}}. \tag{2.2}$$

However, as pointed out in [32], the Casimir energy outside the cylinder does not contribute to the Casimir force acting on the piston since the region outside the cylinder does not have contact with the piston. On the other hand, the computation of the Casimir energy outside the cylinder is a highly nontrivial issue for cylinder of arbitrary cross section. Therefore we would not attempt to carry out this computation here. For the Casimir energies in the left and right chambers, it suffices for us to consider the Casimir energy $E_{\text{Cas}}^{\text{cyl}}(L)$ inside a closed cylinder of the form $\text{cyl} = [0, L] \times \Omega \times \mathcal{N}^n$, where L is the length of the cylinder and Ω is a simply connected domain in \mathbb{R}^{d_1-1} . The Casimir energies of the left and right chambers are then given by $E_{\text{Cas}}^{\text{left}} = E_{\text{Cas}}^{\text{cyl}}(a)$ and $E_{\text{Cas}}^{\text{right}} = E_{\text{Cas}}^{\text{cyl}}(L - a)$ respectively.

In this article, we will consider both the homogeneous and the mixed boundary conditions. For the homogeneous boundary conditions, the walls of the cylinder and the piston both assume Dirichlet boundary conditions or both assume Neumann boundary conditions. In the limit when one end of the cylinder is moved to infinite distance away, we

obtain a configuration equivalent to two Dirichlet plates embedded in a Dirichlet cylinder, or two Neumann plates embedded inside a Neumann cylinder. For the homogeneous boundary conditions, all the walls of the cylinder constituting the left (right) chamber assume the Dirichlet boundary condition or all assume the Neumann boundary conditions. If the field $\varphi(x)$ satisfies Dirichlet boundary conditions on all the walls of the cylinder $\text{cyl} = [0, L] \times \Omega \times \mathcal{N}^n$, the eigenfunctions satisfying (2.1) are given by

$$\varphi_{k,j,l}(x) = e^{-i\omega_{k,j,l}t} \sin \frac{\pi k x^1}{L} \phi_{D;j}(x^2, \dots, x^{d_1}) \Phi_l(y), \quad (2.3)$$

where $k, j \in \mathbb{N}$, $l \in \tilde{\mathbb{N}} = \mathbb{N} \cup \{0\}$. The function $\phi_{D;j}(x^2, \dots, x^{d_1})$ is an eigenfunction of the Laplace operator with Dirichlet boundary conditions on Ω , i.e.,

$$-\sum_{i=2}^{d_1} \frac{\partial^2}{\partial (x^i)^2} \phi_{D;j}(x^2, \dots, x^{d_1}) = \omega_{\Omega,D;j}^2 \phi_{D;j}(x^2, \dots, x^{d_1}),$$

$$\phi_{D;j}|_{\partial\Omega} = 0.$$

The function $\Phi_l(y)$ is an eigenfunction of the operator $-\Delta_G - \xi \mathfrak{R}_G = -\sqrt{G}^{-1} \partial_a \sqrt{G} G^{ab} \partial_b - \xi \mathfrak{R}_G$, ($\mathfrak{R}_G = -\mathfrak{R}$ is the scalar curvature of the metric $G_{ab} dy^a dy^b$) on \mathcal{N}^n with eigenvalue $\omega_{\mathcal{N};l}^2$, i.e.,

$$-(\Delta_G + \xi \mathfrak{R}_G) \Phi_l(y) = \omega_{\mathcal{N};l}^2 \Phi_l(y).$$

The eigenfrequency $\omega_{k,j,l}$ is given by

$$\omega_{k,j,l} = \sqrt{\left(\frac{\pi k}{L}\right)^2 + \omega_{\Omega,D;j}^2 + \omega_{\mathcal{N};l}^2 + m^2}.$$

Since Ω is simply connected, the eigenvalues $\omega_{\Omega,D;j}^2$ are all nonzero.

If the field $\varphi(x)$ satisfies Neumann boundary conditions on all the walls of the cylinder $\text{cyl} = [0, L] \times \Omega \times \mathcal{N}^n$, the eigenfunctions are

$$\varphi_{k,j,l}(x) = e^{-i\omega_{k,j,l}t} \cos \frac{\pi k x^1}{L} \phi_{N;j}(x^2, \dots, x^{d_1}) \Phi_l(y), \quad (2.4)$$

where $k, j, l \in \tilde{\mathbb{N}} = \mathbb{N} \cup \{0\}$. The function $\phi_{N;j}(x^2, \dots, x^{d_1})$ is an eigenfunction of the Laplace operator with Neumann boundary conditions on Ω , i.e.,

$$-\sum_{i=2}^{d_1} \frac{\partial^2}{\partial (x^i)^2} \phi_{N;j}(x^2, \dots, x^{d_1}) = \omega_{\Omega,N;j}^2 \phi_{N;j}(x^2, \dots, x^{d_1}),$$

$$\left. \frac{\partial \phi_{N;j}}{\partial \mathbf{n}} \right|_{\partial\Omega} = 0,$$

where \mathbf{n} is a unit vector perpendicular to $\partial\Omega$. By convention, $\phi_{N;0}(x^2, \dots, x^{d_1})$ is the constant function with zero eigenvalue. For the eigenvalues $\omega_{\mathcal{N};l}^2$, we assume that $\omega_{\mathcal{N};l}^2 \geq 0$ and there are exactly κ of them that are equal to zero.

For the mixed boundary conditions, we consider two types. For the first type, the field satisfies Dirichlet boundary conditions on the walls of the cylinder and Neumann boundary

conditions on the piston. For the second type, the field satisfies Neumann boundary conditions on the walls of the cylinder and Dirichlet boundary conditions on the piston. In the limit where one plate is moved to infinity, we obtain a configuration equivalent to a pair of parallel plates, one assumes Dirichlet boundary condition and one assumes Neumann boundary condition, embedded in a Dirichlet or a Neumann cylinder. For the first type of the mixed boundary conditions, we have to consider the Casimir energy inside the cylinder $\text{cyl} = [0, L] \times \Omega \times \mathcal{N}^n$, where the field $\varphi(x)$ satisfies Dirichlet boundary conditions on the wall $[0, L] \times \partial\Omega \times \mathcal{N}^n$ and the wall $x^1 = 0$, and Neumann boundary condition on the wall $x^1 = L$. In this case, the eigenfunctions are obtained from (2.3) by replacing the function $\sin \frac{\pi kx}{L}$ by $\sin \frac{\pi(k-\frac{1}{2})x}{L}$. For the second type of the mixed boundary conditions, we have to consider the Casimir energy inside the cylinder $\text{cyl} = [0, L] \times \Omega \times \mathcal{N}^n$, where the field $\varphi(x)$ satisfies Neumann boundary conditions on the wall $[0, L] \times \partial\Omega \times \mathcal{N}^n$ and the wall $x^1 = 0$, and Dirichlet boundary condition on the wall $x^1 = L$. In this case, the eigenfunctions are obtained from (2.4) by replacing $\cos \frac{\pi kx}{L}$ by $\cos \frac{\pi(k+\frac{1}{2})x}{L}$.

Before ending this section, we define the variables R and r by $R = \text{Vol}(\Omega)^{1/(d_1-1)}$ and $r = \text{Vol}(\mathcal{N})^{1/n}$ which have the dimension of length to measure the size of the domain Ω and the internal manifold \mathcal{N}^n . Throughout this article, we assume that the size r of the internal manifold is smaller than any measurable length in the $(d_1 + 1)$ -dimensional Minkowski spacetime. The re-scaled variables

$$\begin{aligned} \omega'_{\Omega,*,j} &= R\omega_{\Omega,*,j}, \quad * = D \text{ or } N, \\ \omega'_{\mathcal{N};l} &= r\omega_{\mathcal{N};l}, \end{aligned}$$

are dimensionless variables and are invariant under the re-scaling of the domain Ω and the manifold \mathcal{N}^n .

3 The Casimir energy inside a cylinder

In this section, we compute the Casimir energy inside a closed cylinder of the form $[0, L] \times \Omega \times \mathcal{N}^n$, which is needed for computing the Casimir energy of the left and right chambers of the piston system figure 1.

3.1 Zero temperature Casimir energy

At zero temperature, the Casimir energy is defined naively as the sum of zero point energies:

$$E_{\text{Cas}}^{T=0}(L) = \frac{1}{2} \sum \omega_{k,j,l}.$$

To take into account the homogeneous and mixed boundary conditions, we let

$$\omega_{k,j,l} = \omega_{k,j,l}(\alpha, *, m) = \sqrt{\left(\frac{\pi(k+\alpha)}{L}\right)^2 + \omega_{\Omega,*,j}^2 + \omega_{\mathcal{N};l}^2 + m^2}, \quad k \in \tilde{\mathbb{N}}, j \in J_*, l \in \tilde{\mathbb{N}},$$

where $\alpha = 1$ for homogeneous Dirichlet boundary conditions, $\alpha = 0$ for homogeneous Neumann boundary conditions and $\alpha = 1/2$ for mixed boundary conditions; $*$ = D (resp.

$*$ = N) when $\varphi(x)$ assume Dirichlet (resp. Neumann) boundary conditions on the component $[0, L] \times \partial\Omega \times \mathcal{N}^n$ of the boundary of the cylinder; $J_D = \mathbb{N}$ and $J_N = \tilde{\mathbb{N}}$. Using exponential cut-off regularization, we define the cut-off dependent energy as

$$E_{\text{Cas}}^{T=0}(L; \lambda) = \frac{1}{2} \sum_{k=0}^{\infty} \sum_{j \in J_*} \sum_{l=0}^{\infty} \omega_{k,j,l} e^{-\lambda \omega_{k,j,l}}.$$

Using the same method as the massless case [15], we find that up to the term λ^0 , the small- λ expansion of $E_{\text{Cas}}^{T=0}(L; \lambda)$ is given by

$$E_{\text{Cas}}^{T=0}(L; \lambda) = \sum_{i=0}^{d-1} \frac{\Gamma(d+1-i)}{\Gamma(\frac{d-i}{2})} c_{\text{cyl},\alpha,*,i}(m) \lambda^{i-d-1} - \frac{\psi(1) - \log \lambda}{2\sqrt{\pi}} c_{\text{cyl},\alpha,*,d+1}(m) + \frac{1}{2} \text{FP}_{s=-\frac{1}{2}} \zeta_{\text{cyl},\alpha,*}(s; m), \tag{3.1}$$

where $c_{\text{cyl},\alpha,*,i}(m)$ are the heat kernel coefficients defined so that

$$\sum_{k=0}^{\infty} \sum_{j \in J_*} \sum_{l=0}^{\infty} e^{-t\omega_{k,j,l}^2} = \sum_{i=0}^{M-1} c_{\text{cyl},\alpha,*,i}(m) t^{\frac{i-d}{2}} + O\left(t^{\frac{M-d}{2}}\right) \quad \text{as } t \rightarrow 0^+, \tag{3.2}$$

$\zeta_{\text{cyl},\alpha,*}(s; m)$ is the zeta function

$$\zeta_{\text{cyl},\alpha,*}(s; m) = \sum_{k=0}^{\infty} \sum_{j \in J_*} \sum_{l=0}^{\infty} \omega_{k,j,l}^{-2s},$$

and the finite part of a meromorphic function $f(z)$ with at most simple pole at a point $z = z_0$ is defined by

$$\text{FP}_{z=z_0} f(z) = \lim_{z \rightarrow z_0} \left(f(z) - \frac{\text{Res}_{z=z_0} f(z)}{z - z_0} \right).$$

As $\lambda \rightarrow 0^+$, we see from (3.1) that the divergent part of the Casimir energy contains divergence of order $\log \lambda$ and λ^{-i} , $i = 2, \dots, d+1$, with coefficients depending on the coefficients $c_{\text{cyl},\alpha,*,i}(m)$, which can be expressed in terms of L , m and the geometric invariants of the manifolds Ω and \mathcal{N}^n . In particular, the leading divergence

$$\frac{\Gamma(d+1)}{\Gamma(\frac{d}{2})} c_{\text{cyl},\alpha,*,0}(m) \lambda^{-d-1} = \frac{\Gamma(d+1)}{\Gamma(\frac{d}{2})} \frac{L \text{Vol}(\Omega) \text{Vol}(\mathcal{N}^n)}{2^d \pi^{\frac{d}{2}}} \lambda^{-d-1}$$

is called the bulk divergence and is usually subtracted away in the definition of Casimir energy. The other divergences are called hypersurface divergences and regularization is required to remove these divergences. A conventional method, known as zeta regularization [34], set all the hypersurface divergences to zero and define the regularized zero temperature Casimir energy to be

$$E_{\text{Cas}}^{\text{reg}_{\zeta}, T=0}(L) = \frac{1}{2} \text{FP}_{s=-\frac{1}{2}} \zeta_{\text{cyl},\alpha,*}(s; m) + \frac{1}{2} \log \mu^2 \text{Res}_{s=-\frac{1}{2}} \zeta_{\text{cyl},\alpha,*}(s; m) = \frac{1}{2} \text{FP}_{s=-\frac{1}{2}} \zeta_{\text{cyl},\alpha,*}(s; m) - \frac{\log \mu}{2\sqrt{\pi}} c_{\text{cyl},\alpha,*,d+1}(m), \tag{3.3}$$

where μ is a normalization constant with dimension length^{-1} . This is tantamount to subtracting the divergent terms

$$\sum_{i=0}^{d-1} \frac{\Gamma(d+1-i)}{\Gamma\left(\frac{d-i}{2}\right)} c_{\text{cyl},\alpha,*,i}(m) \lambda^{i-d-1} - \frac{\psi(1) - \log(\lambda\mu)}{2\sqrt{\pi}} c_{\text{cyl},\alpha,*,d+1}(m) \quad (3.4)$$

from the cut-off dependent Casimir energy (3.1) and set $\lambda = 0$. However, for massive scalar fields, this definition is not sufficient. Since a field should not have quantum fluctuations in the limit of infinite mass [33], it is natural to renormalize the Casimir energy so that the renormalized Casimir energy $E_{\text{Cas}}^{\text{ren}}$ vanishes when the mass m approaches infinity, i.e.,

$$E_{\text{Cas}}^{\text{ren}} \xrightarrow{m \rightarrow \infty} 0.$$

To satisfy this condition, one need to extract the leading behavior of the Casimir energy in (3.1) or (3.3) when the mass is large and subtract away those terms that give nontrivial limits when the mass tends to infinity. Using the fact that

$$\sum_{k=0}^{\infty} \sum_{j \in J_*} \sum_{l=0}^{\infty} e^{-t\omega_{k,j,l}^2} = \sum_{k=0}^{\infty} \sum_{j \in J_*} \sum_{l=0}^{\infty} e^{-tm^2} \exp \left\{ -t \left(\left[\frac{\pi(k+\alpha)}{L} \right]^2 + \omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 \right) \right\},$$

it is easy to show that the heat kernel coefficients $c_{\text{cyl},\alpha,*,i}(m)$, $i = 0, 1, 2, \dots$ at any mass can be expressed as a polynomial in m^2 with coefficients in terms of the heat kernel coefficients $c_{\text{cyl},\alpha,*,d+1}(0)$ when $m = 0$. More precisely,

$$c_{\text{cyl},\alpha,*,i}(m) = \sum_{j=0}^{\lfloor \frac{i}{2} \rfloor} \frac{(-1)^j}{j!} c_{\text{cyl},\alpha,*,i-2j}(0) m^{2j}.$$

Therefore, the divergent part of the Casimir energy (3.4) can be expressed as a polynomial in m^2 . As a result, it has to be subtracted away in the renormalization procedure and thus do not contribute to the renormalized Casimir energy. For the regular part of the Casimir energy given by (3.3), the second term that proportional to $c_{\text{cyl},\alpha,*,d+1}(m)$ would not contribute to the renormalized Casimir energy for the same reason above. For the term $(1/2)\text{FP}_{s=-\frac{1}{2}} \zeta_{\text{cyl},\alpha,*}(s; m)$, let

$$\tilde{K}(t) = \sum_{k=0}^{\infty} \sum_{j \in J_*} \sum_{l=0}^{\infty} \exp \left\{ -t \left(\left[\frac{\pi(k+\alpha)}{L} \right]^2 + \omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 \right) \right\} - \sum_{i=0}^{d+1} c_{\text{cyl},\alpha,*,i}(0) t^{\frac{i-d}{2}}.$$

It can be shown that

$$\begin{aligned} \frac{1}{2} \text{FP}_{s=-\frac{1}{2}} \zeta_{\text{cyl},\alpha,*}(s; m) &= \frac{\psi\left(-\frac{1}{2}\right) + \log m^2}{4\sqrt{\pi}} c_{\text{cyl},\alpha,*,d+1}(m) \\ &+ \mathfrak{D}(m) - \frac{1}{4\sqrt{\pi}} \int_0^{\infty} t^{-\frac{3}{2}} e^{-tm^2} \tilde{K}(t) dt, \end{aligned} \quad (3.5)$$

where

$$\mathfrak{D}(m) = -\frac{1}{4\sqrt{\pi}} \sum_{i=0}^{d+1} c_{\text{cyl},\alpha,*,d+1-i}(0) m^i \text{FP}_{s=-\frac{i}{2}} \Gamma(s),$$

and

$$\frac{1}{4\sqrt{\pi}} \int_0^\infty t^{-\frac{3}{2}} e^{-tm^2} \tilde{K}(t) dt \xrightarrow{m \rightarrow \infty} 0.$$

Therefore, as $m \rightarrow \infty$,

$$\frac{1}{2} \text{FP}_{s=-\frac{1}{2}} \zeta_{\text{cyl},\alpha,*}(s; m) = \frac{\psi(-\frac{1}{2}) + \log m^2}{4\sqrt{\pi}} c_{\text{cyl},\alpha,*,d+1}(m) + \mathfrak{D}(m) + o(1).$$

This implies that to obtain the renormalized Casimir energy that vanishes when $m \rightarrow \infty$, we should subtract away the term

$$\sum_{i=0}^{d-1} \frac{\Gamma(d+1-i)}{\Gamma(\frac{d-i}{2})} c_{\text{cyl},\alpha,*,i}(m) \lambda^{i-d-1} + \frac{\psi(-\frac{1}{2}) - 2\psi(1) + \log[\lambda m]^2}{4\sqrt{\pi}} c_{\text{cyl},\alpha,*,d+1}(m) + \mathfrak{D}(m) \quad (3.6)$$

from the cut-off dependent Casimir energy (3.1). From (3.5), we then find that the renormalized Casimir energy is given by

$$\begin{aligned} E_{\text{Cas}}^{\text{ren},T=0}(L) &= \frac{1}{2} \text{FP}_{s=-\frac{1}{2}} \zeta_{\text{cyl},\alpha,*}(s; m) - \frac{\psi(-\frac{1}{2}) + \log m^2}{4\sqrt{\pi}} c_{\text{cyl},\alpha,*,d+1}(m) - \mathfrak{D}(m) \\ &= -\frac{1}{4\sqrt{\pi}} \int_0^\infty t^{-\frac{3}{2}} e^{-tm^2} \tilde{K}(t) dt. \end{aligned} \quad (3.7)$$

To study the behavior of the renormalized Casimir energy with respect to the variation of the plate separation L , we use the fact that for $\alpha = 0, 1/2, 1$,

$$\begin{aligned} \sum_{k=0}^\infty e^{-t \frac{\pi^2(k+\alpha)^2}{L^2}} &= \frac{1}{2} \sum_{k=-\infty}^\infty e^{-t \frac{\pi^2(k+\alpha)^2}{L^2}} + \frac{1}{2} - \alpha \\ &= \frac{L}{2\sqrt{\pi}} \sum_{k=-\infty}^\infty t^{-\frac{1}{2}} e^{2\pi i k \alpha} e^{-\frac{k^2 L^2}{t}} + \frac{1}{2} - \alpha. \end{aligned} \quad (3.8)$$

This implies that

$$\begin{aligned} \zeta_{\text{cyl},\alpha,*}(s; m) &= \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} \sum_{j \in J_*} \sum_{l=0}^\infty e^{-t(\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N};l}^2 + m^2)} \\ &\quad \times \left\{ \frac{L}{2\sqrt{\pi}} \sum_{k=-\infty}^\infty t^{-\frac{1}{2}} e^{2\pi i k \alpha} e^{-\frac{k^2 L^2}{t}} + \frac{1}{2} - \alpha \right\} dt \\ &= \left(\frac{1}{2} - \alpha \right) \zeta_{\Omega \times \mathcal{N},*}(s; m) + \frac{L}{2\sqrt{\pi}} \frac{\Gamma(s - \frac{1}{2})}{\Gamma(s)} \zeta_{\Omega \times \mathcal{N},*}\left(s - \frac{1}{2}; m\right) \\ &\quad + \frac{2L}{\sqrt{\pi} \Gamma(s)} \sum_{k=1}^\infty \sum_{j \in J_*} \sum_{l=0}^\infty e^{2\pi i k \alpha} \left(\frac{kL}{\sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N};l}^2 + m^2}} \right)^{s-\frac{1}{2}} \\ &\quad \times K_{s-\frac{1}{2}} \left(2kL \sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N};l}^2 + m^2} \right), \end{aligned} \quad (3.9)$$

where $K_\nu(z)$ is the modified Bessel function of second kind, and

$$\zeta_{\Omega \times \mathcal{N},*}(s; m) = \sum_{j \in J_*} \sum_{l=0}^{\infty} (\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N};l}^2 + m^2)^{-s}.$$

As a result, we have

$$\begin{aligned} E_{\text{Cas}}^{\text{ren},T=0}(L) &= \frac{1-2\alpha}{4} \text{FP}_{s=-\frac{1}{2}} \zeta_{\Omega \times \mathcal{N},*}(s; m) \\ &+ \frac{L}{8\pi} \left\{ \psi\left(-\frac{1}{2}\right) \text{Res}_{s=-1}(\Gamma(s)\zeta_{\Omega \times \mathcal{N},*}(s)) - \text{FP}_{s=-1}(\Gamma(s)\zeta_{\Omega \times \mathcal{N},*}(s)) \right\} \\ &- \frac{\psi\left(-\frac{1}{2}\right) + \log m^2}{4\sqrt{\pi}} c_{\text{cyl},\alpha,*,d+1}(m) - \mathfrak{D}(m) \\ &- \frac{1}{2\pi} \sum_{k=1}^{\infty} \sum_{j \in J_*} \sum_{l=0}^{\infty} e^{2\pi i k \alpha} \frac{\sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N};l}^2 + m^2}}{k} K_1\left(2kL\sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N};l}^2 + m^2}\right). \end{aligned} \quad (3.10)$$

Using (3.8), we also have

$$\begin{aligned} \sum_{k=0}^{\infty} \sum_{j \in J_*} \sum_{l=0}^{\infty} e^{-t\omega_{k,j,l}^2} &= \left(\frac{L}{2\sqrt{\pi}} t^{-\frac{1}{2}} + \frac{1}{2} - \alpha + \text{e.d.} \right) \\ &\times \left(\sum_{i=0}^{M-1} c_{\Omega \times \mathcal{N},*,i}(m) t^{\frac{i-d+1}{2}} + O\left(t^{\frac{M-d+1}{2}}\right) \right) \quad \text{as } t \rightarrow 0^+, \end{aligned}$$

where e.d. is the exponentially decay terms, and $c_{\Omega \times \mathcal{N},*,i}(m)$ are heat kernel coefficients for an elliptic operator on $\Omega \times \mathcal{N}$. This implies that

$$c_{\text{cyl},\alpha,*,i}(m) = \frac{L}{2\sqrt{\pi}} c_{\Omega \times \mathcal{N},*,i}(m) + \left(\frac{1}{2} - \alpha\right) c_{\Omega \times \mathcal{N},*,i-1}(m). \quad (3.11)$$

Therefore, we read from (3.10) that the renormalized Casimir energy can be written as a linear polynomial in L , plus a term that decays exponentially as L approaches infinity, i.e.,

$$\begin{aligned} E_{\text{Cas}}^{\text{ren},T=0}(L) &= \Sigma_0 + \Sigma_1 L - \frac{1}{2\pi} \sum_{k=1}^{\infty} \sum_{j \in J_*} \sum_{l=0}^{\infty} e^{2\pi i k \alpha} \frac{\sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N};l}^2 + m^2}}{k} \\ &\times K_1\left(2kL\sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N};l}^2 + m^2}\right). \end{aligned} \quad (3.12)$$

3.2 Finite temperature Casimir energy

Now we take into account the temperature correction to the Casimir energy given by

$$T \sum \log\left(1 - e^{-\omega_{k,j,l}/T}\right). \quad (3.13)$$

This summation is finite and no regularization is required. Therefore, the cut-off dependent finite temperature Casimir energy is defined as

$$E_{\text{Cas}}(L; \lambda) = \frac{1}{2} \sum_{k=0}^{\infty} \sum_{j \in J_*} \sum_{l=0}^{\infty} \omega_{k,j,l} e^{-\lambda \omega_{k,j,l}} + T \sum \log\left(1 - e^{-\omega_{k,j,l}/T}\right).$$

It is well known that this can be computed in terms of the finite temperature zeta function given by

$$\zeta_{\text{cyl},\alpha,*,T}(s; m) = \sum_{k=0}^{\infty} \sum_{j \in J_*} \sum_{l=0}^{\infty} \sum_{p=-\infty}^{\infty} e^{-t(\omega_{k,j,l}^2 + (2\pi pT)^2)}.$$

More precisely (see [15]),

$$\begin{aligned} E_{\text{Cas}}(L; \lambda) &= \sum_{i=0}^{d-1} \frac{\Gamma(d+1-i)}{\Gamma\left(\frac{d-i}{2}\right)} c_{\text{cyl},\alpha,*,i}(m) \lambda^{i-d-1} \\ &+ \frac{\log[\lambda\mu] - \psi(1) - \log 2 + 1}{2\sqrt{\pi}} c_{\text{cyl},\alpha,*,d+1}(m) \\ &- \frac{T}{2} (\zeta'_{\text{cyl},\alpha,*,T}(0; m) + \log[\mu^2] \zeta_{\text{cyl},\alpha,*,T}(0; m)). \end{aligned} \quad (3.14)$$

For the renormalization, we observe that the temperature correction to the Casimir energy (3.13) vanishes when the mass m approach infinity. Therefore, the large- m leading behavior of the cut-off dependent finite temperature Casimir energy (3.14) is the same as the large- m leading behavior of the cut-off dependent zero temperature Casimir energy given by (3.6). After subtracting away these large- m non-vanishing terms, we find that the renormalized finite temperature Casimir energy is given by

$$\begin{aligned} E_{\text{Cas}}^{\text{ren}}(L) &= -\frac{T}{2} (\zeta'_{\text{cyl},\alpha,*,T}(0; m) + \log[\mu^2] \zeta_{\text{cyl},\alpha,*,T}(0; m)) \\ &- \frac{\log\left[\frac{m}{\mu}\right]^2 + \psi(1)}{4\sqrt{\pi}} c_{\text{cyl},\alpha,*,d+1}(m) - \mathfrak{D}(m). \end{aligned} \quad (3.15)$$

As a remark, in the zeta regularization scheme, the regularized finite temperature Casimir energy is defined as

$$E_{\text{Cas}}^{\text{reg}\zeta}(L) = -\frac{T}{2} (\zeta'_{\text{cyl},\alpha,*,T}(0; m) + \log[\mu^2] \zeta_{\text{cyl},\alpha,*,T}(0; m)). \quad (3.16)$$

This expression does not go to zero as the mass m approaches infinity. We would like to remark that although the expression (3.15) for the renormalized Casimir energy is derived for the case where the field is confined within a cylindrical cavity, this formula is in fact valid for an arbitrary closed cavity. Similarly, the formula for the renormalized Casimir energy at zero temperature given by (3.7) is also valid for any closed cavity. We would also like to point out that the high temperature expansion of the zeta regularized finite temperature Casimir energy for massive scalar field in manifolds with boundaries have been obtained in [35–39].

Using the same method as in the derivation of (3.9) (see [15]), we find that

$$\begin{aligned} \zeta_{\text{cyl},\alpha,*,T}(0; m) &= \frac{c_{\text{cyl},\alpha,*,d+1}(m)}{2\sqrt{\pi}T}, \\ \zeta'_{\text{cyl},\alpha,*,T}(0; m) &= \Lambda_0 + \Lambda_1 L + \sum_{k=1}^{\infty} \sum_{j \in J_*} \sum_{l=0}^{\infty} \sum_{p=-\infty}^{\infty} \frac{e^{2\pi i k \alpha}}{k} \\ &\quad \times \exp\left(-2kL\sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2 + (2\pi pT)^2}\right), \end{aligned}$$

where Λ_0 and Λ_1 are independent of L , and Λ_1 is given by

$$\begin{aligned} \Lambda_1 = & -\frac{\psi(1)}{4\pi T} c_{\Omega \times \mathcal{N}, *, d+1}(m) + \frac{1}{4\pi T} \text{FP}_{s=-1} \{ \Gamma(s) \zeta_{\Omega \times \mathcal{N}}(s; m) \} \\ & + \frac{2}{\pi} \sum_{j \in J_*} \sum_{l=0}^{\infty} \sum_{p=1}^{\infty} \frac{\sqrt{\omega_{\Omega, *, j}^2 + \omega_{\mathcal{N}; l}^2 + m^2}}{p} K_1 \left(\frac{p \sqrt{\omega_{\Omega, *, j}^2 + \omega_{\mathcal{N}; l}^2 + m^2}}{T} \right). \end{aligned}$$

Together with (3.11), we find that the renormalized Casimir energy (3.15) can be expressed as the sum of a linear polynomial in L plus a term that decays exponentially as L approaches infinity, i.e.,

$$\begin{aligned} E_{\text{Cas}}^{\text{ren}}(L) = & \tilde{\Sigma}_0 + \tilde{\Sigma}_1 L - \frac{T}{2} \sum_{k=1}^{\infty} \sum_{j \in J_*} \sum_{l=0}^{\infty} \sum_{p=-\infty}^{\infty} \frac{e^{2\pi i k \alpha}}{k} \\ & \times \exp \left(-2kL \sqrt{\omega_{\Omega, *, j}^2 + \omega_{\mathcal{N}; l}^2 + m^2 + (2\pi p T)^2} \right), \end{aligned} \quad (3.17)$$

where

$$\begin{aligned} \tilde{\Sigma}_1 = & \frac{1}{8\pi} \sum_{i=0}^{d+1} \left[\text{FP}_{s=-\frac{i}{2}} \Gamma(s) \right] c_{\Omega \times \mathcal{N}, *, d+1-i}(0) m^i \\ & - \frac{1}{8\pi} \{ \text{FP}_{s=-1} \{ \Gamma(s) \zeta_{\Omega \times \mathcal{N}}(s; m) \} + \log[m^2] c_{\Omega \times \mathcal{N}, *, d+1}(m) \} \\ & - \frac{T}{\pi} \sum_{j \in J_*} \sum_{l=0}^{\infty} \sum_{p=1}^{\infty} \frac{\sqrt{\omega_{\Omega, *, j}^2 + \omega_{\mathcal{N}; l}^2 + m^2}}{p} K_1 \left(\frac{p \sqrt{\omega_{\Omega, *, j}^2 + \omega_{\mathcal{N}; l}^2 + m^2}}{T} \right). \end{aligned} \quad (3.18)$$

Before ending this section, we would like to comment that it is easy to deduce from the results above that the Casimir energy for mixed boundary conditions ($\alpha = 1/2$) are related to the Casimir energy for homogeneous boundary conditions ($\alpha = 0$ or 1) by

$$\begin{aligned} E_{\text{Cas}} \left(L; \alpha = \frac{1}{2}, * = D \right) &= E_{\text{Cas}}(2L; \alpha = 1, * = D) - E_{\text{Cas}}(L; \alpha = 1, * = D), \\ E_{\text{Cas}} \left(L; \alpha = \frac{1}{2}, * = N \right) &= E_{\text{Cas}}(2L; \alpha = 0, * = N) - E_{\text{Cas}}(L; \alpha = 0, * = N). \end{aligned} \quad (3.19)$$

3.3 Special case

Here we consider the special case where the cross section of the cylinder Ω is a rectangular region $[0, L_2] \times \dots \times [0, L_{d_1}]$ and the internal manifold \mathcal{N}^n is an n -torus T^n — a product of n circles with radius r_1, \dots, r_n respectively. Notice that the scalar curvature of the torus T^n is zero. The spectrum $\omega_{\Omega, *, j}^2$ of the Laplace operator with Dirichlet or Neumann boundary conditions on Ω is given by

$$\left(\frac{\pi j_2}{L_2} \right)^2 + \dots + \left(\frac{\pi j_{d_1}}{L_{d_1}} \right)^2, \quad \mathbf{j} = (j_2, \dots, j_{d_1}) \in \begin{cases} \mathbb{N}^{d_1-1}, & \text{if } * = D, \\ \tilde{\mathbb{N}}^{d_1-1}, & \text{if } * = N, \end{cases}$$

and the spectrum $\omega_{\mathcal{N};i}^2$ of the Laplace operator on T^n is given by

$$\left(\frac{l_1}{r_1}\right)^2 + \dots + \left(\frac{l_n}{r_n}\right)^2, \quad \mathbf{l} = (l_1, \dots, l_n) \in \mathbb{Z}^n.$$

For simplicity, we only consider the cases where $\alpha = 0$ or 1 , i.e., all the walls of the rectangular cavity $[0, L_1] \times \dots \times [0, L_{d_1}]$ assume Dirichlet boundary conditions or all the walls assume Neumann boundary conditions. The results for the case where $\alpha = 1/2$ can be obtained from the results for $\alpha = 0$ or 1 by (3.19). The finite temperature zeta function $\zeta_{\text{cyl},D/N;T}(m)$ can be written as a sum of inhomogeneous Epstein zeta function, i.e.,

$$\zeta_{\text{cyl},D/N;T}(s; m) = 2^{-d_1} \sum_{i=0}^{d_1} (\mp 1)^{d_1-i} \sum_{1 \leq \sigma_1 < \dots < \sigma_i \leq d_1} \times Z_{i+n+1} \left(s; \frac{\pi}{L_{\sigma_1}}, \dots, \frac{\pi}{L_{\sigma_i}}, \frac{1}{r_1}, \dots, \frac{1}{r_n}, 2\pi T; m \right),$$

where

$$Z_j(s; a_1, \dots, a_j; m) = \sum_{(k_1, \dots, k_j) \in \mathbb{Z}^j} \frac{1}{([k_1 a_1]^2 + \dots + [k_j a_j]^2 + m^2)^s}.$$

The properties of the inhomogeneous Epstein zeta function $Z_j(s; c_1, \dots, c_j; m)$ have been discussed quite extensively (see e.g. [40–43]). Using those properties, we find that the renormalized Casimir energy $E_{\text{Cas};D/N}^{\text{ren}}(L_1)$ is given by

$$\begin{aligned} E_{\text{Cas};D/N}^{\text{ren}}(L_1) &= - \frac{\prod_{l=1}^n r_l}{2^{d_1+1}} \sum_{i=0}^{d_1} \frac{(\mp 1)^{d_1-i}}{\pi^{\frac{i+1-n}{2}}} m^{\frac{i+n+1}{2}} \\ &\times \sum_{1 \leq \sigma_1 < \dots < \sigma_i \leq d_1} \left[\prod_{j=1}^i L_{\sigma_j} \right] \sum_{(k_{\sigma_1}, \dots, k_{\sigma_i}, l_1, \dots, l_n, p) \in \mathbb{Z}^{i+n+1} \setminus \{0\}} \\ &\times \left(\sum_{j=1}^i [L_{\sigma_j} k_{\sigma_j}]^2 + \sum_{j=1}^n [\pi r_j l_j]^2 + \left[\frac{p}{2T} \right]^2 \right)^{-\frac{i+n+1}{4}} \\ &\times K_{\frac{i+n+1}{2}} \left(2m \sqrt{\sum_{j=1}^i [L_{\sigma_j} k_{\sigma_j}]^2 + \sum_{j=1}^n [\pi r_j l_j]^2 + \left[\frac{p}{2T} \right]^2} \right). \end{aligned} \tag{3.20}$$

The terms with $p = 0$ in (3.20) give the zero temperature Casimir energy. The sum of the terms with $\sigma_1 \geq 2$ corresponds to the term $\tilde{\Sigma}_0$ in (3.17) which is independent of L_1 . The sum of the terms with $\sigma_1 = 1$ and $k_{\sigma_1} = k_1 = 0$ corresponds to the term $\tilde{\Sigma}_1$ in (3.17) which is proportional to L_1 . The rest of the terms decay exponentially when $L_1 \rightarrow \infty$.

4 The Casimir force acting on a piston or two parallel plates

4.1 The Casimir force acting on a piston embedded in a closed cylinder or on two parallel plates embedded in an infinitely long cylinder

The Casimir force acting the piston is obtained by differentiating the Casimir energy of the piston system (2.2) with respect to a , the position of the piston. As explained in section 2, the Casimir energy of the region outside the cylinder does not contribute. The Casimir force acting on the piston is the sum of the Casimir forces arise from the variations of the Casimir energies in the left and right chambers:

$$F_{\text{Cas}}(a; L) = -\frac{\partial}{\partial a} (E_{\text{Cas}}(a) + E_{\text{Cas}}(L - a)). \quad (4.1)$$

From (3.17), we find that the contribution to the (renormalized) Casimir force from the left chamber is given by

$$F_{\text{Cas}}^{\text{ren, left}}(a) = -\tilde{\Sigma}_1 - (-1)^{2\alpha} T \sum_{j \in J_*} \sum_{l=0}^{\infty} \sum_{p=-\infty}^{\infty} \frac{\sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2 + (2\pi pT)^2}}{\exp\left(2a\sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2 + (2\pi pT)^2}\right) - (-1)^{2\alpha}}, \quad (4.2)$$

where $\tilde{\Sigma}_1$ is a term independent of a given by (3.18). The contribution to the (renormalized) Casimir force from the right chamber is negative the contribution from the left chamber, with a replaced by $L - a$. As a result, the term $\tilde{\Sigma}_1$ cancels out and we find that the Casimir force acting on the piston is given by

$$F_{\text{Cas}}(a; L) = F_{\text{Cas}}^{\infty}(a) - F_{\text{Cas}}^{\infty}(L - a), \quad (4.3)$$

where

$$F_{\text{Cas}}^{\infty}(a) = -(-1)^{2\alpha} T \sum_{j \in J_*} \sum_{l=0}^{\infty} \sum_{p=-\infty}^{\infty} \frac{\sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2 + (2\pi pT)^2}}{\exp\left(2a\sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2 + (2\pi pT)^2}\right) - (-1)^{2\alpha}} \quad (4.4)$$

can be interpreted as the limit of the Casimir force when the right chamber is infinitely long.

Since the difference between the cut-off dependent Casimir energy (3.14) and the renormalized Casimir energy (3.17) is given by (3.6), which by (3.11) is a linear function in L , this implies that we will obtain the same result for the Casimir force whether we use the cut-off dependent Casimir energy or the renormalized Casimir energy. In other words, the Casimir force acting on the piston is independent of the regularization and renormalization procedures employed.

It is easy to deduce from (4.4) that the Casimir force for $\alpha = 1/2$ is related to the Casimir force for $\alpha = 0$ or 1 by

$$F_{\text{Cas}} \left(a; L; \alpha = \frac{1}{2}; * = D/N \right) = 2F_{\text{Cas}}(2a; 2L; \alpha = 1/0, * = D/N) - F_{\text{Cas}}(a; L; \alpha = 1/0, * = D/N), \quad (4.5)$$

which also follows from (3.19).

The expression (4.4) is negative if $\alpha = 0$ or 1 and positive if $\alpha = 1/2$. Moreover, its absolute value is a monotonically decreasing function of a . Therefore, if the quantum field assumes either Dirichlet or Neumann boundary conditions on both the piston and the walls of the cylinder, the Casimir force acting on the piston is an attractive force tending to pull the piston towards the closer wall. If the quantum field assumes Dirichlet (resp. Neumann) boundary conditions on the piston and Neumann (resp. Dirichlet) boundary conditions on the walls of the cylinder, then the Casimir force acting on the piston is a repulsive force tending to restore the piston to its equilibrium position $x^1 = L/2$. In both cases, the magnitude of the Casimir force increases as the piston is moving away from the equilibrium position.

Eq. (4.4) can also be interpreted as the Casimir force acting on two parallel plates embedded in an infinitely long cylinder with cross section $\Omega \times \mathcal{N}^n$. It shows that the Casimir force between the plates is attractive if the field assumes the same boundary conditions on the plates and is repulsive if the field assumes different boundary conditions on the plates, regardless of the boundary conditions assumed on the surrounding transversal wall. Writing $\omega_{\Omega,*,j} = \omega'_{\Omega,*,j}/R$ and $\omega_{\mathcal{N},l} = \omega'_{\mathcal{N},l}/r$, eq. (4.4) shows that when a/r or a/R or am is large, the magnitude of the Casimir force decays exponentially. At high temperature T , eq. (4.4) shows that the leading term of the Casimir force is given by a term linear in T :

$$F_{\text{Cas}}^{\infty, T \gg 1}(a) \sim -(-1)^{2\alpha} T \sum_{j \in J_*} \sum_{l=0}^{\infty} \frac{\sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2}}{\exp\left(2a\sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2}\right) - (-1)^{2\alpha}}, \quad (4.6)$$

and the remaining term decays exponentially. Notice that if we consider the contribution to the Casimir force from the left chamber (4.2), then the result (A.1) of appendix A shows that when $T \gg 1$,

$$F_{\text{Cas}}^{\text{ren, left}}(a) \sim \frac{1}{4\pi} \sum_{i=0}^{d-1} \Gamma\left(\frac{i+2}{2}\right) \zeta_R(i+2) c_{\Omega \times \mathcal{N}, *, d-1-i}(m) (2T)^{i+2} + O(T). \quad (4.7)$$

We observe that in general, there are terms of order T^2, T^3, \dots, T^{d+1} . In particular, the leading term is

$$\begin{aligned} F_{\text{Cas}}^{\text{ren, left}}(a) &\sim \frac{2^{d-1} \Gamma\left(\frac{d+1}{2}\right) \zeta_R(d+1)}{\pi} c_{\Omega \times \mathcal{N}, *, 0}(m) T^{d+1} \\ &= \frac{\Gamma\left(\frac{d+1}{2}\right) \zeta_R(d+1)}{\pi^{\frac{d+1}{2}}} \text{Vol}(\Omega \times \mathcal{N}) T^{d+1}, \end{aligned}$$

which is the Stefan-Boltzmann term. Usually this term is subtracted away in the regularization of the Casimir energy since it can be interpreted as the contribution to the vacuum

energy in the absence of boundaries. However, besides the limiting case of infinite parallel plates, there are still terms with order T^2, \dots, T^d . Nevertheless, since these terms are independent of a , they cancel the corresponding contributions from the right chamber and the high temperature leading term of the Casimir force acting on the piston is a term linear in T given by (4.6). This is usually called the classical term [45–48] due to the absence of the Planck constant \hbar in this term. In a recent work on Casimir effect of electromagnetic field in three dimensional ideal metal rectangular box [49], it has been argued that the terms of order T^2, \dots, T^{d+1} have to be subtracted away in order to be consistent with thermodynamics. Here we find that in the piston scenario, such terms are naturally absent due to the cancelation between the two sides of the plate.

For the low temperature asymptotic expansion of the Casimir force, we use the formula

$$\begin{aligned} & \frac{(-1)^{2\alpha-1} \sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2 + (2\pi pT)^2}}{\exp\left(2a\sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2 + (2\pi pT)^2}\right) - (-1)^{2\alpha}} \\ &= \frac{1}{2\sqrt{\pi}} \frac{\partial}{\partial a} \left\{ a \int_0^\infty t^{-\frac{1}{2}} \sum_{k=1}^\infty e^{2\pi i k \alpha} \exp\left\{-\frac{1}{t} (\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + [2\pi pT]^2 + m^2) - tk^2 a^2\right\} dt \right\} \end{aligned} \quad (4.8)$$

and the formula

$$\sum_{k=-\infty}^\infty \exp(-tk^2) = \sqrt{\pi t^{-\frac{1}{2}}} \sum_{k=-\infty}^\infty \exp\left(-\frac{\pi^2 k^2}{t}\right). \quad (4.9)$$

These give

$$\begin{aligned} F_{\text{Cas}}^\infty(a) &= \frac{1}{2\pi} \sum_{k=1}^\infty \sum_{j \in J_*} \sum_{l=0}^\infty \sum_{p=-\infty}^\infty \\ &\times e^{2\pi i k \alpha} \left\{ \sqrt{\frac{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2}{[ka]^2 + \left[\frac{p}{2T}\right]^2}} K_1 \left(2\sqrt{(\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2) \left([ka]^2 + \left[\frac{p}{2T}\right]^2\right)} \right) \right. \\ &\left. - 2k^2 a^2 \frac{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2}{[ka]^2 + \left[\frac{p}{2T}\right]^2} K_2 \left(2\sqrt{(\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2) \left([ka]^2 + \left[\frac{p}{2T}\right]^2\right)} \right) \right\}. \end{aligned} \quad (4.10)$$

In the zero temperature limit, we obtain the zero temperature Casimir force from the terms with $p = 0$:

$$\begin{aligned} F_{\text{Cas}}^{\infty, T=0}(a) &= \frac{1}{2\pi} \sum_{k=1}^\infty \sum_{j \in J_*} \sum_{l=0}^\infty e^{2\pi i k \alpha} \left\{ \frac{\sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2}}{ka} K_1 \left(2ka \sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2} \right) \right. \\ &\left. - 2(\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2) K_2 \left(2ka \sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2} \right) \right\}, \end{aligned} \quad (4.11)$$

which can also be derived directly from eq. (3.12). The temperature correction to the Casimir force is the sum of the terms with $p \neq 0$ in (4.10). It goes to zero exponentially

fast when $T \rightarrow 0$. Using the identity $K_2(z) = K_0(z) + 2K_1(z)/z$, we can rewrite the zero temperature Casimir force (4.11) as

$$F_{\text{Cas}}^{\infty, T=0}(a) = -\frac{1}{2\pi} \sum_{k=1}^{\infty} \sum_{j \in J_*} \sum_{l=0}^{\infty} e^{2\pi i k \alpha} \left\{ \frac{\sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2}}{ka} K_1 \left(2ka \sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2} \right) + 2 \left(\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2 \right) K_0 \left(2ka \sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2} \right) \right\}, \quad (4.12)$$

which shows manifestly that the zero temperature Casimir force is attractive if $\alpha = 0$ or 1 and is repulsive if $\alpha = 1/2$.

For the behavior of the Casimir force with respect to the variation of mass, we observe that the function

$$x \mapsto \frac{x}{e^x - 1}$$

is a decreasing function. Therefore the magnitude of the Casimir force acting between a pair of parallel plates embedded in an infinitely long cylinder (of arbitrary cross section) is a decreasing function of m when $\alpha = 0$ or 1. In other words, for a pair of parallel plates with identical boundary conditions, the increase in mass reduces the Casimir effect. This property cannot be obviously inferred from the expressions for the zero temperature Casimir force given by (4.11) or (4.12). We see here the advantage of considering the Casimir force at any finite temperature. It enables us to derive some properties of the Casimir force from the expression (4.4), which cannot be derived directly from the expression for zero temperature Casimir force (4.11) or (4.12).

Notice that when the mass m decreases to zero, (4.4) naively shows that the Casimir force tends to the Casimir force for massless scalar fields [15]. To be more careful, we need to discuss the cases where the field assumes Dirichlet boundary conditions and Neumann boundary conditions on the surrounding walls $[0, L] \times \partial\Omega \times \mathcal{N}^n$ separately. If the field assumes Dirichlet boundary conditions on the wall $[0, L] \times \partial\Omega \times \mathcal{N}^n$, then the Dirichlet eigenvalues $\omega_{\Omega,D;j}^2$ are all nonzero. In this case, we can immediately set $m = 0$ in (4.4) and obtain the Casimir force for massless scalar field. However, if the field assumes Neumann boundary conditions on the wall $[0, L] \times \partial\Omega \times \mathcal{N}^n$, there is exactly one zero Neumann eigenvalue $\omega_{\Omega,N;0}^2$. In this case, there are κ pairs of (j, l) such that $\omega_{\Omega,N;j}^2 + \omega_{\mathcal{N};l}^2 = 0$. Separating the sum over $(j, l) \in J_N \times \tilde{N}$ in (4.4) as a sum over those (j, l) with $\omega_{\Omega,N;j}^2 + \omega_{\mathcal{N};l}^2 = 0$ and those (j, l) with $\omega_{\Omega,N;j}^2 + \omega_{\mathcal{N};l}^2 \neq 0$, we find that we can immediately set $m = 0$ in the latter sum to obtain the contribution of the modes with $\omega_{\Omega,N;j}^2 + \omega_{\mathcal{N};l}^2 \neq 0$ to the Casimir force due to massless scalar field. For the sum over (j, l) with $\omega_{\Omega,N;j}^2 + \omega_{\mathcal{N};l}^2 = 0$ given by

$$-(-1)^{2\alpha} \kappa T \sum_{p=-\infty}^{\infty} \frac{\sqrt{m^2 + (2\pi p T)^2}}{\exp \left(2a \sqrt{m^2 + (2\pi p T)^2} \right) - (-1)^{2\alpha}}, \quad (4.13)$$

we have to be careful when taking the massless limit for the term with $p = 0$. Using the fact that

$$\frac{x}{e^x - 1} = 1 + O(x), \quad \frac{x}{e^x + 1} = O(x) \quad \text{as } x \rightarrow 0,$$

we find that the massless limit of (4.13) is given by

$$\kappa \left\{ -|2\alpha - 1| \frac{T}{2a} + 4\pi T^2 \sum_{p=1}^{\infty} \frac{(-1)^{2\alpha-1} p}{\exp(4\pi p T a) - (-1)^{2\alpha}} \right\}. \quad (4.14)$$

Except for the factor κ , this is the finite temperature Casimir force between a pair of parallel plates in (1+1)-dimensional Minkowski spacetime due to massless scalar field with Dirichlet or Neumann boundary conditions on both plates (for $\alpha = 0$ or 1) and with Dirichlet boundary condition on one plate and Neumann boundary condition on the other plate (for $\alpha = 1/2$) (see [50]). As in [50], the Casimir force due to the modes with $\omega_{\Omega, N; j}^2 + \omega_{\mathcal{N}; l}^2 = 0$ (4.14) has an alternative expression given by

$$\kappa \left(-\frac{\pi}{24a^2} - \frac{\pi T^2}{6} + \frac{\pi}{a^2} \sum_{k=1}^{\infty} \frac{k}{\exp\left(\frac{\pi k}{T a}\right) - 1} \right), \quad \text{if } \alpha = 0, 1; \quad (4.15)$$

and

$$\kappa \left(\frac{\pi}{48a^2} - \frac{\pi T^2}{6} + \frac{\pi}{a^2} \sum_{k=1}^{\infty} \frac{k + \frac{1}{2}}{\exp\left(\frac{\pi(k + \frac{1}{2})}{T a}\right) - 1} \right), \quad \text{if } \alpha = \frac{1}{2}. \quad (4.16)$$

The first term in (4.15) and (4.16) give the respective zero temperature Casimir force. Notice that in contrast to the massive case where the force decays exponentially, they decay in the order $1/a^2$ when a is large. This is a long range force which is the subject of study in the context of electromagnetic fields in the recent work [51]. At finite temperature, (4.14) shows that this long range force is present if and only if $\alpha \neq 1/2$, i.e., if and only if the boundary conditions assumed on the two plates and the walls of the cylinder are both Neumann conditions. In this case, the long range force is of order T/a . From this analysis, we find that long range force may exist only in the massless case when Neumann boundary conditions are assumed on the wall of the cylinder and the two plates or when the temperature is zero and Dirichlet boundary conditions are assumed on the two plates. The transition from massless to massive field will change the nature of the force from long range to short range.

To investigate the dependence of the Casimir force $F_{\text{Cas}}^{\infty}(a)$ on the size R of the cross section Ω , we rewrite $\omega_{\Omega, *; j}$ as $\omega'_{\Omega, *; j}/R$, where the re-scaled frequency $\omega'_{\Omega, *; j}$ is independent of the size R of the cross section Ω . Using the same argument about the dependence of the Casimir force on mass, we see that if $\alpha = 0$ or 1, the Casimir force increases when the size R increases. The asymptotic behaviors of the Casimir force when the plate separation a is much smaller than the size R of the cross section Ω is derived in appendix B. We read

from (B.2) and (B.5) that if $a \ll R$, then if $aT \gg 1$,

$$\begin{aligned}
 F_{\text{Cas}}^\infty(a) &\sim \frac{T}{a\sqrt{\pi}} \sum_{i=0}^{d_1-1} c_{\Omega/R,*,i} \left(\frac{R}{a}\right)^{d_1-i-1} \sum_{k=1}^{\infty} \sum_{l=0}^{\infty} \sum_{p=-\infty}^{\infty} e^{2\pi i k \alpha} \left\{ \left(\frac{a\sqrt{\omega_{\mathcal{N};l}^2 + (2\pi p T)^2 + m^2}}{k} \right)^{\frac{d_1-i}{2}} \right. \\
 &\quad \times K_{\frac{d_1-i}{2}} \left(2ka\sqrt{\omega_{\mathcal{N};l}^2 + (2\pi p T)^2 + m^2} \right) - 2 \frac{\left(a\sqrt{\omega_{\mathcal{N};l}^2 + (2\pi p T)^2 + m^2} \right)^{\frac{d_1+2-i}{2}}}{k^{\frac{d_1-2-i}{2}}} \\
 &\quad \left. \times K_{\frac{d_1+2-i}{2}} \left(2ka\sqrt{\omega_{\mathcal{N};l}^2 + (2\pi p T)^2 + m^2} \right) \right\} + O\left(\left(\frac{R}{a}\right)^{-1}\right), \tag{4.17}
 \end{aligned}$$

and if $aT \ll 1$,

$$\begin{aligned}
 F_{\text{Cas}}^\infty(a) &\sim -\frac{1}{a^2} \sum_{i=0}^{d_1-1} c_{\Omega/R,*,i} \left(\frac{R}{a}\right)^{d_1-1-i} \left\{ \frac{1}{2\pi} \sum_{k=1}^{\infty} e^{2\pi i k \alpha} \sum_{l=0}^{\infty} \left[(d_1-i) \left(\frac{a\sqrt{\omega_{\mathcal{N};l}^2 + m^2}}{k} \right)^{\frac{d_1+1-i}{2}} \right. \right. \\
 &\quad \left. \left. \times K_{\frac{d_1+1-i}{2}} \left(2ka\sqrt{\omega_{\mathcal{N};l}^2 + m^2} \right) + 2 \frac{\left(a\sqrt{\omega_{\mathcal{N};l}^2 + m^2} \right)^{\frac{d_1+3-i}{2}}}{k^{\frac{d_1-1-i}{2}}} K_{\frac{d_1-1-i}{2}} \left(2ka\sqrt{\omega_{\mathcal{N};l}^2 + m^2} \right) \right] \right. \\
 &\quad \left. - 2^{\frac{d_1-i}{2}} \pi^{\frac{3}{2}} (aT)^{\frac{d_1-i-2}{2}} \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \sum_{p=1}^{\infty} (k+\alpha)^2 \left(\frac{\sqrt{[\pi(k+\alpha)]^2 + (a\omega_{\mathcal{N};l})^2 + (am)^2}}{p} \right)^{\frac{d_1-i-2}{2}} \right. \\
 &\quad \left. \times K_{\frac{d_1-i-2}{2}} \left(\frac{p}{Ta} \sqrt{[\pi(k+\alpha)]^2 + (a\omega_{\mathcal{N};l})^2 + (am)^2} \right) \right. \\
 &\quad \left. + \frac{(2aT)^{\frac{d_1-i+1}{2}}}{2\pi} \sum_{l=0}^{\infty} \sum_{p=1}^{\infty} \left(\frac{a\sqrt{\omega_{\mathcal{N};l}^2 + m^2}}{p} \right)^{\frac{d_1-i+1}{2}} K_{\frac{d_1-i+1}{2}} \left(\frac{p}{T} \sqrt{\omega_{\mathcal{N};l}^2 + m^2} \right) \right\} \\
 &\quad + O\left(\left(\frac{R}{a}\right)^{-1}\right). \tag{4.18}
 \end{aligned}$$

Here $c_{\Omega/R,*,i}$ are the heat kernel coefficients of the Laplace operator with Dirichlet ($* = D$) or Neumann ($* = N$) boundary conditions on Ω/R . Notice that when $r \ll a \ll R$, the large- R non-vanishing terms of the Casimir force can be written as a polynomial of order $d_1 - 1$ in R with coefficients depending on the geometric invariants $c_{\Omega,*,i}$ of Ω , and Bessel series that depend on the geometry of the internal manifold \mathcal{N}^n , and the plate separation a . From these expressions, it is easy to read that if the mass m is also very large, the Casimir force decays exponentially. In the case the mass m is small, or more precisely if $am \ll 1 \ll Rm$, we obtain from (B.4) and (B.5) that if $\alpha = 0$ or 1 ,

$$F_{\text{Cas}}^\infty(a) \sim \frac{\kappa T}{2\sqrt{\pi}a} \sum_{j=1}^{\lfloor \frac{d_1}{2} \rfloor} c_{\Omega/R,*,d_1-2j} \left(\frac{R}{a}\right)^{2j-1} \left\{ \frac{1}{2} \frac{(-1)^j}{j!} (am)^{2j} \left(\log\left(\frac{am}{2\pi}\right)^2 + 2 - \psi(j+1) - \psi(1) \right) \right\}$$

$$\begin{aligned}
 & + \sqrt{\pi} \sum_{q=0}^{j-1} \frac{(-1)^q}{q!} (am)^{2q} (2q - 2j + 1) \pi^{2j-2q-1} \Gamma\left(-j + \frac{1}{2} + q\right) \zeta_R(-2j + 1 + 2q) \Big\} \\
 & + \frac{\kappa T}{2\sqrt{\pi}a} \sum_{j=0}^{\lfloor \frac{d_1-1}{2} \rfloor} c_{\Omega/R,*,d_1-2j-1} \left(\frac{R}{a}\right)^{2j} \times \left\{ -\sqrt{\pi} \frac{(-1)^j}{j!} (am)^{2j} - \frac{1}{2} \Gamma\left(-j - \frac{1}{2}\right) (am)^{2j+1} \right. \\
 & \quad \left. + \sqrt{\pi} \sum_{q=0}^{j-1} \frac{(-1)^q}{q!} (am)^{2q} (2q - 2j) \pi^{2j-2q} \Gamma(-j + q) \zeta_R(-2j + 2q) \right\} \\
 & + \frac{T}{\sqrt{\pi}a} \sum_{i=0}^{d_1-1} c_{\Omega,*,i} \left(\frac{R}{a}\right)^{d_1-1-i} \sum_{q=0}^{d_1+1-i} \frac{(-1)^q}{q!} (am)^{2q} \sum_{k=1}^{\infty} \sum_{\substack{(l,p) \in \tilde{\mathbb{N}} \times \mathbb{Z} \\ \omega_{\mathcal{N};l}^2 + (2\pi pT)^2 \neq 0}} \\
 & \times \left\{ \left(\frac{a \sqrt{\omega_{\mathcal{N};l}^2 + (2\pi pT)^2}}{k} \right)^{\frac{d_1-i-2q}{2}} K_{\frac{d_1-i-2q}{2}} \left(2ka \sqrt{\omega_{\mathcal{N};l}^2 + (2\pi pT)^2} \right) \right. \\
 & \quad \left. - 2 \frac{\left(a \sqrt{\omega_{\mathcal{N};l}^2 + (2\pi pT)^2} \right)^{\frac{d_1+2-i-2q}{2}}}{k^{\frac{d_1-2-i-2q}{2}}} K_{\frac{d_1+2-i-2q}{2}} \left(2ka \sqrt{\omega_{\mathcal{N};l}^2 + (2\pi pT)^2} \right) \right\}, \quad (4.19)
 \end{aligned}$$

if $aT \gg 1$; and

$$\begin{aligned}
 F_{\text{Cas}}^{\infty}(a) & \sim \frac{\kappa}{4\pi a^2} \sum_{j=1}^{\lfloor \frac{d_1+1}{2} \rfloor} c_{\Omega/R,*,d_1+1-2j} \left(\frac{R}{a}\right)^{2j-2} \left\{ \frac{1}{2} \frac{(-1)^j}{j!} (am)^{2j} \left(\log\left(\frac{am}{2\pi}\right)^2 + 2 - \psi(j+1) - \psi(1) \right) \right. \\
 & + \sqrt{\pi} \sum_{q=0}^{j-1} \frac{(-1)^q}{q!} (am)^{2q} (2q - 2j + 1) \pi^{2j-2q-1} \Gamma\left(-j + \frac{1}{2} + q\right) \zeta_R(-2j + 1 + 2q) \Big\} \\
 & + \frac{\kappa}{4\pi a^2} \sum_{j=0}^{\lfloor \frac{d_1}{2} \rfloor} c_{\Omega/R,*,d_1-2j} \left(\frac{R}{a}\right)^{2j-1} \left\{ -\sqrt{\pi} \frac{(-1)^j}{j!} (am)^{2j} - \frac{1}{2} \Gamma\left(-j - \frac{1}{2}\right) (am)^{2j+1} \right. \\
 & \left. + \sqrt{\pi} \sum_{q=0}^{j-1} \frac{(-1)^q}{q!} (am)^{2q} (2q - 2j) \pi^{2j-2q} \Gamma(-j + q) \zeta_R(-2j + 2q) \right\} \\
 & - \frac{1}{2\pi a^2} \sum_{i=0}^{d_1-1} c_{\Omega,*,i} \left(\frac{R}{a}\right)^{d_1-1-i} \sum_{q=0}^{d_1+1-i} \frac{(-1)^q}{q!} (am)^{2q} \sum_{k=1}^{\infty} \sum_{\substack{l \in \tilde{\mathbb{N}} \\ \omega_{\mathcal{N};l}^2 \neq 0}} \\
 & \times \left\{ (d_1 - i - 2q) \left(\frac{\omega_{\mathcal{N};l}}{k}\right)^{\frac{d_1+1-i-2q}{2}} K_{\frac{d_1+1-i-2q}{2}}(2ka\omega_{\mathcal{N};l}) + 2 \frac{\omega_{\mathcal{N};l}}{k^{\frac{d_1-1-i-2q}{2}}} K_{\frac{d_1-1-i-2q}{2}}(2ka\omega_{\mathcal{N};l}) \right\} \quad (4.20)
 \end{aligned}$$

if $T = 0$. The omitted terms goes to zero when $am \rightarrow 0$ or $R/a \rightarrow \infty$. Notice that these asymptotic behaviors contain logarithmic terms in am which goes to zero when the mass m approaches zero. From (4.19) and (4.20), we find that when m is small, then as $a \rightarrow 0^+$,

the leading order term of the Casimir force is given by

$$F_{\text{Cas}}^\infty(a) \sim -(d_1 - 1)\Gamma\left(\frac{d_1}{2}\right) \zeta_R(d_1) \frac{\text{Vol}(\Omega)}{(4\pi)^{\frac{d_1}{2}}} \frac{\kappa T}{a^{d_1+1}}, \quad \text{if } aT \gg 1,$$

and

$$F_{\text{Cas}}^\infty(a) \sim -d_1\Gamma\left(\frac{d_1+1}{2}\right) \zeta_R(d_1+1) \frac{\text{Vol}(\Omega)}{(4\pi)^{\frac{d_1+1}{2}}} \frac{\kappa}{a^{d_1}}, \quad \text{if } aT \ll 1,$$

respectively, which are independent of the mass m . This shows that when $am \ll 1$, the mass correction to the Casimir force is not significant.

If $d_1 = 3$ and $\alpha = 0$ or 1 , (4.19) and (4.20) give respectively

$$F_{\text{Cas}}^\infty(a) \sim \kappa T \left\{ -\frac{c_{\Omega/R,*;0} R^2}{2a^3} \zeta_R(3) - \frac{\pi^{\frac{3}{2}} c_{\Omega/R,*;1} R}{12a^2} - \frac{c_{\Omega/R,*;2}}{2a} + \frac{m^2 c_{\Omega/R,*;0} R^2}{2a} - \frac{m^2 c_{\Omega/R,*;1} R}{4\sqrt{\pi}} \log(am)^2 \right\} + O(a^0) \quad (4.21)$$

as $r \ll a \rightarrow 0^+$

if $T \gg 1$, and

$$F_{\text{Cas}}^\infty(a) \sim \kappa \left\{ -\frac{\pi^3 c_{\Omega/R,*;0} R^2}{120a^4} - \frac{c_{\Omega/R,*;1} R}{4\sqrt{\pi} a^3} \zeta_R(3) - \frac{\pi c_{\Omega/R,*;2}}{24a^2} + \frac{m^2 c_{\Omega/R,*;1} R}{4\sqrt{\pi} a} - \frac{c_{\Omega/R,*;2}}{8\pi} m^2 \log(am)^2 + \frac{m^4 c_{\Omega/R,*;0} R^2}{16\pi} \log(am)^2 \right\} + O(a^0) \quad (4.22)$$

as $r \ll a \rightarrow 0^+$,

if $T \ll 1$. When $\Omega = [0, L_2] \times [0, L_3]$ is a rectangle, (4.22) gives the correct behavior of the Casimir force when $a \ll 1$ which was derived in [44].

For the influence of the extra dimensions, we rewrite $\omega_{\mathcal{N};l}$ as $\omega'_{\mathcal{N};l}/r$, where the re-scaled frequency $\omega'_{\mathcal{N};l}$ is independent of the size r of the extra dimensions \mathcal{N}^n . We find from each of the formulas and asymptotic expansions for the Casimir force $F_{\text{Cas}}^\infty(a)$ derived above that, as the size r is very small compared to the plate separation a , the terms with nonzero $\omega_{\mathcal{N};l}^2$ contribute Casimir force that are exponentially small. In the limit the internal manifold \mathcal{N}^n vanishes, i.e. $r \rightarrow 0^+$, only the κ terms corresponding to $\omega_{\mathcal{N};l}^2 = 0$ remain and they give κ times the Casimir force in $(d_1 + 1)$ -dimensional Minkowski spacetime. In other words, if $\kappa = 0$, the Casimir force goes to zero in the limit of vanishing internal space. This is definitely not a desired physical situation. The situation that is of physical interest is the recovery of the Casimir force in $(d_1 + 1)$ -dimensional Minkowski spacetime in the limit of vanishing extra dimensions, or equivalently $\kappa = 1$. This happens in particular when \mathcal{N}^n has zero scalar curvature or when $\xi = 0$ (minimal coupling) and \mathcal{N}^n is connected. In this case, (4.4) shows that the presence of extra dimensions enhances the Casimir effect. In case $\alpha = 0$ or 1 , the expression (4.4) shows that the magnitude of the Casimir force becomes larger when the size of the internal manifold r is larger.

When the size r of the internal manifold \mathcal{N}^n becomes comparable to the plate separation a , the correction to the Casimir force in $(d_1 + 1)$ -dimensional Minkowski spacetime due to the extra dimensions (i.e. the contributions from the terms with $\omega_{\mathcal{N};l}^2 \neq 0$) can become substantial and it depends on the geometry of the internal manifold \mathcal{N}^n .

4.2 The Casimir force density acting on a pair of infinite parallel plates

In this section, we consider the limit where the size R of the cross section Ω goes to infinity, which is tantamount to two parallel plates in a $(d_1 + 1)$ -dimensional Minkowski spacetime M^{d_1+1} , with an n -dimensional internal manifold compactified to \mathcal{N}^n at every point of M^{d_1+1} . In this case, we should consider the Casimir force density $\mathcal{F}_{\text{Cas}}^{\parallel}(a)$ on the plates $x^1 = 0$ and $x^1 = a$ which is defined as the limit

$$\mathcal{F}_{\text{Cas}}^{\parallel}(a) = \lim_{R \rightarrow \infty} \frac{F_{\text{Cas}}^{\infty}(a)}{\text{Vol}(\Omega)} = \lim_{R \rightarrow \infty} \frac{F_{\text{Cas}}^{\infty}(a)}{R^{d_1-1}}.$$

From (4.17) and (4.18), we observe that when R is large, the leading term of $F_{\text{Cas}}^{\infty}(a)$ is of order R^{d_1-1} coming from the term with $i = 0$. All the remaining terms are of order smaller than R^{d_1-1} . Using the fact that $c_{\Omega/R,*;0} = 1/(2\sqrt{\pi})^{d_1-1}$, we obtain immediately the following high and low temperature expansions for the Casimir force density $\mathcal{F}_{\text{Cas}}^{\parallel}(a)$:

$$\begin{aligned} \mathcal{F}_{\text{Cas}}^{\parallel}(a) &= \frac{1}{2^{d_1-1} \pi^{\frac{d_1}{2}}} \frac{T}{a^{d_1}} \sum_{k=1}^{\infty} \sum_{l=0}^{\infty} \sum_{p=-\infty}^{\infty} e^{2\pi i k \alpha} \left(\frac{a \sqrt{\omega_{\mathcal{N};l}^2 + (2\pi p T)^2 + m^2}}{k} \right)^{\frac{d_1}{2}} \\ &\quad \times K_{\frac{d_1}{2}} \left(2ka \sqrt{\omega_{\mathcal{N};l}^2 + (2\pi p T)^2 + m^2} \right) \\ &\quad - \frac{1}{2^{d_1-2} \pi^{\frac{d_1}{2}}} \frac{T}{a^{d_1}} \sum_{k=1}^{\infty} \sum_{l=0}^{\infty} \sum_{p=-\infty}^{\infty} e^{2\pi i k \alpha} \frac{\left(a \sqrt{\omega_{\mathcal{N};l}^2 + (2\pi p T)^2 + m^2} \right)^{\frac{d_1+2}{2}}}{k^{\frac{d_1-2}{2}}} \\ &\quad \times K_{\frac{d_1+2}{2}} \left(2ka \sqrt{\omega_{\mathcal{N};l}^2 + (2\pi p T)^2 + m^2} \right), \\ \mathcal{F}_{\text{Cas}}^{\parallel}(a) &= - \frac{d_1}{2^{d_1} \pi^{\frac{d_1+1}{2}}} \frac{1}{a^{d_1+1}} \sum_{k=1}^{\infty} \sum_{l=0}^{\infty} e^{2\pi i k \alpha} \left(\frac{a \sqrt{\omega_{\mathcal{N};l}^2 + m^2}}{k} \right)^{\frac{d_1+1}{2}} \\ &\quad \times K_{\frac{d_1+1}{2}} \left(2ka \sqrt{\omega_{\mathcal{N};l}^2 + m^2} \right) - \frac{1}{2^{d_1-1} \pi^{\frac{d_1+1}{2}}} \frac{1}{a^{d_1+1}} \\ &\quad \times \sum_{k=1}^{\infty} \sum_{l=0}^{\infty} e^{2\pi i k \alpha} \frac{\left(a \sqrt{\omega_{\mathcal{N};l}^2 + m^2} \right)^{\frac{d_1+3}{2}}}{k^{\frac{d_1-1}{2}}} K_{\frac{d_1-1}{2}} \left(2ka \sqrt{\omega_{\mathcal{N};l}^2 + m^2} \right) \\ &\quad + \frac{(aT)^{\frac{d_1-2}{2}}}{2^{\frac{d_1-2}{2}} \pi^{\frac{d_1-4}{2}} a^{d_1+1}} \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \sum_{p=1}^{\infty} (k+\alpha)^2 \left(\frac{\sqrt{(\pi[k+\alpha])^2 + (a\omega_{\mathcal{N};l})^2 + (am)^2}}{p} \right)^{\frac{d_1-2}{2}} \\ &\quad \times K_{\frac{d_1-2}{2}} \left(\frac{p}{Ta} \sqrt{(\pi[k+\alpha])^2 + (a\omega_{\mathcal{N};l})^2 + (am)^2} \right) \end{aligned}$$

$$- \frac{(aT)^{\frac{d_1+1}{2}}}{2^{\frac{d_1-1}{2}} \pi^{\frac{d_1+1}{2}} a^{d_1+1}} \sum_{l=0}^{\infty} \sum_{p=1}^{\infty} \left(\frac{a \sqrt{\omega_{\mathcal{N};l}^2 + m^2}}{p} \right)^{\frac{d_1+1}{2}} K_{\frac{d_1+1}{2}} \left(\frac{p}{T} \sqrt{\omega_{\mathcal{N};l}^2 + m^2} \right). \tag{4.23}$$

The first two terms in (4.23) give the zero temperature Casimir force density and it agrees with the result obtained in [16]. Notice that since the Casimir force density $\mathcal{F}_{\text{Cas}}^{\parallel}(a)$ is derived as a limit of the Casimir force $F_{\text{Cas}}^{\infty}(a)$, it follows that for a pair of infinite parallel plates with Dirichlet or Neumann boundary conditions on both plates, the Casimir force is attractive. For a pair of infinite parallel plates with Dirichlet boundary condition on one plate and Neumann boundary condition on the other plate, the Casimir force is repulsive. For either homogeneous or mixed boundary conditions, the magnitude of the Casimir force is a decreasing function of the plate separation a , but it is enhanced by the presence of the extra dimensions. If both the plates assume the same (Dirichlet or Neumann) boundary conditions, the Casimir force density $\mathcal{F}_{\text{Cas}}^{\parallel}(a)$ is also a decreasing function of mass. In the high temperature regime, the leading term of the Casimir force is linear in temperature. In the low temperature regime, the Casimir effect is dominated by the zero temperature term.

4.3 Special case

Here we consider the special case as in section 3.3, where the cross section of the cylinder Ω is a rectangular region $[0, L_2] \times \dots \times [0, L_{d_1}]$ and the internal manifold \mathcal{N}^n is an n -torus T^n — a product of n circles with radius r_1, \dots, r_n respectively. For simplicity, we only consider the Casimir force $F_{\text{Cas}}^{\infty}(a)$ acting on a rectangular piston moving freely inside a semi-infinite long rectangular box when both the piston and the walls of the rectangular box assume the same (Dirichlet or Neumann) boundary conditions. The result for the case where they assume different boundary conditions can be obtained using (4.5). From (3.20) and (4.2), we find that the left closed chamber contributes the Casimir force

$$F_{\text{Cas}}^{\text{left}}(a; D/N) = -\tilde{\Sigma}_1 - T \sum_{\mathbf{j} \in \mathbb{N}^{d_1-1} / \tilde{\mathbb{N}}^{d_1-1}} \sum_{l \in \mathbb{Z}^n} \sum_{p=-\infty}^{\infty} \frac{\sqrt{\sum_{i=2}^{d_1} \left(\frac{\pi j_i}{L_i}\right)^2 + \sum_{i=1}^n \left(\frac{l_i}{r_i}\right)^2 + m^2 + (2\pi pT)^2}}{\exp\left(2a \sqrt{\sum_{i=2}^{d_1} \left(\frac{\pi j_i}{L_i}\right)^2 + \sum_{i=1}^n \left(\frac{l_i}{r_i}\right)^2 + m^2 + (2\pi pT)^2}\right) - 1},$$

and the right infinitely long chamber contributes

$$F_{\text{Cas}}^{\text{right}}(D/N) = \tilde{\Sigma}_1 = - \frac{\prod_{l=1}^n r_l}{2^{d_1+1}} \sum_{i=0}^{d_1-1} \frac{(\mp 1)^{d_1-i-1}}{\pi^{\frac{i-n+2}{2}}} m^{\frac{i+n+2}{2}} \times \sum_{2 \leq \sigma_1 < \dots < \sigma_i \leq d_1} \left[\prod_{j=1}^i L_{\sigma_j} \right] \sum_{(k_{\sigma_1}, \dots, k_{\sigma_i}, l_1, \dots, l_n, p) \in \mathbb{Z}^{i+n+1} \setminus \{0\}} \tag{4.24}$$

$$\begin{aligned} & \times \left(\sum_{j=1}^i [L_{\sigma_j} k_{\sigma_j}]^2 + \sum_{j=1}^n [\pi r_j l_j]^2 + \left[\frac{p}{2T} \right]^2 \right)^{-\frac{i+n+2}{4}} \\ & \times K_{\frac{i+n+2}{2}} \left(2m \sqrt{\sum_{j=1}^i [L_{\sigma_j} k_{\sigma_j}]^2 + \sum_{j=1}^n [\pi r_j l_j]^2 + \left[\frac{p}{2T} \right]^2} \right). \end{aligned}$$

For Neumann boundary conditions, it is obvious that $\tilde{\Sigma}_1$ is negative. Using (4.9), we can rewrite $\tilde{\Sigma}_1$ in the form (3.18), where the terms with $p = 0$ in (4.24) are the T -independent terms correspond to the first two terms in (3.18). As is shown in appendix A, when T is large, $\tilde{\Sigma}_1$ is negative and dominated by a term proportional T^{d+1} . Therefore, when T is large, there is a large repulsive force due to the vacuum fluctuations of the scalar field in the left chamber that tend to push the piston away from the closed end of the rectangular box. However, the vacuum fluctuations of the field in the right chamber give rise to a larger force in the opposite direction. The sum of these two forces is

$$\begin{aligned} F_{\text{Cas}}^\infty(a; D/N) = & -T \sum_{j \in \mathbb{N}^{d_1-1} / \tilde{\mathbb{N}}^{d_1-1}} \sum_{l \in \mathbb{Z}^n} \sum_{p=-\infty}^{\infty} \\ & \frac{\sqrt{\sum_{i=2}^{d_1} \left(\frac{\pi j_i}{L_i} \right)^2 + \sum_{i=1}^n \left(\frac{l_i}{r_i} \right)^2 + m^2 + (2\pi p T)^2}}{\exp \left(2a \sqrt{\sum_{i=2}^{d_1} \left(\frac{\pi j_i}{L_i} \right)^2 + \sum_{i=1}^n \left(\frac{l_i}{r_i} \right)^2 + m^2 + (2\pi p T)^2} \right) - 1}, \end{aligned} \quad (4.25)$$

which tends to move the piston towards the closed end of the rectangular box. The expression (4.25) shows that the Casimir force $F_{\text{Cas}}^\infty(a; D/N)$ is exponentially small when any of the parameters a, m, T is large or any of the parameters $L_2, \dots, L_{d_1}, r_1, \dots, r_n$ is small. There are a few alternative expressions for the Casimir force $F_{\text{Cas}}^\infty(a; D/N)$ which can be used to study the behaviors of the Casimir force at other limits. Using the formula (4.10), we find that if a and m are large and $L_2, \dots, L_{d_1}, r_1, \dots, r_n, T$ are small, the Casimir force can be computed using the formula

$$\begin{aligned} F_{\text{Cas}}^\infty(a; D/N) = & \frac{1}{2\pi} \sum_{k=1}^{\infty} \sum_{j \in \mathbb{N}^{d_1-1} / \tilde{\mathbb{N}}^{d_1-1}} \sum_{l \in \mathbb{Z}^n} \sum_{p=-\infty}^{\infty} \\ & \times \left\{ \sqrt{\frac{\sum_{i=2}^{d_1} \left(\frac{\pi j_i}{L_i} \right)^2 + \sum_{i=1}^n \left(\frac{l_i}{r_i} \right)^2 + m^2}{[ka]^2 + \left[\frac{p}{2T} \right]^2}} \right. \\ & \times K_1 \left(2 \sqrt{\left(\sum_{i=2}^{d_1} \left(\frac{\pi j_i}{L_i} \right)^2 + \sum_{i=1}^n \left(\frac{l_i}{r_i} \right)^2 + m^2 \right) \left([ka]^2 + \left[\frac{p}{2T} \right]^2 \right)} \right) \\ & \left. - 2k^2 a^2 \frac{\sum_{i=2}^{d_1} \left(\frac{\pi j_i}{L_i} \right)^2 + \sum_{i=1}^n \left(\frac{l_i}{r_i} \right)^2 + m^2}{[ka]^2 + \left[\frac{p}{2T} \right]^2} \right) \end{aligned}$$

$$\times K_2 \left(2 \sqrt{\left(\sum_{i=2}^{d_1} \left(\frac{\pi j_i}{L_i} \right)^2 + \sum_{i=1}^n \left(\frac{l_i}{r_i} \right)^2 + m^2 \right) \left([ka]^2 + \left[\frac{p}{2T} \right]^2 \right)} \right) \Bigg\}.$$

In the zero temperature limit, we find that the zero temperature Casimir force is given by

$$\begin{aligned} & F_{\text{Cas}}^\infty(a; T = 0; D/N) \\ &= \frac{1}{2\pi} \sum_{k=1}^{\infty} \sum_{\mathbf{j} \in \mathbb{N}^{d_1-1} / \mathbb{N}^{d_1-1}} \sum_{\mathbf{l} \in \mathbb{Z}^n} \left\{ \frac{\sqrt{\sum_{i=2}^{d_1} \left(\frac{\pi j_i}{L_i} \right)^2 + \sum_{i=1}^n \left(\frac{l_i}{r_i} \right)^2 + m^2}}{ka} \right. \\ & \times K_1 \left(2ka \sqrt{\sum_{i=2}^{d_1} \left(\frac{\pi j_i}{L_i} \right)^2 + \sum_{i=1}^n \left(\frac{l_i}{r_i} \right)^2 + m^2} \right) \\ & \left. - 2 \left(\sum_{i=2}^{d_1} \left(\frac{\pi j_i}{L_i} \right)^2 + \sum_{i=1}^n \left(\frac{l_i}{r_i} \right)^2 + m^2 \right) K_2 \left(2ka \sqrt{\sum_{i=2}^{d_1} \left(\frac{\pi j_i}{L_i} \right)^2 + \sum_{i=1}^n \left(\frac{l_i}{r_i} \right)^2 + m^2} \right) \right\}. \end{aligned}$$

This expression shows that the zero temperature Casimir force is exponentially small when any of the parameters a, m is large or any of the parameters $L_2, \dots, L_{d_1}, r_1, \dots, r_n$ is small. For the situation that we are more interested in, i.e., the case where $r_i \ll a \ll L_j, 1 \leq i \leq n, 2 \leq j \leq d_1$, the Casimir force $F_{\text{Cas}}^\infty(a; D/N)$ can be written as the sum of two terms, where the first term

$$\begin{aligned} & - \frac{T}{2^{d_1-1}} \sum_{i=0}^{d_1-1} (\mp 1)^{d_1-1-i} \frac{\tilde{S}_i}{\pi^{\frac{i+1}{2}}} \sum_{k=1}^{\infty} \sum_{\mathbf{l} \in \mathbb{Z}^n} \sum_{p=-\infty}^{\infty} \left\{ i \left(\frac{\sqrt{\sum_{j=1}^n \left(\frac{l_j}{r_j} \right)^2 + (2\pi p T)^2 + m^2}}{ka} \right)^{\frac{i+1}{2}} \right. \\ & \times K_{\frac{i+1}{2}} \left(2ka \sqrt{\sum_{j=1}^n \left(\frac{l_j}{r_j} \right)^2 + (2\pi p T)^2 + m^2} \right) + 2 \frac{\left(\sum_{j=1}^n \left(\frac{l_j}{r_j} \right)^2 + (2\pi p T)^2 + m^2 \right)^{\frac{i+3}{4}}}{(ka)^{\frac{i-1}{2}}} \\ & \left. \times K_{\frac{i-1}{2}} \left(2ka \sqrt{\sum_{j=1}^n \left(\frac{l_j}{r_j} \right)^2 + (2\pi p T)^2 + m^2} \right) \right\} \end{aligned} \quad (4.26)$$

is the dominating term, and the second term

$$\begin{aligned} & - \frac{T}{2^{d_1}} \sum_{i=0}^{d_1-1} \frac{(\mp 1)^{d_1-1-i}}{\pi^{\frac{i+1}{2}}} \sum_{\mathbf{l} \in \mathbb{Z}^n} \sum_{p=-\infty}^{\infty} \sum_{2 \leq \sigma_1 < \dots < \sigma_i \leq d_1} \left[\prod_{j=1}^i L_{\sigma_j} \right] \sum_{(k_{\sigma_1}, \dots, k_{\sigma_i}) \in \mathbb{Z}^i \setminus \{\mathbf{0}\}} \\ & \times \left\{ \left(\frac{\sum_{j=1}^n \left(\frac{l_j}{r_j} \right)^2 + (2\pi p T)^2 + m^2}{\sum_{j=1}^i [k_{\sigma_j} L_{\sigma_j}]^2} \right)^{\frac{i+1}{4}} \right. \\ & \left. \times K_{\frac{i+1}{2}} \left(2 \sqrt{\left(\sum_{j=1}^n \left(\frac{l_j}{r_j} \right)^2 + (2\pi p T)^2 + m^2 \right) \left(\sum_{j=1}^i [k_{\sigma_j} L_{\sigma_j}]^2 \right)} \right) \right\} \end{aligned} \quad (4.27)$$

$$\begin{aligned}
 & - \frac{4\pi^{\frac{5}{2}}}{a^3} \sum_{k_1=1}^{\infty} k_1^2 \left(\frac{\left(\frac{\pi k_1}{a}\right)^2 + \sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + (2\pi p T)^2 + m^2}{\sum_{j=1}^i [k_{\sigma_j} L_{\sigma_j}]^2} \right)^{\frac{i-2}{4}} \\
 & \times K_{\frac{i-2}{2}} \left(2 \sqrt{\left(\frac{\pi k_1}{a}\right)^2 + \sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + (2\pi p T)^2 + m^2} \left(\sum_{j=1}^i [k_{\sigma_j} L_{\sigma_j}]^2 \right) \right) \Big\}
 \end{aligned}$$

decays exponentially when the area of the cross section $L_2 \dots L_{d_1}$ is large. The term \tilde{S}_i in (4.26) is equal to the sum

$$\tilde{S}_i = \sum_{2 \leq \sigma_1 < \dots < \sigma_i \leq d_1} \left[\prod_{j=1}^i L_{\sigma_j} \right].$$

The expressions (4.26) and (4.27) are suitable for investigating the high temperature behavior. The high temperature leading term is linear in T , which is equal to the sum of the terms with $p = 0$. At low temperature, we have the following alternative expressions for (4.26) and (4.27):

$$\begin{aligned}
 & - \frac{1}{2^{d_1}} \sum_{i=0}^{d_1-1} (\mp 1)^{d_1-1-i} \frac{\tilde{S}_i}{\pi^{\frac{i+2}{2}}} \sum_{l \in \mathbb{Z}^n} \left\{ - \sum_{k=1}^{\infty} \left(\frac{\sqrt{\sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + m^2}}{ka} \right)^{\frac{i+2}{2}} \right. \\
 & \times K_{\frac{i+2}{2}} \left(2ka \sqrt{\sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + m^2} \right) + 2 \sum_{k=1}^{\infty} \frac{\left(\sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + m^2\right)^{\frac{i+4}{4}}}{(ka)^{\frac{i}{2}}} \\
 & \times K_{\frac{i+4}{2}} \left(2ka \sqrt{\sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + m^2} \right) + (2T)^{\frac{i+2}{2}} \sum_{p=1}^{\infty} \left(\frac{\sqrt{\sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + m^2}}{p} \right)^{\frac{i+2}{2}} \\
 & \times K_{\frac{i+2}{2}} \left(\frac{p}{T} \sqrt{\sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + m^2} \right) - \frac{4\pi^{\frac{5}{2}} (2T)^{\frac{i-1}{2}}}{a^3} \sum_{p=1}^{\infty} \sum_{k=1}^{\infty} k^2 \left(\frac{\sqrt{\left(\frac{\pi k}{a}\right)^2 + \sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + m^2}}{p} \right)^{\frac{i-1}{2}} \\
 & \left. \times K_{\frac{i-1}{2}} \left(\frac{p}{T} \sqrt{\left(\frac{\pi k}{a}\right)^2 + \sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + m^2} \right) \right\} \tag{4.28}
 \end{aligned}$$

and

$$- \frac{1}{2^{d_1+1}} \sum_{i=0}^{d_1-1} \frac{(\mp 1)^{d_1-1-i}}{\pi^{\frac{i+2}{2}}} \sum_{l \in \mathbb{Z}^n} \sum_{p=-\infty}^{\infty} \sum_{2 \leq \sigma_1 < \dots < \sigma_i \leq d_1} \left[\prod_{j=1}^i L_{\sigma_j} \right]$$

$$\begin{aligned}
 & \times \sum_{(k_{\sigma_1}, \dots, k_{\sigma_i}) \in \mathbb{Z}^i \setminus \{0\}} \left\{ \left(\frac{\sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + m^2}{\left(\frac{p}{2T}\right)^2 + \sum_{j=1}^i [k_{\sigma_j} L_{\sigma_j}]^2} \right)^{\frac{i+2}{4}} \right. \\
 & \times K_{\frac{i+2}{2}} \left(2 \sqrt{\left(\sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + m^2 \right) \left(\left(\frac{p}{2T}\right)^2 + \sum_{j=1}^i [k_{\sigma_j} L_{\sigma_j}]^2 \right)} \right) \\
 & - \frac{4\pi^{\frac{5}{2}}}{a^3} \sum_{k_1=1}^{\infty} \left(\frac{\left(\frac{\pi k_1}{a}\right)^2 + \sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + m^2}{\left(\frac{p}{2T}\right)^2 + \sum_{j=1}^i [k_{\sigma_j} L_{\sigma_j}]^2} \right)^{\frac{i-1}{4}} \\
 & \left. \times K_{\frac{i-1}{2}} \left(2 \sqrt{\left(\left(\frac{\pi k_1}{a}\right)^2 + \sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + m^2 \right) \left(\left(\frac{p}{2T}\right)^2 + \sum_{j=1}^i [k_{\sigma_j} L_{\sigma_j}]^2 \right)} \right) \right\}. \tag{4.29}
 \end{aligned}$$

The $p = 0$ terms in (4.28) and (4.29) give the zero temperature Casimir force. In the limit the cross section is large, i.e. $L_i \rightarrow \infty$ for $i = 2, \dots, d_1$, we find from the term with $i = d_1 - 1$ in (4.26) and (4.28) that the Casimir force density acting on a pair of infinite parallel plates is given by

$$\begin{aligned}
 & \mathcal{F}_{\text{Cas}}^{\parallel}(a) \\
 & = -\frac{T}{2^{d_1-1} \pi^{\frac{d_1}{2}}} \sum_{k=1}^{\infty} \sum_{l \in \mathbb{Z}^n} \sum_{p=-\infty}^{\infty} \left\{ (d_1 - 1) \left(\frac{\sqrt{\sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + (2\pi p T)^2 + m^2}}{ka} \right)^{\frac{d_1}{2}} \right. \\
 & \times K_{\frac{d_1}{2}} \left(2ka \sqrt{\sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + (2\pi p T)^2 + m^2} \right) \\
 & \left. + 2 \frac{\left(\sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + (2\pi p T)^2 + m^2 \right)^{\frac{d_1+2}{4}}}{(ka)^{\frac{d_1-2}{2}}} K_{\frac{d_1-2}{2}} \left(2ka \sqrt{\sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + (2\pi p T)^2 + m^2} \right) \right\},
 \end{aligned}$$

or alternatively

$$\begin{aligned}
 \mathcal{F}_{\text{Cas}}^{\parallel}(a) & = -\frac{1}{2^{d_1} \pi^{\frac{d_1+1}{2}}} \sum_{l \in \mathbb{Z}^n} \left\{ -\sum_{k=1}^{\infty} \left(\frac{\sqrt{\sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + m^2}}{ka} \right)^{\frac{d_1+1}{2}} K_{\frac{d_1+1}{2}} \left(2ka \sqrt{\sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + m^2} \right) \right. \\
 & \left. + 2 \sum_{k=1}^{\infty} \frac{\left(\sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + m^2 \right)^{\frac{d_1+3}{4}}}{(ka)^{\frac{d_1-1}{2}}} K_{\frac{d_1+3}{2}} \left(2ka \sqrt{\sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + m^2} \right) \right\}
 \end{aligned}$$




Figure 2. Left: The Casimir force $F_{\text{Cas}}^\infty(a)$ as a function of r/a for different values of m and T . Right: The Casimir force $F_{\text{Cas}}^\infty(a)$ as a function of temperature for different values of r/a and m .




Figure 3. The Casimir force $F_{\text{Cas}}^\infty(a)$ as a function of mass for different values of r/a and T .

$$\begin{aligned}
 & + (2T)^{\frac{d_1+1}{2}} \sum_{p=1}^{\infty} \left(\frac{\sqrt{\sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + m^2}}{p} \right)^{\frac{d_1+1}{2}} K_{\frac{d_1+1}{2}} \left(\frac{p}{T} \sqrt{\sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + m^2} \right) \\
 & - \frac{4\pi^{\frac{5}{2}} (2T)^{\frac{d_1-2}{2}}}{a^3} \sum_{p=1}^{\infty} \sum_{k=1}^{\infty} \left(\frac{\sqrt{\left(\frac{\pi k_1}{a}\right)^2 + \sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + m^2}}{p} \right)^{\frac{d_1-2}{2}} \\
 & \times K_{\frac{d_1-2}{2}} \left(\frac{p}{T} \sqrt{\left(\frac{\pi k_1}{a}\right)^2 + \sum_{j=1}^n \left(\frac{l_j}{r_j}\right)^2 + m^2} \right) \Bigg\}.
 \end{aligned}$$

In figures 2 and 3, we show graphically the behavior of the Casimir force $F_{\text{Cas}}^\infty(a)$ acting on two parallel plates embedded in an infinitely long rectangular cylinder in a $(4 + 1)$ -dimensional background spacetime with an extra dimension curled up to a circle of radius

r . We assume that the massive scalar field is subjected to Dirichlet boundary conditions on the walls of the cylinder and the two plates. Figure 2(a) shows the variations of the Casimir force as a function of the size r of the extra dimension, or more precisely, the ratio of the size r to the plate separation a . It shows that the magnitude of the Casimir force is increased if the size of the extra dimension is increased. When $r/a > 0.4$, the existence of extra dimension can contribute substantially to the Casimir force. Figure 2(b) shows that the magnitude of the Casimir force is an increasing function of temperature. For $a = 6\mu\text{m}$, the graph shows that the Casimir force depends linearly on temperature when $T > 200\text{K}$. Figure 3 shows that the magnitude of the Casimir force decreases when the mass m increases.

5 Conclusion

In this paper, we have investigated the Casimir effect for massive scalar field with general curvature coupling in $(d+1)$ -dimensional spacetime with $n = d - d_1$ extra dimensions. We consider the cases that the field assumes Dirichlet or Neumann boundary conditions on two parallel plates embedded in an infinitely long cylinder. We derive a general expression for the renormalized Casimir energy that vanishes in the infinite mass limit. A lots of the properties of the Casimir force are similar to the massless case. In particular, if the field assumes Dirichlet or Neumann boundary conditions on both plates, then the Casimir force acting on the plates are attractive. If the field assumes Dirichlet boundary condition on one plate and Neumann boundary condition on the other plate, then the Casimir force is repulsive. Passing from massless to massive, we find that the strength of the Casimir force is reduced if both plates assume the same boundary conditions.

For the influence of the extra dimensions, we find that the presence of extra dimensions enhances the Casimir effect. When the size of the internal manifold shrinks to zero, one obtains the Casimir force in the (d_1+1) -dimensional spacetime if and only if a certain elliptic operator on the internal manifold has exactly one zero mode. This happens in particular in the case of minimal coupling or zero scalar curvature.

For the extension of this work, it would be interesting to consider the general case of Robin boundary conditions as in [16]. This will be important for considering the finite temperature Casimir effect in the braneworld model especially for the radion field stabilization mechanism, as has been shown in [16] for the zero temperature case. The work along this direction will be reported elsewhere.

Acknowledgments

This project is funded by Ministry of Science, Technology and Innovation, Malaysia under e-Science fund 06-02-01-SF0080.

A High temperature asymptotic behavior of $\tilde{\Sigma}_1$

In this section, we consider the asymptotic behavior of $\tilde{\Sigma}_1$ (3.18) at high temperature. The first two terms in (3.18) are independent of T . For the third term, we have

$$\begin{aligned}
& -\frac{T}{\pi} \sum_{j \in J_*} \sum_{l=0}^{\infty} \sum_{p=1}^{\infty} \frac{\sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N};l}^2 + m^2}}{p} K_1 \left(\frac{p \sqrt{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N};l}^2 + m^2}}{T} \right) \\
&= -\frac{1}{4\pi} \int_0^{\infty} \sum_{j \in J_*} \sum_{l=0}^{\infty} \sum_{p=1}^{\infty} \exp \left\{ -\frac{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N};l}^2 + m^2}{t} - \frac{tp^2}{4T^2} \right\} dt \\
&= -\frac{1}{4\pi} \int_0^{\infty} \sum_{p=1}^{\infty} \exp \left\{ -\frac{tp^2}{4T^2} \right\} \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \Gamma(z) t^z \zeta_{\Omega \times \mathcal{N},*}(z; m) dz dt \quad (\text{A.1}) \\
&= -\frac{1}{4\pi} \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \Gamma(z) \Gamma(z+1) (2T)^{2z+2} \zeta_R(2z+2) \zeta_{\Omega \times \mathcal{N},*}(z; m) dz \\
&\sim -\frac{1}{4\pi} \sum_{i=0}^{d-1} \Gamma \left(\frac{i+2}{2} \right) \zeta_R(i+2) c_{\Omega \times \mathcal{N},*;d-1-i}(m) (2T)^{i+2} + O(T).
\end{aligned}$$

B The asymptotic behavior of the Casimir force $F_{\text{Cas}}^{\infty}(a)$ in different limits

In this section, we derive the asymptotic behavior of the Casimir force $F_{\text{Cas}}^{\infty}(a)$ in different limits. We only consider the case where $\alpha = 0$ or 1 . The case $\alpha = 1/2$ can be derived analogously.

First we consider the asymptotic behavior when $r < a < R$. In the high temperature regime, it follows from (4.8) that

$$\begin{aligned}
F_{\text{Cas}}^{\infty}(a) &= \frac{T}{2\sqrt{\pi}} \frac{\partial}{\partial a} \left\{ a \int_0^{\infty} t^{-\frac{1}{2}} \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \Gamma(z) t^z \zeta_{\Omega,*}(z) \sum_{k=1}^{\infty} \sum_{l=0}^{\infty} \sum_{p=-\infty}^{\infty} \right. \\
&\quad \left. \times \exp \left(-\frac{\omega_{\mathcal{N};l}^2 + (2\pi pT)^2 + m^2}{t} - tk^2 a^2 \right) dz dt \right\} \\
&\sim \frac{T}{2\sqrt{\pi}} \sum_{i=0}^{\infty} c_{\Omega,*,i} \frac{\partial}{\partial a} \left\{ a \int_0^{\infty} t^{\frac{d_1-i}{2}-1} \sum_{k=1}^{\infty} \sum_{l=0}^{\infty} \sum_{p=-\infty}^{\infty} \right. \\
&\quad \left. \times \exp \left(-\frac{\omega_{\mathcal{N};l}^2 + (2\pi pT)^2 + m^2}{t} - tk^2 a^2 \right) dt \right\}, \quad (\text{B.1})
\end{aligned}$$

where $\zeta_{\Omega,*}(z)$ is the zeta function

$$\zeta_{\Omega,*}(s) = \sum_{j \in J_*} \omega_{\Omega,*,j}^{-2s},$$

and $c_{\Omega,*,i}$ is the heat kernel coefficient of the Laplace operator with Dirichlet/Neumann boundary conditions on Ω , i.e.,

$$\sum_{j \in J_*} e^{-t\omega_{\Omega,*,j}^2} = \sum_{i=0}^{M-1} c_{\Omega,*,i} t^{\frac{i-d_1+1}{2}} + O\left(t^{\frac{M-d_1+1}{2}}\right) \quad \text{as } t \rightarrow 0^+.$$

In terms of the measure R of the size of Ω ,

$$c_{\Omega,*,i} = R^{d_1-i-1} c_{\Omega/R,*,i} \propto R^{d_1-i-1}.$$

Therefore when $r < a < R$,

$$\begin{aligned} & F_{\text{Cas}}^\infty(a) \\ & \sim \frac{T}{a\sqrt{\pi}} \sum_{i=0}^{M-1} c_{\Omega/R,*,i} \left(\frac{R}{a}\right)^{d_1-i-1} \sum_{k=1}^{\infty} \sum_{l=0}^{\infty} \sum_{p=-\infty}^{\infty} \left\{ \left(\frac{a\sqrt{\omega_{\mathcal{N};l}^2 + (2\pi pT)^2 + m^2}}{k} \right)^{\frac{d_1-i}{2}} \right. \\ & \quad \times K_{\frac{d_1-i}{2}} \left(2ka\sqrt{\omega_{\mathcal{N};l}^2 + (2\pi pT)^2 + m^2} \right) \\ & \quad \left. - 2 \frac{\left(a\sqrt{\omega_{\mathcal{N};l}^2 + (2\pi pT)^2 + m^2} \right)^{\frac{d_1+2-i}{2}}}{k^{\frac{d_1-2-i}{2}}} K_{\frac{d_1+2-i}{2}} \left(2ka\sqrt{\omega_{\mathcal{N};l}^2 + (2\pi pT)^2 + m^2} \right) \right\} \\ & \quad + O\left(\left(\frac{R}{a}\right)^{d_1-M-1}\right). \end{aligned} \tag{B.2}$$

This expansion only gives the behavior when $am \gg 1$. For the expansion when $am \ll 1$, we go back to (B.1) and consider the expansion of

$$\mathfrak{I}(i; a) = \frac{\partial}{\partial a} \left\{ a \int_0^\infty t^{\frac{d_1-i}{2}-1} \sum_{k=1}^{\infty} \sum_{l=0}^{\infty} \sum_{p=-\infty}^{\infty} \exp\left(-\frac{\omega_{\mathcal{N};l}^2 + (2\pi pT)^2 + m^2}{t} - tk^2 a^2\right) dt \right\} \tag{B.3}$$

when $am \ll 1$. Recall that there are κ zero eigenvalues $\omega_{\mathcal{N};l}^2$. We write (B.3) as the sum of $\mathfrak{I}_1(i; a)$ and $\mathfrak{I}_2(i; a)$, where

$$\mathfrak{I}_1(i; a) = \kappa \frac{\partial}{\partial a} \left\{ a \int_0^\infty t^{-\frac{i+2-d_1}{2}} \sum_{k=1}^{\infty} \exp\left(-\frac{m^2}{t} - tk^2 a^2\right) dt \right\}$$

and

$$\mathfrak{I}_2(i; a) = \frac{\partial}{\partial a} \left\{ a \int_0^\infty t^{\frac{d_1-i}{2}-1} \sum_{k=1}^{\infty} \sum_{\substack{(l,p) \in \mathbb{N} \times \mathbb{Z} \\ \omega_{\mathcal{N};l}^2 + (2\pi pT)^2 \neq 0}} \exp\left(-\frac{\omega_{\mathcal{N};l}^2 + (2\pi pT)^2 + m^2}{t} - tk^2 a^2\right) dt \right\}.$$

For the term $\mathfrak{I}_2(i; a)$, we can use the Taylor expansion of $e^{-m^2/t}$ to write

$$\begin{aligned} \mathfrak{I}_2(i; a) = & 2a^{i-d_1} \sum_{q=0}^{\infty} \frac{(-1)^q}{q!} (am)^{2q} \left\{ \left(\frac{a\sqrt{\omega_{\mathcal{N};l}^2 + (2\pi pT)^2}}{k} \right)^{\frac{d_1-i-2q}{2}} \right. \\ & \times K_{\frac{d_1-i-2q}{2}} \left(2ka\sqrt{\omega_{\mathcal{N};l}^2 + (2\pi pT)^2} \right) \\ & \left. - 2 \frac{\left(a\sqrt{\omega_{\mathcal{N};l}^2 + (2\pi pT)^2} \right)^{\frac{d_1+2-i-2q}{2}}}{(k)^{\frac{d_1-2-i-2q}{2}}} K_{\frac{d_1+2-i-2q}{2}} \left(2ka\sqrt{\omega_{\mathcal{N};l}^2 + (2\pi pT)^2} \right) \right\}. \end{aligned}$$

For $\mathfrak{I}_1(i; a)$, notice that when s is large enough, and $am \ll 1$,

$$\begin{aligned} \mathfrak{I}_1(s; a) &= \int_0^\infty t^{-s} \sum_{k=1}^{\infty} \exp\left(-\frac{m^2}{t} - tk^2 a^2\right) dt = \int_0^\infty t^{s-2} \sum_{k=1}^{\infty} \exp\left(-tm^2 - \frac{k^2 a^2}{t}\right) dt \\ &= -\frac{1}{2} \Gamma(s-1) m^{-2s+2} + \frac{\sqrt{\pi}}{2a} \Gamma\left(s - \frac{1}{2}\right) m^{-2s+1} \\ &\quad + \frac{\sqrt{\pi}}{a} \sum_{q=0}^{\infty} \frac{(-1)^q}{q!} m^{2q} \Gamma\left(s - \frac{1}{2} + q\right) \zeta_R(2s-1+2q) \left(\frac{\pi}{a}\right)^{-2s+1-2q}. \end{aligned} \tag{B.4}$$

By analytic continuation, this formula holds for all s . From these, we find that when $r < a < R$ and $am \ll 1 \ll Rm$, the Casimir force $F_{\text{Cas}}^\infty(a)$ has the asymptotic expansion (4.19).

In the low temperature regime, we use (3.8) to transform (4.10) to

$$\begin{aligned} F_{\text{Cas}}^\infty(a) &= \frac{1}{4\pi} \frac{\partial}{\partial a} \left\{ a \int_0^\infty \sum_{k=1}^{\infty} \sum_{j \in J_*} \sum_{l=0}^{\infty} \exp\left\{-\frac{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2}{t} - tk^2 a^2\right\} dt \right. \\ &\quad + 2\sqrt{\pi} \int_0^\infty t^{-\frac{1}{2}} \sum_{k=1}^{\infty} \sum_{j \in J_*} \sum_{l=0}^{\infty} \sum_{p=1}^{\infty} \\ &\quad \times \exp\left\{-\frac{1}{t} \left(\left[\frac{\pi k}{a} \right]^2 + \omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2 \right) - t \left(\frac{p}{2T} \right)^2\right\} dt \Big\} \\ &\quad - \frac{1}{4\pi} \int_0^\infty \sum_{j \in J_*} \sum_{l=0}^{\infty} \sum_{p=1}^{\infty} \exp\left\{-\frac{\omega_{\Omega,*,j}^2 + \omega_{\mathcal{N},l}^2 + m^2}{t} - t \left(\frac{p}{2T} \right)^2\right\} dt. \end{aligned}$$

Using the same method as we derive (B.2), we find that

$$\begin{aligned} F_{\text{Cas}}^\infty(a) \sim & -\frac{1}{a^2} \sum_{i=0}^{\infty} c_{\Omega/R,*,i} \left(\frac{R}{a}\right)^{d_1-1-i} \left\{ \frac{1}{2\pi} \sum_{k=1}^{\infty} \sum_{l=0}^{\infty} \left[(d_1-i) \left(\frac{a\sqrt{\omega_{\mathcal{N};l}^2 + m^2}}{k} \right)^{\frac{d_1+1-i}{2}} \right. \right. \\ & \left. \left. \times K_{\frac{d_1+1-i}{2}} \left(2ka\sqrt{\omega_{\mathcal{N};l}^2 + m^2} \right) + 2 \frac{\left(a\sqrt{\omega_{\mathcal{N};l}^2 + m^2} \right)^{\frac{d_1+3-i}{2}}}{k^{\frac{d_1-1-i}{2}}} K_{\frac{d_1-1-i}{2}} \left(2ka\sqrt{\omega_{\mathcal{N};l}^2 + m^2} \right) \right] \right\} \end{aligned}$$

$$\begin{aligned}
& - 2^{\frac{d_1-i}{2}} \pi^{\frac{3}{2}} (aT)^{\frac{d_1-i-2}{2}} \sum_{k=1}^{\infty} \sum_{l=0}^{\infty} \sum_{p=1}^{\infty} k^2 \left(\frac{\sqrt{(\pi k)^2 + (a\omega_{\mathcal{N};l})^2 + (am)^2}}{p} \right)^{\frac{d_1-i-2}{2}} \\
& \times K_{\frac{d_1-i-2}{2}} \left(\frac{p}{Ta} \sqrt{(\pi k)^2 + (a\omega_{\mathcal{N};l})^2 + (am)^2} \right) + \frac{(2aT)^{\frac{d_1-i+1}{2}}}{2\pi} \sum_{l=0}^{\infty} \sum_{p=1}^{\infty} \left(\frac{a\sqrt{\omega_{\mathcal{N};l}^2 + m^2}}{p} \right)^{\frac{d_1-i+1}{2}} \\
& \times K_{\frac{d_1-i+1}{2}} \left(\frac{p}{T} \sqrt{\omega_{\mathcal{N};l}^2 + m^2} \right) \Bigg\}. \tag{B.5}
\end{aligned}$$

References

- [1] J. Polchinski, *String theory*, Cambridge Monographs on Mathematical Physics, Cambridge U.K. (1998).
- [2] H.-B. Cheng, *The asymptotic behavior of Casimir force in the presence of compactified universal extra dimensions*, *Phys. Lett. B* **643** (2006) 311 [[hep-th/0609099](#)] [[SPIRES](#)].
- [3] H.-B. Cheng, *The Casimir force on a piston in the spacetime with extra compactified dimensions*, *Phys. Lett. B* **668** (2008) 72 [[arXiv:0801.2810](#)] [[SPIRES](#)].
- [4] S.A. Fulling and K. Kirsten, *Comment on: “The Casimir force on a piston in the spacetime with extra compactified dimensions”* [*Phys. Lett. B* **668** (2008) 72], *Phys. Lett. B* **671** (2009) 179 [[arXiv:0811.0779](#)] [[SPIRES](#)].
- [5] K. Kirsten and S.A. Fulling, *Kaluza-Klein models as pistons*, *Phys. Rev. D* **79** (2009) 065019 [[arXiv:0901.1902](#)] [[SPIRES](#)].
- [6] M. Frank, I. Turan and L. Ziegler, *The Casimir force in Randall Sundrum models*, *Phys. Rev. D* **76** (2007) 015008 [[arXiv:0704.3626](#)] [[SPIRES](#)].
- [7] R. Linares, H.A. Morales-Tecotl and O. Pedraza, *Casimir force for a scalar field in warped brane worlds*, *Phys. Rev. D* **77** (2008) 066012 [[arXiv:0712.3963](#)] [[SPIRES](#)].
- [8] M. Frank, N. Saad and I. Turan, *The Casimir force in Randall Sundrum models with $q + 1$ dimensions*, *Phys. Rev. D* **78** (2008) 055014 [[arXiv:0807.0443](#)] [[SPIRES](#)].
- [9] K. Poppenhaefer, S. Hossenfelder, S. Hofmann and M. Bleicher, *The Casimir effect in the presence of compactified universal extra dimensions*, *Phys. Lett. B* **582** (2004) 1 [[hep-th/0309066](#)] [[SPIRES](#)].
- [10] L. Perivolaropoulos, *Vacuum energy, the cosmological constant and compact extra dimensions: constraints from Casimir effect experiments*, *Phys. Rev. D* **77** (2008) 107301 [[arXiv:0802.1531](#)] [[SPIRES](#)].
- [11] F. Pascoal, L.F.A. Oliveira, F.S.S. Rosa and C. Farina, *Estimate for the size of the compactification radius of a one extra dimension universe*, *Braz. J. Phys.* **38** (2008) 581 [[hep-th/0701181](#)] [[SPIRES](#)].
- [12] A. Edery and V. Marachevsky, *Compact dimensions and the Casimir effect: the Proca connection*, *JHEP* **12** (2008) 035 [[arXiv:0810.3430](#)] [[SPIRES](#)].
- [13] C. Ccapa Ttira, C.D. Fosco, A.P.C. Malbouisson and I. Roditi, *Quantum effects in QED_{3+1} with an extra compactified dimension*, [arXiv:0812.4746](#) [[SPIRES](#)].

- [14] L.P. Teo, *Finite temperature Casimir effect in spacetime with extra compactified dimensions*, *Phys. Lett. B* **672** (2009) 190 [[arXiv:0812.4641](#)] [[SPIRES](#)].
- [15] L.P. Teo, *Finite temperature Casimir effect in Kaluza-Klein spacetime*, *Nucl. Phys. B* **819** (2009) 431 [[arXiv:0901.2195](#)] [[SPIRES](#)].
- [16] E. Elizalde, S.D. Odintsov and A.A. Saharian, *Repulsive Casimir effect from extra dimensions and Robin boundary conditions: from branes to pistons*, *Phys. Rev. D* **79** (2009) 065023 [[arXiv:0902.0717](#)] [[SPIRES](#)].
- [17] A.A. Saharian, *Casimir densities in brane models with compact internal spaces*, [arXiv:0811.4031](#) [[SPIRES](#)].
- [18] F.S. Accetta and E.W. Kolb, *Finite temperature instability for compactification*, *Phys. Rev. D* **34** (1986) 1798 [[SPIRES](#)].
- [19] W.D. Goldberger and I.Z. Rothstein, *Quantum stabilization of compactified AdS_5* , *Phys. Lett. B* **491** (2000) 339 [[hep-th/0007065](#)] [[SPIRES](#)].
- [20] S.-K. Nam, *Casimir force in compact noncommutative extra dimensions and radius stabilization*, *JHEP* **10** (2000) 044 [[hep-th/0008083](#)] [[SPIRES](#)].
- [21] W.-H. Huang, *Finite-temperature Casimir effect on the radius stabilization of noncommutative torus*, *JHEP* **11** (2000) 041 [[hep-th/0011037](#)] [[SPIRES](#)].
- [22] E. Ponton and E. Poppitz, *Casimir energy and radius stabilization in five and six dimensional orbifolds*, *JHEP* **06** (2001) 019 [[hep-ph/0105021](#)] [[SPIRES](#)].
- [23] R. Hofmann, P. Kanti and M. Pospelov, *(De-)stabilization of an extra dimension due to a Casimir force*, *Phys. Rev. D* **63** (2001) 124020 [[hep-ph/0012213](#)] [[SPIRES](#)].
- [24] W.-H. Huang, *Casimir effect on the radius stabilization of the noncommutative torus*, *Phys. Lett. B* **497** (2001) 317 [[hep-th/0010160](#)] [[SPIRES](#)].
- [25] E. Elizalde, *Matching the observational value of the cosmological constant*, *Phys. Lett. B* **516** (2001) 143 [[hep-th/0007094](#)] [[SPIRES](#)].
- [26] C.L. Gardner, *Primordial inflation and present-day cosmological constant from extra dimensions*, *Phys. Lett. B* **524** (2002) 21 [[hep-th/0105295](#)] [[SPIRES](#)].
- [27] K.A. Milton, *Dark energy as evidence for extra dimensions*, *Grav. Cosmol.* **9** (2003) 66 [[hep-ph/0210170](#)] [[SPIRES](#)].
- [28] M.R. Setare, *Stress-energy tensor for parallel plate on background of conformally flat brane-world geometries and cosmological constant problem*, *Eur. Phys. J. C* **38** (2004) 373 [[SPIRES](#)].
- [29] G. Mahajan, S. Sarkar and T. Padmanabhan, *Casimir effect confronts cosmological constant*, *Phys. Lett. B* **641** (2006) 6 [[astro-ph/0604265](#)] [[SPIRES](#)].
- [30] M. Doran and J. Jaeckel, *What measurable zero point fluctuations can(not) tell us about dark energy*, *JCAP* **08** (2006) 010 [[astro-ph/0605711](#)] [[SPIRES](#)].
- [31] V.N. Marachevsky, *Casimir interaction of two plates inside a cylinder*, *Phys. Rev. D* **75** (2007) 085019 [[hep-th/0703158](#)] [[SPIRES](#)].
- [32] R.M. Cavalcanti, *Casimir force on a piston*, *Phys. Rev. D* **69** (2004) 065015 [[quant-ph/0310184](#)] [[SPIRES](#)].
- [33] M. Bordag, U. Mohideen and V.M. Mostepanenko, *New developments in the Casimir effect*, *Phys. Rept.* **353** (2001) 1 [[quant-ph/0106045](#)] [[SPIRES](#)].

- [34] S. Blau, M. Visser and A. Wipf, *Zeta functions and the Casimir energy*, *Nucl. Phys. B* **310** (1988) 163 [[arXiv:0906.2817](#)] [[SPIRES](#)].
- [35] J.S. Dowker and G. Kennedy, *Finite temperature and boundary effects in static space-times*, *J. Phys. A* **11** (1978) 895 [[SPIRES](#)].
- [36] J.S. Dowker, *Finite temperature and vacuum effects in higher dimensions*, *Class. Quant. Grav.* **1** (1984) 359 [[SPIRES](#)].
- [37] J.S. Dowker and J.P. Schofield, *Chemical potentials in curved space*, *Nucl. Phys. B* **327** (1989) 267 [[SPIRES](#)].
- [38] K. Kirsten, *Grand thermodynamic potential in a static space-time with boundary*, *Class. Quant. Grav.* **8** (1991) 2239 [[SPIRES](#)].
- [39] K. Kirsten, *Casimir effect at finite temperature*, *J. Phys. A* **24** (1991) 3281 [[SPIRES](#)].
- [40] E. Elizalde, *A very simple computation of the Casimir effect*, *Nuovo Cim. B* **104** (1989) 685 [[SPIRES](#)].
- [41] K. Kirsten, *Inhomogeneous multidimensional Epstein zeta functions*, *J. Math. Phys.* **32** (1991) 3008 [[SPIRES](#)].
- [42] E. Elizalde, *An extension of the Chowla-Selberg formula useful in quantizing with the Wheeler-DeWitt equation*, *J. Phys. A* **27** (1994) 3775 [[hep-th/9402155](#)] [[SPIRES](#)].
- [43] E. Elizalde, *Multidimensional extension of the generalized Chowla-Selberg formula*, *Commun. Math. Phys.* **198** (1998) 83 [[hep-th/9707257](#)] [[SPIRES](#)].
- [44] S.C. Lim and L.P. Teo, *Three dimensional Casimir piston for massive scalar fields*, to appear in *Annals Phys.* [[arXiv:0807.3613](#)] [[SPIRES](#)].
- [45] J. Feinberg, A. Mann and M. Revzen, *Casimir effect: the classical limit*, *Annals Phys.* **288** (2001) 103 [[hep-th/9908149](#)] [[SPIRES](#)].
- [46] I. Klich, J. Feinberg, A. Mann and M. Revzen, *Casimir energy of a dilute dielectric ball with uniform velocity of light at finite temperature*, *Phys. Rev. D* **62** (2000) 045017 [[hep-th/0001019](#)] [[SPIRES](#)].
- [47] M. Schaden and L. Spruch, *Classical Casimir effect: the interaction of ideal parallel walls at a finite temperature*, *Phys. Rev. A* **65** (2002) 034101.
- [48] A. Scardicchio and R.L. Jaffe, *Casimir effects: an optical approach II. Local observables and thermal corrections*, *Nucl. Phys. B* **743** (2006) 249 [[quant-ph/0507042](#)] [[SPIRES](#)].
- [49] B. Geyer, G.L. Klimchitskaya and V.M. Mostepanenko, *Thermal Casimir effect in ideal metal rectangular boxes*, *Eur. Phys. J. C* **57** (2008) 823 [[arXiv:0808.3754](#)] [[SPIRES](#)].
- [50] S.C. Lim and L.P. Teo, *Finite-temperature Casimir effect in piston geometry and its classical limit*, *Eur. Phys. J. C* **60** (2009) 323 [[SPIRES](#)].
- [51] E. Álvarez and F.D. Mazzitelli, *Long range Casimir force induced by transverse electromagnetic modes*, *Phys. Rev. D* **79** (2009) 045019 [[arXiv:0901.2641](#)] [[SPIRES](#)].