

## Improved constraints on $Z$ bosons from electroweak precision data

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

JHEP08(2009)017

(<http://iopscience.iop.org/1126-6708/2009/08/017>)

[The Table of Contents](#) and [more related content](#) is available

Download details:

IP Address: 80.92.225.132

The article was downloaded on 03/04/2010 at 10:22

Please note that [terms and conditions apply](#).

## Improved constraints on $Z'$ bosons from electroweak precision data

---

Jens Erler,<sup>a</sup> Paul Langacker,<sup>b</sup> Shoaib Munir<sup>a</sup> and Eduardo Rojas<sup>a</sup>

<sup>a</sup>*Departamento de Física Teórica, Instituto de Física,  
Universidad Nacional Autónoma de México, 04510 México D.F., México*

<sup>b</sup>*School of Natural Sciences, Institute for Advanced Study,  
Einstein Drive, Princeton, NJ 08540, U.S.A.*

*E-mail:* [erler@fisica.unam.mx](mailto:erler@fisica.unam.mx), [pgl@ias.edu](mailto:pgl@ias.edu), [smunir@fisica.unam.mx](mailto:smunir@fisica.unam.mx),  
[eduardor@fisica.unam.mx](mailto:eduardor@fisica.unam.mx)

**ABSTRACT:** We analyze various models with an extra  $U(1)$  gauge symmetry in addition to the Standard Model (SM) gauge group at low energies, and impose limits on the mass of the neutral  $Z'$  boson,  $M_{Z'}$ , predicted in all such models, and on the  $Z - Z'$  mixing angle,  $\theta_{ZZ'}$ . The precision electroweak (EW) data strongly constrain  $\theta_{ZZ'}$  to very small values and for most models we find lower limits on  $M_{Z'}$  of  $\mathcal{O}(1 \text{ TeV})$ . In one case we obtain a somewhat better fit than in the SM (although this is only marginally statistically significant) and here we find a weak upper limit at the 90% C.L.

**KEYWORDS:** Beyond Standard Model, GUT, Standard Model

**ARXIV EPRINT:** [0906.2435](https://arxiv.org/abs/0906.2435)

---

## Contents

<b>1</b>	<b>Overview</b>	<b>1</b>
<b>2</b>	<b>Extended Higgs sectors and exotics</b>	<b>4</b>
<b>3</b>	<b>Details of the analyses</b>	<b>6</b>
<b>4</b>	<b>Results and discussion</b>	<b>9</b>

---

## 1 Overview

With the advent of the LHC, particle physics has entered a new exciting era. Within a few years of data accumulation, the LHC should be able to test and constrain many types of new physics beyond the SM. In particular, the discovery reach for extra neutral gauge bosons is exceptional. Searches for a high invariant dilepton mass peak in about  $100\text{ fb}^{-1}$  of accumulated data will find or exclude  $Z'$  bosons up to about 5 TeV, and a luminosity upgraded LHC (by roughly a factor of 10) can extend the reach by another TeV [1]. After a potential discovery, the LHC will have some diagnostic means to narrow down the underlying  $Z'$  model [2] by studying, for example, leptonic forward-backward asymmetries (for reviews, see refs. [3]–[5]) and heavy quark final states [6, 7]. Furthermore, angular distributions of Drell-Yan leptons may help to discriminate a  $Z'$  against spin-0 (sneutrino) and spin-2 (Kaluza-Klein graviton) resonances [8]. However, the hadronic LHC environment will make it difficult to specify the  $Z'$  properties completely or with satisfactory precision. Electroweak precision measurements will therefore play an important complementary rôle in this context. Already these data give rise to strong constraints on possible  $Z'$  and in many cases yield the best limits. In this paper we will revisit the EW precision data in the presence of  $Z'$  bosons. This is motivated by much higher precision in several measurements (such as from the Tevatron) compared to previous studies [9, 10] and also by significant shifts and new measurements at low energies, a sector with increasing impact on global analyses of this type.

Neutral gauge sectors with an additional U(1) symmetry in addition to the SM hypercharge  $U(1)_Y$  and an associated  $Z'$  gauge boson are among the best motivated extensions of the SM. They are predicted in most Grand Unified Theories (GUTs) and appear copiously in superstring theories. An especially compelling motivation for extended gauge theories came from the development of GUTs larger than the original SU(5) model, such as SO(10) or  $E_6$ , which allow the SM gauge group to be embedded into them [11]. There is an extensive range of models with an extra U(1) symmetry (for a review, see [5] and references therein). Among these, models based on the  $E_6$  GUT group and left-right symmetry

groups have been extensively pursued in the literature and are particularly significant from the point of view of LHC phenomenology. In the context of supersymmetry, this class of models also arises [12] from requiring the extra U(1) symmetry to provide a solution to the  $\mu$ -problem [13], to forbid rapid (dimension 4) proton decay, to protect all fields by chirality and supersymmetry from acquiring high scale masses, and to be consistent with anomaly cancellation, gauge coupling unification and family universality (to avoid the strong constraints from the flavor-changing neutral current (FCNC) sector [14]). The models studied here include:

**$E_6$  based models:** extra U(1) gauge symmetries appear in the decomposition of the SO(10) [15] or  $E_6$  [16, 17] GUT groups.  $E_6$  contains the maximal subgroup  $\text{SO}(10) \times \text{U}(1)_\psi$ , and SO(10) can be further decomposed into its  $\text{SU}(5) \times \text{U}(1)_\chi$  maximal subgroup. We are considering models in which the linear combination,

$$\text{U}(1)' = \cos \beta \text{U}(1)_\chi + \sin \beta \text{U}(1)_\psi, \tag{1.1}$$

survives down to the EW scale, using a convention in which the mixing angle in eq. (1.1) satisfies<sup>1</sup>  $-90^\circ < \beta \leq 90^\circ$ . The full  $E_6$  symmetry would impose strong constraints on these models, which are often unrealistic. For the purpose of our analyses, we are, however, mostly interested in the effects, phenomenology, and constraints associated directly with the  $Z'$  bosons and not in other aspects of these models<sup>2</sup>.

$Z_\chi$ :  $\beta = 0^\circ \implies Z' = Z_\chi$ , which is also the unique solution to the conditions of (i) family universality, (ii) no extra matter other than right handed neutrino, (iii) absence of gauge and mixed gauge/gravitational anomalies and (iv) orthogonality to the hypercharge generator.

$Z_\psi$ :  $\beta = 90^\circ \implies Z' = Z_\psi$ , possessing only axial vector couplings to ordinary fermions. As discussed in section 4, it is among the least constrained by the precision data.

$Z_\eta$ :  $\beta = -\arctan \sqrt{5/3} \approx -52.2^\circ \implies Z' = \sqrt{3/8} Z_\chi - \sqrt{5/8} Z_\psi \equiv Z_\eta$ , occurring in Calabi-Yau compactifications [19] of the heterotic string [20] if  $E_6$  breaks directly to a rank 5 subgroup [21] via the Hosotani mechanism [22].

$Z_I$ :  $\beta = \arctan \sqrt{3/5} \approx 37.8^\circ \implies Z' = \sqrt{5/8} Z_\chi + \sqrt{3/8} Z_\psi \equiv -Z_I$ , which is orthogonal to the  $Z_\eta$ . This boson [16] has the defining property of vanishing couplings to up-type quarks. Its production is thus suppressed at hadron colliders, especially at the Tevatron since in high-energy  $p\bar{p}$  collisions  $Z'$  production through down quarks is suppressed by a factor of 25 relative to up quarks [3].

$Z_S$ : a supersymmetric model with a secluded U(1)' breaking sector and a large supersymmetry breaking  $A$ -term was introduced (i) to provide an approximately

---

<sup>1</sup>Note, that by restricting  $\beta$  to a semi-circle makes the sign of the mixing angle  $\theta_{ZZ'}$  physically meaningful.

<sup>2</sup>There are classes of  $Z'$  models closely related to the  $E_6$  ones motivated by minimal gauge unification [5, 12] which generically require more than one kind of U(1)' breaking singlet field. The issue of obtaining a suitable symmetry breaking pattern is discussed in [18].

flat potential allowing the generation of a  $Z$ - $Z'$  mass hierarchy [23] and (ii) to produce a strong first order EW phase transition for EW baryogenesis [24]. Such a sector is obtained if the right-handed  $\nu$  has  $-2\times$  the  $U(1)'$  charge of the other SM singlet in a **27** representation of  $E_6$ , giving  $\beta = \arctan \sqrt{5/27} \approx 23.3^\circ \implies Z' = \sqrt{27/32} Z_\chi + \sqrt{5/32} Z_\psi \equiv Z_S$ , which is numerically close to the  $Z_I$ .

$Z_N$ :  $\beta = \arctan \sqrt{15} \approx 75.5^\circ \implies Z' = (Z_\chi + \sqrt{15} Z_\psi)/4 \equiv Z_N$ , which is a consequence of choosing the right-handed neutrinos to have zero  $U(1)'$  charges so that they can acquire very heavy Majorana masses [25]–[27] and are thus suitable to take part in the standard seesaw mechanism [28] with three naturally light neutrinos. The  $Z_N$  boson also appears in a model referred to as the ESSM [29] or the  $E_6$ SSM [30].

$Z_R$ : all models discussed so far assume negligible kinetic mixing, i.e., the absence of a cross term,

$$-\frac{\sin \chi}{2} F'_{\mu\nu} F_Y^{\mu\nu}, \tag{1.2}$$

between the gauge kinetic terms for the  $U(1)'$  and the  $U(1)_Y$  gauge bosons [31]–[34]. This is motivated by the orthogonality of all  $U(1)$  subgroups within a simple GUT group like  $E_6$ . A usually very small kinetic term arises at the two-loop level from the renormalization group evolution of the gauge couplings [35]. An exception in the case in which incomplete GUT multiplets survive below the unification scale and in which a relatively large kinetic mixing term can be generated [36]. In any case, there is no general reason to ignore kinetic mixing and we now address it in a different but equivalent formalism: one can always redefine the gauge boson fields to remove any term of the form (1.2). The effects due to  $\sin \chi \neq 0$  then manifest themselves in the  $U(1)'$  charges which will in general have a non-trivial hypercharge component. In the  $E_6$  context, for example, one can write the  $Z'$  as the general (family-universal) combination [10],

$$Z' = \cos \alpha \cos \beta Z_\chi + \sin \alpha \cos \beta Z_Y + \sin \beta Z_\psi. \tag{1.3}$$

The restriction to  $\beta = 0^\circ$  corresponds to general  $SO(10)$  based models. Specifying further to  $\alpha = \arctan \sqrt{3/2} \approx 50.8^\circ \implies Z' = \sqrt{2/5} Z_\chi + \sqrt{3/5} Z_Y \equiv Z_R$ , which couples to charges proportional to the diagonal (third) component of right-handed isospin,  $SU(2)_R$ . We are unaware of this case having been introduced in the literature. But as we will discuss in section 4, the resulting  $Z'$  gives a reasonably good fit, and technically even a finite 90% C.L. upper limit can be set on its mass.

$Z_{LR}$ : models with left-right symmetry (reviewed in ref. [37]) are based on the gauge group  $SU(3)_C \times SU(2)_L \times SU(2)_R \times U(1)_{B-L} \subset SO(10)$  and contain a boson,  $Z_{LR} \equiv \sqrt{3/5} (\bar{\alpha} Z_R - Z_{B-L}/2\bar{\alpha})$ . Here  $B$  and  $L$  coincide, respectively, with baryon and lepton number for the ordinary fermions, and  $Z' = \sqrt{3/5} Z_\chi - \sqrt{2/5} Z_Y \equiv -\sqrt{3/8} Z_{B-L}$  is obtained by choosing  $\alpha = -\arctan \sqrt{2/3} \approx -39.2^\circ$  ( $B$  or  $L$  individually cannot be parametrized in this way). The parameter  $\bar{\alpha} =$

$\sqrt{g_R^2/g_L^2 \cot^2 \theta_W - 1}$ , with  $\theta_W$  the weak mixing angle, gives the coupling strength in terms of the  $SU(2)_{L,R}$  gauge couplings,  $g_{L,R}$ . Manifest left-right symmetry (which we will assume) requires  $g_L = g_R$ , while the very strong coupling limit ( $\bar{\alpha}, g_R/g_L \rightarrow \infty$ ) implies  $Z_{LR} \rightarrow Z_R$ .

$Z_\psi$ : a leptophobic  $Z'$  has vanishing  $U(1)'$  charges to charged leptons and left-handed neutrinos. One version of this idea [36] is a variation of the  $Z_\eta$  model with kinetic mixing added. The choice  $(\alpha, \beta) = (\arctan \sqrt{8/27}, -\arctan \sqrt{9/7}) \approx (28.6^\circ, -48.6^\circ) \implies Z' = \sqrt{27/80} Z_\chi + 1/\sqrt{10} Z_Y - 3/4 Z_\psi \equiv Z_\psi$ . The effects of a leptophobic  $Z'$  are very difficult to observe but it can be searched for in the dijet [38] and  $t\bar{t}$  [39] channels at hadron colliders. Moreover, mixing effects at LEP 1 strongly constrain  $\theta_{ZZ'}$  even in this case.

**Sequential  $Z'$ :** the  $Z_{SM}$  boson is defined to have the same couplings to fermions as the SM  $Z$ . Such a boson is not expected in the context of gauge theories unless it has different couplings to exotic fermions than the ordinary  $Z$ . However, it serves as a useful reference case when comparing constraints from various sources. It could also play the role of an excited state of the ordinary  $Z$  in models of compositeness or with extra dimensions at the weak scale.

**A superstring  $Z'$ :** there is a family non-universal  $Z_{string}$  boson appearing in a specific model [40] based on the free fermionic string construction with real fermions. This model has been investigated in considerable detail [41, 42] with the goal of understanding some of the characteristics of (weakly coupled) string theories, and of contrasting them with the more conventional ideas such as GUTs. While this specific model itself is not realistic (for example, it fails to produce an acceptable fermion mass spectrum) the predicted  $Z_{string}$  it contains is itself not ruled out (ignoring issues related to CP violation and FCNCs [14]). Its coupling strength is predicted and so are its fermion couplings. Such a  $Z_{string}$  can be naturally at the electroweak scale [43, 44].

## 2 Extended Higgs sectors and exotics

The incorporation of (one or more) extra gauge group(s) in the models listed above generally warrants an extended fermionic sector for two main reasons: (i) cancellation of gauge and mixed gauge-gravitational anomalies to assure quantum consistency of the theory, and (ii) in the context of low-energy supersymmetry, the unification of gauge couplings at high energies. Provided that all fermions belong to complete  $E_6$  representations the anomalies are cancelled automatically. In a bottom-up approach, however, the condition of anomaly cancellation restricts the  $U(1)'$  charge assignments of the SM fermions and the exotics [12, 45].

The structure of the Higgs sector of the underlying model is important as it may affect the  $\rho_0$  parameter,

$$\rho_0 \equiv \frac{\sum_i (t_i^2 - t_{3i}^2 + t_i) |\langle \phi_i \rangle|^2}{\sum_i 2t_{3i}^2 |\langle \phi_i \rangle|^2}, \tag{2.1}$$

$Z'$	$C$	range	restricted range ( $\omega = 0, \tau \geq \frac{1}{2}$ )
$Z_\chi$	$\frac{2}{\sqrt{10}} (1 - \frac{5}{2}\omega)$	$\frac{2}{\sqrt{10}} [-\frac{3}{2}, +1]$	$\frac{2}{\sqrt{10}}$
$Z_\psi$	$\sqrt{\frac{2}{3}} (1 - 2\tau - \frac{3}{2}\omega)$	$\sqrt{\frac{2}{3}} [-1, +1]$	$\sqrt{\frac{2}{3}} [-1, 0]$
$Z_\eta$	$-\frac{1}{\sqrt{15}} (1 - 5\tau)$	$\frac{1}{\sqrt{15}} [-1, +4]$	$\frac{1}{\sqrt{15}} [\frac{3}{2}, +4]$
$Z_I$	$\tau + 2\omega - 1$	$[-1, 1]$	$[-\frac{1}{2}, 0]$
$Z_S$	$\frac{7}{\sqrt{60}} (1 - \frac{5}{7}\tau - \frac{15}{7}\omega)$	$\frac{7}{\sqrt{60}} [-\frac{8}{7}, +1]$	$\frac{1}{\sqrt{15}} [+1, +\frac{9}{4}]$
$Z_N$	$\frac{3}{\sqrt{10}} (1 - \frac{5}{3}(\tau + \omega))$	$\frac{3}{\sqrt{10}} [-\frac{2}{3}, +1]$	$\frac{1}{\sqrt{10}} [-2, +\frac{1}{2}]$
$Z_R$	$1 - \omega$	$[0, +1]$	$1$
$Z_{LR}$	$\sqrt{\frac{3}{5}}\bar{\alpha} [1 - (1 + \frac{1}{\bar{\alpha}^2})\omega]$	$\sqrt{\frac{3}{5}} [-\frac{1}{\bar{\alpha}}, +\bar{\alpha}]$	$\sqrt{\frac{3}{5}}\bar{\alpha}$
$Z_\psi$	$\sqrt{\frac{3}{2}}\tau$	$\sqrt{\frac{3}{2}} [0, +1]$	$\sqrt{\frac{3}{2}} [\frac{1}{2}, +1]$

**Table 1.** Special Higgs sectors for  $E_6$  based models. The third column shows the most general range for  $C$  if all three Higgs doublets in a **27** representation participate in spontaneous symmetry breaking. The last column corresponds to the restricted range appropriate for supersymmetry inspired models.

where  $t_i$  ( $t_{3i}$ ) is (the third component of) the weak isospin of the Higgs field  $\phi_i$ , and which enters the neutral and charged ( $M_W$ ) gauge boson mass interdependence,

$$M_0 = \frac{M_W}{\sqrt{\rho_0} \cos \theta_W}. \quad (2.2)$$

$\rho_0 = 1$  corresponds to a Higgs sector with only SU(2) doublets and singlets. In that case the mass parameter  $M_0$  (the ordinary  $Z$  mass in the absence of  $Z - Z'$  mixing) is predicted. In general there is mixing between the mass eigenstates of the  $Z'$  and the  $Z$  given by [46],

$$\tan^2 \theta_{ZZ'} = \frac{M_0^2 - M_Z^2}{M_{Z'}^2 - M_0^2}. \quad (2.3)$$

Allowing  $\rho_0$  as an additional fit parameter means that the Higgs sector of the model is arbitrary and may include higher-dimensional Higgs representations. In addition, the presence of non-degenerate multiplets of heavy fermions or scalars will affect the  $W$  and  $Z$  self energies at the loop level, and therefore contribute to the  $T$  parameter [47]. With the current data set, the phenomenological consequences of  $\rho_0$  and  $T$  are indistinguishable and values quoted for  $\rho_0$  really apply to the combination  $\rho_0/(1 - \alpha T)$ .

If the U(1)' charge assignments of the Higgs fields,  $Q'_i$ , are known in a specific model, then there exists an additional constraint [48],

$$\theta_{ZZ'} = C \frac{g_2 M_Z^2}{g_1 M_{Z'}^2}, \quad (2.4)$$

where  $g_1 = g_L / \cos \theta_W$  and where  $g_2 = \sqrt{5/3} g_1 \sin \theta_W \sqrt{\lambda}$  is the  $U(1)'$  gauge coupling. The latter is given in terms of  $\lambda$  which is of order unity (we will set  $\lambda = 1$  as is conventionally done), and in fact  $\lambda \sim 1$  in GUT models breaking directly to  $SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)'$ .  $C$  is a function of vacuum expectation values (VEVs) of the Higgs fields and the  $Q'_i$ ,

$$C = -\frac{\sum_i t_{3i} Q'_i |\langle \phi_i \rangle|^2}{\sum_i t_{3i}^2 |\langle \phi_i \rangle|^2}. \quad (2.5)$$

As an illustration, for the  $E_6$  based models one may restrict oneself to the case where the Higgs fields arise from a **27** representation. The  $U(1)'$  quantum numbers are then predicted and eq. (2.5) receives contributions from the VEVs of three Higgs doublets,  $x \equiv \langle \phi_\nu \rangle$ ,  $v \equiv \langle \phi_N \rangle$  and  $\bar{v} \equiv \langle \phi_{\bar{N}} \rangle$ , respectively, in correspondence with the standard lepton doublet, as well as the two doublets contained in the  $\bar{\mathbf{5}}$  and  $\mathbf{5}$  of  $SU(5) \subset E_6$ . They satisfy the sum rule,  $|v|^2 + |\bar{v}|^2 + |x|^2 = (\sqrt{2} G_F)^{-1} = (246.22 \text{ GeV})^2$ , and we introduce the ratios,

$$\tau = \frac{|\bar{v}|^2}{|v|^2 + |\bar{v}|^2 + |x|^2} \quad (0 \leq \tau \leq 1), \quad (2.6)$$

$$\omega = \frac{|x|^2}{|v|^2 + |\bar{v}|^2 + |x|^2} \quad (0 \leq \omega \leq 1). \quad (2.7)$$

In supersymmetric models one usually assumes  $x = \omega = 0$  to avoid spontaneous breaking of lepton number and problems with charged current universality, as well as  $\bar{v} \geq v$  (implying  $\tau \geq 1/2$ ) to avoid non-perturbative values for the top quark Yukawa coupling. The resulting ranges for  $C$  are shown in table 1.

### 3 Details of the analyses

The theoretical evaluation uses the special purpose FORTRAN package GAPP [49] dedicated to the Global Analysis of Particle Properties. All experimental and theoretical uncertainties are included and their correlations accounted for. All errors have been added in quadrature and in most (but not all) cases been treated as Gaussian. The effects of the  $Z'$  bosons are taken into account as first order perturbations to the SM expressions.

The most stringent indirect constraints on  $M_{Z'}$  come from low-energy weak neutral current experiments displayed in table 2 together with other non  $Z$ -pole observables. The first set shown are the most recent combinations of  $M_W$  [50] and the top quark mass,  $m_t$  [51].

The second set are effective four-Fermi operator coefficients ( $g_{L,R}^2$ ) and cross section ratios ( $\kappa, R^\nu, R^{\bar{\nu}}$ ) from neutrino and anti-neutrino deep inelastic scattering ( $\nu$ -DIS) at FNAL [52, 53] and CERN [54, 55]. The NuTeV [53] results are very preliminary. We have updated ref. [53] to account for the recently measured strange quark asymmetry [56]. The incorporation of other effects like more recent QED radiative corrections [57, 58] and parton distribution functions [59] (allowing some level of charge symmetry violation) are likely to decrease the  $2\sigma$  deviation in  $g_L^2$  shown in the table. On the other hand, the world average [60] of the  $K_{e3}$  branching fraction has been corrected upwards several times in the previous years, making for a larger correction for the  $\nu_e$  ( $\bar{\nu}_e$ ) contamination of the



Quantity	Group(s)	Value	Standard Model	pull
$m_t$ [GeV]	Tevatron	$173.1 \pm 1.4$	$173.1 \pm 1.4$	0.0
$M_W$ [GeV]	Tevatron	$80.432 \pm 0.039$	$80.380 \pm 0.015$	1.3
$M_W$ [GeV]	LEP 2	$80.376 \pm 0.033$		-0.1
$g_L^2$	NuTeV	$0.3010 \pm 0.0015$	$0.3039 \pm 0.0002$	-2.0
$g_R^2$	NuTeV	$0.0308 \pm 0.0011$	0.0300	0.7
$\kappa$	CCFR	$0.5820 \pm 0.0041$	$0.5831 \pm 0.0003$	-0.3
$R^\nu$	CDHS	$0.3096 \pm 0.0043$	$0.3091 \pm 0.0002$	0.1
$R^\nu$	CHARM	$0.3021 \pm 0.0041$		-1.7
$R^{\bar{\nu}}$	CDHS	$0.384 \pm 0.018$	$0.3861 \pm 0.0001$	-0.1
$R^{\bar{\nu}}$	CHARM	$0.403 \pm 0.016$		1.1
$R^{\bar{\nu}}$	CDHS 1979	$0.365 \pm 0.016$	$0.3815 \pm 0.0001$	-1.0
$g_V^{\nu e}$	CHARM II + older	$-0.040 \pm 0.015$	$-0.0397 \pm 0.0003$	0.0
$g_A^{\nu e}$	CHARM II + older	$-0.507 \pm 0.014$	$-0.5064 \pm 0.0001$	0.0
$Q_W(\text{Ti})$	Oxford + Seattle	$-116.4 \pm 3.6$	-116.8	0.1
$Q_W(\text{Cs})$	Boulder	$-73.16 \pm 0.35$	$-73.16 \pm 0.03$	0.0
$Q_W(e)$	SLAC E158	$-0.0403 \pm 0.0053$	$-0.0472 \pm 0.0005$	1.3
$\cos \gamma C_{1d} - \sin \gamma C_{1u}$	Young et al.	$0.342 \pm 0.063$	$0.3885 \pm 0.0002$	-0.7
$\sin \gamma C_{1d} + \cos \gamma C_{1u}$	Young et al.	$-0.0285 \pm 0.0043$	$-0.0335 \pm 0.0001$	1.2
CKM unitarity	KLOE dominated	$1.0000 \pm 0.0006$	1	0.0
$(g_\mu - 2 - \alpha/\pi)/2$	BNL E821	$4511.07 \pm 0.74$	$4509.04 \pm 0.09$	2.7

**Table 2.** Non  $Z$ -pole precision observables from FNAL, CERN, SLAC, JLab, and elsewhere. Shown are the experimental results, the SM predictions, and the pulls. The SM errors are from the parametric uncertainties in the Higgs boson and quark masses and in the strong and electromagnetic coupling constants evaluated at  $M_Z$ .

dominantly  $\nu_\mu$  ( $\bar{\nu}_\mu$ ) beams, and which by itself would be indicative of an increase in the deviation. More precise statements about the size and the sign of the net effect of these corrections will only be possible after the completion of the re-analysis of the NuTeV result, which is currently in progress [61]. The  $g_{V,A}^{\nu e}$  in the third set are effective four-Fermi couplings for elastic  $\nu$ - $e$  scattering [62].

$Q_W$  denote so-called weak charges measured in atomic parity violation [63]–[65] and polarized Møller scattering [66]. The extracted value for  $Q_W(\text{Cs})$  has shifted very recently from a  $1\sigma$  deviation to perfect agreement with the SM. This is due to the state of the art atomic structure calculation of ref. [67] which also brought the atomic theory uncertainty below the measurement error. This is of significant importance for  $Z'$  studies, since they easily affect and conversely are strongly constrained by precision weak charges. A previous  $2\sigma$  deviation in  $Q_W(\text{Cs})$  based on the same measurement [65] but a different evaluation of the atomic physics [68] even indicated the presence of a  $Z'$  [10]. Related to nuclear weak charges are the two linear combinations of four-Fermi couplings  $C_{1u}$  and  $C_{1d}$  (with  $\tan \gamma \approx 0.445$  [69]) which are the result of a global analysis of parity violating electron scattering experiments on nuclear fixed targets [70].

Quantity	Group(s)	Value	Standard Model	pull
$M_Z$ [GeV]	LEP 1	$91.1876 \pm 0.0021$	$91.1874 \pm 0.0021$	0.1
$\Gamma_Z$ [GeV]	LEP 1	$2.4952 \pm 0.0023$	$2.4954 \pm 0.0010$	-0.1
$\sigma_{\text{had}}$ [nb]	LEP 1	$41.541 \pm 0.058$	$41.483 \pm 0.008$	1.6
$R_e$	LEP 1	$20.804 \pm 0.050$	$20.736 \pm 0.010$	1.4
$R_\mu$	LEP 1	$20.785 \pm 0.033$	$20.736 \pm 0.010$	1.5
$R_\tau$	LEP 1	$20.764 \pm 0.045$	$20.782 \pm 0.010$	-0.4
$A_{\text{FB}}(e)$	LEP 1	$0.0145 \pm 0.0025$	$0.0163 \pm 0.0002$	-0.7
$A_{\text{FB}}(\mu)$	LEP 1	$0.0169 \pm 0.0013$		0.5
$A_{\text{FB}}(\tau)$	LEP 1	$0.0188 \pm 0.0017$		1.5
$R_b$	LEP 1 + SLD	$0.21629 \pm 0.00066$	$0.21578 \pm 0.00005$	0.8
$R_c$	LEP 1 + SLD	$0.1721 \pm 0.0030$	$0.17224 \pm 0.00003$	0.0
$R_{s,d}/R_{(d+u+s)}$	OPAL	$0.371 \pm 0.022$	0.3592	0.5
$A_{\text{FB}}(b)$	LEP 1	$0.0992 \pm 0.0016$	$0.1033 \pm 0.0007$	-2.5
$A_{\text{FB}}(c)$	LEP 1	$0.0707 \pm 0.0035$	$0.0738 \pm 0.0006$	-0.9
$A_{\text{FB}}(s)$	DELPHI + OPAL	$0.098 \pm 0.011$	$0.1034 \pm 0.0001$	-0.5
$A_b$	SLD	$0.923 \pm 0.020$	$0.9347 \pm 0.0001$	-0.6
$A_c$	SLD	$0.670 \pm 0.027$	$0.6679 \pm 0.0004$	0.1
$A_s$	SLD	$0.895 \pm 0.091$	$0.9357 \pm 0.0001$	-0.4
$Q_{\text{FB}}$	LEP 1	$0.0403 \pm 0.0026$	$0.0423 \pm 0.0003$	-0.8
$A_{\text{LR}}$ (hadrons)	SLD	$0.1514 \pm 0.0022$	$0.1473 \pm 0.0010$	1.9
$A_{\text{LR}}$ (leptons)	SLD	$0.1544 \pm 0.0060$		1.2
$A_\mu$	SLD	$0.142 \pm 0.015$		-0.4
$A_\tau$	SLD	$0.136 \pm 0.015$		-0.8
$A_e(Q_{\text{LR}})$	SLD	$0.162 \pm 0.043$		0.3
$A_\tau(\mathcal{P}_\tau)$	LEP 1	$0.1439 \pm 0.0043$		-0.8
$A_e(\mathcal{P}_\tau)$	LEP 1	$0.1498 \pm 0.0049$		0.5
$\sin^2 \theta_W^{\text{eff}}(e)$	Tevatron	$0.2316 \pm 0.0018$	$0.2315 \pm 0.0001$	0.1

**Table 3.**  $Z$ -pole precision observables from LEP 1, the SLC, and the Tevatron. The SM errors are parametric as in table 2.

Finally, the constraints in the last two lines from first row CKM matrix unitarity [60, 71, 72] and from the anomalous magnetic moment of the muon [73] are affected by  $Z'$  bosons at the one-loop level. These loop diagrams are finite and give rise to rather small but not necessarily negligible effects. For example, a  $W - Z'$  box contribution could violate quark-lepton universality in the charged-current sector (and therefore apparently violate CKM unitarity) and is logarithmically enhanced for large  $M_{Z'}$  [74]. We included the analogous effect in the parameter  $\Delta r$  [75] describing the relation between the Fermi constant and  $M_W$ .

The size of the mixing angle  $\theta_{ZZ'}$  is strongly constrained by the very high precision  $Z$ -pole experiments [76] at LEP and SLC shown in table 3. The first set of measurements is from the  $Z$  line shape, from the (inverse) leptonic branching ratios normalized to the total hadronic  $Z$  decay width and from leptonic forward-backward asymmetries,  $A_{\text{FB}}(\ell)$ .

$Z'$	$M_{Z'} [\text{GeV}]$				$\sin \theta_{ZZ'}$			$\chi_{\min}^2$
	EW (this work)	CDF	DØ	LEP 2	$\sin \theta_{ZZ'}$	$\sin \theta_{ZZ'}^{\min}$	$\sin \theta_{ZZ'}^{\max}$	
$Z_\chi$	1,141	892	640	673	-0.0004	-0.0016	0.0006	47.3
$Z_\psi$	147	878	650	481	-0.0005	-0.0018	0.0009	46.5
$Z_\eta$	427	982	680	434	-0.0015	-0.0047	0.0021	47.7
$Z_I$	1,204	789	575		0.0003	-0.0005	0.0012	47.4
$Z_S$	1,257	821			-0.0003	-0.0013	0.0005	47.3
$Z_N$	623	861			-0.0004	-0.0015	0.0007	47.4
$Z_R$	442				-0.0003	-0.0015	0.0009	46.1
$Z_{LR}$	998	630		804	-0.0004	-0.0013	0.0006	47.3
$Z_{\underline{L}}$	(803)	(740)			-0.0015	-0.0094	0.0081	47.7
$Z_{SM}$	1,403	1,030	780	1,787	-0.0008	-0.0026	0.0006	47.2
$Z_{\text{string}}$	1,362				0.0002	-0.0005	0.0009	47.7
SM	$\infty$				0			48.5

**Table 4.** 95% C.L. lower mass limits on extra  $Z'$  bosons for various models from EW precision data and constraints on  $\sin \theta_{ZZ'}$  assuming  $\rho_0 = 1$  (fixed). For comparison, we show (where applicable) in the third, fourth and fifth column the limits obtained by CDF, DØ and LEP 2. In the following columns we give, respectively, the central value and the 95% C.L. lower and upper limits for  $\sin \theta_{ZZ'}$ . Also indicated is the  $\chi^2$  minimum for each model. The last row is included for comparison with the standard case of only one  $Z$  boson.

The second set represents similarly defined quantities of the quark sector. While all  $A_{FB}$  are practically sensitive only to the effective weak mixing angle defined for the initial state,  $\sin^2 \theta_W^{\text{eff}}(e)$ , the quantities  $A_q$  are functions of the effective weak mixing angle of the respective quark flavor,  $q$  ( $Q_{FB}$  is a similar observable for light quarks). The third set is a variety of cross section asymmetries sensitive to  $\sin^2 \theta_W^{\text{eff}}(e)$ ,  $\sin^2 \theta_W^{\text{eff}}(\mu)$ , or  $\sin^2 \theta_W^{\text{eff}}(\tau)$ . For details, see references [60, 76, 77]. The most recent result is the determination of  $\sin^2 \theta_W^{\text{eff}}(e)$  by the CDF [78] and DØ [79] Collaborations and is obtained from the forward-backward asymmetry for  $e^+e^-$  final states. Many of the entries in table 3 are of much higher precision than typical low-energy observables. The  $Z'$  amplitude, however, is almost entirely out of phase with and therefore negligible compared to the resonating  $Z$  amplitude. The  $Z'$  enters here mainly through a modification of the couplings of the ordinary  $Z$  to fermions, as well as through eq. (2.2), and indirectly by affecting the extracted value of the QCD coupling.

## 4 Results and discussion

In table 4 we present our limits on the  $Z'$  parameters for the models introduced in section 1. In this table we specify our results for the case  $\rho_0 = 1$  fixed but make no further assumptions regarding the Higgs sector except that the Higgs boson mass,  $M_H$ , is restricted to  $114.4 \text{ GeV} \leq M_H \leq 1 \text{ TeV}$ , where the upper end is from requiring perturbativity. The lower limit is the SM bound from LEP 2 [80] although we recall that this does not necessarily apply in the presence of new physics. Also shown in table 4 are the current limits on various

$Z'$	$M_{Z'} [\text{GeV}]$	$\sin \theta_{ZZ'}$	$\sin \theta_{ZZ'}^{\min}$	$\sin \theta_{ZZ'}^{\max}$	$\rho_0$	$\rho_0^{\min}$	$\rho_0^{\max}$	$\chi_{\min}^2$
$Z_\psi$	147	-0.0004	-0.0018	0.0010	1.0002	0.9996	1.0035	46.1
$Z_R$	439	-0.0003	-0.0015	0.0012	1.0003	0.9996	1.0035	45.3
SM	$\infty$	0			1.0003	0.9996	1.0035	47.9


**Table 5.** 95% C.L. limits on  $M_{Z'}$ ,  $\sin \theta_{ZZ'}$  and  $\rho_0$  when the latter is allowed to float freely.

$Z'$  boson masses from the Tevatron and LEP 2. The CDF limits [81] are from a search for a dimuon invariant mass peak. Notice that the  $Z_I$  and  $Z_{\cancel{L}}$  bosons face the weakest limits as is expected from their hadrophobic and leptophobic characters, respectively (no limit on the  $Z_{LR}$  is available from Run II at the Tevatron; the entry shown in the table is the CDF Run I result [82] from the combined dimuon and dielectron channels). Not shown are the dielectron channel search limits from CDF Run II [83] which are similar but slightly lower. There is a significant excess at a dielectron invariant mass of 240 GeV, but this is not confirmed in the  $\mu^+\mu^-$  channel. The results from DØ [84] are based on the dielectron final state. The mass limits at the Tevatron assume that no decay channels into exotic fermions or superpartners are open to the  $Z'$ ; otherwise the limits would be moderately weaker. LEP 2 constrains virtual  $Z'$  bosons by their effects on cross sections and angular distributions of dileptons, hadrons,  $b\bar{b}$  and  $c\bar{c}$  final states [85]. The table shows that the mass limits from the EW precision data are generally competitive with and in many cases stronger than those from colliders. We stress that these classes of limits are highly complementary. The result for the leptophobic  $Z_{\cancel{L}}$  (in parentheses) in the EW column is for the special Higgs sector with  $\tau = 1/2$ , i.e., for the lower end of the restricted range in table 1. For the upper end ( $\tau = 1$ ) we find a limit of 1.32 TeV. The CDF number [38] refers to the  $Z_{SM}$  limit from the dijet channel and should give a rough estimate of the sensitivity to our specific  $Z_{\cancel{L}}$ .

In the most general situation  $\rho_0$  is allowed to differ from 1 and is treated as a free fit parameter. We give the results of this case for the  $Z_\psi$  and the  $Z_R$  models in table 5. The comparison with table 4 shows that the presence of the extra fit parameter has little impact on the extracted  $Z'$  constraints.

Note, that all weak charges and the  $C_{1q}$  are proportional to some vector coupling,  $v$ , and hence blind to the  $Z_\psi$  which has only axial-vector couplings,  $a$ , to ordinary fermions. This is why the EW data give very weak constraints on its mass. The loop effects on the last two observables in table 2 gain therefore relative importance. In fact,  $Z'$  effects on  $g_\mu - 2$  are proportional to  $v_\mu^2 - 5a_\mu^2$  so that there is an additional enhancement (of an otherwise very small effect).


Figures 1, 2 and 3 show 90% C.L. exclusion contours for all models except for the  $Z_{\cancel{L}}$  (since its mass is in general unbounded). The solid (black) lines specify use of the constraint  $\rho_0 = 1$  while the dashed (blue) lines are for  $\rho_0$  free. We also show the extra constraints for the specific Higgs sectors described in section 2. These are represented by the dotted (red) lines unless they belong to the restricted range in table 1 in which case they are long-dashed (green). The numbers in the plots refer to the values of  $\tau$  or  $\omega$ , whichever carries the larger coefficient in table 1. The best fit locations (for  $\rho_0 = 1$ ) are indicated by



**Figure 1.** 95% C.L. contours in  $M_{Z'}$  vs.  $\sin \theta_{ZZ'}$  for various models. See the text for details.

an "x". The lower limits from CDF (dot-dashed and black),  $D\bar{O}$  (double-dot-dashed and magenta) and LEP 2 (dot-double-dashed and orange) given in table 4 are also shown.


In all figures and in tables 4 and 5 we used the  $M_H$  window mentioned above. However, the SM best fit value,  $M_H = 96^{+29}_{-24}$  GeV, is below this range. It is interesting to note [10, 86]



**Figure 2.** 95% C.L. contours in  $M_{Z'}$  vs.  $\sin \theta_{ZZ'}$  for various models. See the text for details.

that the presence of a  $Z'$  often moves the central value up to the allowed region. Table 6 shows the best fit values and  $1\sigma$  errors for  $M_H$  when the LEP 2 bound is removed.

Some  $Z'$  models have a fairly low minimum  $\chi^2$ , especially the  $Z_\psi$  and the  $Z_R$ . Table 6 shows the  $\chi^2$  minimum of the  $Z_R$  model about 3 units below the SM value, technically



**Figure 3.** 95% C.L. contours in  $M_{Z'}$  vs.  $\sin \theta_{ZZ'}$  for the sequential  $Z'$  boson and the  $Z_{\text{string}}$  model. See the text for details.

$Z'$	$Z_\chi$	$Z_\psi$	$Z_\eta$	$Z_I$	$Z_S$	$Z_N$
$M_H$ [GeV]	$171^{+493}_{-89}$	$97^{+31}_{-25}$	$423^{+577}_{-350}$	$141^{+304}_{-61}$	$149^{+353}_{-68}$	$117^{+222}_{-40}$
$\chi^2_{\min}$	47.3	46.1	47.7	47.4	47.3	47.4
$Z'$	$Z_R$	$Z_{LR}$	$Z_{\cancel{L}}$	$Z_{\text{SM}}$	$Z_{\text{string}}$	SM
$M_H$ [GeV]	$84^{+31}_{-24}$	$110^{+174}_{-35}$	$126^{+276}_{-52}$	$331^{+669}_{-246}$	$134^{+299}_{-58}$	$96^{+29}_{-25}$
$\chi^2_{\min}$	45.1	47.3	47.7	47.2	47.7	48.0

**Table 6.**  $1\sigma$  ranges of  $M_H$  allowed by each model and the best fit  $\chi^2$  values.

implying an *upper* bound on the  $Z_R$  mass of about 29 TeV at the 90% C.L. This is actually the reason why we included the  $Z_R$  in this paper in the first place. Of course, at present there is little significance to this observation since we have two additional fit parameters ( $M'_Z$  and  $\sin \theta_{ZZ'}$ ) and various parameters for the charges (like the angles  $\alpha$  and  $\beta$ ) to adjust. Nevertheless, this is somewhat surprising given that the SM fit is quite good with  $\chi^2_{\min} = 48.0$  for 45 effective degrees of freedom. It may be useful to note that the improvement in  $\chi^2$  arises mainly through  $\sigma_{\text{had}}$ ,  $Q_W(e)$ , and the  $e^-$ -DIS observables, where the latter two are of special interest in view of proposed and approved experiments to be performed at JLab.

## Acknowledgments

We thank the Institute for Nuclear Theory at the University of Washington for its hospitality and the Department of Energy for partial support during the early stages of this work. Work at IF-UNAM is supported by CONACyT project 82291-F. The work of P.L. is supported by the IBM Einstein Fellowship and by NSF grant PHY-0503584.

## References

- [1] S. Godfrey, *Update of discovery limits for extra neutral gauge bosons at hadron colliders*, in the proceedings of the *APS/DPF/DPB Summer Study on the Future of Particle Physics*, June 30–July 21, Snowmass, U.S.A. (2001), [hep-ph/0201093](#) [[SPIRES](#)].
- [2] F. Petriello and S. Quackenbush, *Measuring  $Z'$  couplings at the CERN LHC*, *Phys. Rev. D* **77** (2008) 115004 [[arXiv:0801.4389](#)] [[SPIRES](#)].
- [3] A. Leike, *The phenomenology of extra neutral gauge bosons*, *Phys. Rept.* **317** (1999) 143 [[hep-ph/9805494](#)] [[SPIRES](#)].
- [4] T.G. Rizzo,  *$Z'$  phenomenology and the LHC*, in the proceedings of *Exploring new frontiers using colliders and neutrinos (TASI2006)*, June 4–30, Boulder, U.S.A. (2006), [hep-ph/0610104](#) [[SPIRES](#)].
- [5] P. Langacker, *The physics of heavy  $Z'$  gauge bosons*, [arXiv:0801.1345](#) [[SPIRES](#)].
- [6] V. Barger, T. Han and D.G.E. Walker, *Top quark pairs at high invariant mass: a model-independent discriminator of new physics at the LHC*, *Phys. Rev. Lett.* **100** (2008) 031801 [[hep-ph/0612016](#)] [[SPIRES](#)].
- [7] S. Godfrey and T.A.W. Martin, *Identification of extra neutral gauge bosons at the LHC using  $b$ - and  $t$ -quarks*, *Phys. Rev. Lett.* **101** (2008) 151803 [[arXiv:0807.1080](#)] [[SPIRES](#)].
- [8] P. Osland, A.A. Pankov, A.V. Tsytrinov and N. Paver, *Spin and model identification of  $Z'$  bosons at the LHC*, *Phys. Rev. D* **79** (2009) 115021 [[arXiv:0904.4857](#)] [[SPIRES](#)].
- [9] J. Erler and P. Langacker, *Constraints on extended neutral gauge structures*, *Phys. Lett. B* **456** (1999) 68 [[hep-ph/9903476](#)] [[SPIRES](#)].
- [10] J. Erler and P. Langacker, *Indications for an extra neutral gauge boson in electroweak precision data*, *Phys. Rev. Lett.* **84** (2000) 212 [[hep-ph/9910315](#)] [[SPIRES](#)].
- [11] H. Georgi and S.L. Glashow, *Unity of all elementary particle forces*, *Phys. Rev. Lett.* **32** (1974) 438 [[SPIRES](#)].
- [12] J. Erler, *Chiral models of weak scale supersymmetry*, *Nucl. Phys. B* **586** (2000) 73 [[hep-ph/0006051](#)] [[SPIRES](#)].
- [13] J.E. Kim and H.P. Nilles, *The  $\mu$  problem and the strong CP problem*, *Phys. Lett. B* **138** (1984) 150 [[SPIRES](#)].
- [14] P. Langacker and M. Plümacher, *Flavor changing effects in theories with a heavy  $Z'$  boson with family nonuniversal couplings*, *Phys. Rev. D* **62** (2000) 013006 [[hep-ph/0001204](#)] [[SPIRES](#)].
- [15] R.W. Robinett and J.L. Rosner, *Mass scales in grand unified theories*, *Phys. Rev. D* **26** (1982) 2396 [[SPIRES](#)].
- [16] P. Langacker, R.W. Robinett and J.L. Rosner, *New heavy gauge bosons in  $pp$  and  $p\bar{p}$  collisions*, *Phys. Rev. D* **30** (1984) 1470 [[SPIRES](#)].



- [17] J.L. Hewett and T.G. Rizzo, *Low-energy phenomenology of superstring inspired  $E_6$  models*, *Phys. Rept.* **183** (1989) 193 [SPIRES].
- [18] P. Langacker, G. Paz and I. Yavin, *Scalar potentials and accidental symmetries in supersymmetric  $U(1)'$  models*, *Phys. Lett.* **B 671** (2009) 245 [arXiv:0811.1196] [SPIRES].
- [19] P. Candelas, G.T. Horowitz, A. Strominger and E. Witten, *Vacuum configurations for superstrings*, *Nucl. Phys.* **B 258** (1985) 46 [SPIRES].
- [20] D.J. Gross, J.A. Harvey, E.J. Martinec and R. Rohm, *The heterotic string*, *Phys. Rev. Lett.* **54** (1985) 502 [SPIRES].
- [21] E. Witten, *Symmetry breaking patterns in superstring models*, *Nucl. Phys.* **B 258** (1985) 75 [SPIRES].
- [22] Y. Hosotani, *Dynamical gauge symmetry breaking as the Casimir effect*, *Phys. Lett.* **B 129** (1983) 193 [SPIRES].
- [23] J. Erler, P. Langacker and T.-j. Li, *The  $Z$ - $Z'$  mass hierarchy in a supersymmetric model with a secluded  $U(1)'$  breaking sector*, *Phys. Rev.* **D 66** (2002) 015002 [hep-ph/0205001] [SPIRES].
- [24] J. Kang, P. Langacker, T.-j. Li and T. Liu, *Electroweak baryogenesis in a supersymmetric  $U(1)'$  model*, *Phys. Rev. Lett.* **94** (2005) 061801 [hep-ph/0402086] [SPIRES].
- [25] E. Ma, *Neutrino masses in an extended gauge model with  $E_6$  particle content*, *Phys. Lett.* **B 380** (1996) 286 [hep-ph/9507348] [SPIRES].
- [26] V. Barger, P. Langacker and H.-S. Lee, *Primordial nucleosynthesis constraints on  $Z'$  properties*, *Phys. Rev.* **D 67** (2003) 075009 [hep-ph/0302066] [SPIRES].
- [27] J.-h. Kang, P. Langacker and T.-j. Li, *Neutrino masses in supersymmetric  $SU(3)(C) \times SU(2)(L) \times U(1)(Y) \times U(1)'$  models*, *Phys. Rev.* **D 71** (2005) 015012 [hep-ph/0411404] [SPIRES].
- [28] M. Gell-Mann, P. Ramond and R. Slansky, *Complex spinors and unified theories*, in the proceedings of the *Supergravity workshop*, September 27–29, Stony Brook, U.S.A. (1979) [SPIRES].
- [29] S.F. King, S. Moretti and R. Nevzorov, *Theory and phenomenology of an exceptional supersymmetric standard model*, *Phys. Rev.* **D 73** (2006) 035009 [hep-ph/0510419] [SPIRES]; *Exceptional supersymmetric standard model*, *Phys. Lett.* **B 634** (2006) 278 [hep-ph/0511256] [SPIRES].
- [30] S. F. King, S. Moretti and R. Nevzorov,  *$E_6$ Ssm*, *AIP Conf. Proc.* **881** (2007) 138 [hep-ph/0610002] [SPIRES].
- [31] B. Holdom, *Two  $U(1)$ 's and epsilon charge shifts*, *Phys. Lett.* **B 166** (1986) 196 [SPIRES].
- [32] T. Matsuoka and D. Suematsu, *Low-energy gauge interactions from the  $E_8 \times E_8'$  superstring theory*, *Prog. Theor. Phys.* **76** (1986) 901 [SPIRES].
- [33] F. del Aguila, G.D. Coughlan and M. Quirós, *Gauge coupling renormalization with several  $U(1)$  factors*, *Nucl. Phys.* **B 307** (1988) 633 [SPIRES].
- [34] R. Foot and X.-G. He, *Comment on  $ZZ'$  mixing in extended gauge theories*, *Phys. Lett.* **B 267** (1991) 509 [SPIRES].
- [35] P. Langacker and J. Wang,  *$U(1)'$  symmetry breaking in supersymmetric  $E_6$  models*, *Phys. Rev.* **D 58** (1998) 115010 [hep-ph/9804428] [SPIRES].
- [36] K.S. Babu, C.F. Kolda and J. March-Russell, *Leptophobic  $U(1)$ s and the  $R_b$ - $R_c$  crisis*, *Phys. Rev.* **D 54** (1996) 4635 [hep-ph/9603212] [SPIRES]; *Implications of generalized  $ZZ'$  mixing*, *Phys. Rev.* **D 57** (1998) 6788 [hep-ph/9710441] [SPIRES].

- [37] R.N. Mohapatra, *Unification and supersymmetry*, Springer-Verlag, Berlin Germany (1986).
- [38] CDF collaboration, T. Aaltonen et al., *Search for new particles decaying into dijets in proton- antiproton collisions at  $\sqrt{s} = 1.96$  TeV*, *Phys. Rev. D* **79** (2009) 112002 [[arXiv:0812.4036](#)] [[SPIRES](#)].
- [39] CDF collaboration, T. Aaltonen et al., *Search for resonant  $t\bar{t}$  production in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV*, *Phys. Rev. Lett.* **100** (2008) 231801 [[arXiv:0709.0705](#)] [[SPIRES](#)].
- [40] S. Chaudhuri, S.W. Chung, G. Hockney and J.D. Lykken, *String consistency for unified model building*, *Nucl. Phys. B* **456** (1995) 89 [[hep-ph/9501361](#)] [[SPIRES](#)].
- [41] G. Cleaver, M. Cvetič, J.R. Espinosa, L.L. Everett and P. Langacker, *Classification of flat directions in perturbative heterotic superstring vacua with anomalous U(1)*, *Nucl. Phys. B* **525** (1998) 3 [[hep-th/9711178](#)] [[SPIRES](#)].
- [42] G. Cleaver et al., *Physics implications of flat directions in free fermionic superstring models. I: mass spectrum and couplings*, *Phys. Rev. D* **59** (1999) 055005 [[hep-ph/9807479](#)] [[SPIRES](#)]; *Physics implications of flat directions in free fermionic superstring models. II: renormalization group analysis*, *Phys. Rev. D* **59** (1999) 115003 [[hep-ph/9811355](#)] [[SPIRES](#)].
- [43] M. Cvetič and P. Langacker, *Implications of abelian extended gauge structures from string models*, *Phys. Rev. D* **54** (1996) 3570 [[hep-ph/9511378](#)] [[SPIRES](#)]; *New gauge bosons from string models*, *Mod. Phys. Lett. A* **11** (1996) 1247 [[hep-ph/9602424](#)] [[SPIRES](#)].
- [44] M. Cvetič, D.A. Demir, J.R. Espinosa, L.L. Everett and P. Langacker, *Electroweak breaking and the  $\mu$  problem in supergravity models with an additional U(1)*, *Phys. Rev. D* **56** (1997) 2861 [[hep-ph/9703317](#)] [[SPIRES](#)].
- [45] T. Appelquist, B.A. Dobrescu and A.R. Hopper, *Nonexotic neutral gauge bosons*, *Phys. Rev. D* **68** (2003) 035012 [[hep-ph/0212073](#)] [[SPIRES](#)].
- [46] P. Langacker, *Bounds on mixing between light and heavy gauge bosons*, *Phys. Rev. D* **30** (1984) 2008 [[SPIRES](#)].
- [47] M.E. Peskin and T. Takeuchi, *A new constraint on a strongly interacting Higgs sector*, *Phys. Rev. Lett.* **65** (1990) 964 [[SPIRES](#)].
- [48] P. Langacker and M.-x. Luo, *Constraints on additional Z bosons*, *Phys. Rev. D* **45** (1992) 278 [[SPIRES](#)].
- [49] J. Erler, *Global fits to electroweak data using GAPP*, in the proceedings of *Physics at RUN II: QCD and Weak Boson Physics Workshop: 2<sup>nd</sup> General Meeting*, June 3–4, Batavia, U.S.A. (1999), [hep-ph/0005084](#) [[SPIRES](#)].
- [50] ALEPH collaboration, CDF collaboration, DØ collaboration, DELPHI collaboration, L3 collaboration, OPAL collaboration and SLD collaboration, LEP, TEVATRON AND SLD EW working groups and SLD HEAVY FLAVOUR GROUP, *Precision electroweak measurements and constraints on the standard model*, [arXiv:0811.4682](#) [[SPIRES](#)].
- [51] CDF collaboration and DØ collaborations, TEVATRON ELECTROWEAK WORKING GROUP, *Combination of CDF and D0 results on the mass of the top quark*, [arXiv:0903.2503](#) [[SPIRES](#)].
- [52] CCFR collaboration, K.S. McFarland et al., *A precision measurement of electroweak parameters in neutrino nucleon scattering*, *Eur. Phys. J. C* **1** (1998) 509 [[hep-ex/9701010](#)] [[SPIRES](#)].
- [53] NUTeV collaboration, G.P. Zeller et al., *A precise determination of electroweak parameters in neutrino nucleon scattering*, *Phys. Rev. Lett.* **88** (2002) 091802 [[hep-ex/0110059](#)] [[SPIRES](#)].

- [54] CHDS collaboration, A. Blondel et al., *Electroweak parameters from a high statistics neutrino nucleon scattering experiment*, *Z. Phys. C* **45** (1990) 361 [SPIRES].
- [55] CHARM collaboration, J.V. Allaby et al., *A precise determination of the electroweak mixing angle from semileptonic neutrino scattering*, *Z. Phys. C* **36** (1987) 611 [SPIRES].
- [56] D. Mason et al., *Measurement of the nucleon strange-antistrange asymmetry at next-to-leading order in QCD from NuTeV dimuon data*, *Phys. Rev. Lett.* **99** (2007) 192001 [SPIRES].
- [57] K.P.O. Diener, S. Dittmaier and W. Hollik, *Electroweak radiative corrections to deep-inelastic neutrino scattering: implications for NuTeV?*, *Phys. Rev. D* **69** (2004) 073005 [hep-ph/0310364] [SPIRES]; *Electroweak higher-order effects and theoretical uncertainties in deep-inelastic neutrino scattering*, *Phys. Rev. D* **72** (2005) 093002 [hep-ph/0509084] [SPIRES].
- [58] A.B. Arbuzov, D.Y. Bardin and L.V. Kalinovskaya, *Radiative corrections to neutrino deep inelastic scattering revisited*, *JHEP* **06** (2005) 078 [hep-ph/0407203] [SPIRES].
- [59] A.D. Martin, R.G. Roberts, W.J. Stirling and R.S. Thorne, *Uncertainties of predictions from parton distributions. I: theoretical errors*, *Eur. Phys. J. C* **35** (2004) 325 [hep-ph/0308087] [SPIRES].
- [60] PARTICLE DATA GROUP collaboration, C. Amsler et al., *Review of particle physics*, *Phys. Lett. B* **667** (2008) 1 [SPIRES].
- [61] G.P. Zeller, private communication.
- [62] CHARM-II collaboration, P. Vilain et al., *Precision measurement of electroweak parameters from the scattering of muon-neutrinos on electrons*, *Phys. Lett. B* **335** (1994) 246 [SPIRES].
- [63] N.H. Edwards et al., *Precise measurement of parity nonconserving optical rotation in atomic Thallium*, *Phys. Rev. Lett.* **74** (1995) 2654.
- [64] P.A. Vetter, D.M. Meekhof, P.K. Majumder, S.K. Lamoreaux and E.N. Fortson, *Precise test of electroweak theory from a new measurement of parity nonconservation in atomic thallium*, *Phys. Rev. Lett.* **74** (1995) 2658 [SPIRES].
- [65] C.S. Wood et al., *Measurement of parity nonconservation and an anapole moment in Cesium*, *Science* **275** (1997) 1759 [SPIRES].
- [66] SLAC E158 collaboration, P.L. Anthony et al., *Precision measurement of the weak mixing angle in Moeller scattering*, *Phys. Rev. Lett.* **95** (2005) 081601 [hep-ex/0504049] [SPIRES].
- [67] S.G. Porsev, K. Beloy and A. Derevianko, *Precision determination of electroweak coupling from atomic parity violation and implications for particle physics*, *Phys. Rev. Lett.* **102** (2009) 181601 [arXiv:0902.0335] [SPIRES].
- [68] J.S.M. Ginges and V.V. Flambaum, *Violations of fundamental symmetries in atoms and tests of unification theories of elementary particles*, *Phys. Rept.* **397** (2004) 63 [physics/0309054].
- [69] R.D. Young, private communication.
- [70] R.D. Young, R.D. Carlini, A.W. Thomas and J. Roche, *Testing the standard model by precision measurement of the weak charges of quarks*, *Phys. Rev. Lett.* **99** (2007) 122003 [arXiv:0704.2618] [SPIRES].
- [71] J.C. Hardy and I.S. Towner, *Superaligned  $0^+$  to  $0^+$  nuclear beta decays: a new survey with precision tests of the conserved vector current hypothesis and the standard model*, *Phys. Rev. C* **79** (2009) 055502 [arXiv:0812.1202] [SPIRES].
- [72] KLOE collaboration, F. Bossi et al., *Precision kaon and hadron physics with KLOE*, *Riv. Nuovo Cim.* **031** (2008) 531 [arXiv:0811.1929] [SPIRES].

- [73] MUON G-2 collaboration, G.W. Bennett et al., *Measurement of the negative muon anomalous magnetic moment to 0.7 ppm*, *Phys. Rev. Lett.* **92** (2004) 161802 [[hep-ex/0401008](#)] [[SPIRES](#)].
- [74] W.J. Marciano and A. Sirlin, *Constraint on additional neutral gauge bosons from electroweak radiative corrections*, *Phys. Rev. D* **35** (1987) 1672 [[SPIRES](#)].
- [75] A. Sirlin, *Radiative corrections in the  $SU(2)_L \times U(1)$  theory: a simple renormalization framework*, *Phys. Rev. D* **22** (1980) 971 [[SPIRES](#)].
- [76] ALEPH collaboration, DELPHI collaboration, L3 collaboration, OPAL collaboration and SLD collaboration, LEP EW WORKING GROUP and SLD EW AND HEAVY FLAVOUR GROUPS, *Precision electroweak measurements on the Z resonance*, *Phys. Rept.* **427** (2006) 257 [[hep-ex/0509008](#)] [[SPIRES](#)].
- [77] J. Erler and P. Langacker, *Status of the standard model*, in the proceedings of the 5<sup>th</sup> *International Wein Symposium: A Conference on Physics Beyond the Standard Model (WEIN98)*, June 14–21, Santa Fe, U.S.A. (1998), [hep-ph/9809352](#) [[SPIRES](#)].
- [78] CDF collaboration, D.E. Acosta et al., *Measurement of the forward-backward charge asymmetry of electron positron pairs in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV*, *Phys. Rev. D* **71** (2005) 052002 [[hep-ex/0411059](#)] [[SPIRES](#)].
- [79] D0 collaboration, V.M. Abazov et al., *Measurement of the forward-backward charge asymmetry and extraction of  $\sin^2 \theta_W^{\text{eff}}$  in  $p\bar{p} \rightarrow Z/\gamma^* + X \rightarrow e^+e^- + X$  events produced at  $\sqrt{s} = 1.96$  TeV*, *Phys. Rev. Lett.* **101** (2008) 191801 [[arXiv:0804.3220](#)] [[SPIRES](#)].
- [80] ALEPH collaboration, DELPHI COLLABORATION, L3 collaboration, OPAL collaboration, LEP WORKING GROUP FOR HIGGS BOSON SEARCHES, R. Barate et al., *Search for the standard model Higgs boson at LEP*, *Phys. Lett. B* **565** (2003) 61 [[hep-ex/0306033](#)] [[SPIRES](#)].
- [81] CDF collaboration, T. Aaltonen et al., *A search for high-mass resonances decaying to dimuons at CDF*, *Phys. Rev. Lett.* **102** (2009) 091805 [[arXiv:0811.0053](#)] [[SPIRES](#)].
- [82] CDF collaboration, F. Abe et al., *Search for new gauge bosons decaying into dileptons in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8$  TeV*, *Phys. Rev. Lett.* **79** (1997) 2192 [[SPIRES](#)].
- [83] CDF collaboration, T. Aaltonen et al., *Search for High-Mass  $e^+ e^-$  Resonances in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV*, *Phys. Rev. Lett.* **102** (2009) 031801 [[arXiv:0810.2059](#)] [[SPIRES](#)].
- [84] D0 collaboration, *Search for heavy Z bosons in the dielectron channel with 200  $pb^{-1}$  of data with the D0 detector*, DO Note 4375-conf (2004), <http://www-d0.fnal.gov/Run2Physics/WWW/results/prelim/NP/N03/N03.pdf>
- [85] ALEPH collaboration, DELPHI collaboration, L3 collaboration and OPAL collaboration, LEP ELECTROWEAK WORKING GROUP, J. Alcaraz et al., *A combination of preliminary electroweak measurements and constraints on the standard model*, *Phys. Rept.* **427** (2006) 257 [[hep-ex/0612034](#)] [[SPIRES](#)].
- [86] M.S. Chanowitz, *A  $Z'$  boson and the higgs boson mass*, [arXiv:0806.0890](#) [[SPIRES](#)].