

Topology and higher dimensional representations

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

JHEP08(2009)084

(<http://iopscience.iop.org/1126-6708/2009/08/084>)

[The Table of Contents](#) and [more related content](#) is available

Download details:

IP Address: 80.92.225.132

The article was downloaded on 03/04/2010 at 09:25

Please note that [terms and conditions apply](#).

Topology and higher dimensional representations

Zoltán Fodor,^{a,b,c,d} Kieran Holland,^e Julius Kuti,^f Dániel Nógrádi^f and Chris Schroeder^f

^aDepartment of Theoretical Physics, University of Wuppertal,
Gaussstrasse 20, Wuppertal 42119, Germany

^bNIC/DESY Zeuthen Forschungsgruppe,
Platanenallee 6, Zeuthen 15738, Germany

^cForschungszentrum Jülich,
D-52425 Jülich, Germany

^dInstitute for Theoretical Physics, Eötvös University,
Pázmány Péter sétány 1/A, Budapest 1117, Hungary

^eDepartment of Physics, University of the Pacific,
3601 Pacific Avenue, Stockton, CA 95211, U.S.A.

^fDepartment of Physics, University of California San Diego,
9500 Gilman Dr, La Jolla, CA 92093, U.S.A.

E-mail: fodor@bodri.elte.hu, kholland@pacific.edu, jkuti@ucsd.edu,
nogradi@bodri.elte.hu, crs@physics.ucsd.edu

ABSTRACT: SU(3) gauge theory in the 2-index symmetric (sextet) and fundamental representations is considered in symmetric and periodic boxes. Using the overlap formulation in the quenched approximation it is shown that the topological charge obtained from the sextet index theorem always leads to an integer value and agrees with the charge obtained from the fundamental index theorem in the continuum. At larger lattice spacing configurations exist with fractional topological charge if the sextet index is used but these are lattice artifacts and the probability of finding such a configuration rapidly approaches zero. By considering the decomposition of the sextet representation with respect to an SU(2) subgroup it is shown that the SU(2) adjoint index theorem leads to integer charge as well. We conclude that the non-zero value of the bilinear gaugino condensate in $\mathcal{N} = 1$ super-Yang-Mills theory cannot be attributed to configurations with fractional topological charge once periodic boundary conditions are imposed.

KEYWORDS: Lattice QCD, Lattice Gauge Field Theories

ARXIV EPRINT: [0905.3586](https://arxiv.org/abs/0905.3586)

Contents

1	Introduction	1
2	Topology and index theorem	2
2.1	Sextet representation of SU(3)	3
2.2	Adjoint representation of SU(2)	7
3	Topological susceptibility	10
4	Discussion	10

1 Introduction

The chirally invariant overlap formulation is known to have advantageous topological and continuum properties in the fundamental representation [1–5], but much less is known in higher dimensional representations. A naive use of the Atiyah-Singer index theorem at finite lattice spacing may lead to fractional topological charge $Q = I/2T$ where T is the trace normalization factor of the representation and I is the index of the Dirac operator, if I is not an integer multiple of $2T$.

Here we study the 2-index symmetric (sextet) representation of SU(3) in the quenched approximation. In this case $2T = 5$. This model has been considered previously in [6–11].

Our motivations for this work are twofold. First, the theory of $N_f = 2$ fermions in the sextet of SU(3) is a promising candidate for a walking technicolor model [12]. The $N_f = 0$ theory certainly breaks chiral symmetry and is a good testing ground for chiral and/or topological properties of sextet overlap fermions before addressing dynamical questions such as the infrared properties of the $N_f = 2$ theory. Results on non-abelian gauge theories with fermions in higher dimensional representations have been reported in [13–20].

Second, in SU(N) $\mathcal{N} = 1$ super-Yang-Mills theory the gaugino condensate involving $2N$ gaugino fields is non-zero due to the contribution of instantons to this correlation function since there are $2N$ adjoint zero modes for a charge 1 instanton. The general expectation is that the bilinear gaugino condensate is non-zero as well [21]. However it is not clear if this non-zero value can be attributed to some special class of configurations if periodic boundary conditions are used. Since the fermionic definition of the topological charge can lead to fractional values at finite lattice spacing as mentioned above, it is worth investigating what happens to these configurations in the continuum limit. If the fractional value $1/N$ does survive the continuum limit with periodic boundary conditions they potentially can explain the non-zero value of the bilinear gaugino condensate on R^4 . This would then be similar to the case of $S^1 \times R^3$ where fractional topological charge configurations do exist and it has been argued that these can explain the gaugino condensate [22].

The main result of our paper is that once periodic boundary conditions are imposed, configurations where $Q = I/2T$ is not an integer disappear in the continuum limit. In the continuum all $SU(3)$ configurations respect the sextet index theorem, i.e. zero modes come in 5-tuples, furthermore the topological charge defined by the sextet index agrees with the one from the fundamental index and is an integer.

Although we have only measured the sextet and fundamental index for $SU(3)$ with periodic boundary conditions the following property of continuum gauge fields provides an opportunity to address the supersymmetric case of $SU(2)$ as well. An $SU(N)$ gauge field can always be deformed to lie in an $SU(2)$ subgroup and the representation then can be decomposed with respect to this subgroup. In our case the decomposition of the sextet of $SU(3)$ contains both the fundamental and adjoint of $SU(2)$. This observation leads to the conclusion that in $SU(2)$ gauge theory the fundamental and adjoint index theorems both give the same definition of the topological charge in the continuum. In particular, the charge is an integer.

Consequently, the bilinear gaugino condensate cannot be attributed to fractionally charged configurations subject to periodic boundary conditions.

Although we have not investigated other higher dimensional representations, we conjecture that our findings hold more generally. Specifically, in a large periodic box, (1) the fermionic definition of the topological charge $Q = I/2T$ will always be an integer and (2) two definitions from two different representations will agree in the continuum limit.

The adjoint representation of $SU(2)$ was investigated directly in [13] and a slight increase has been reported in the probability of finding a configuration with adjoint index 2, suggesting $Q = 1/2$, as the lattice spacing decreased at fixed physical volume. This raised the possibility that the probability stays finite in the continuum limit. We will show however that if the lattice spacing is too large the decrease in this probability with decreasing lattice spacing is quite small and can easily be confused with a slight increase at low statistics. A more consistent decrease of the probability with decreasing lattice spacing does however set in for sufficiently small lattice spacing.

2 Topology and index theorem

The Atiyah-Singer index theorem in the continuum states that $I_R = 2T_R Q$ where I_R is the index of the Dirac operator in representation R , T_R is its trace normalization factor and Q is the topological charge of the underlying gauge field. At finite lattice spacing a gluonic definition of Q is not available in general but the above relation can be employed to define it using the overlap operator which possesses exact chiral zero modes [23]. Clearly, two different representations may lead to different charges for the same configuration. Also, if $2T_R > 1$ the fermionic definition may lead to non-integer charge. In the fundamental representation this problem does not arise because $2T = 1$ and the fermionic definition always gives an integer charge (different choices for the negative Wilson mass may still lead to different integers but in the continuum limit the charge will not be sensitive to this choice).

β	L/a	r_0/a	a [fm]	#config
5.9500	12	4.9122	0.1018	441
6.0384	14	5.7306	0.0873	543
6.1210	16	6.5496	0.0763	408
6.2719	20	8.1867	0.0611	211
6.4064	24	9.8239	0.0509	105

Table 1. Simulation parameters using the Wilson plaquette gauge action. The physical volume is kept fixed at $L = 1.2215$ fm.

For gauge group $SU(N)$ the T factors for some of the most frequently used representations are given by,

$$T_F = \frac{1}{2}, \quad T_{adj} = N, \quad T_S = \frac{N+2}{2}, \quad T_A = \frac{N-2}{2}, \quad (2.1)$$

where F , adj , S and A refer to the fundamental, adjoint, 2-index symmetric and 2-index anti-symmetric representations, respectively. The index for $SU(N)$ will be denoted by I_R^N . Unless otherwise indicated Q and I will stand for the absolute value of these quantities.

2.1 Sextet representation of $SU(3)$

For the sextet of $SU(3)$ we have $2T = 5$ thus the continuum index theorem suggests that zero modes of the Dirac operator ought to come in 5-tuples.

We have measured the index of the Dirac operator in both the fundamental and sextet representation in the quenched approximation at 5 different lattice spacings at the same physical volume. For a similar study in the fundamental representation only, see [24]. The index was obtained by calculating the first few eigenvalues of the Dirac operator until a non-zero eigenvalue is found and checking whether they are chiral or not. Zero modes are clearly identified by their chirality being ± 1 whereas non-zero modes have 0 chirality just as in the continuum. The non-zero eigenvalues carry important chiral information encoded in the Banks-Casher relation and random matrix theory. These findings will be reported elsewhere [25].

The Wilson plaquette action was used on symmetrical L^4 lattices with periodic boundary conditions and the Sommer scale $r_0 = 0.5$ fm (in the fundamental representation) was used to set the scale [26]. The negative Wilson mass in the fundamental representation was chosen at -1.4 while for the sextet it was set to -1.7 . In both cases 2 steps of stout smearing was applied with smearing parameter 0.15 [27]. The implementation of the overlap operator is the same as in [28]. The simulation parameters are summarized in table 1. The fundamental and sextet indices are tabulated in tables 2–4 together with the number of configurations that have been observed with the given values.

First, let us analyze to what extent sextet zero modes come in 5-tuples. The distribution of the sextet index is shown on figure 1 without taking the absolute value. Clearly, as the lattice spacing decreases the number of configurations that would lead to fractional charge is suppressed. The probability that one finds such a configuration is shown on figure 2.

I_F^3	I_S^3	$5I_F^3 - I_S^3$	$I_S^3 - I_F^3$	#configs	I_F^3	I_S^3	$5I_F^3 - I_S^3$	$I_S^3 - I_F^3$	#configs
1	5	0	4	52	1	0	5	-1	4
0	2	-2	2	35	0	6	-6	6	4
1	3	2	2	32	3	15	0	12	3
0	0	0	0	30	2	5	5	3	3
1	7	-2	6	29	4	18	2	14	2
2	10	0	8	26	4	17	3	13	2
2	8	2	6	21	3	19	-4	16	2
1	4	1	3	18	3	17	-2	14	2
0	1	-1	1	18	3	12	3	9	2
1	6	-1	5	17	3	11	4	8	2
2	12	-2	10	12	4	20	0	16	1
2	11	-1	9	12	4	19	1	15	1
1	1	4	0	12	4	16	4	12	1
1	2	3	1	11	4	15	5	11	1
0	3	-3	3	10	3	20	-5	17	1
2	9	1	7	8	3	18	-3	15	1
2	7	3	5	8	3	14	1	11	1
2	6	4	4	8	3	10	5	7	1
1	9	-4	8	8	3	7	8	4	1
0	4	-4	4	8	2	13	-3	11	1
3	13	2	10	6	1	12	-7	11	1
0	5	-5	5	6	1	10	-5	9	1
2	14	-4	12	5	0	9	-9	9	1
1	8	-3	7	5	0	7	-7	7	1
3	16	-1	13	4					

Table 2. The sextet and fundamental indices for $L/a = 12$.

Fitting this probability to a function of a requires some care. Unfortunately we do not have a solid theoretical expectation for the a -dependence as a whole and neither for the limit $a \rightarrow 0$. In principle the $a \rightarrow 0$ limit could be worked out analytically and we hope to return to this question in the future. Currently, however, the only guide is the data itself. The chiral symmetry of overlap fermions ensure $O(a)$ improvement in on-shell correlation functions of local operators [2], but the probability plotted on figure 2 is not such a quantity. Hence as $a \rightarrow 0$ we do not expect $O(a)$ terms to be absent.

Although the linear fit to the data does not go through zero at $a = 0$, the actual linear fit does seem to work very well. We cannot completely rule out the scenario that the probability of finding a configuration with sextet indices that are not multiples of 5 will only vanish at zero lattice spacing, deviating from the fitted straight line for smaller a values than the simulation points. However, our current data set is consistent with the linear fit, reaching zero at finite lattice spacing.

I_F^3	I_S^3	$5I_F^3 - I_S^3$	$I_S^3 - I_F^3$	#configs	I_F^3	I_S^3	$5I_F^3 - I_S^3$	$I_S^3 - I_F^3$	#configs
1	5	0	4	95	1	5	0	4	94
0	0	0	0	66	0	0	0	0	58
0	2	-2	2	49	2	10	0	8	54
2	10	0	8	48	1	3	2	2	30
1	3	2	2	37	1	7	-2	6	19
2	8	2	6	27	0	2	-2	2	19
1	7	-2	6	24	1	4	1	3	15
0	1	-1	1	21	3	13	2	10	14
1	4	1	3	18	2	8	2	6	14
3	15	0	12	17	0	1	-1	1	14
1	1	4	0	12	3	15	0	12	12
3	13	2	10	11	2	9	1	7	8
1	6	-1	5	11	1	6	-1	5	6
0	3	-3	3	11	0	4	-4	4	6
2	12	-2	10	9	3	17	-2	14	4
2	9	1	7	9	2	12	-2	10	4
2	6	4	4	8	2	11	-1	9	4
0	4	-4	4	8	0	3	-3	3	4
1	8	-3	7	7	4	20	0	16	3
3	14	1	11	6	2	6	4	4	3
2	7	3	5	6	1	9	-4	8	3
3	17	-2	14	5	1	1	4	0	3
2	11	-1	9	5	4	16	4	12	2
1	2	3	1	4	3	14	1	11	2
4	20	0	16	3	2	5	5	3	2
3	16	-1	13	3	2	4	6	2	2
2	13	-3	11	3	4	22	-2	18	1
4	22	-2	18	2	4	18	2	14	1
4	16	4	12	2	3	18	-3	15	1
3	12	3	9	2	3	12	3	9	1
3	11	4	8	2	2	7	3	5	1
2	14	-4	12	2	1	10	-5	9	1
2	0	10	-2	2	1	8	-3	7	1
5	20	5	15	1	1	2	3	1	1
4	18	2	14	1	1	0	5	-1	1
3	18	-3	15	1					
2	5	5	3	1					
2	4	6	2	1					
1	9	-4	8	1					
1	0	5	-1	1					
0	5	-5	5	1					

Table 3. The sextet and fundamental indices for $L/a = 14$ (left) and $L/a = 16$ (right).

I_F^3	I_S^3	$5I_F^3 - I_S^3$	$I_S^3 - I_F^3$	#configs	I_F^3	I_S^3	$5I_F^3 - I_S^3$	$I_S^3 - I_F^3$	#configs
1	5	0	4	64	1	5	0	4	41
0	0	0	0	51	0	0	0	0	28
2	10	0	8	32	2	10	0	8	14
3	15	0	12	14	3	15	0	12	3
1	7	-2	6	10	2	12	-2	10	3
0	2	-2	2	6	1	3	2	2	3
2	8	2	6	5	4	20	0	16	2
1	4	1	3	5	1	2	3	1	2
1	3	2	2	5	1	7	-2	6	2
0	1	-1	1	5	0	1	-1	1	1
2	12	-2	10	4	0	2	-2	2	1
3	13	2	10	3	0	3	-3	3	1
4	18	2	14	1	1	4	1	3	1
3	17	-2	14	1	1	6	-1	5	1
3	14	1	11	1	2	8	2	6	1
2	9	1	7	1	2	9	1	7	1
1	6	-1	5	1					
1	2	3	1	1					
1	1	4	0	1					

Table 4. The sextet and fundamental indices for $L/a = 20$ (left) and $L/a = 24$ (right).

Keeping in mind these cautionary remarks we proceed with the linear fit. It implies that at a certain small but finite lattice spacing all configurations that are not consistent with the sextet index theorem and integer charge are gone. The extrapolated value is $a_c/r_0 = 0.075(8)$ corresponding to $a_c = 0.038(4) fm$. At our physical volume this corresponds to approximately $\beta = 6.64$ and $L/a = 32$. The precise value of a_c presumably depends on the fixed finite physical volume, as well as on the details of the fermion matrix such as the number of smearing steps and the smearing parameter. Naturally, it would be most interesting to calculate the sextet and fundamental indices on lattices finer than a_c to verify the validity of the extrapolation used, but currently we do not have sufficient resources.

Second, let us investigate how many configurations violate the continuum expectation $5I_F^3 = I_S^3$. Their number is actually almost the same as the number of configurations with I_S^3 not a multiple of 5, see tables 2–4. Correspondingly, the extrapolated lattice spacing value a_c where all configurations with the property $5I_F^3 \neq I_S^3$ disappear is the same, within errors, as the one obtained above. Our comments about the $a \rightarrow 0$ behaviour above naturally apply to this case as well.

Summarizing this section we conclude that if $a < a_c$, or perhaps only in the continuum limit, the only remaining configurations are the ones where I_S^3 is an integer multiple of 5 and in addition the topological charge defined from the fundamental representation coincides with the definition from the sextet representation. In particular it is an integer.




Figure 1. Normalized distribution of the SU(3) sextet index at fixed physical volume and various lattice spacings.

2.2 Adjoint representation of SU(2)

Although we have only studied the fundamental and sextet representations of SU(3) our results have implications for the adjoint representation of SU(N) as well. This is because in the continuum an SU(N) gauge field can always be deformed to lie in an SU(2) subgroup only. The fermion representation can then be decomposed with respect to this SU(2) subgroup. For the sextet of SU(3) this decomposition is

$$\mathbf{6}_{\text{SU}(3)} = \mathbf{3}_{\text{SU}(2)} \oplus \mathbf{2}_{\text{SU}(2)} \oplus \mathbf{1}_{\text{SU}(2)} , \quad (2.2)$$

containing the adjoint and fundamental of SU(2).

From the relations (2.1) we find that an SU(2) charge 1 gauge field will carry 4 adjoint and 1 fundamental zero modes, $I_{\text{adj}}^2 = 4Q$ and $I_F^2 = Q$. This also confirms that for SU(3)




Figure 2. Probability of finding a configuration where the number of sextet zero modes is not a multiple of 5.

there are $4 + 1 = 5$ sextet zero modes. These 5 sextet zero modes of a charge 1 SU(3) gauge field can be grouped into a group of 4 which are the 4 adjoint SU(2) zero modes and a group containing just a single fundamental SU(2) zero mode.

This is the reason why our sextet SU(3) zero modes know about adjoint SU(2) zero modes.

Since we have seen in the previous section that the SU(3) sextet index is compatible with integer charge if the lattice spacing is small enough, it strongly suggests that the SU(2) adjoint index is as well. To make this claim more quantitative, we have $I_S^3 - I_F^3 = I_{\text{adj}}^2$ for an SU(2) gauge field embedded in SU(3) using the decomposition (2.2), since in this case $I_F^2 = I_F^3$. If the SU(2) adjoint index theorem is not compatible with integer charge, $I_S^3 - I_F^3$ should not be a multiple of 4. Inspecting the tables 2–4 we find that once $I_S^3 - I_F^3$ is not a multiple of 4, the remainder is almost always 2, i.e. $I_S^3 - I_F^3 = 4k + 2$ suggesting $Q = k + 1/2$. In [13] the reported violation of the SU(2) adjoint index theorem was also found in this channel.

On figure 3 we plot the probability of finding a configuration with the property $I_S^3 - I_F^3 = 4k + 2$. It is worth noting that the decrease in this probability from the largest to the second largest lattice spacing is consistent with zero. At smaller lattice spacing a clear decreasing trend sets in which again can be fitted with a linear a -dependence, omitting the largest lattice spacing. The extrapolated a_c value where all such configurations disappear is $a_c/r_0 = 0.068(12)$, compatible with the one obtained in the previous section. Naturally, all the comments in the previous section about the $a \rightarrow 0$ limit and the fitting procedure itself apply to this case as well.

There are also other ways of measuring how the SU(2) index theorem is violated. For




Figure 3. Probability of finding configurations with $I_S^3 - I_F^3 = 4k + 2$ for integer k . These can be interpreted as embedded SU(2) configurations with $I_{\text{adj}}^2 = 4k + 2$ suggesting $Q = k + 1/2$.

example, configurations with $I_{\text{adj}}^2 = 4k + 2$ presumably have $I_F^2 = k$ or $k + 1$. Consequently, we have $|5I_F^3 - I_S^3| = 2$ or 3 for all k . The probability of finding configurations with this property behaves very similarly to figure 3. Or the probability of finding configurations with the property $I_S^3 = 5k + 2$ or $5k + 3$ again behaves similarly. The decrease in all of these probabilities from the largest to the second largest lattice spacing is either consistent with zero or just barely outside 1 standard deviation. All of these observations lead to the conclusion that the low statistics is the most probable reason why the absence of configurations with $I_{\text{adj}}^2 = 2$ was not ruled out in the continuum limit in [13]. The lattice spacings are difficult to compare since that simulation was directly for the gauge group SU(2) though.

From our analysis we see however that in the continuum the adjoint SU(2) index theorem is compatible with integer charge and configurations with fractional charge are lattice artifacts. In this respect the heuristic arguments in [29] suggesting that the configurations in [13] might have been too rough may actually be correct.

Instantons give rise to a non-zero condensate involving $2N$ gauginos in SU(N) $\mathcal{N} = 1$ super-Yang-Mills theory because each instanton carries $2N$ zero modes. The general expectation is that the bilinear gaugino condensate is non-zero as well. Non-periodic boundary conditions are capable of supporting gauge fields with fractional topological charge and it has been argued that these configurations are responsible for the non-zero bilinear condensate [22]. However periodic boundary conditions, at least in the continuum, only allow for smooth gauge fields of integer charge. Since in general the path integral has contributions from non-smooth gauge fields as well, the configurations to which the adjoint index assigns a fractional charge at finite lattice spacing would have been good candidates for explaining the bilinear condensate in a periodic box.

a/r_0	0.2036	0.1745	0.1527	0.1222	0.1018
$r_0^4 \chi_F$	0.062(4)	0.066(5)	0.066(3)	0.059(7)	0.049(11)
$r_0^4 \chi_S$	0.060(5)	0.061(5)	0.062(3)	0.058(7)	0.050(12)
$r_0^4 \chi'_S$	0.052(7)	0.060(8)	0.059(6)	0.055(7)	0.050(13)

Table 5. Topological susceptibility from both the fundamental and sextet representations. In the last row for the sextet representation only configurations with integer $Q = I_S/5$ were considered.

One consequence of our finding is that this is not the case. The bilinear gaugino condensate cannot be attributed to configurations with fractional topological charge in a periodic box because such configurations do not survive the continuum limit.

3 Topological susceptibility

The topological susceptibility, $\chi = \langle Q^2 \rangle / V$, can straightforwardly be calculated from the distribution of indices and the corresponding charges. The continuum expectation is certainly that the representation used to define the topological charge is irrelevant and any two representations should lead to the same topological susceptibility. The fundamental index always leads to integer charges even at finite lattice spacing, but the sextet index gives fractional charges whenever it is not a multiple of 5. In any case, the charge will be defined as $Q = I/2T$ for both representations.

Table 5 shows the measured susceptibilities for both representations. They are consistent with [30] at comparable lattice spacings. Disregarding the largest lattice spacing a linear fit in a^2 gives $r_0^4 \chi_F = 0.050(8)$ and $r_0^4 \chi_S = 0.054(6)$ for the continuum values. Clearly, they are compatible with each other and also with the continuum extrapolated value in physical volume $L = 1.49 fm$ in [24] and the combined infinite volume and continuum extrapolated value in [31]. Our volume is $L = 1.2215 fm$ but the volume dependence is expected to be very small as long as L is larger than about $1 fm$.

We have seen that in the sextet representation configurations where $I/2T$ is not an integer are lattice artifacts. Consequently one might define the topological susceptibility via restricting the vacuum expectation value $\langle Q^2 \rangle / V$ to those configurations only where $I/2T$ is an integer. This would reduce the statistics and increase errors. Nevertheless the topological susceptibility could be continuum extrapolated in this way as well, omitting the largest lattice spacing similarly to above. The extrapolated topological susceptibility is $r_0^4 \chi'_S = 0.048(3)$ (although with a very small χ^2/dof) which, within errors, agrees with the one obtained above, involving all configurations.

4 Discussion

Strong numerical evidence has been presented that the sextet and fundamental representation give the same integral definition of the topological charge in the continuum limit in a large periodic box. What is known rigorously is that the purely geometric definition of the charge gives an integer once a smoothness condition on the plaquettes, $\|1 - U_p\| < \varepsilon$, is satisfied for some ε . This is a sufficient but not necessary condition [32].

Although the plaquette condition precludes the existence of a positive transfer matrix at finite lattice spacing [33], it also enters the rigorous treatment of the overlap operator. Its locality can be established rigorously if this condition is imposed on all plaquettes [34]. However for locality the link variables need to be much smoother than for having an integer geometric charge, i.e. the ε needed for locality is much smaller than the one needed for an integer geometric charge.

We conjecture that if the stronger plaquette condition is imposed on the link variables, any two representations will lead to identical and integral topological charge via the index theorem and that this integer will agree with the geometric definition. This stronger plaquette condition is probably only sufficient but not necessary.

One byproduct of our analysis was a result on $\mathcal{N} = 1$ super-Yang-Mills theory on R^4 . The non-zero value of the bilinear gaugino condensate is certainly due to non-perturbative effects but it is entirely possible that it does not come from semi-classical gauge fields. For example instantons are most probably not the only contribution to the non-zero multi-linear condensate involving $2N$ gauginos either [35, 36]. In this case instantons of course do contribute but most probably there are other non-perturbative contributions as well. The situation seems similar for the bilinear condensate the only difference being that no contribution can be attributed to semi-classical fields and the entire quantity comes from other non-perturbative effects, at least in a large periodic box, or equivalently on R^4 .

It should be noted that of course on $S^1 \times R^3$ objects with fractional topological charge are well known to exist [37–41] and have been argued to give rise to a non-zero bilinear condensate under suitable boundary conditions [42]. However here we have considered the case of periodic boundary conditions which is appropriate for R^4 .

Acknowledgments

DN would like to acknowledge useful discussions with David Adams, Tom DeGrand, Urs Heller and Ken Intriligator. The computations were carried out on the USQCD clusters at Fermilab and the GPU clusters at Wuppertal University. This work is supported by the NSF under grant 0704171, by the DOE under grants DOE-FG03-97ER40546, DOE-FG-02-97ER25308, by the DFG under grant FO 502/1 and by SFB-TR/55.

References

- [1] P. Hasenfratz, V. Laliena and F. Niedermayer, *The index theorem in QCD with a finite cut-off*, *Phys. Lett. B* **427** (1998) 125 [[hep-lat/9801021](#)] [[SPIRES](#)].
- [2] F. Niedermayer, *Exact chiral symmetry, topological charge and related topics*, *Nucl. Phys. Proc. Suppl.* **73** (1999) 105 [[hep-lat/9810026](#)] [[SPIRES](#)].
- [3] D.H. Adams, *Axial anomaly and topological charge in lattice gauge theory with overlap-Dirac operator*, *Annals Phys.* **296** (2002) 131 [[hep-lat/9812003](#)] [[SPIRES](#)].
- [4] D.H. Adams, *General bounds on the Wilson-Dirac operator*, *Phys. Rev. D* **68** (2003) 065009 [[hep-lat/9907005](#)] [[SPIRES](#)].

- [5] D.H. Adams, *On the continuum limit of fermionic topological charge in lattice gauge theory*, *J. Math. Phys.* **42** (2001) 5522 [[hep-lat/0009026](#)] [[SPIRES](#)].
- [6] J.B. Kogut, J. Shigemitsu and D.K. Sinclair, *Chiral symmetry breaking with octet and sextet quarks*, *Phys. Lett. B* **145** (1984) 239 [[SPIRES](#)].
- [7] Y. Shamir, B. Svetitsky and T. DeGrand, *Zero of the discrete β -function in SU(3) lattice gauge theory with color sextet fermions*, *Phys. Rev. D* **78** (2008) 031502 [[arXiv:0803.1707](#)] [[SPIRES](#)].
- [8] B. Svetitsky, Y. Shamir and T. DeGrand, *Nonperturbative infrared fixed point in sextet QCD*, [arXiv:0809.2885](#) [[SPIRES](#)].
- [9] T. DeGrand, Y. Shamir and B. Svetitsky, *Exploring the phase diagram of sextet QCD*, [arXiv:0809.2953](#) [[SPIRES](#)].
- [10] Z. Fodor, K. Holland, J. Kuti, D. Nogradi and C. Schroeder, *Nearly conformal electroweak sector with chiral fermions*, [arXiv:0809.4888](#) [[SPIRES](#)].
- [11] T. DeGrand, Y. Shamir and B. Svetitsky, *Phase structure of SU(3) gauge theory with two flavors of symmetric-representation fermions*, *Phys. Rev. D* **79** (2009) 034501 [[arXiv:0812.1427](#)] [[SPIRES](#)].
- [12] D.D. Dietrich and F. Sannino, *Walking in the SU(N)*, *Phys. Rev. D* **75** (2007) 085018 [[hep-ph/0611341](#)] [[SPIRES](#)].
- [13] R.G. Edwards, U.M. Heller and R. Narayanan, *Evidence for fractional topological charge in SU(2) pure Yang-Mills theory*, *Phys. Lett. B* **438** (1998) 96 [[hep-lat/9806011](#)] [[SPIRES](#)].
- [14] P.H. Damgaard, U.M. Heller, R. Niclasen and B. Svetitsky, *Patterns of spontaneous chiral symmetry breaking in vectorlike gauge theories*, *Nucl. Phys. B* **633** (2002) 97 [[hep-lat/0110028](#)] [[SPIRES](#)].
- [15] L. Del Debbio, A. Patella and C. Pica, *Higher representations on the lattice: numerical simulations. SU(2) with adjoint fermions*, [arXiv:0805.2058](#) [[SPIRES](#)].
- [16] S. Catterall, J. Giedt, F. Sannino and J. Schneible, *Phase diagram of SU(2) with 2 flavors of dynamical adjoint quarks*, *JHEP* **11** (2008) 009 [[arXiv:0807.0792](#)] [[SPIRES](#)].
- [17] A. Hietanen, J. Rantaharju, K. Rummukainen and K. Tuominen, *Spectrum of SU(2) gauge theory with two fermions in the adjoint representation*, *PoS(LATTICE 2008)065* [[arXiv:0810.3722](#)] [[SPIRES](#)].
- [18] L. Del Debbio, A. Patella and C. Pica, *Fermions in higher representations. Some results about SU(2) with adjoint fermions*, [arXiv:0812.0570](#) [[SPIRES](#)].
- [19] A.J. Hietanen, J. Rantaharju, K. Rummukainen and K. Tuominen, *Spectrum of SU(2) lattice gauge theory with two adjoint Dirac flavours*, *JHEP* **05** (2009) 025 [[arXiv:0812.1467](#)] [[SPIRES](#)].
- [20] A.J. Hietanen, K. Rummukainen and K. Tuominen, *Evolution of the coupling constant in SU(2) lattice gauge theory with two adjoint fermions*, [arXiv:0904.0864](#) [[SPIRES](#)].
- [21] G. Veneziano and S. Yankielowicz, *An effective Lagrangian for the pure $N = 1$ supersymmetric Yang-Mills theory*, *Phys. Lett. B* **113** (1982) 231 [[SPIRES](#)].
- [22] E. Cohen and C. Gomez, *Chiral symmetry breaking in supersymmetric Yang-Mills*, *Phys. Rev. Lett.* **52** (1984) 237 [[SPIRES](#)].

- [23] H. Neuberger, *Exactly massless quarks on the lattice*, *Phys. Lett. B* **417** (1998) 141 [[hep-lat/9707022](#)] [[SPIRES](#)].
- [24] L. Giusti, M. Lüscher, P. Weisz and H. Wittig, *Lattice QCD in the ϵ -regime and random matrix theory*, *JHEP* **11** (2003) 023 [[hep-lat/0309189](#)] [[SPIRES](#)].
- [25] Z. Fodor, K. Holland, J. Kuti, D. Nogradi and C. Schroeder, *Chiral properties of SU(3) sextet fermions*, [arXiv:0908.2466](#) [[SPIRES](#)].
- [26] S. Necco and R. Sommer, *The $N_f = 0$ heavy quark potential from short to intermediate distances*, *Nucl. Phys. B* **622** (2002) 328 [[hep-lat/0108008](#)] [[SPIRES](#)].
- [27] C. Morningstar and M.J. Peardon, *Analytic smearing of SU(3) link variables in lattice QCD*, *Phys. Rev. D* **69** (2004) 054501 [[hep-lat/0311018](#)] [[SPIRES](#)].
- [28] Z. Fodor, S.D. Katz and K.K. Szabo, *Dynamical overlap fermions, results with hybrid Monte-Carlo algorithm*, *JHEP* **08** (2004) 003 [[hep-lat/0311010](#)] [[SPIRES](#)].
- [29] M. Garcia Perez, A. Gonzalez-Arroyo and A. Sastre, *Adjoint zero-modes as a tool to understand the Yang-Mills vacuum*, *PoS(LATTICE 2007)328* [[arXiv:0710.0455](#)] [[SPIRES](#)].
- [30] T.G. Kovacs, *Locality and topology with fat link overlap actions*, *Phys. Rev. D* **67** (2003) 094501 [[hep-lat/0209125](#)] [[SPIRES](#)].
- [31] S. Dürr, Z. Fodor, C. Hölbling and T. Kurth, *Precision study of the SU(3) topological susceptibility in the continuum*, *JHEP* **04** (2007) 055 [[hep-lat/0612021](#)] [[SPIRES](#)].
- [32] M. Lüscher, *Topology of lattice gauge fields*, *Commun. Math. Phys.* **85** (1982) 39 [[SPIRES](#)].
- [33] M. Creutz, *Positivity and topology in lattice gauge theory*, *Phys. Rev. D* **70** (2004) 091501 [[hep-lat/0409017](#)] [[SPIRES](#)].
- [34] P. Hernández, K. Jansen and M. Lüscher, *Locality properties of Neuberger's lattice Dirac operator*, *Nucl. Phys. B* **552** (1999) 363 [[hep-lat/9808010](#)] [[SPIRES](#)].
- [35] J. Fuchs and M.G. Schmidt, *Instanton induced Green functions in the superfield formalism*, *Z. Phys. C* **30** (1986) 161 [[SPIRES](#)].
- [36] T.J. Hollowood, V.V. Khoze, W.-J. Lee and M.P. Mattis, *Breakdown of cluster decomposition in instanton calculations of the gluino condensate*, *Nucl. Phys. B* **570** (2000) 241 [[hep-th/9904116](#)] [[SPIRES](#)].
- [37] T.C. Kraan and P. van Baal, *Periodic instantons with non-trivial holonomy*, *Nucl. Phys. B* **533** (1998) 627 [[hep-th/9805168](#)] [[SPIRES](#)].
- [38] T.C. Kraan and P. van Baal, *Monopole constituents inside SU(n) calorons*, *Phys. Lett. B* **435** (1998) 389 [[hep-th/9806034](#)] [[SPIRES](#)].
- [39] F. Bruckmann, D. Nogradi and P. van Baal, *Constituent monopoles through the eyes of fermion zero-modes*, *Nucl. Phys. B* **666** (2003) 197 [[hep-th/0305063](#)] [[SPIRES](#)].
- [40] F. Bruckmann, D. Nogradi and P. van Baal, *Higher charge calorons with non-trivial holonomy*, *Nucl. Phys. B* **698** (2004) 233 [[hep-th/0404210](#)] [[SPIRES](#)].
- [41] E. Poppitz and M. Ünsal, *Index theorem for topological excitations on $R^3 \times S^1$ and Chern-Simons theory*, *JHEP* **0903** (2009) 027 [[arXiv:0812.2085](#)] [[SPIRES](#)].
- [42] N.M. Davies, T.J. Hollowood, V.V. Khoze and M.P. Mattis, *Gluino condensate and magnetic monopoles in supersymmetric gluodynamics*, *Nucl. Phys. B* **559** (1999) 123 [[hep-th/9905015](#)] [[SPIRES](#)].