

On sl_3 Knizhnik-Zamolodchikov equations and \mathcal{W}_3 null-vector equations

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

JHEP10(2009)002

(<http://iopscience.iop.org/1126-6708/2009/10/002>)

[The Table of Contents](#) and [more related content](#) is available

Download details:

IP Address: 80.92.225.132

The article was downloaded on 01/04/2010 at 13:40

Please note that [terms and conditions apply](#).

On sl_3 Knizhnik-Zamolodchikov equations and \mathcal{W}_3 null-vector equations

Sylvain Ribault

*Laboratoire de Physique Théorique et Astroparticules, UMR5207 CNRS-UM2,
Université Montpellier II, Place E. Bataillon, CC 070,
34095 Montpellier Cedex 05, France*

E-mail: ribault@lpta.univ-montp2.fr

ABSTRACT: Starting from Sklyanin's separation of variables for the sl_3 Yangian model, we derive the separation of variables for the quantum sl_3 Gaudin model. We use the resulting new variables for rewriting the sl_3 Knizhnik-Zamolodchikov equations, and comparing them with certain null-vector equations in conformal field theories with \mathcal{W}_3 -algebra symmetry. The two sets of equations are remarkably similar, but become identical only in the critical level limit. This is in contrast to the sl_2 Knizhnik-Zamolodchikov equations, which are known to be equivalent to Belavin-Polyakov-Zamolodchikov equations for all values of the level.

KEYWORDS: Conformal and W Symmetry, Integrable Field Theories

ARXIV EPRINT: [0811.4587](https://arxiv.org/abs/0811.4587)

Contents

1	Introduction and conjecture	1
2	Gaudin hamiltonians in conformal field theory	3
2.1	Knizhnik-Zamolodchikov equations	4
2.2	Ward identities for the cubic field	5
3	Review of the sl_2 case	6
3.1	Separation of variables for the sl_2 Gaudin model	7
3.2	The sl_2 Knizhnik-Zamolodchikov equations in Sklyanin variables	9
3.3	Comparison with Virasoro null-vector equations	10
4	The sl_3 case	12
4.1	Separation of variables for the sl_3 Gaudin model	12
4.2	The sl_3 Knizhnik-Zamolodchikov equations in Sklyanin variables	15
4.3	\mathcal{W}_3 null-vector equations	17
4.4	Comparing sl_3 Knizhnik-Zamolodchikov equations with \mathcal{W}_3 null-vector equations	21
5	Conclusion	23
A	A few technical results	24
A.1	Helpful identities	24
A.2	A characterization of $\mathcal{F}_2(y)$	25
A.3	Study of the differential operator D_1	26

1 Introduction and conjecture

Many interesting models of two-dimensional conformal field theories are based on affine Lie algebras \widehat{sl}_N and their cosets, starting with Wess-Zumino-Witten models. To solve such theories is an interesting challenge, whose difficulty depends more from the choice of the underlying Lie algebra sl_N , than from the particular coset or real form chosen.

For example, the sl_2 family includes string theory in AdS_3 and in the $SL(2, \mathbb{R})/U(1)$ 2d black hole, as well as the H_3^+ model; the simplest non-rational nontrivial model of the family is however Liouville theory, also known as conformal sl_2 Toda theory. In several of the other theories in the sl_2 family, it turns out that arbitrary correlation functions have a simple relation to certain Liouville theory correlation functions [1, 2]. This relation entails a relation between the Knizhnik-Zamolodchikov equations which follow from \widehat{sl}_2 symmetry,

and the Belavin-Polyakov-Zamolodchikov equations which follow from the conformal symmetry of Liouville theory [3]. The relation to Liouville theory is helpful in solving certain models in the sl_2 family, by disentangling the particular details of a model from its general sl_2 -based properties. For example, the H_3^+ -Liouville relation was very helpful in solving the H_3^+ model on a disc [4]. Moreover, playing with the Liouville side of the relation leads to the discovery of new conformal field theories which generalize the H_3^+ model [5], and which can be considered as members of an extended sl_2 family.

The intuitive reason why such a relation exists is that $\widehat{sl_2}$ representations are parametrized by just one number, their spin. So it is not very surprising that the dynamics of say the H_3^+ model, a theory of three interacting bosons, are in some sense effectively one-dimensional. Applied to a theory with an $\widehat{sl_{N>2}}$ symmetry algebra, which may involve as many as $N^2 - 1$ bosons, this reasoning suggests that it could be related to a theory of only $N - 1$ bosons. Such a theory is present in the sl_N family: namely, conformal sl_N Toda theory, which can be described by the Lagrangian $L = (\partial\phi, \bar{\partial}\phi) + \sum_{i=1}^{N-1} e^{b(e_i, \phi)}$ where the field $\phi(z, \bar{z})$ and the simple roots e_i live in the $N - 1$ -dimensional root space of sl_N . (See for example [6] for details.) It is therefore natural to investigate whether correlation functions of that theory have a simple relation to correlation functions of other models in the family. Such a relation would be a welcome simplification: for instance, in the sl_3 family, we would trade 8 bosons of the $SL(3, \mathbb{R})$ WZW model for the 2 bosons of sl_3 conformal Toda theory.

The investigation of the $sl_{N>2}$ families is motivated both from the appearance of groups of rank higher than one in many interesting string theory backgrounds, and from the observation that theories in the $sl_{N>2}$ families are qualitatively more difficult, and more generic, than theories in the sl_2 family. This is due to features like: infinite fusion multiplicities, correlation functions involving degenerate fields without obeying nontrivial differential equations, and structures constants which can probably not be written in terms of known special functions [6]. These are serious obstacles in the way of solving such theories. Nevertheless, we do know a strong explicit constraint on the correlation functions of all models which have the full $\widehat{sl_N}$ symmetry: they obey KZ equations. The aim of the present article is therefore to determine whether the sl_3 KZ equations are related to some null-vector equations in conformal sl_3 Toda theory, which follow from its symmetry algebra \mathcal{W}_3 .

In analogy with the sl_2 case, we will look for a relation based on Sklyanin's separation of variables [7]. As the KZ equations are closely related to the Gaudin Hamiltonians, we will use Sklyanin's separation of variables for the quantum sl_3 Gaudin model. Before using it, we will actually have to work it out, as this has apparently not been fully done in the existing literature. A rather close starting point is available though: the separation of variables for the sl_3 Yangian model [8].

Let us now sketch the correlation functions we are interested in and the relation we are aiming at. Consider a theory with an $\widehat{sl_3}$ symmetry algebra at level k . We are interested in correlation functions of generic $\widehat{sl_3}$ affine primary fields $\Phi^j(x|z)$, where the spin j labels sl_3 representations, the variable x is a generic isospin coordinate (a triplet of complex numbers), and z is a coordinate on the complex plane where the field lives. We denote an

n -point function of such fields as

$$\Omega_n \equiv \left\langle \prod_{i=1}^n \Phi^{j_i}(x_i|z_i) \right\rangle. \quad (1.1)$$

We will seek to relate such correlation functions to fairly particular correlation functions in a theory with a \mathcal{W}_3 symmetry algebra at parameter $b = (k - 3)^{-\frac{1}{2}}$, which involve not only n generic \mathcal{W}_3 -primary fields $V_{\alpha_i}(z_i)$ corresponding to $\Phi^{j_i}(x_i|z_i)$, but also $3n - 6$ degenerate fields $V_{-b^{-1}\omega_1}(y_a)$ with the special value $-b^{-1}\omega_1$ for their \mathcal{W}_3 momentum:

$$\tilde{\Omega}_n \equiv \left\langle \prod_{a=1}^{3n-6} V_{-b^{-1}\omega_1}(y_a) \prod_{i=1}^n V_{\alpha_i}(z_i) \right\rangle. \quad (1.2)$$

The number of degenerate fields is of the order of $3n$, which allows their worldsheet positions y_a to (approximately) correspond to the $3n$ components of the isospin variables $x_1 \cdots x_n$. This will also allow $\tilde{\Omega}_n$ to obey some differential equations which may be related to the KZ equations for Ω_n . Moreover, the tentative relation between Ω_n and $\tilde{\Omega}_n$ will involve a simple twist function

$$\Theta_n = \prod_{a < b} (y_a - y_b)^\lambda \prod_i \prod_a (y_a - z_i)^\mu \prod_{i < j} (z_i - z_j)^\nu, \quad (1.3)$$

for some constants λ, μ, ν to be determined in terms of the level k of our \widehat{sl}_3 algebra; and the integral transformation \mathcal{K} with integration kernel $K(\{x_i|\{y_a\}, U|\{z_i\})$ which implements Sklyanin's separation of variables, and may therefore depend on the spins j_i but not on the level k . We will then investigate the validity of the conjecture $\Omega_n \stackrel{?}{\sim} \mathcal{K} \cdot \Theta_n \tilde{\Omega}_n \equiv \int dU \prod_a dy_a K \cdot \Theta_n \tilde{\Omega}_n$, or more explicitly

$$\Omega_n(\{x_i|\{z_i\}) \stackrel{?}{\sim} \int dU \prod_a dy_a K(\{x_i|\{y_a\}, U|\{z_i\}) \cdot \Theta_n(\{y_a|\{z_i\}) \tilde{\Omega}_n(\{y_a|\{z_i\}). \quad (1.4)$$

The meaning of the equivalence \sim here is that both sides obey the same differential equations. If true, this equivalence may then be promoted to a relation between physical correlation function of specific models, like the relation between the H_3^+ model and Liouville theory [1], but this is not the focus of the present article. This is why we do not worry about such details as the dependence of the correlation functions on antiholomorphic variables.

The article will start with a brief review of the KZ equations and other Ward identities in conformal field theories with \widehat{sl}_N symmetries, where we will explain how the Gaudin Hamiltonians appear in such equations. We will then review the KZ-BPZ relation in the sl_2 case; the reader is not advised to skip that section as the KZ-BPZ relation is presented in a form suitable for generalization to sl_3 . In the sl_3 case, we will then find that the conjecture (1.4) holds only in the critical level limit $k \rightarrow 3$.

2 Gaudin hamiltonians in conformal field theory

We will review how the Gaudin Hamiltonians appear in Ward identities obeyed by correlation functions in conformal field theories with an \widehat{sl}_N symmetry algebra. The Ward

identities associated to the stress-energy tensor $T^J(z)$ lead to the KZ equations, which involve the ordinary Gaudin Hamiltonians. The Ward identities associated to the cubic field $W^J(z)$ involve higher Gaudin Hamiltonians.

2.1 Knizhnik-Zamolodchikov equations

The affine Lie algebra \widehat{sl}_N is an infinite-dimensional extension of the simple Lie algebra sl_N . The generators t^a of sl_N , its structure constants f_c^{ab} , and its metric κ^{ab} are defined by the relations

$$[t^a, t^b] = f_c^{ab} t^c \quad , \quad \kappa^{ab} \equiv \text{Tr } t^a t^b \quad , \quad f_c^{ab} f_b^{cd} = 2N \kappa^{ad} \quad , \quad (2.1)$$

where here and in the following the trace Tr is taken in the fundamental representation, so that our metric κ^{ab} coincides with the renormalized Killing form of [9](13.13). The affine Lie algebra \widehat{sl}_N can be formulated as the algebra of currents $J^a(z)$ with the operator product expansion

$$J^a(z) J^b(w) = -\frac{k \kappa^{ab}}{(z-w)^2} + f_c^{ab} \frac{J^c(w)}{z-w} + (J^a J^b)(w) + \mathcal{O}(z-w) \quad , \quad (2.2)$$

where the parameter k is called the level, and the normal-ordered product $(J^a J^b)(w)$ is defined by the present formula. Conformal symmetry follows from the existence of a Virasoro algebra with central charge $c = \frac{k(N^2-1)}{k-N}$, generated by the Sugawara stress-energy tensor

$$T^J(z) \equiv -\frac{1}{2(k-N)} (J^a J^a)(z) \quad , \quad (2.3)$$

where $J^a J^a$ is a shorthand for $\kappa_{ab} J^a J^b$. The identification of $T^J(z)$ with the generator of conformal transformations will be at the origin of the KZ equations. These equations are satisfied by any correlation function (1.1) of n affine primary fields $\Phi^{j_i}(x_i|z_i)$ on the complex z -plane, where the spins j_i label representations of sl_N , the isospin variables x_i label the states in a given representation, and the complex numbers z_i are positions on the Euclidean two-dimensional spacetime. The affine primary fields are defined by their operator product expansions with the currents $J^a(z)$,

$$J^a(z) \Phi^j(x|w) = \frac{D^a \Phi^j(x|w)}{z-w} + \mathcal{O}(1) \quad , \quad (2.4)$$

where D^a provides a realization of the representation of spin j in terms of differential operators acting on the isospin variables x , so that $[D^a, D^b] = f_c^{ab} D^c$. We will keep this realization arbitrary, without committing to any particular choice of isospin variables. Let us however give an example of such a choice in the sl_2 case:

$$D^- = \frac{\partial}{\partial x} \quad , \quad D^3 = x \frac{\partial}{\partial x} - j \quad , \quad D^+ = x^2 \frac{\partial}{\partial x} - 2jx \quad . \quad (2.5)$$

The KZ equations are now obtained by inserting $T^J(z)$ into the correlation function Ω_n , and using the conformal Ward identity for $T^J(z)$ on the one hand, and the affine Ward

identities for $(J^a J^a)(z)$ on the other hand:

$$\begin{aligned} \left\langle T^J(z) \prod_{i=1}^n \Phi^{j_i}(x_i|z_i) \right\rangle &= \sum_{i=1}^n \left(\frac{L_{0,(i)}^J}{(z-z_i)^2} + \frac{L_{-1,(i)}^J}{z-z_i} \right) \Omega_n \\ &= -\frac{1}{2(k-N)} \sum_{i=1}^n \frac{D_{(i)}^a}{z-z_i} \sum_{\ell=1}^n \frac{D_{(\ell)}^a}{z-z_\ell} \Omega_n, \end{aligned} \quad (2.6)$$

where the subscript (i) in $D_{(i)}^a$ indicates that it acts on the isospin variables x_i , and by definition $L_{p,(i)}^J$ is the p -th mode of $T^J(z)$ acting on $\Phi^{j_i}(x_i|z_i)$, according to

$$L_p^J \Phi^j(x|z) \equiv \frac{1}{2\pi i} \oint_z dw (w-z)^{p+1} T^J(w) \Phi^j(x|z). \quad (2.7)$$

Calling Δ^J the eigenvalues of L_0^J , such that $L_{0,(i)}^J \Omega_n = \Delta_{j_i}^J \Omega_n$, we first deduce from eq. (2.6) the expression for Δ_j in terms of the quadratic Casimir $C_2(j) \equiv D^a D^a$ of the $s\ell_N$ representation with spin j ,

$$\Delta_j^J \equiv -\frac{C_2(j)}{2(k-N)}. \quad (2.8)$$

Now $T^J(z)$ is assumed to generate conformal transformations, and in particular $L_{-1,(i)}^J \Omega_n = \frac{\delta}{\delta z_i} \Omega_n$. (We define $\frac{\delta}{\delta z_i} \equiv \left. \frac{\partial}{\partial z_i} \right|_{x_i}$ as a derivative at fixed isospin variables.) Together with eq. (2.6), this implies the KZ equations [10]

$$(k-N) \frac{\delta}{\delta z_i} \Omega_n = -H_i \Omega_n \quad , \quad H_i \equiv \sum_{\ell \neq i} \frac{D_{(i)}^a D_{(\ell)}^a}{z_i - z_\ell}, \quad (2.9)$$

The n commuting differential operators H_i are called the Gaudin Hamiltonians. Through its dependence on $D_{(i)}^a$ and $D_{(\ell)}^a$, each one of the n Hamiltonians involves all of the n isospin variables x_i , which makes the problem of their simultaneous diagonalization difficult. This difficulty will be solved by Sklyanin's separation of variables, which replaces the isospins x_i with new variables y_i , and combines the Gaudin eigenvalue equations into an essentially equivalent set of equations, each of which involves only one of the new variables.

2.2 Ward identities for the cubic field

In addition to the quadratic invariant tensor $\kappa^{ab} = \text{Tr } t^a t^b$, it is possible to define the fully symmetric cubic invariant tensor

$$d^{abc} \equiv \text{Tr } (t^a t^b t^c + t^a t^c t^b). \quad (2.10)$$

This tensor vanishes in the case of $s\ell_2$, but not in the cases of $s\ell_{N \geq 3}$. It can then be used for constructing the invariant cubic field

$$W^J(z) \equiv \frac{1}{6} \rho \, d_{abc} (J^a (J^b J^c))(z) \quad , \quad \rho \equiv \frac{i}{(k-N)^{\frac{3}{2}}}. \quad (2.11)$$

This generalizes the Sugawara construction (2.3), with however two substantial differences. First, while the field $T^J(z)$ is interpreted as the generator of conformal transformations, there is no such geometrical interpretation for $W^J(z)$. Second, while the field $T^J(z)$ obeys a Virasoro algebra, the field $W^J(z)$ does not obey the higher \mathcal{W}_3 algebra [11]. In other words, while the Virasoro algebra can be realized as either a coset of \widehat{sl}_2 or a subalgebra of the enveloping algebra of $\widehat{sl}_{N \geq 2}$ (albeit with differing central charges), the \mathcal{W}_3 algebra is a coset of \widehat{sl}_3 but not a subalgebra of the enveloping algebra of $\widehat{sl}_{N \geq 3}$.

In analogy with eq. (2.6) we now have

$$\begin{aligned} \left\langle W^J(z) \prod_{i=1}^n \Phi^{j_i}(x_i|z_i) \right\rangle &= \sum_{i=1}^n \left(\frac{W_{0,(i)}^J}{(z-z_i)^3} + \frac{W_{-1,(i)}^J}{(z-z_i)^2} + \frac{W_{-2,(i)}^J}{z-z_i} \right) \Omega_n \\ &= \frac{1}{6} \rho \, d_{abc} \sum_{i=1}^n \frac{D_{(i)}^a}{z-z_i} \sum_{\ell=1}^n \frac{D_{(\ell)}^b}{z-z_\ell} \sum_{m=1}^n \frac{D_{(m)}^c}{z-z_m} \Omega_n, \end{aligned} \quad (2.12)$$

where by definition $W_{p,(i)}^J$ is the p -th mode of $W^J(z)$ acting on $\Phi^{j_i}(x_i|z_i)$, according to

$$W_p^J \Phi^j(x|z) \equiv \frac{1}{2\pi i} \oint_z dw (w-z)^{p+2} W^J(w) \Phi^j(x|z). \quad (2.13)$$

Calling q^J the eigenvalues of W_0^J , such that $W_{0,(i)}^J \Omega_n = q_{j_i}^J \Omega_n$, we first deduce from eq. (2.12) the expression for q_j^J in terms of the cubic Casimir $C_3(j) \equiv d_{abc}(D^a D^b D^c + D^a D^c D^b)$ of the sl_N representation with spin j ,

$$q_j^J = \frac{1}{6} \rho \, C_3(j). \quad (2.14)$$

We further deduce

$$W_{-1,(i)}^J \Omega_n = \frac{1}{2} \rho \, H'_i \Omega_n, \quad (2.15)$$

$$W_{-2,(i)}^J \Omega_n = \frac{1}{2} \rho \, H''_i \Omega_n, \quad (2.16)$$

where the differential operators H'_i and H''_i are higher Gaudin Hamiltonians, whose explicit expressions in terms of $D_{(i)}^a$ can easily be derived from eq. (2.12). But, in contrast to L_{-1}^J , the operators W_{-1}^J and W_{-2}^J are not interpreted as differential operators with respect to z . The equations (2.15) and (2.16), which generalize the KZ equations, are therefore not differential equations, and they will therefore not help us test our conjecture. Nevertheless, they will naturally appear in certain formulas.

3 Review of the sl_2 case

In this section we will review the relation between the sl_2 KZ equations and BPZ equations. This was originally found by Feigin, Frenkel and Stoyanovsky [3], using Sklyanin's separation of variables for the sl_2 Gaudin model [7]. However, the original derivation relied on a particular choice of the isospin variables. This choice of isospin variables makes the result

remarkably simple, but has no analog in the sl_3 case, as we will show. We will therefore reanalyze the sl_2 case, using whenever possible objects which do have analogs in the sl_3 or even sl_N cases. We will present systematic derivations of their relevant properties, which will help clarify whether and how they can be generalized to the sl_3 case.

3.1 Separation of variables for the sl_2 Gaudin model

Let us consider a system of n representations of sl_2 with spins $j_1 \cdots j_n$. Consider the associated quantum variables $D_{(i)}^a$ such that $[D_{(i)}^a, D_{(j)}^b] = \delta_{ij} f_c^{ab} D_{(i)}^c$ with $D_{(i)}^a D_{(i)}^a = C_2(j_i)$. The system comes with parameters $z_1 \cdots z_n$. Sklyanin's separation of variables for this system involves three ingredients:

1. A function $B(u)$ of an arbitrary variable u (the spectral parameter), whose zeroes are the separated variables y_i , so that $B(y_i) = 0$;
2. Another function $A(u)$ such that $p_i = A(y_i)$ is the conjugate momenta to y_i ;
3. A kinematical identity, called the characteristic equation, which for any given i relates y_i and p_i .

We now briefly review the construction of these three objects in the sl_2 case. They are built from the sl_2 Lax matrix

$$I(u) \equiv - \sum_{i=1}^n \frac{t^a D_{(i)}^a}{u - z_i}, \tag{3.1}$$

whose matrix elements $I_\alpha^\beta(u)$ obey the identity

$$(u - v)[I_\alpha^\gamma(u), I_\beta^\epsilon(v)] = \delta_\alpha^\epsilon I_\beta^\gamma(u) - \delta_\beta^\gamma I_\alpha^\epsilon(u) - \delta_\alpha^\epsilon I_\beta^\gamma(v) + \delta_\beta^\gamma I_\alpha^\epsilon(v). \tag{3.2}$$

With the particular choice eq. (2.5) for the sl_2 isospin variable x , the sl_2 Lax matrix is explicitly

$$I(u) = - \begin{bmatrix} \frac{1}{2} \sum_{i=1}^n \frac{1}{u-z_i} \left(x_i \frac{\partial}{\partial x_i} - j_i \right) & \sum_{i=1}^n \frac{1}{u-z_i} \frac{\partial}{\partial x_i} \\ \sum_{i=1}^n \frac{1}{u-z_i} \left(x_i^2 \frac{\partial}{\partial x_i} - 2j_i x_i \right) & -\frac{1}{2} \sum_{i=1}^n \frac{1}{u-z_i} \left(x_i \frac{\partial}{\partial x_i} - j_i \right) \end{bmatrix}. \tag{3.3}$$

Now choosing

$$B(u) \equiv I_1^2(u) \quad , \quad A(u) \equiv I_1^1(u), \tag{3.4}$$

it is easy to check that

$$[B(u), B(v)] = 0 \quad , \quad [A(u), A(v)] = 0, \tag{3.5}$$

$$(u - v)[A(u), B(v)] = B(v) - B(u). \tag{3.6}$$

These relations ensure that the operators y_i defined as the zeroes of $B(u)$, and $p_i = A(y_i)$, do satisfy

$$[y_i, y_j] = 0 \quad , \quad [p_i, y_j] = \delta_{ij} \quad , \quad [p_i, p_j] = 0. \tag{3.7}$$

In particular, $[p_i, B(v)] = \frac{B(v)}{y_i - v}$ agrees with $B(v) \propto \frac{\prod_i (v - y_i)}{\prod_j (v - z_j)}$. There is however a problem of operator ordering in the expressions $A(y_i)$ and $B(y_i)$, because the separated variables y_i are operators. This problem is dealt with in reference [7]. We will ignore it in the forthcoming heuristic derivation of the characteristic equation. Let us start with $\det(A(y_i)\text{id} - I(y_i)) = 0$, where id is the identity matrix. (The determinant of a matrix whose first line vanishes is zero.) This implies $p_i^2 - \frac{1}{2}(I_\alpha^\beta I_\beta^\alpha)(y_i) = 0$. This characteristic equation can easily be rewritten as

$$p_i^2 - \frac{1}{2} \sum_\ell \frac{C_2(j_\ell)}{(y_i - z_\ell)^2} - \sum_\ell \frac{1}{y_i - z_\ell} H_\ell = 0, \quad (3.8)$$

where H_ℓ is of course a Gaudin Hamiltonian (2.9), and $C_2(j)$ is the quadratic Casimir of a spin- j representation.

Functional space interpretation. We now wish to consider the quantum variables $D_{(i)}^a$ as differential operators acting on functions $\Psi(\{x_i\})$ of isospin variables x_i . (An example of such a realization was given in eq. (2.5).) Similarly, the separated variables y_ℓ and their associated momenta p_ℓ may act on functions $\tilde{\Psi}(\{y_\ell\})$, in particular $p_\ell \tilde{\Psi} = \frac{\partial}{\partial y_\ell} \tilde{\Psi}$. The separation of variables $\{x_i\} \rightarrow \{y_\ell\}, U$ (where the extra variable U will be defined shortly) is then interpreted as an integral transformation \mathcal{K} such that

$$\Psi(\{x_i\}) = \mathcal{K} \tilde{\Psi}(\{y_\ell\}, U) = \int dU \int \prod_\ell dy_\ell K(\{x_i\}|\{y_\ell\}, U) \tilde{\Psi}(\{y_\ell\}, U), \quad (3.9)$$

where the kernel K is characterized as a common eigenvector of the commuting operators $B(u)$

$$\left(B(u) - U \frac{\prod_\ell (u - y_\ell)}{\prod_i (u - z_i)} \right) K(\{x_i\}|\{y_\ell\}, U) = 0. \quad (3.10)$$

The simultaneous diagonalization of the Gaudin Hamiltonians H_j , namely the set of equations $(H_\ell - E_\ell)\Psi = 0$, can now be reformulated using the characteristic equation (3.8), which implies

$$\left(\frac{\partial^2}{\partial y_i^2} - \frac{1}{2} \sum_\ell \frac{C_2(j_\ell)}{(y_i - z_\ell)^2} - \sum_\ell \frac{E_\ell}{y_i - z_\ell} \right) \tilde{\Psi} = 0, \quad (3.11)$$

The solutions of this equation can be found in factorized form $\tilde{\Psi} = \prod_i \tilde{\psi}(y_i)$. This justifies the name ‘‘separation of variables’’ attributed to the change of variables $x_i \rightarrow y_i$.

Some remarks. Finding the kernel K by the simultaneous diagonalization of the operators $B(u)$ is easy in the $s\ell_2$ case because $B(u) = I_1^2(u)$ is a sum of n commuting operators, so that we have $K(\{x_i\}|\{y_\ell\}, U) = \prod_{i=1}^n k_i(x_i|\{y_\ell\}, U)$ where the equation on k_i is obtained from eq. (3.10) in the limit $u \rightarrow z_i$:

$$\left((t^a)_1^2 D_{(i)}^a + \mu_i \right) k_i(x_i|\{y_\ell\}, U) = 0 \quad , \quad \mu_i \equiv U \frac{\prod_\ell (z_i - y_\ell)}{\prod_{j \neq i} (z_i - z_j)}. \quad (3.12)$$

For example, if the isospin variables are chosen as in eq. (2.5), then we find $k_i = e^{-\mu_i x_i}$. This suggests that we could use other isospin variables $\hat{\mu}_i$ such that $D_{(i)}^a (t^a)_1^2 = -\hat{\mu}_i$, then we would find $k_i \propto \delta(\hat{\mu}_i - \mu_i)$, so that we could explicitly perform the integrals in eq. (3.9). This would lead to $\Psi(\{\hat{\mu}_i\}) \propto \Psi(\{y_\ell\}, U)$ with simple proportionality factors, as the change of variables $\{\hat{\mu}_i\} \rightarrow (\{y_\ell\}, U)$ would now be local and described by the functions $\mu_i(\{y_\ell\}, U)$. More generally, for any choice of isospin variables, the kernel K will be of the type

$$K(\{x_i\}|\{y_\ell\}, U|\{z_j\}) = \prod_{i=1}^n k_i(x_i|\{\mu_j\}), \quad (3.13)$$

where $\mu_j(\{y_\ell\}, U|\{z_j\})$ is defined in eq. (3.12), and we made the z_j -dependence explicit. Thus, in the sl_2 case, the kernel K can be determined explicitly, and this is because the operator $B(u)$ is a linear function of the Lax matrix $I(u)$.

Let us finally be more precise about the number of variables y_ℓ . They are defined as the zeroes of a rational function $B(u)$ which, barring extra constraints, has n poles and degree -1 . Therefore we must have $n - 1$ such variables, and the n th variable U is the eigenvalue of $-(t^a)_1^2 \sum_{i=1}^n D_{(i)}^a$. In conformal field theory applications, we however impose the extra constraint $\sum_{i=1}^n D_{(i)}^a = 0$, so that $B(u)$ has degree -2 . This yields $n - 2$ variables $\{y_\ell\}_{\ell=1 \dots n-2}$, and U is the eigenvalue of $-(t^a)_1^2 \sum_{i=1}^n z_i D_{(i)}^a$.

3.2 The sl_2 Knizhnik-Zamolodchikov equations in Sklyanin variables

We just saw that Sklyanin's separation of variables is useful tool for simultaneously diagonalizing the sl_2 Gaudin Hamiltonians. This problem is closely related to the problem of solving the KZ equations (2.9), which are obtained by replacing the eigenvalues of the Gaudin Hamiltonians H_i with $-(k - 2) \frac{\delta}{\delta z_i}$. This suggests that it may be interesting to rewrite the KZ equations in terms of Sklyanin's variables. To do this, we will use the characteristic equation (3.8) which such variables obey, and apply it to $\mathcal{K}^{-1} \Omega_n$, which is a function of $\{y_i\}$, so that $p_i \mathcal{K}^{-1} \Omega_n = \frac{\partial}{\partial y_i} \mathcal{K}^{-1} \Omega_n$. While itself just a kinematical identity, the characteristic equation then allows us to reorganize the KZ equations as

$$\left(\frac{1}{k-2} \frac{\partial^2}{\partial y^2} + \sum_{\ell=1}^n \frac{1}{y-z_\ell} \mathcal{K}^{-1} \frac{\delta}{\delta z_\ell} \mathcal{K} + \sum_{\ell=1}^n \frac{\Delta_{j_\ell}^J}{(y-z_\ell)^2} \right) \mathcal{K}^{-1} \Omega_n = 0, \quad (3.14)$$

where we drop the index from y_i , and we use $\Delta_j^J = -\frac{C_2(j)}{2(k-2)}$ from eq. (2.8). We still have to perform the change of variables on the z_ℓ -derivatives at fixed isospins, i.e. to rewrite $\mathcal{K}^{-1} \frac{\delta}{\delta z_\ell} \mathcal{K}$ in terms of $\frac{\partial}{\partial z_\ell} \Big|_{y_a}$. This is rather easy because of the particular form of the kernel (3.13), where the dependences on $\{y_a\}, U$ and $\{z_\ell\}$ are channeled through the particular functions $\{\mu_i\}$. This implies that the integral transformation (3.9) just adds first-order differential operators $\frac{\partial}{\partial y_a}, \frac{\partial}{\partial U}$ to $\frac{\delta}{\delta z_\ell}$, so that

$$\mathcal{K}^{-1} \frac{\delta}{\delta z_\ell} \mathcal{K} = \frac{\partial}{\partial z_\ell} + \sum_a \frac{\partial y_a}{\partial z_\ell} \Big|_{\mu_i} \frac{\partial}{\partial y_a} + \frac{\partial U}{\partial z_\ell} \Big|_{\mu_i} \frac{\partial}{\partial U}. \quad (3.15)$$

Denoting $\{y_a\} = \{y, \{y_b\}\}$, we obtain the KZ equations in Sklyanin variables,

$$\left(\frac{1}{k-2} \frac{\partial^2}{\partial y^2} + \sum_{\ell=1}^n \frac{1}{y-z_\ell} \left(\frac{\partial}{\partial z_\ell} + \frac{\partial}{\partial y} \right) + \sum_b \frac{1}{y-y_b} \left(\frac{\partial}{\partial y_b} - \frac{\partial}{\partial y} \right) + \sum_{\ell=1}^n \frac{\Delta_{j_\ell}^J}{(y-z_\ell)^2} \right) \mathcal{K}^{-1} \Omega_n = 0 . \quad (3.16)$$

In this equation the variables are no longer separated, as the variables y_b appear in addition to y .

3.3 Comparison with Virasoro null-vector equations

In the previous subsection, we have studied the KZ equations in a CFT with an \widehat{sl}_2 symmetry algebra at level k . We will now compare them with null-vector equations in a CFT with a Virasoro symmetry algebra at central charge $c = 1 + 6(b + b^{-1})^2$ where $b^2 \equiv \frac{1}{k-2}$. This is the Virasoro algebra which would be obtained from our \widehat{sl}_2 algebra by quantum Hamiltonian reduction (see for instance [11]), although that reduction does not explain the relation between differential equations which we are about to review.

The Virasoro algebra can be formulated in terms of the stress-energy tensor $T(z)$, which obeys

$$T(z)T(w) = \frac{\frac{1}{2}c}{(z-w)^4} + \frac{2T(w)}{(z-w)^2} + \frac{\partial T(w)}{z-w} + \mathcal{O}(1) . \quad (3.17)$$

Primary fields $V_\alpha(w)$ of momentum α and conformal dimension $\Delta_\alpha = \alpha(b + b^{-1} - \alpha)$ are defined by

$$T(z)V_\alpha(w) = \frac{\Delta_\alpha V_\alpha(w)}{(z-w)^2} + \frac{\partial V_\alpha(w)}{z-w} + \mathcal{O}(1) . \quad (3.18)$$

This definition does not distinguish the primary fields V_α and $V_{b+b^{-1}-\alpha}$, which have the same conformal dimension. These fields are therefore assumed to be proportional, with a proportionality constant called the reflection coefficient. This \mathbb{Z}_2 symmetry can be understood as the action of the Weyl group of sl_2 on the space of the momenta α .

The Virasoro representation generated by the degenerate field $V_{-\frac{1}{2b}}$ is known to have a null-vector at level two Namely, $(L_{-2} + b^2 L_{-1}^2)V_{-\frac{1}{2b}} = 0$, where the modes L_p are defined as in eq. (2.7). This implies that correlation functions involving such a degenerate field obey the Belavin-Polyakov-Zamolodchikov equation [12]

$$\left[b^2 \frac{\partial^2}{\partial y^2} + \sum_{i=1}^n \frac{1}{y-z_i} \frac{\partial}{\partial z_i} + \sum_{i=1}^n \frac{\Delta_{\alpha_i}}{(y-z_i)^2} \right] \left\langle V_{-\frac{1}{2b}}(y) \prod_{i=1}^n V_{\alpha_i}(z_i) \right\rangle = 0 . \quad (3.19)$$

Curiously, this equation is formally identical to the variable-separated KZ equation (3.14). The meaning of this formal similarity is not clear to us. The KZ equations in Sklyanin

variables (3.16) actually involve $n - 2$ variables $y_1 \cdots y_{n-2}$, therefore we should rather consider correlation functions of the type

$$\tilde{\Omega}_n \equiv \left\langle \prod_{a=1}^{n-2} V_{-\frac{1}{2b}}(y_a) \prod_{i=1}^n V_{\alpha_i}(z_i) \right\rangle . \quad (3.20)$$

We then expect such correlation functions to be related to Ω_n (1.1) as in equation (1.4). That equation means that the twisted BPZ equations satisfied by $\Theta_n \tilde{\Omega}_n$ are identical to the KZ equations in Sklyanin variables (3.16). This can indeed be checked by explicit calculation, provided we correctly specify the function Θ_n as well as the relation between sl_2 spins j_i and Virasoro momenta α_i . Requiring that the $\alpha - j$ relation is compatible with the respective Weyl symmetries $j \rightarrow -j - 1$ and $\alpha \rightarrow b + b^{-1} - \alpha$, and that conformal dimensions $\Delta_j^J = -\frac{j(j+1)}{k-2}$ eq. (2.8) and Δ_α are related by a constant shift, determines the relation

$$\alpha = b(j + 1) + \frac{1}{2b} \quad , \quad \Delta_\alpha = \Delta_j^J + \frac{1}{2} + \frac{1}{4b^2} . \quad (3.21)$$

We still have to specify the values of the parameters λ, μ, ν in the ansatz (1.3) for the function Θ_n . We could determine these values by requiring the twisted BPZ equations to agree with eq. (3.16), and we would find

$$\lambda = \frac{1}{2b^2} \quad , \quad \mu = -\frac{1}{2b^2} \quad , \quad \nu = \frac{1}{2b^2} . \quad (3.22)$$

There are simple concurring arguments for the values of λ and ν . First, the value of λ is determined by the requirement of continuity of $\Theta_n \tilde{\Omega}_n$ at $y_a = y_b$. This requirement plays an important role in the boundary H_3^+ model [4]. Second, the value of ν follows from checking equation (1.4) in the simplest case $n = 2$, when there are no y_a variables and no BPZ equations.

Let us now comment on this twist function Θ_n and its relation to free field correlation functions. In this paragraph we will consider full correlation functions with dependences on both holomorphic and antiholomorphic variables, and the full twist factor which is thus $|\Theta_n|^2$. With the above values (3.22) for λ, μ, ν , we observe that the inverse twist factor $|\Theta_n|^{-2}$ coincides with the free field correlation function formally obtained from $\tilde{\Omega}_n$ by taking the fields $V_{\alpha_i}(z_i)$ to have momenta $\alpha_i = \frac{1}{2b}$ instead of $\alpha_i = b(j_i + 1) + \frac{1}{2b}$. This means

$$|\Theta_n|^{-2} = \left\langle \prod_{a=1}^{n-2} V_{-\frac{1}{2b}}(y_a) \prod_{i=1}^n V_{\frac{1}{2b}}(z_i) \right\rangle^{\text{free}} . \quad (3.23)$$

This interpretation of Θ_n plays a role in a recent proof of the FZZ conjecture [13], see also [14]. For now, let us explain the origin of this observed relation by studying the $b \rightarrow 0$ limit of the H_3^+ -Liouville relation. This relation can be written as $\Omega_n \sim \mathcal{K}\bar{\mathcal{K}}|\Theta_n|^2\tilde{\Omega}_n$, whose factors we now analyze:

- The Liouville correlation function $\tilde{\Omega}_n$ reduces to $\left\langle \prod_{a=1}^{n-2} V_{-\frac{1}{2b}}(y_a) \prod_{i=1}^n V_{\frac{1}{2b}}(z_i) \right\rangle$ as $b \rightarrow 0$. And it turns out that this coincides with a free field correlation function, be-

cause the momentum conservation condition is obeyed. Namely, the sum of the momenta is $(n-2) \times -\frac{1}{2b} + n \times \frac{1}{2b} = \frac{1}{b}$ which coincides with the dominant term in the Liouville background charge $\frac{1}{b} + b$. Therefore, according to standard path-integral reasoning in Liouville theory [15], we have $\tilde{\Omega}_n \underset{b \rightarrow 0}{\sim} R_n \left\langle \prod_{a=1}^{n-2} V_{-\frac{1}{2b}}(y_a) \prod_{i=1}^n V_{\frac{1}{2b}}(z_i) \right\rangle^{\text{free}} = R_n \left| \frac{\prod(y_a - z_i)}{\prod(y_a - y_b) \prod(z_i - z_j)} \right|^{\frac{1}{b^2}}$ where R_n is b -independent.

- The H_3^+ correlation function Ω_n is expected to have a finite “minisuperspace” limit [16] as $b \rightarrow 0$ which is equivalent to $k \rightarrow \infty$ where k is the level.
- The separation of variables \mathcal{K} is b -independent by definition.
- So the twist factor $|\Theta_n|^2$ must absorb the $b \rightarrow 0$ divergence of the Liouville correlation function $\tilde{\Omega}_n$, which implies the relation (3.23) and the values (3.22) for the parameters λ, μ, ν . (This reasoning does not exclude the presence of extra terms in λ, μ, ν which would be finite in the $b \rightarrow 0$ limit.)

This concludes our reminder of the KZ-BPZ relation in the sl_2 case. In the next section we will analyze the sl_3 KZ equations along the same lines.

4 The sl_3 case

4.1 Separation of variables for the sl_3 Gaudin model

To the best of our knowledge, the full quantum separation of variables for the sl_3 Gaudin model has not been derived yet. By the full separation of variables we mean the determination of $A(u)$, $B(u)$ and a characteristic equation, like in the sl_2 case.¹ Sklyanin did however derive the full separation of variables for the classical sl_3 Gaudin model [18]. In order to derive the quantum version, we will use Sklyanin’s separation of variables for models with an sl_3 Yangian symmetry [8], see also [19] for a generalization to sl_N . This Yangian symmetry is present in the Gaudin model, which will allow us to derive its quantum characteristic equation from the Yangian’s.

sl_3 Yangian symmetry. As in the sl_2 case, the variables of the sl_N Gaudin model can be combined into an sl_N Lax matrix $I(u)$ (3.1) obeying the relation (3.2). It is however possible to combine the variables into another sl_N matrix, which depends on an extra parameter η ,

$$Y(u) \equiv \left(\text{id} - \frac{\eta}{u - z_1} t^a D_{(1)}^a \right) \left(\text{id} - \frac{\eta}{u - z_2} t^a D_{(2)}^a \right) \cdots \left(\text{id} - \frac{\eta}{u - z_n} t^a D_{(n)}^a \right) \quad (4.1)$$

$$= \text{id} + \eta I(u) + \frac{1}{2} \eta^2 : I^2 : (u) + \frac{1}{6} \eta^3 : I^3 : (u) + \cdots, \quad (4.2)$$

where the definition of the normal ordering in $: I^2 : (u)$ and $: I^3 : (u)$ follows from the chosen ordering of the factors of $Y(u)$. This object can be shown to obey the Yangian

¹A different approach was proposed in [17], which consists in trying to use the sl_2 separation of variables in the sl_3 case. This approach requires a particular choice of isospin variables. The results are complicated.

algebra

$$(u - v)Y_\alpha^\gamma(u)Y_\beta^\epsilon(v) + \eta Y_\alpha^\epsilon(u)Y_\beta^\gamma(v) = (u - v)Y_\beta^\epsilon(v)Y_\alpha^\gamma(u) + \eta Y_\alpha^\epsilon(v)Y_\beta^\gamma(u). \quad (4.3)$$

Sklyanin's separated variables y_ℓ for the Yangian [8] are defined as the zeroes of a function

$$B^Y(u) = Y_3^2(u)Y_2^1(u)Y_3^2(u - \eta) - Y_3^2(u)Y_3^1(u)Y_2^2(u - \eta) + Y_3^1(u)Y_3^2(u)Y_1^1(u - \eta) - Y_3^1(u)Y_1^2(u)Y_3^1(u - \eta), \quad (4.4)$$

while the conjugate variables are given by $X_i = A^Y(y_i)$ where

$$A^Y(u) = Y_1^1(u) - Y_3^2(u - \eta)^{-1}Y_3^1(u - \eta)Y_1^2(u), \quad (4.5)$$

Let us point out that interesting structural insight into these formulas for $A^Y(u)$ and $B^Y(u)$ was obtained in [20], based on general properties of matrices with non-commuting elements. The functions $A^Y(u)$ and $B^Y(u)$ obey the commutation relations

$$[A^Y(u), A^Y(v)] = 0, \quad [B^Y(u), B^Y(v)] = 0, \quad (4.6)$$

$$\frac{u - v}{\eta} [A^Y(u), B^Y(v)] = B^Y(u)A^Y(v) Y_3^2(u - \eta)^{-1}Y_3^2(u)^{-1} \cdot Y_3^2(v - \eta)Y_3^2(v) - B^Y(v)A^Y(u), \quad (4.7)$$

so that

$$[y_i, y_j] = 0 \quad , \quad [X_i, y_j] = -\eta\delta_{ij}X_i \quad , \quad [X_i, X_j] = 0. \quad (4.8)$$

The quantum characteristic equation is then

$$X_i^3 - X_i^2 t_1(y_i) + X_i t_2(y_i - \eta) - d(y_i - 2\eta) = 0, \quad (4.9)$$

with the invariant operators $t_1(u)$, $t_2(u)$ and $d(u)$ defined as [8]

$$t_1(u) = \text{Tr } Y(u) \quad , \quad t_2(u) = \text{Tr } \tilde{Y}(u) \quad , \quad d(u)\delta_\alpha^\gamma = Y_\alpha^\beta(u)\tilde{Y}_\beta^\gamma(u + \eta), \quad (4.10)$$

where the matrix \tilde{Y} is constructed by transposing the quantum comatrix of Y . For instance, $\tilde{Y}_3^2(u) = -Y_3^2(u)Y_1^1(u + \eta) + Y_3^1(u)Y_1^2(u + \eta)$, where the η -shifts are the manifestation of the quantum character of the comatrix whose $\frac{3}{2}$ matrix element we just wrote. Operator ordering issues in expressions like $t_2(y_i - \eta)$ are resolved by inserting the operator y_i from the left.

From the Yangian to the Gaudin model. We will now construct objects $A(u)$, $B(u)$ and a quantum characteristic equation for the sl_3 Gaudin model. Such η -independent functions of the matrix $I(u)$ will be obtained by expanding the corresponding objects for the sl_3 Yangian algebra in powers of η . We find

$$A^Y(u) = 1 - \eta A(u) + \mathcal{O}(\eta^2), \quad A(u) = -I_1^1 + \frac{I_3^1 I_1^2}{I_3^2}, \quad (4.11)$$

$$B^Y(u) = \eta^3 B(u) + \mathcal{O}(\eta^4), \quad B(u) = I_2^1 I_3^2 I_3^2 - I_3^2 I_3^1 I_2^2 + I_3^1 I_3^2 I_1^1 - I_1^2 I_3^1 I_3^1, \quad (4.12)$$

where we omitted the spectral parameter u in $I_\alpha^\beta(u)$, and we point out that our formula for $A(u)$ is free of ordering ambiguities because $I_3^2(u)$ commutes with both $I_1^2(u)$ and $I_3^1(u)$. The commutation relations (4.6) for $A^Y(u)$ and $B^Y(u)$ imply the analogous relations

$$[A(u), A(v)] = 0 \quad , \quad [B(u), B(v)] = 0, \quad (4.13)$$

$$(u-v)[A(u), B(v)] = B(v) - B(u) \frac{I_3^2(v)I_3^2(v)}{I_3^2(u)I_3^2(u)}, \quad (4.14)$$

which may be compared to the corresponding relations in the sl_2 case eq. (3.6).

Let us rewrite the characteristic equation (4.9) as:

$$(X_i - 1)^3 - (X_i - 1)^2 [t_1(y_i) - 3] + (X_i - 1) [t_2(y_i - \eta) - 2t_1(y_i) + 3] + [1 - t_1(y_i) + t_2(y_i - \eta) - d(y_i - 2\eta)] = 0. \quad (4.15)$$

The leading behaviour of this equation as $\eta \rightarrow 0$ will turn out to be $\mathcal{O}(\eta^3)$. To compute this behaviour, we of course need to compute the behaviours of X_i and y_i as $\eta \rightarrow 0$. It turns out that we only need the $\mathcal{O}(\eta)$ behaviour of X_i . We therefore define the variable p_i by $X_i = 1 - \eta p_i + \mathcal{O}(\eta^2)$. As for y_i we only need need the leading $\mathcal{O}(1)$ behaviour. To this leading order, the zeroes of $B^Y(u)$ coincide with those of $B(u)$, so that we do not need distinct notations and call them all y_i . The most complicated part of the calculation however does not involve such subtleties, but rather deals with the last term in eq. (4.15),

$$\begin{aligned} & 1 - t_1(u) + t_2(u - \eta) - d(u - 2\eta) \quad (4.16) \\ &= (1 - Y_1^1(u - 2\eta))(Y_3^3(u - \eta) - 1)Y_2^2(u) + (Y_1^1(u - \eta) - Y_1^1(u - 2\eta))Y_2^2(u) \\ & \quad + (Y_1^1(u - 2\eta) - 1)Y_3^3(u - \eta)Y_2^2(u) + (Y_1^1(u - \eta) - 1)Y_3^3(u) + (1 - Y_1^1(u)) \\ & \quad + (Y_3^3(u - 2\eta) - 1)Y_1^1(u - \eta)Y_2^2(u) - Y_3^2(u - 2\eta)Y_1^3(u - \eta)Y_2^1(u) \\ & \quad - Y_1^2(u - 2\eta)Y_3^1(u - \eta)Y_2^3(u) + Y_1^3(u - 2\eta)Y_3^1(u - \eta)Y_2^2(u) - Y_1^3(u - \eta)Y_3^1(u) \\ &= \eta^3 [-I_1^1 I_3^3 I_2^2 + I_1^1 I_3^2 I_2^3 + I_3^3 I_1^2 I_2^1 + I_2^2 I_1^3 I_3^1 - I_3^2 I_1^3 I_2^1 - I_1^2 I_3^3 I_2^3 \\ & \quad + I_1^1 (I')_3^3 - I_1^1 (I')_1^1 - I_1^3 (I')_3^1 - (I')_1^3 I_3^1 - (I'')_1^1] + \mathcal{O}(\eta^4), \end{aligned}$$

where we omitted the spectral parameter u in $I_\alpha^\beta(u)$, and used the sl_3 -defining relation $I_1^1 + I_2^2 + I_3^3 = 0$. We then obtain the following quantum characteristic equation of the sl_3 Gaudin model:

$$p_i^3 - p_i \cdot \frac{1}{2} (I_\alpha^\beta I_\beta^\alpha)(y_i) + \frac{1}{4} (I_\alpha^\beta I_\beta^\alpha)'(y_i) + \frac{1}{6} \left(I_\alpha^\beta I_\beta^\gamma I_\gamma^\alpha + I_\beta^\alpha I_\gamma^\beta I_\alpha^\gamma \right) (y_i) = 0. \quad (4.17)$$

Notice that the particular cubic invariant which appears in this formula is related to the fully symmetric invariant tensor d_{abc} eq. (2.10). Using the definition (3.1) of $I(u)$, we indeed have

$$\left(I_\alpha^\beta I_\beta^\gamma I_\gamma^\alpha + I_\beta^\alpha I_\gamma^\beta I_\alpha^\gamma \right) (u) = -d_{abc} \sum_{i=1}^n \frac{D_{(i)}^a}{u - z_i} \sum_{\ell=1}^n \frac{D_{(\ell)}^b}{u - z_\ell} \sum_{m=1}^n \frac{D_{(m)}^c}{u - z_m}. \quad (4.18)$$

This could further be expressed in terms of the higher Gaudin Hamiltonians of section 2.2, so that the characteristic equation could help simultaneously diagonalize these Hamiltonians.

Some remarks. Like in the sl_2 case, Sklyanin's change of variables can be interpreted as an integral transformation \mathcal{K} (3.9) acting on a functional space. The kernel K of \mathcal{K} now obeys

$$\left(B(u) - U \frac{\prod_{\ell}(u - y_{\ell})}{\prod_i(u - z_i)^3} \right) K(\{x_i\}|\{y_{\ell}\}, U) = 0 . \tag{4.19}$$

However, the simultaneous diagonalization of the commuting operators $B(u)$ is now a difficult problem, as $B(u)$ is now cubic and not linear in $I(u)$, and thus no longer a sum of n commuting operators. Therefore, the kernel K is no longer of the form (3.13). Certainly, no choice of isospin variables exists such that the kernel K has a simple expression. Another difference with the sl_2 case is the counting of variables: generic functions of the sl_3 isospin coordinates x_i should correspond to functions of not only y_i and U , but also of two extra variables. These extra variables are necessary for the transformation \mathcal{K} to be invertible. We will neglect this issue,² as well as the issue of precisely defining the relevant functional spaces, and we will assume \mathcal{K} to be invertible.

Let us finally determine the number of separated variables y_i – that is, the number of zeroes of $B(u)$. Barring extra constraints, this is of course $3n - 3$. In conformal field theory applications, we however impose the extra constraints $\sum_{i=1}^n D_{(i)}^a = 0$, so that $I(u)$ has degree -2 . This does not immediately imply that $B(u)$ (eq. (4.12)), which is cubic in $I(u)$, has degree -6 , because $\sum_{i=1}^n D_{(i)}^a = 0$ only holds when directly applied to a physical correlation function, and the matrix elements of $I(u)$ generically do not commute with each other. Rather, the degree of $B(u)$ depends on its precise form and should be evaluated by explicit calculation. We find that each one of the four terms of $B(u)$ has degree -5 , while $B(u)$ itself has degree -6 . This means that there are $3n - 6$ separated variables. Therefore, as in the sl_2 case, the number of separated variables vanishes for $n = 2$.

4.2 The sl_3 Knizhnik-Zamolodchikov equations in Sklyanin variables

Let us consider a conformal field theory with an $\widehat{sl_3}$ symmetry algebra. The Ward identities consist in the n KZ differential equations (2.9), plus $2n$ extra non-differential relations (2.15) and (2.16), which express $W_{-1,(i)}^J$ and $W_{-2,(i)}^J$ in terms of differential operators acting on isospin variables. Let us reorganize all these relations by injecting them into the characteristic equation of the quantum sl_3 Gaudin model (4.17). The result is schematically of the form

$$\left[\frac{\partial^3}{\partial y^3} + (k - 3) \frac{\partial}{\partial y} \cdot T^J(y) - \frac{1}{2}(k - 3) \partial T^J(y) - \frac{1}{\rho} W^J(y) \right] \mathcal{K}^{-1} \Omega_n = 0, \tag{4.20}$$

²A construction of the extra variables seems to be available in the article [21].

where the constant ρ was defined in eq. (2.11). Explicitly,

$$\left[\frac{\partial^3}{\partial y^3} + (k-3) \frac{\partial}{\partial y} \cdot \sum_{i=1}^n \left(\frac{1}{y-z_i} \mathcal{K}^{-1} \frac{\delta}{\delta z_i} \mathcal{K} + \frac{\Delta_{j_i}^J}{(y-z_i)^2} \right) + \frac{1}{2} (k-3) \sum_{i=1}^n \left(\frac{1}{(y-z_i)^2} \mathcal{K}^{-1} \frac{\delta}{\delta z_i} \mathcal{K} + \frac{2\Delta_{j_i}^J}{(y-z_i)^3} \right) - \frac{1}{\rho} \sum_{i=1}^n \left(\frac{\mathcal{K}^{-1} W_{-2,(i)}^J \mathcal{K}}{y-z_i} + \frac{\mathcal{K}^{-1} W_{-1,(i)}^J \mathcal{K}}{(y-z_i)^2} + \frac{q_{j_i}^J}{(y-z_i)^3} \right) \right] \mathcal{K}^{-1} \Omega_n = 0, \quad (4.21)$$

where Ω_n is still an n -point function of the type (1.1).

In this equation, the terms involving $W_{-1,(i)}^J$ and $W_{-2,(i)}^J$ refer to correlation functions involving descendents of the primary fields $\Phi^j(\mu|z)$. We have little control over such non-differential terms, and we would like to ignore them in the following. This could be done by considering appropriate linear combinations of our $3n - 6$ equations. (Remember that the variable y spans the $3n - 6$ separated variables $\{y_a\}$). We will for simplicity adopt the alternative approach of working modulo the unwanted terms. Let us make this precise by defining the space \mathcal{D}_S of differential operators in y_a, z_i (including functions of y_a, z_i) which are symmetric under permutations of $\{y_1, y_2 \cdots y_{3n-6}\}$. For any choice $\{y_a\} = \{y, y_b\}$ of a distinguished variable y we further define

$$\mathcal{F}_2(y) \equiv \sum_{i=1}^n \frac{1}{y-z_i} \mathcal{D}_S + \sum_{i=1}^n \frac{1}{(y-z_i)^2} \mathcal{D}_S. \quad (4.22)$$

By a simple counting of variables it can be realized that any differential operator which is symmetric under permutations of $\{y_b\}$ does belong to $\mathcal{F}_3(y) \equiv \sum_{i=1}^n \frac{1}{y-z_i} \mathcal{D}_S + \sum_{i=1}^n \frac{1}{(y-z_i)^2} \mathcal{D}_S + \sum_{i=1}^n \frac{1}{(y-z_i)^3} \mathcal{D}_S$. But it does not always belong to $\mathcal{F}_2(y)$, so we can define a nontrivial equivalence \sim as the equality modulo $\mathcal{F}_2(y)$. Thus, equation (4.21) simplifies to

$$\left[\frac{\partial^3}{\partial y^3} + \frac{\partial}{\partial y} \cdot \sum_{i=1}^n \frac{k-3}{y-z_i} \mathcal{K}^{-1} \frac{\delta}{\delta z_i} \mathcal{K} + \sum_{i=1}^n \frac{(k-3)\Delta_{j_i}^J}{(y-z_i)^2} \frac{\partial}{\partial y} - \sum_{i=1}^n \frac{\frac{1}{\rho} q_{j_i}^J + (k-3)\Delta_{j_i}^J}{(y-z_i)^3} \right] \mathcal{K}^{-1} \Omega_n \sim 0 \quad (4.23)$$

Having thus eliminated $W_{-1,(i)}^J$ and $W_{-2,(i)}^J$, we are left with operators $\frac{\delta}{\delta z_i}$, which we recall are z_i -derivatives at fixed isospin variables. We expect $\mathcal{K}^{-1} \frac{\delta}{\delta z_i} \mathcal{K}$ to be a combination of the operators $\frac{\partial}{\partial z_i}$, $\frac{\partial}{\partial y_a}$ and $\frac{\partial}{\partial U}$, although we do not know how to compute it. And it is not clear whether $\mathcal{K}^{-1} \frac{\delta}{\delta z_i} \mathcal{K}$ is a first-order differential operator, as happened in the $s\ell_2$ case (see eq. (3.15)). Nevertheless, we do know that $\mathcal{K}^{-1} \frac{\delta}{\delta z_i} \mathcal{K}$ is independent from the level k , which is a parameter of our conformal field theory but neither of the Gaudin model nor of its separation of variables. Therefore, we will still be able to extract useful information from eq. (4.23), a sum of terms with various power-like dependences on $(k-3)$, by considering all terms which are not linear in $(k-3)$.

4.3 \mathcal{W}_3 null-vector equations

Let us first briefly explain why we try to relate conformal field theories with an \widehat{sl}_3 symmetry at level k to theories with a \mathcal{W}_3 symmetry at central charge $c = 2 + 24(b + b^{-1})^2$ where

$$b^2 = \frac{1}{k-3} . \tag{4.24}$$

A theory with an \widehat{sl}_3 symmetry like the $sl_3(\mathbb{R})$ WZW model can be written in terms of eight quantum fields, as sl_3 is eight-dimensional. However, affine \widehat{sl}_3 highest-weight representations are parametrized by just two numbers, namely the two components of the sl_3 spin j . This suggests that the non-trivial dynamics of the theory really take place in a two-dimensional space, where j would play the role of the momentum. There exists such an sl_3 -based theory which involves just two interacting quantum fields: the conformal sl_3 Toda theory, which has a \mathcal{W}_3 symmetry algebra. The correct parameter b for this algebra is suggested by the Drinfeld-Sokolov reduction, which realizes \mathcal{W}_3 as a kind of coset of the \widehat{sl}_3 algebra.

\mathcal{W}_3 algebra and primary fields. Referring to the review article [11] for more details, we recall that the \mathcal{W}_3 algebra is spanned by the modes of the fields $T(z) = \sum_{n \in \mathbb{Z}} L_n z^{-n-2}$ and $W(z) = \sum_{n \in \mathbb{Z}} W_n z^{-n-3}$. Let us write the defining relations of the \mathcal{W}_3 algebra in the form of commutation relations for the modes L_n, W_n rather than operator product expansions for the fields $T(z), W(z)$, as this form is more convenient for finding null vectors in representations:

$$[L_m, L_n] = (m-n)L_{m+n} + \frac{c}{12}m(m^2-1)\delta_{m+n,0} , \tag{4.25}$$

$$[L_m, W_n] = (2m-n)W_{m+n} , \tag{4.26}$$

$$[W_m, W_n] = \frac{(22+5c)}{48} \frac{c}{360} m(m^2-1)(m^2-4)\delta_{m+n,0} + \frac{(22+5c)}{48} \frac{m-n}{15} \left(m^2+n^2 - \frac{1}{2}mn - 4 \right) L_{m+n} + \frac{1}{3}(m-n)\Lambda_{m+n} , \tag{4.27}$$

where we introduce, using the normal ordering $: L_m L_n := L_m L_n$ if $m \leq n$,

$$\Lambda_m = \sum_{n \in \mathbb{Z}} : L_n L_{m-n} : + \frac{1}{5} x_m L_m \quad \text{with} \quad \begin{cases} x_{2\ell} = (1+\ell)(1-\ell) \\ x_{2\ell+1} = (\ell+2)(1-\ell) \end{cases} . \tag{4.28}$$

A primary fields V_α of the \mathcal{W}_3 algebra of momentum α , conformal dimension Δ_α and charge q_α is defined by its operator product expansions with $T(z)$ eq. (3.18) and $W(z)$:

$$W(z)V_\alpha(w) = \frac{q_\alpha V_\alpha(w)}{(z-w)^3} + \frac{W_{-1}V_\alpha(w)}{(z-w)^2} + \frac{W_{-2}V_\alpha(w)}{z-w} + \mathcal{O}(1) . \tag{4.29}$$

The momenta α now belong to the two-dimensional root space of the Lie algebra sl_3 . A basis of this space is provided by the simple roots e_1, e_2 whose scalar products appear in the Cartan matrix $\begin{pmatrix} (e_1, e_1) & (e_1, e_2) \\ (e_2, e_1) & (e_2, e_2) \end{pmatrix} = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$. We may also use the dual basis $\omega_1 = \frac{2}{3}e_1 + \frac{1}{3}e_2, \omega_2 = \frac{1}{3}e_1 + \frac{2}{3}e_2$ such that $(e_i, \omega_j) = \delta_{ij}$. We decompose the momenta along this

dual basis: $\alpha = \alpha_1\omega_1 + \alpha_2\omega_2$, and we introduce the vector $Q = (b + b^{-1})(e_1 + e_2)$. The conformal dimension and charge are parametrized in terms of the momentum as

$$\Delta_\alpha = \frac{1}{2}(\alpha, 2Q - \alpha), \tag{4.30}$$

$$q_\alpha = \frac{i}{27}[\alpha_1 - \alpha_2][2\alpha_1 + \alpha_2 - 3(b + b^{-1})][\alpha_1 + 2\alpha_2 - 3(b + b^{-1})]. \tag{4.31}$$

\mathcal{W}_3 degenerate fields. Let us now justify the choice of the field $V_{-b^{-1}\omega_1}$ in the correlator $\tilde{\Omega}_n$ (1.2) which appears in our conjecture. We wish $\tilde{\Omega}_n$ to obey third-order differential equations, which would correspond to the sl_3 KZ equations in Sklyanin variables. This suggests that we use the simplest non-trivial degenerate fields, which have null vectors at levels 1, 2 and 3. But there are actually four such degenerate fields, with $\alpha \in \{-b\omega_1, -b\omega_2, -b^{-1}\omega_1, -b^{-1}\omega_2\}$, whereas we want only one of them to appear in $\tilde{\Omega}_n$, because the original isospin variables are invariant under permutations of the Sklyanin variables.

By analogy with the sl_2 case, we focus on the fields $V_{-b^{-1}\omega_1}$ and $V_{-b^{-1}\omega_2}$, whose momenta go to zero in the critical level limit $k \rightarrow 3$. They are related to the other two fields by the \mathcal{W}_3 algebra self-duality $b \rightarrow b^{-1}$, which is however not an invariance of the \widehat{sl}_3 algebra. And they are related to each other by the Dynkin diagram automorphism $\omega_1 \leftrightarrow \omega_2$ of sl_3 , which acts on general primary fields V_α as $(\Delta_\alpha, q_\alpha) \rightarrow (\Delta_\alpha, -q_\alpha)$. This symmetry does have a counterpart in the separation of variables for the sl_3 Gaudin model. The construction of the separated variables was indeed based on the introduction of an sl_3 Lax matrix $I(u)$ (3.1), so that sl_3 generators act in the fundamental representation. But we could alternatively have used the antifundamental representation, which is related to the fundamental by the Dynkin diagram automorphism. With our conventions, our choice of the fundamental representation will turn out to correspond to the choice of the degenerate field $V_{-b^{-1}\omega_1}$ of the \mathcal{W}_3 algebra. The three corresponding null-vector equations are [22]

$$\left[iW_{-1} + \left(\frac{b}{2} + \frac{5}{6b} \right) L_{-1} \right] V_{-b^{-1}\omega_1} = 0, \tag{4.32}$$

$$\left[iW_{-2} - \frac{2}{3b}L_{-2} - bL_{-1}^2 \right] V_{-b^{-1}\omega_1} = 0, \tag{4.33}$$

$$\left[iW_{-3} - \left(\frac{b}{2} + \frac{1}{6b} \right) L_{-3} + bL_{-1}L_{-2} + b^3L_{-1}^3 \right] V_{-b^{-1}\omega_1} = 0. \tag{4.34}$$

The last null-vector equation implies that any correlation function with one degenerate field obeys $E_1 \langle V_{-b^{-1}\omega_1}(y) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle = 0$, where

$$\begin{aligned} E_1 \equiv & \frac{\partial^3}{\partial y^3} + \frac{1}{b^2} \frac{\partial}{\partial y} \cdot \sum_{i=1}^n \left(\frac{1}{y - z_i} \frac{\partial}{\partial z_i} + \frac{\Delta_{\alpha_i}}{(y - z_i)^2} \right) \\ & + \left(\frac{1}{2b^2} + \frac{1}{6b^4} \right) \sum_{i=1}^n \left(\frac{1}{(y - z_i)^2} \frac{\partial}{\partial z_i} + \frac{2\Delta_{\alpha_i}}{(y - z_i)^3} \right) \\ & + \frac{i}{b^3} \sum_{i=1}^n \left(\frac{W_{-2,(i)}}{y - z_i} + \frac{W_{-1,(i)}}{(y - z_i)^2} + \frac{q_{\alpha_i}}{(y - z_i)^3} \right). \end{aligned} \tag{4.35}$$

This may be compared with eq. (4.21), which is formally similar, or even identical if the term with coefficient $\frac{1}{6b^4}$ is absorbed into the other terms by redefining $W_{-1,(i)}$ and q_{α_i} . Like in the $s\ell_2$ case, the meaning of this formal similarity is not clear.

Now the equations obeyed by correlation functions with several degenerate fields like $\tilde{\Omega}_n$ eq. (1.2) are significantly more complicated than E_1 , because eliminating W_{-1}, W_{-2} descendants of the degenerate fields requires the use of the first two null-vector equations (4.32), (4.33). Still denoting $\{y_a\} = \{y, y_b\}$, we obtain the equation $E_2\tilde{\Omega}_n = 0$ with

$$\begin{aligned}
 E_2 \equiv E_1 &+ \frac{1}{b^2} \sum_b \frac{1}{y-y_b} \frac{\partial^2}{\partial y_b^2} + \frac{1}{b^2} \frac{\partial}{\partial y} \cdot \sum_b \left(\frac{1}{y-y_b} \frac{\partial}{\partial y_b} + \frac{\Delta_{-b^{-1}\omega_1}}{(y-y_b)^2} \right) \\
 &+ \frac{2}{3b^4} \sum_{b,i} \frac{1}{(y-y_b)(y_b-z_i)} \left(\frac{\partial}{\partial z_i} + \frac{\Delta_{\alpha_i}}{y_b-z_i} \right) \\
 &+ \frac{2}{3b^4} \sum_{b \neq c} \frac{1}{(y-y_b)(y_b-y_c)} \left(\frac{\partial}{\partial y_c} + \frac{\Delta_{-b^{-1}\omega_1}}{y_b-y_c} \right) \\
 &- \frac{2}{3b^4} \sum_b \frac{1}{(y-y_b)^2} \left(\frac{\partial}{\partial y_b} + \frac{\partial}{\partial y} \right) \\
 &\quad + \left(\left(\frac{1}{b^2} + \frac{1}{b^4} \right) \Delta_{-b^{-1}\omega_1} + \frac{i}{b^3} q_{-b^{-1}\omega_1} \right) \sum_b \frac{1}{(y-y_b)^3},
 \end{aligned} \tag{4.36}$$

where

$$\Delta_{-b^{-1}\omega_1} = -1 - \frac{4}{3b^2}, \quad q_{-b^{-1}\omega_1} = -\frac{i}{27b^3}(4+3b^2)(5+3b^2). \tag{4.37}$$

Relating \mathcal{W}_3 momenta to $\widehat{s\ell_3}$ spins. In order to compare the equation $E_2\tilde{\Omega}_n = 0$ with the KZ equations in Sklyanin variables (4.23), we should specify how we relate $\widehat{s\ell_3}$ primary fields $\Phi^j(\mu|z)$ to \mathcal{W}_3 primary fields $V_\alpha(z)$. We are looking for a relation between α and j which translates into a simple relation between $(\Delta_\alpha, q_\alpha)$ and (Δ_j^J, q_j^J) . We propose

$$\alpha = -bj + b^{-1}(e_1 + e_2) \Rightarrow \begin{cases} \Delta_\alpha = \Delta_j^J + 2 + b^{-2} \\ q_\alpha = q_j^J \end{cases}, \tag{4.38}$$

where we use the following expressions for (Δ_j^J, q_j^J) defined in eqs. (2.8) and (2.14)

$$\Delta_j^J = -\frac{1}{k-3} \frac{1}{2} (j, j + 2e_1 + 2e_2), \tag{4.39}$$

$$q_j^J = \frac{1}{(k-3)^{\frac{3}{2}}} \frac{i}{27} [j_1 - j_2] [2(j_1 + 1) + (j_2 + 1)] [(j_1 + 1) + 2(j_2 + 1)], \tag{4.40}$$

where the components (j_1, j_2) of the spin j are defined as $j = j_1\omega_1 + j_2\omega_2$. Notice that our relation between α and j maps the principal unitary series of $s\ell_3$ representations $j \in -e_1 - e_2 + i\mathbb{R}^2$ to the \mathcal{W}_3 representations which appear in the physical spectrum of conformal $s\ell_3$ Toda theory [6] $\alpha \in Q + i\mathbb{R}^2$. Such choices of α or j lead to real values of (Δ, q) if $k > 3$.

However, there does not need to be any relation between the \widehat{sl}_3 creation operators W_{-1}^J, W_{-2}^J and their \mathcal{W}_3 counterparts W_{-1}, W_{-2} . While relating $L_{-1}^J = \frac{\delta}{\delta z}$ to $L_{-1} = \frac{\partial}{\partial z}$, though difficult in practice, is in principle a simple matter of performing the change of variables, there is apparently no principle which would determine how W_{-1}^J, W_{-2}^J would behave through the change of variables. This is why we work modulo $\mathcal{F}_2(y)$, ignoring the non-differential terms which involve such operators, and being left with differential equations. Now the presence of degenerate fields in correlation functions of \mathcal{W}_3 fields does not necessarily lead to differential equations, a fact which makes conformal sl_3 Toda theory much more complicated than Liouville theory [6]. Differential equations actually appear provided the number of degenerate fields is large enough. We are inserting $3n-6$ degenerate fields $V_{-b^{-1}\omega_1}$ together with the n generic fields V_{α_i} , which is enough for eliminating the $2n$ terms $W_{-1,(i)}, W_{-2,(i)}$ and being left with $n-6$ differential equations.

Twisting \mathcal{W}_3 null-vector equations. Finally, we should determine the twist factor Θ_n which appears in the conjecture (1.4), so as to be able to compute

$$E_3 \equiv \Theta_n E_2 \Theta_n^{-1} \quad \text{such that} \quad E_3 \cdot \Theta_n \tilde{\Omega}_n = 0 . \quad (4.41)$$

The values of the parameters λ, ν can be derived as in the sl_2 case. Requiring continuity of $\Theta_n \tilde{\Omega}_n$ at $y_a = y_b$ implies $\lambda = 2\Delta_{-b^{-1}\omega_1} - \Delta_{-2b^{-1}\omega_1} = \frac{2}{3b^2}$, and requiring that the conjecture (1.4) holds in the case $n = 2$ implies $\nu = 2\Delta_\alpha - 2\Delta_j^J = \frac{2}{b^2} + 4$, see eq. (4.38). Notice however that this only determines ν up to b -independent terms, as the unknown b -independent kernel K may also contribute.

These constraints leave the parameter μ arbitrary. We will obtain an ansatz for μ , and confirm the values of λ and ν , by generalizing the relation (3.23) between Θ_n and free field correlation functions which was observed in the sl_2 case. In the sl_3 case the analogous relation is

$$|\Theta_n|^{-2} = \left\langle \prod_{a=1}^{3n-6} V_{-b^{-1}\omega_1}(y_a) \prod_{i=1}^n V_{b^{-1}(e_1+e_2)}(z_i) \right\rangle^{\text{free}} . \quad (4.42)$$

This ansatz leads to the values

$$\lambda = \frac{2}{3b^2} \quad , \quad \mu = -\frac{1}{b^2} \quad , \quad \nu = \frac{2}{b^2} . \quad (4.43)$$

These values will turn out to be the only ones such that, modulo $\mathcal{F}_2(y)$, the only non-differential terms in E_3 are of the type $\frac{c_i}{(y-z_i)^3}$. This is a rather non-trivial requirement as many non-differential terms can potentially appear (cf appendix A.1). Working modulo $\mathcal{F}_2(y)$ eq. (4.22), and using the relation (4.38) between \widehat{sl}_3 and \mathcal{W}_3 representation data, we indeed compute

$$E_3 \sim \frac{\partial^3}{\partial y^3} + \frac{1}{b^2} D_2 + \frac{1}{b^4} D_1 + \frac{1}{b^2} \sum_{i=1}^n \frac{\Delta_{j_i}^J}{(y-z_i)^2} \frac{\partial}{\partial y} + \sum_{i=1}^n \frac{\frac{i}{b^3} q_{j_i}^J - \frac{1}{b^2} \Delta_{j_i}^J}{(y-z_i)^3} , \quad (4.44)$$

where we introduced two differential operators D_1 and D_2 of respective orders 1 and 2, which depend neither on the field momenta α_i nor on the model parameter b ,

$$D_1 \equiv - \sum_i \frac{1}{(y - z_i)^2} \frac{\partial}{\partial y} + 2 \left(\sum_i \frac{1}{y - z_i} \right)^2 \frac{\partial}{\partial y} + 3 \sum_i \frac{1}{y - z_i} \sum_b \frac{1}{y - y_b} \left(\frac{\partial}{\partial y_b} - \frac{\partial}{\partial y} \right) - 2 \sum_{b \neq c} \frac{1}{y - y_b} \frac{1}{y_b - y_c} \left(\frac{\partial}{\partial y_b} - \frac{\partial}{\partial y} \right), \quad (4.45)$$

$$D_2 \equiv \sum_i \frac{1}{y - z_i} \frac{\partial}{\partial y} \left(\frac{\partial}{\partial z_i} + 3 \frac{\partial}{\partial y} \right) + \sum_b \frac{1}{y - y_b} \left(\frac{\partial}{\partial y_b} - \frac{\partial}{\partial y} \right) \left(\frac{\partial}{\partial y_b} + 2 \frac{\partial}{\partial y} \right) + \sum_b \frac{1}{(y - y_b)^2} \frac{\partial}{\partial y}. \quad (4.46)$$

4.4 Comparing sl_3 Knizhnik-Zamolodchikov equations with \mathcal{W}_3 null-vector equations

We are now in a position to test the conjecture (1.4) by comparing the KZ equations in Sklyanin variables (4.23), which apply to $\mathcal{K}^{-1}\Omega_n$, with the twisted \mathcal{W}_3 null-vector equations (4.44), which apply to $\Theta_n\tilde{\Omega}_n$. We will first do the comparison for general values of b , and then explain in more detail what happens in the particular limits $b \rightarrow \infty$ and $b \rightarrow 0$.

The comparison for general b . To start with, the non-differential terms agree. This is actually a very non-trivial statement, as we started with complicated non-differential terms in eq. (4.37) and then generated more terms by twisting with Θ_n . The freedoms to choose the three parameters λ, μ, ν of Θ_n and to ignore terms belonging to $\mathcal{F}_2(y)$ is a priori not sufficient to ensure the dozens of required cancellations, which nevertheless occur as can be seen in explicit calculations. These calculations use some helpful identities which are gathered in appendix A.1. The existence of a simple twist which simplifies the differential equations obeyed by correlation functions involving many identical degenerate fields might well be a general phenomenon in conformal field theory, as we now see that it happens for the simplest degenerate field in theories with \mathcal{W}_3 symmetry, in addition to the already known cases of the two simplest degenerate fields in theories with Virasoro symmetry [3, 5].

Let us then examine the term $\frac{1}{b^2}D_2$ in eq. (4.44). Agreement with the corresponding term in eq. (4.23) would occur provided

$$\frac{\partial}{\partial y} \cdot \sum_i \frac{1}{y - z_i} \mathcal{K}^{-1} \frac{\delta}{\delta z_i} \mathcal{K} \stackrel{?}{\sim} D_2. \quad (4.47)$$

It seems technically challenging to check this identity. But remember that our inability to explicitly perform Sklyanin's change of variables for $\frac{\delta}{\delta z_i}$ does not contaminate the other terms in our equations, as we do know that the change of variables must be independent from the parameter $b = (k - 3)^{-\frac{1}{2}}$.

Let us now examine the term $\frac{1}{b^4}D_1$. We would like this term to vanish modulo $\mathcal{F}_2(y)$, as no such term is present in eq. (4.23). However, it is rather obvious that D_1 does not belong to $\mathcal{F}_2(y)$, although it has quite a few remarkable properties. This is explained in detail in the appendix A.3. As a result, the conjecture cannot hold for general values of b .

The critical level limit $b \rightarrow \infty$. We notice that the term $\frac{1}{b^4}D_1$, which is responsible for the failure of our conjecture, vanishes in the $b \rightarrow \infty$ limit. Therefore, the conjecture has better chances to hold in that limit. To completely prove that it does, we still need to clear one subtlety with the term $\frac{1}{b^2}D_2$. This term seems to vanish in the $b \rightarrow \infty$ limit but actually it does not. This is because near $b \rightarrow \infty$ our correlation functions do not have finite limits. Rather, the Toda correlation function $\tilde{\Omega}_n = \langle \prod_a V_{-b^{-1}\omega_1}(y_a) \prod_i V_{-bj_i+b^{-1}(e_1+e_2)}(z_i) \rangle$ involves “heavy” fields $V_{-bj_i+b^{-1}(e_1+e_2)}(z_i)$ whose momenta grow as b . On general grounds (see for instance [6]), it is therefore expected that $\tilde{\Omega}_n \underset{b \rightarrow \infty}{\sim} e^{b^2 S(\{z_i\})} T_n$ where S and T_n are b -independent functions, and S depends only on $\{z_i\}$ and not on $\{y_a\}$. The differential operator $\frac{1}{b^2}D_2$, which contains derivatives with respect to z_i , may yield a finite contribution when such derivatives act on $e^{b^2 S(\{z_i\})}$.

We should therefore check whether eq. (4.47) holds to the leading order in b^2 when acting on functions of the type $e^{b^2 S(\{z_i\})} T_n$. This is actually the case, because the only term in D_2 with z_i -derivatives is $\frac{\partial}{\partial y} \cdot \sum_i \frac{1}{y-z_i} \frac{\partial}{\partial z_i}$, and $\mathcal{K}^{-1} \frac{\delta}{\delta z_i} \mathcal{K} S(\{z_i\}) = \frac{\partial}{\partial z_i} S(\{z_i\})$. This completes the proof of the conjecture (1.4) in the critical level limit $b \rightarrow \infty \Leftrightarrow k \rightarrow 3$.

Notice that this $b \rightarrow \infty$ limit is not sensitive to the twist function Θ_n . This is because the exponents λ, μ, ν (4.43) vanish in this limit so that $\Theta_n \underset{b \rightarrow \infty}{\rightarrow} 1$.

The minisuperspace limit $b \rightarrow 0$. In this limit, the discrepant term $\frac{1}{b^4}D_1$, which is responsible for the failure of the conjecture (1.4) for general b , grows larger. We may therefore obtain some insights on the reasons for this failure.

As in the sl_2 case, we will consider full correlation functions (with both holomorphic and antiholomorphic dependences) and use path-integral reasonings in sl_3 Toda theory. For full correlation functions, the conjecture reads $\Omega_n \sim \mathcal{K} \bar{\mathcal{K}} |\Theta_n|^2 \tilde{\Omega}_n$. As in the sl_2 case, the transformation \mathcal{K} is b -independent, Ω_n is expected to have a finite limit, and the Toda correlation function $\tilde{\Omega}_n$ behaves as $\tilde{\Omega}_n \underset{b \rightarrow 0}{\sim} R_n \langle \prod_{a=1}^{3n-6} V_{-b^{-1}\omega_1}(y_a) \prod_{i=1}^n V_{b^{-1}(e_1+e_2)}(z_i) \rangle$ where R_n is b -independent.

Therefore $\tilde{\Omega}_n$ simplifies in the $b \rightarrow 0$ limit but, in contrast to the sl_2 case, its leading behaviour does not reduce to a free field correlation function. This is because the simplified correlation function $\langle \prod_{a=1}^{3n-6} V_{-b^{-1}\omega_1}(y_a) \prod_{i=1}^n V_{b^{-1}(e_1+e_2)}(z_i) \rangle$ does not obey momentum conservation, given the value $2Q = 2(b+b^{-1})(e_1+e_2)$ of the background charge in sl_3 Toda theory. However, momentum conservation can be restored by inserting $n-2$ screening operators $V_{b^{-1}e_1}$. (See [6] for similar reasonings and calculations in sl_3 Toda theory.) Thus,

$$\tilde{\Omega}_n \underset{b \rightarrow 0}{\sim} R_n \left\langle \prod_{a=1}^{3n-6} V_{-b^{-1}\omega_1}(y_a) \prod_{i=1}^n V_{b^{-1}(e_1+e_2)}(z_i) \prod_{\ell=1}^{n-2} \int d^2 x_\ell V_{b^{-1}e_1}(x_\ell) \right\rangle^{\text{free}}. \quad (4.48)$$

This free correlation function is the product of the free correlation function (4.42), which we took as our ansatz for $|\Theta_n|^2$, and an integral over x_ℓ , leading to

$$|\Theta_n|^2 \tilde{\Omega}_n \underset{b \rightarrow 0}{\sim} R_n \int \prod_{\ell} d^2 x_\ell \prod_{a,\ell} |x_\ell - y_a|^{\frac{2}{b^2}} \prod_{i,\ell} |x_\ell - z_i|^{-\frac{2}{b^2}} \prod_{\ell \neq \ell'} |x_\ell - x_{\ell'}|^{-\frac{4}{b^2}}. \quad (4.49)$$

The integral in this formula is expected to be dominated by a saddle point, where the x_ℓ s are solutions of

$$2 \sum_{\ell' \neq \ell} \frac{1}{x_\ell - x_{\ell'}} + \sum_i \frac{1}{x_\ell - z_i} - \sum_a \frac{1}{x_\ell - y_a} = 0 . \tag{4.50}$$

(Curiously, these are the Bethe equations for the sl_2 Jaynes-Cummings-Gaudin model at infinite coupling and with spins $\pm \frac{1}{2}$ [23].) The dominant behaviour of the integral is expected to be of the form $|\theta_n|^{\frac{2}{b^2}}$ as $b \rightarrow 0$, with θ_n a b -independent quantity. This $|\theta_n|^{\frac{2}{b^2}}$ factor contradicts the existence of a finite limit for $|\Theta_n|^2 \tilde{\Omega}_n$ as $b \rightarrow 0$, which follows from the conjecture.

One may be tempted to modify the conjecture by adding a factor $\theta_n^{-\frac{1}{b^2}}$ to the twist function Θ_n . This would not only correct the leading behaviour in the $b \rightarrow 0$ limit, but also make the conjecture compatible with global conformal symmetry. We have not mentioned global conformal symmetry until now because this subject is independent from the differential equations in terms of which the conjecture was formulated. It is however easy to see that the conjecture is incompatible with the behaviour of correlation functions under scaling transformations $(z_i, y_a) \rightarrow (\lambda z_i, \lambda y_a)$, except in the $b \rightarrow \infty$ limit.

However, adding the factor $\theta_n^{-\frac{1}{b^2}}$ would spoil the agreement between most terms of the KZ equations (4.23) and the twisted \mathcal{W}_3 null-vector equations (4.44), in particular the terms depending on the spins j_i . The modified conjecture would only hold at the level of the j_i -independent dominant factors in the $b \rightarrow 0$ limit, which would not be interesting.

5 Conclusion

The comparison of sl_3 KZ equations in Sklyanin variables (4.23) with \mathcal{W}_3 null-vector equations (4.44) does not support the conjecture (1.4) in its general form. Nevertheless, the KZ equations are very similar to the null-vector equations: many terms agree nontrivially, and the disagreement is confined to a term which does not depend on the spins j_i of the fields. This remarkable quasi-agreement makes it unlikely that a full agreement can be obtained by modifying the conjecture.

In the critical level limit $k \rightarrow 3 \Leftrightarrow b \rightarrow \infty$, the disagreement disappears and the conjecture (1.4) is true. This limit plays an important role in the Langlands correspondence [24], which might possibly explain why the conjecture (1.4) holds for sl_2 and not for sl_3 , and why in the sl_3 case it holds only in the critical level limit. Another hopeful source of insights is the recent work on conformal Toda theories [6], where the $sl_{N \geq 3}$ cases are understood to be qualitatively different from the sl_2 case. Of course, we already pointed out a significant qualitative difference, namely the failure of the \widehat{sl}_3 cubic field $W^J(z)$ (2.11) to obey the \mathcal{W}_3 algebra. It is not clear how this is related to our problem.

Our results in the sl_3 case lead to natural conjectures in $sl_{N > 3}$ cases, where we expect the KZ equations in Sklyanin variables to agree with \mathcal{W}_N null-vector equations only in the critical level limit $k \rightarrow N$. Let us tentatively perform a counting of equations. There are $\frac{1}{2}N(N - 1)$ isospin variables on the lhs of eq. (1.4), and on the rhs we expect $\frac{1}{2}N(N -$

$1)(n-2)$ Sklyanin variables y_a plus $N(N-1)$ extra variables, which may be collectively included in the symbol U . Differential equations for the sl_N Toda correlation function which generalizes $\tilde{\Omega}_n$ are obtained by eliminating $\frac{1}{2}(N-2)(N+1)n$ non-differential terms from the $\frac{1}{2}N(N-1)(n-2)$ null-vector equations. Thus, we have $n-N(N-1)$ differential equations. When it comes to $\mathcal{K}\Theta_n\tilde{\Omega}_n$, we should presumably add an equation for each one of the extra variables, reaching n differential equations. This precisely the number of KZ equations for the lhs Ω_n of eq. (1.4). In addition, we have the same number of global Ward identities on both sides of eq. (1.4), namely N^2-1 .

A A few technical results

A.1 Helpful identities

The following identities are used in computing the non-differential terms of the operator $E_3 \equiv \Theta_n E_2 \Theta_n^{-1}$ eq. (4.44). Some identities are written modulo terms in $\mathcal{F}_2(y)$ (4.22), as indicated by the relation sign \sim . All identities are proved by elementary manipulations, using observations of the type $\frac{1}{(y-z_i)^2} \sum_b \frac{1}{y_b-z_i} = \frac{1}{(y-z_i)^2} \left(\sum_a \frac{1}{y_a-z_i} - \frac{1}{y-z_i} \right) \sim -\frac{1}{(y-z_i)^3}$.

$$\left(\sum_i \frac{1}{y-z_i} \right)^3 \sim \sum_i \frac{1}{y-z_i} \sum_j \frac{1}{(y-z_j)^2} \sim \sum_i \frac{1}{(y-z_i)^3}, \quad (\text{A.1})$$

$$\sum_b \frac{1}{y-y_b} \frac{1}{(y_b-z_i)^2} \sim \frac{2}{(y-z_i)^3} + \frac{1}{(y-z_i)^2} \sum_b \frac{1}{y-y_b}, \quad (\text{A.2})$$

$$\sum_b \frac{1}{y-y_b} \left(\sum_i \frac{1}{y_b-z_i} \right)^2 \sim \sum_i \frac{-2}{(y-z_i)^3} + \sum_b \frac{1}{y-y_b} \left(\sum_i \frac{1}{y-z_i} \right)^2, \quad (\text{A.3})$$

$$\sum_{bij} \frac{1}{y-y_b} \frac{1}{y-z_j} \frac{1}{y_b-z_i} \sim \sum_i \frac{-1}{(y-z_i)^3} + \sum_b \frac{1}{y-y_b} \left(\sum_i \frac{1}{y-z_i} \right)^2, \quad (\text{A.4})$$

$$\sum_b \frac{1}{y-y_b} \frac{1}{y_b-z_i} \frac{1}{y_b-z_j} \sim \frac{1}{(y-z_i)(y-z_j)} \sum_b \frac{1}{y-y_b} \quad (\text{A.5})$$

$$\sum_b \frac{1}{(y-y_b)^2} \frac{1}{y_b-z_i} \sim \frac{1}{(y-z_i)^3} + \frac{1}{(y-z_i)^2} \sum_b \frac{1}{y-y_b} + \frac{1}{y-z_i} \sum_b \frac{1}{(y-y_b)^2}, \quad (\text{A.6})$$

$$\sum_{b \neq c} \frac{1}{y-y_b} \frac{1}{y_b-y_c} \frac{1}{y_b-z_i} \sim \frac{1}{(y-z_i)^3} + \frac{1}{2} \frac{1}{y-z_i} \left[\left(\sum_b \frac{1}{y-y_b} \right)^2 - \sum_b \frac{1}{(y-y_b)^2} \right], \quad (\text{A.7})$$

$$\sum_{b \neq c} \frac{1}{y-y_b} \frac{1}{y_b-y_c} \frac{1}{y_c-z_i} \sim \frac{2}{(y-z_i)^3} + \frac{1}{2} \frac{1}{y-z_i} \left[\left(\sum_b \frac{1}{y-y_b} \right)^2 - \sum_b \frac{1}{(y-y_b)^2} \right] \quad (\text{A.8})$$

$$-\frac{1}{(y-z_i)^2} \sum_b \frac{1}{y-y_b} + \frac{1}{y-z_i} \sum_b \frac{1}{y-y_b} \sum_c \frac{1}{y_c-z_i}, \quad (\text{A.9})$$

$$\sum_{b \neq c} \frac{1}{(y-y_b)^2} \frac{1}{y_b-y_c} = \sum_b \frac{1}{(y-y_b)^2} \sum_c \frac{1}{y-y_c} - \sum_b \frac{1}{(y-y_b)^3}, \quad (\text{A.10})$$

$$\sum_{\substack{b \neq c \\ d \neq c}} \frac{1}{y-y_b} \frac{1}{y_b-y_c} \frac{1}{y_c-y_d} = - \sum_{b \neq c} \frac{1}{y-y_b} \frac{1}{(y_b-y_c)^2} + \frac{1}{6} \sum_{b \neq c \neq d} \frac{1}{y-y_b} \frac{1}{y-y_c} \frac{1}{y-y_d}, \quad (\text{A.11})$$

$$\sum_{\substack{c \neq b \\ d \neq b}} \frac{1}{y-y_b} \frac{1}{y_b-y_c} \frac{1}{y_b-y_d} = \sum_{b \neq c} \frac{1}{y-y_b} \frac{1}{(y_b-y_c)^2} + \frac{1}{3} \sum_{b \neq c \neq d} \frac{1}{y-y_b} \frac{1}{y-y_c} \frac{1}{y-y_d}, \quad (\text{A.12})$$

$$\sum_{b \neq c \neq d} \frac{1}{y-y_b} \frac{1}{y-y_c} \frac{1}{y-y_d} = \left(\sum_b \frac{1}{y-y_b} \right)^3 + \sum_b \frac{2}{(y-y_b)^3} - 3 \sum_b \frac{1}{y-y_b} \sum_c \frac{1}{(y-y_c)^2}. \quad (\text{A.13})$$

A.2 A characterization of $\mathcal{F}_2(y)$

Here we will justify the characterisation (A.19) of the space $\mathcal{F}_2(y)$ defined in eq. (4.22).

For pedagogical reasons we will begin with the simpler problem of characterizing the space of permutation-symmetric functions of m variables $\{y_a\}$. More precisely, given a function $f(t, \{y_a\})$ which is permutation-symmetric in $\{y_a\}$, depends on an additional variable t , and is regular at $t = y_a$, we want to determine whether $f(y, \{y_a\})$ is actually permutation-symmetric although it apparently depends on y . This amounts to determining whether $f(y_{a'}, \{y_a\})$ actually depends on the choice of a' . If it does not, then for any polynomial $P(t)$ of degree $m-2$ we have

$$\sum_{a'} \oint_{y_{a'}} dt \frac{P(t)f(t)}{\prod_a(t-y_a)} = f(y) \sum_{a'} \oint_{y_{a'}} dt \frac{P(t)}{\prod_a(t-y_a)} = f(y) \oint_{\infty} dt \frac{P(t)}{\prod_a(t-y_a)} = 0. \quad (\text{A.14})$$

So we have transformed the $m-1$ conditions $f(y_1) = f(y_2) = \dots = f(y_m)$ into the condition $\sum_{a'} \oint_{y_{a'}} dt \frac{P(t)f(t)}{\prod_a(t-y_a)} = 0$, which can then be evaluated by moving the integration contours, if the analytic properties of $f(t)$ permit.

Let us apply a similar reasoning to the characterization of $\mathcal{F}_2(y)$. If $f(y) \in \mathcal{F}_2(y)$, for instance $f(y) = \frac{1}{(y-z_{i_0})^2} \tilde{f}(y)$ where $\tilde{f}(y)$ is actually permutation-symmetric, then given any polynomial $P(t)$ of degree $n-7$ we have

$$\sum_{a'} \oint_{y_{a'}} dt P(t) \frac{\prod_{i=1}^n (t-z_i)^2}{\prod_{a=1}^{3n-6} (t-y_a)} f(t) = \tilde{f}(y) \oint_{\infty} dt P(t) \frac{\prod_{i \neq i_0} (t-z_i)^2}{\prod_a (t-y_a)} = 0. \quad (\text{A.15})$$

Thus, to know whether $f(y) \in \mathcal{F}_2(y)$, we only need to evaluate the left hand-side of this equality. To do this we can use the assumed analytic properties of $f(t)$: namely, that it is meromorphic with singularities only at $t = z_i$, and goes to zero as $t \rightarrow \infty$. This implies

$$\sum_{a'} \oint_{y_{a'}} dt P(t) \frac{\prod_i (t-z_i)^2}{\prod_a (t-y_a)} f(t) = - \sum_{i=1}^n \oint_{z_i} dt P(t) \frac{\prod_i (t-z_i)^2}{\prod_a (t-y_a)} f(t), \quad (\text{A.16})$$

which proves $f(y) \in \mathcal{F}_2(y) \Rightarrow \langle P, f \rangle = 0$ as in eq. (A.19). The reverse implication follows from a simple counting of variables: the space of polynomials of degree $n-7$ has dimension $n-6$, which is precisely the number of constraints which we expect for characterizing the space $\mathcal{F}_2(y)$.

A.3 Study of the differential operator D_1

As explained in section 4.4, our conjecture (1.4) implies the relation $D_1 \stackrel{?}{\sim} 0$ or equivalently $D_1 \stackrel{?}{\in} \mathcal{F}_2(y)$, where D_1 is the first-order differential operator written explicitly in eq. (4.45). Here we provide a rigorous argument that this relation is not true, which implies that the conjecture cannot hold for general values of the parameter b .

To start with, let us reduce the study of the first-order differential operator D_1 to the study of mere functions. The operator D_1 , like all our differential equations, is assumed to act on functions which are symmetric under permutations of the $3n - 6$ variables $\{y_a\}$. The space of such functions is algebraically generated by the $3n$ functions

$$\rho_i \equiv \sum_a \log(y_a - z_i), \quad \sigma_i \equiv \sum_a \frac{1}{y_a - z_i}, \quad \tau_i \equiv \sum_a \frac{1}{(y_a - z_i)^2}, \quad i = 1 \cdots n. \quad (\text{A.17})$$

Therefore, $D_1 \sim 0 \Leftrightarrow D_1 \rho_i \sim D_1 \sigma_i \sim D_1 \tau_i \sim 0$. Direct calculations show

$$\begin{aligned} D_1 \rho_i &\sim 0, \\ D_1 \sigma_i &\sim -\frac{1}{(y - z_i)^4} + \frac{\sigma_i}{(y - z_i)^3} + \frac{2}{(y - z_i)^3} \sum_{j \neq i} \frac{1}{y - z_j}, \\ D_1 \tau_i &\sim -\frac{8}{(y - z_i)^5} + \frac{6\sigma_i}{(y - z_i)^4} + \frac{10}{(y - z_i)^4} \sum_{j \neq i} \frac{1}{y - z_j} + \frac{4\tau_i}{(y - z_i)^3} - \frac{2\sigma_i^2}{(y - z_i)^3} \\ &\quad - \frac{6\sigma_i}{(y - z_i)^3} \sum_{j \neq i} \frac{1}{y - z_j} + \frac{2}{(y - z_i)^3} \sum_{j \neq i} \frac{1}{(y - z_j)^2} - \frac{4}{(y - z_i)^3} \left(\sum_{j \neq i} \frac{1}{y - z_j} \right)^2. \end{aligned} \quad (\text{A.18})$$

So $D_1 \sigma_i$ and $D_1 \tau_i$ do not manifestly vanish modulo $\mathcal{F}_2(y)$. Let us however study them further. They may be considered as values at $t = y$ of functions $f(t) = f(t, \{y_a\}, \{z_i\})$ which are invariant under permutations of $\{y_a\}$ but depend on the additional variable t . Let us consider the space of such functions, which we in addition assume to be meromorphic in t with no singularities besides $t = z_i$, and to go to zero as $t \rightarrow \infty$. Let us moreover introduce the space \mathcal{P}_{n-7} of polynomials $P(t)$ of degree $n - 7$. As we show in appendix A.2,

$$f(y) \in \mathcal{F}_2(y) \Leftrightarrow \forall P \in \mathcal{P}_{n-7}, \quad \langle P, f \rangle \equiv \sum_{i=1}^n \oint_{z_i} dt P(t) \frac{\prod_{i=1}^n (t - z_i)^2}{\prod_{a=1}^{3n-6} (t - y_a)} f(t) = 0. \quad (\text{A.19})$$

Then, explicit calculations yields

$$\langle P, D_1 \sigma_i \rangle = 2\pi i \frac{\prod_{k \neq i} (z_i - z_k)^2}{\prod_a (z_i - y_a)} P'(z_i), \quad (\text{A.20})$$

$$\langle P, D_1 \tau_i \rangle = 2\pi i \frac{\prod_{k \neq i} (z_i - z_k)^2}{\prod_a (z_i - y_a)} \left[4P''(z_i) + \left(2\sigma_i + 6 \sum_{k \neq i} \frac{1}{z_i - z_k} \right) P'(z_i) \right]. \quad (\text{A.21})$$

This explicitly demonstrates that $D_1 \notin \mathcal{F}_2(y)$.

However, D_1 still has remarkable properties with respect to the constant polynomial $P = 1$, namely $\langle 1, D_1 \sigma_i \rangle = \langle 1, D_1 \tau_i \rangle = 0$. These non-trivial identities sensitively depend on the general structure of D_1 and on the particular values of λ, μ, ν which determine its

coefficients. This implies that, whereas arbitrary differential operators belong to $\mathcal{F}_2(y)$ for $n \leq 6$, $D_1 \in \mathcal{F}_2(y)$ for $n \leq 7$. The significance of these properties of D_1 is not clear. When combined with $D_1 \rho_i \sim 0$, they suggest that $D_1 \sim 0$ when applied to a special class of permutation-symmetric function of y_a (and z_i), and one might wonder whether $\Theta_n \tilde{\Omega}_n$ actually belongs to this class. Given the freedom to choose $y \in \{y_a\}$, this would imply that $\tilde{\Omega}_n$ satisfies $n - 6$ further differential equations. But $\tilde{\Omega}_n$ is not expected to satisfy any further differential equations besides the global Ward identities, whose number is n -independent. So the supposition $D_1 \cdot \Theta_n \tilde{\Omega}_n \stackrel{?}{\sim} 0$ certainly fails for $n > 7$, and so does our conjecture (1.4).

Acknowledgments

I wish to thank Alexander Chervov and Alexey Litvinov for interesting discussions and very helpful comments on this manuscript. I am also grateful to Nicolas Crampé, Vladimir Fateev, Philippe Roche, Volker Schomerus and Joerg Teschner for stimulating discussions. I acknowledge the hospitality of DESY, Hamburg while part of this work was done.

References

- [1] S. Ribault and J. Teschner, H_3^+ WZNW correlators from Liouville theory, *JHEP* **06** (2005) 014 [[hep-th/0502048](#)] [[SPIRES](#)].
- [2] Y. Hikida and V. Schomerus, H_3^+ WZNW model from Liouville field theory, *JHEP* **10** (2007) 064 [[arXiv:0706.1030](#)] [[SPIRES](#)].
- [3] A.V. Stoyanovsky, A relation between the Knizhnik-Zamolodchikov and Belavin-Polyakov-Zamolodchikov systems of partial differential equations, [math-ph/0012013](#) [[SPIRES](#)].
- [4] K. Hosomichi and S. Ribault, Solution of the H_3^+ model on a disc, *JHEP* **01** (2007) 057 [[hep-th/0610117](#)] [[SPIRES](#)].
- [5] S. Ribault, A family of solvable non-rational conformal field theories, *JHEP* **05** (2008) 073 [[arXiv:0803.2099](#)] [[SPIRES](#)].
- [6] V.A. Fateev and A.V. Litvinov, Correlation functions in conformal Toda field theory I, *JHEP* **11** (2007) 002 [[arXiv:0709.3806](#)] [[SPIRES](#)].
- [7] E.K. Sklyanin, Quantum inverse scattering method. Selected topics, [hep-th/9211111](#) [[SPIRES](#)].
- [8] E.K. Sklyanin, Separation of variables in the quantum integrable models related to the Yangian $Y[sl(3)]$, *J. Math. Sci.* **80** (1996) 1861 [[hep-th/9212076](#)] [[SPIRES](#)].
- [9] P. Di Francesco, P. Mathieu and D. Senechal, *Conformal field theory*, Springer, New York U.S.A. (1997), pag. 890.
- [10] V.G. Knizhnik and A.B. Zamolodchikov, Current algebra and Wess-Zumino model in two dimensions, *Nucl. Phys.* **B 247** (1984) 83 [[SPIRES](#)].
- [11] P. Bouwknegt and K. Schoutens, *W symmetry in conformal field theory*, *Phys. Rept.* **223** (1993) 183 [[hep-th/9210010](#)] [[SPIRES](#)].

- [12] A.A. Belavin, A.M. Polyakov and A.B. Zamolodchikov, *Infinite conformal symmetry in two-dimensional quantum field theory*, *Nucl. Phys. B* **241** (1984) 333 [SPIRES].
- [13] Y. Hikida and V. Schomerus, *The FZZ-duality conjecture — A proof*, *JHEP* **03** (2009) 095 [[arXiv:0805.3931](#)] [SPIRES].
- [14] G. Giribet and M. Leoni, *A twisted FZZ-like dual for the 2D black hole*, *Rept. Math. Phys.* **61** (2008) 151 [[arXiv:0706.0036](#)] [SPIRES].
- [15] A.B. Zamolodchikov and A.B. Zamolodchikov, *Structure constants and conformal bootstrap in Liouville field theory*, *Nucl. Phys. B* **477** (1996) 577 [[hep-th/9506136](#)] [SPIRES].
- [16] J. Teschner, *The mini-superspace limit of the $SL(2, C)/SU(2)$ WZNW model*, *Nucl. Phys. B* **546** (1999) 369 [[hep-th/9712258](#)] [SPIRES].
- [17] E. Mukhin, V. Schechtman, V. Tarasov, A. Varchenko, *On the new form of Bethe ansatz equations and separation of variables in the sl_3 Gaudin model*, [math/0609428](#).
- [18] E.K. Sklyanin, *Separation of variables in the classical integrable $SL(3)$ magnetic chain*, *Commun. Math. Phys.* **150** (1992) 181 [[hep-th/9211126](#)] [SPIRES].
- [19] F.A. Smirnov, *Separation of variables for quantum integrable models related to $U_q(\widehat{sl}_N)$* , [math-ph/0109013](#).
- [20] A. Chervov and G. Falqui, *Manin matrices and Talalaev's formula*, *J. Phys. A* **41** (2008) 194006 [[arXiv:0711.2236](#)] [SPIRES].
- [21] M.R. Adams, J.P. Harnad and J. Hurtubise, *Darboux coordinates on coadjoint orbits of Lie algebras*, *Lett. Math. Phys.* **40** (1997) 41 [[solv-int/9603001](#)] [SPIRES].
- [22] G.M.T. Watts, *Fusion in the $W(3)$ algebra*, *Commun. Math. Phys.* **171** (1995) 87 [[hep-th/9403163](#)] [SPIRES].
- [23] O. Babelon and D. Talalaev, *On the Bethe ansatz for the Jaynes-Cummings-Gaudin model*, *J. Stat. Mech.* (2007) P06013 [[hep-th/0703124](#)] [SPIRES].
- [24] E. Frenkel, *Lectures on the Langlands program and conformal field theory*, [hep-th/0512172](#) [SPIRES].