

Noncommutative $\mathcal{N} = 1$ super Yang-Mills, the Seiberg-Witten map and UV divergences

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

JHEP11(2009)092

(<http://iopscience.iop.org/1126-6708/2009/11/092>)

[The Table of Contents](#) and [more related content](#) is available

Download details:

IP Address: 80.92.225.132

The article was downloaded on 01/04/2010 at 13:31

Please note that [terms and conditions apply](#).

RECEIVED: August 3, 2009

REVISED: October 5, 2009

ACCEPTED: October 30, 2009

PUBLISHED: November 20, 2009

Noncommutative $\mathcal{N} = 1$ super Yang-Mills, the Seiberg-Witten map and UV divergences

C.P. Martín^a and C. Tamarit^b

^a*Departamento de Física Teórica I, Facultad de Ciencias Físicas,
Universidad Complutense de Madrid,
28040 Madrid, Spain*

^b*Kavli Institute for Theoretical Physics, University of California,
Santa Barbara, CA, 93106-4030, U.S.A.*

E-mail: carmelo@elbereth.fis.ucm.es, tamarit@kitp.ucsb.edu

ABSTRACT: Classically, the dual under the Seiberg-Witten map of noncommutative $U(N)$, $\mathcal{N} = 1$ super Yang-Mills theory is a field theory with ordinary gauge symmetry whose fields carry, however, a θ -deformed nonlinear realisation of the $\mathcal{N} = 1$ supersymmetry algebra in four dimensions. For the latter theory we work out at one-loop and first order in the noncommutative parameter matrix $\theta^{\mu\nu}$ the UV divergent part of its effective action in the background-field gauge, and, for $N \neq 1$, we show that for finite values of N the gauge sector fails to be renormalisable; however, in the large N limit the full theory is renormalisable, in keeping with the expectations raised by the quantum behaviour of the theory's noncommutative classical dual. We also obtain –for $N \geq 3$, the case with $N = 2$ being trivial– the UV divergent part of the effective action of the $SU(N)$ noncommutative theory in the enveloping-algebra formalism that is obtained from the previous ordinary $U(N)$ theory by removing the $U(1)$ degrees of freedom. This noncommutative $SU(N)$ theory is also renormalisable.

KEYWORDS: Supersymmetric gauge theory, Renormalization Regularization and Renormalons, Non-Commutative Geometry

ARXIV EPRINT: [0907.2437](https://arxiv.org/abs/0907.2437)

Contents

1	Introduction	1
2	The model and the background field method	5
3	Computation of the divergent part of the effective action	8
3.1	Commutative limit	12
3.2	Noncommutative contributions to $\tilde{\Gamma}^{(3,0)}$	12
3.3	Noncommutative contributions to $\tilde{\Gamma}^{(0,1)}$	13
3.4	Noncommutative contributions to $\tilde{\Gamma}^{(1,1)}$	13
3.5	Noncommutative contributions to $\tilde{\Gamma}^{(0,2)}$	14
3.6	Final expression	15
4	Analysing renormalisability	16
4.1	Commutative renormalisation	18
4.2	Renormalisation of the noncommutative bosonic sector	18
4.3	Renormalisation of the noncommutative gaugino sector	20
5	Conclusions	22
A	Lie algebra, Dirac algebra, Majorana spinors	23
B	Feynman rules for $\alpha = 1$, $\mathfrak{S}_\mu^{\text{tree}} = \mathfrak{L}^{\text{tree}} = 0$	24

1 Introduction

Noncommutative gauge theories are known to arise as low energy limits of (super)string theory [1, 2], and they are interesting on their own as examples of nonlocal theories. One of their intriguing features is that noncommutative $U(N)$ gauge theories, considered as effective descriptions of the dynamics of D-branes with Neveu-Schwarz backgrounds, are known to have a dual description in terms of fields with ordinary gauge invariance [1]. This equivalence, which can be traced back to the possibility of choosing different yet equivalent regularisations of the D-Brane effective action, can be formulated by means of a map which relates noncommutative and ordinary gauge fields in a way consistent with their respective gauge symmetries, so that orbits of noncommutative gauge transformations are mapped into orbits of ordinary gauge transformations. These maps are called Seiberg-Witten maps. Their role linking different DBI actions has also been shown to hold, at least to a certain approximation, in the $\mathcal{N} = 1$ supersymmetric case [3]. In principle, this equivalence holds for the D-Brane effective actions, but one may wonder whether it also holds, at the quantum

level, for the noncommutative gauge theories that do not involve the higher order terms present in the DBI actions.

The idea of mapping noncommutative to ordinary gauge symmetries was the starting point for the formulation of noncommutative gauge theories for arbitrary gauge groups by means of Seiberg-Witten maps pioneered in refs. [4–6]. In the “standard” formalism, closure under gauge transformations restricts the gauge groups to be $U(N)$ and the representations to be (anti-)fundamental or bi-(anti)-fundamental, while the formalism which makes use of Seiberg-Witten maps, also referred to as the enveloping algebra formalism, makes it possible to consider arbitrary gauge groups and representations by mapping the enveloping-algebra valued noncommutative gauge fields to ordinary Lie-algebra valued gauge fields.

The quantum properties of noncommutative gauge theories, both in the standard and enveloping algebra approaches, have been analysed in many works. Concerning the standard approach, nonsupersymmetric noncommutative $U(N)$ Yang-Mills theories are plagued by pathological IR divergences coming from the UV/IR mixing effect [7], which are suppressed in the large N limit, in which only planar diagrams contribute and the sole effect of noncommutativity is producing phase factors depending on the external momenta which can be taken out of the loop integrals. Noncommutative supersymmetric gauge theories [8] exhibit a better behaviour in the infrared, as the problematic divergences are milder or altogether absent [9–11]. These milder noncommutative IR divergences are logarithmic and can be integrated leading to a consistent renormalisable supersymmetric noncommutative Wess-Zumino [12] and most likely to consistent renormalisable, or even UV finite, supersymmetric noncommutative $U(N)$ theories [13, 14]. A noncommutative extension of the MSSM has been put forward in ref. [15], which contains more “particle” states than the ordinary MSSM due to the noncommutative anomaly cancellation conditions [16, 17] and other noncommutative requirements.

On the other hand, concerning the theories defined by means of Seiberg-Witten maps, they are known to have gauge anomaly cancellation conditions identical to their commutative counterparts [18], and their renormalisability properties have been studied in a wide number of papers [19–27]. The results can be summarised as follows: pure gauge theories, $U(1)$ or $SU(N)$, are one-loop renormalisable at least to first order in the noncommutativity parameters. The introduction of matter fields in the form of Dirac fermions or complex scalars in arbitrary representations (but such that the matter Lagrangian in terms of noncommutative fields does not involve a covariant derivative with a star-product commutator), does not spoil the renormalisability of the gauge sector of the theory; however, the full theory seems to be nonrenormalisable in all cases analysed. These cases for which the renormalisability of the matter sector has been addressed are: Dirac fermions with gauge groups $U(1)$ [20, 21] or $SU(2)$ in the fundamental representation [22], and $U(1)$ complex scalars [25]. Renormalisability is spoiled by the appearance of divergences in matter field Green functions which cannot be removed by multiplicative renormalisations or field redefinitions. There is still no definitive answer concerning whether other types of matter fields or representations could overcome this problem, despite promising results concerning chiral fermions [27]. Still, the renormalisability properties of theories with Majorana fermions or/and covariant derivatives involving a star-product commutator have not been studied.

Moreover, supersymmetry could be expected to make some divergences go away. However, though generally supersymmetry is associated with a cancellation of divergences between bosonic and fermionic degrees of freedom, and noncommutative $U(N)$ theories defined by means of Seiberg-Witten maps have been shown to be compatible with supersymmetry, it turns out that the latter is realised nonlinearly in the ordinary fields [3], and thus it is not clear how it will affect divergences.

Comparing the quantum properties of noncommutative theories in both the standard and enveloping algebra approaches raises interesting questions regarding their equivalence for $U(N)$ gauge groups, for which the Seiberg-Witten map establishes a classical equivalence. The different gauge anomaly cancellation conditions makes this equivalence doubtful in the presence of chiral fermions, at least when noncommutativity is treated perturbatively. In the case of theories without matter, the equivalence has been found to hold for noncommutative Chern-Simons [28] –a theory which is UV finite–, whereas for other gauge theories with or without matter there is no concluding evidence, since on the side of the enveloping algebra approach the theories studied have exclusively $U(1)$ and $SU(N)$ gauge groups, while to make contact with the standard formalism one should consider $U(N)$ in the large N limit, in which the theories, at least at the one-loop level, are supposed to be well behaved and renormalisable for infinitesimal noncommutativity.

We have so far identified several issues that need further investigation, and, indeed, there are two goals that one can achieve by studying the renormalisability of the $\mathcal{N} = 1$ super Yang-Mills θ -expanded theory. On one hand, the study of the renormalisability properties, both for the gauge sector and the full theory, of noncommutative theories defined by means of Seiberg-Witten maps (i.e., theories defined within the enveloping-algebra formalism of refs. [4–6]) with Majorana fermions and/or involving a covariant derivative with star-product commutators and/or supersymmetry. This study is much needed since, within the enveloping-algebra formalism, the inclusion of fermions in noncommutative $SU(N)$ theories in a nonsupersymmetric way leads to nonrenormalisable theories: the fermionic part of nonsupersymmetric noncommutative $SU(N)$ is nonrenormalisable –see ref. [22]. On the other hand, to check in a highly-nontrivial setting the equivalence at the quantum level of the standard and enveloping algebra (θ -expanded) approaches for supersymmetric noncommutative $U(N)$ gauge theories in the large N limit, i.e., the quantum duality of supersymmetric noncommutative $U(N)$ formulated in terms of noncommutative fields and the supersymmetric theory, whose fields are ordinary gauge fields carrying a nonlinear realisation of supersymmetry, obtained from the former by using the Seiberg-Witten map. In this regard, it should be stressed that the Seiberg-Witten map is not a field redefinition of the ordinary field –it maps ordinary gauge orbits to noncommutative gauge orbits– and, hence, it is far from clear that renormalisability of the planar sector of the standard noncommutative supersymmetric $U(N)$ theory implies renormalisability of the large- N part of its classical ordinary dual under the Seiberg-Witten map. Besides, the Seiberg-Witten map is defined by products of fields at the same point and renormalisation of the Green functions of elementary fields does not imply renormalisation of Green functions of composite operators.

It is plain that the final and more ambitious goal regarding the perturbative renormalisability of the planar (and $SU(N)$) sector of the theta-expanded $U(N)$ SYM theory

would be to prove renormalisability at any order in $\theta^{\mu\nu}$ and the number of loops. Here perturbative renormalisability is understood in the physical sense that once the coupling constants and mass parameters that occur in the classical action –these coupling constants and mass parameters being a finite set– have been renormalized, the so remaining (if any) UV divergences can be accounted for by field redefinitions which are parametrised by an infinite number of UV divergent parameters, which constitute the physically irrelevant part of the Seiberg-Witten map. This all-order perturbative renormalisability proof seems to be technically out of reach at the moment and to carry it out will require a great deal of ingenuity –though some partial all-order-in-theta proofs have been given in refs. [19, 26]–, in this regard some improvement of the techniques introduced in ref. [29] may be of help. This all-order renormalisability proof is a preliminary requirement to fully establish quantum Seiberg-Witten map duality, on the one hand, and, to fully make sense out of any noncommutative field theory formulated within the enveloping algebra formalism. However, as it has happened in the past in the history of quantum field theory, in the absence of general proofs explicit one-loop computations at first order in $\theta^{\mu\nu}$ are always welcome since they may serve as a guidance on how to achieve the ultimate goal and to reveal whether some properties such as supersymmetry maybe useful to attain that goal –U(N) YM theory in the large-N limit in the case at hand–. Besides, these computations show in which cases the ultimate goal cannot be accomplished–for finite N, the U(N) YM theory in the case analysed here–. All in all, a full perturbative proof of the renormalizability of the kind of θ -expanded theories studied in this paper requires far more work and a better understanding of both the technical issues and formal ideas involved in their formulation.

The aim of this paper is to take the first steps in the study of the two open issues mentioned above by analysing the renormalisability properties of $\mathcal{N} = 1$ U(N) super Yang-Mills in the enveloping algebra approach, with the ordinary fields taking values in the fundamental representation of the gauge group. First, the theory has a Majorana fermion with a covariant derivative involving a star-product commutator; supersymmetry is also present for the noncommutative fields, and it is inherited by the ordinary fields albeit in a nonlinear fashion. Secondly, since we have a U(N) gauge group in the fundamental representation, the theory can also be formulated in the standard approach, in which case, in the large N limit, it is renormalisable and well-behaved for small noncommutativity. We will analyse whether one-loop renormalisability in the background field gauge is achieved at least for large N. Further, in order to complement previous research regarding theories with simple gauge groups, we will study the renormalisability properties of the SU(N) model that results from eliminating the U(1) degrees of freedom in the U(N) theory, with the goal of seeing whether the modified field content and interactions yield a better behaviour at the quantum level. To tackle these problems, we will compute the divergent part of the one-loop effective action at first order in the noncommutative parameters $\theta^{\mu\nu}$, using the background field method in the background field gauge and dimensional regularisation, and we will study whether the divergences can be removed by appropriate multiplicative renormalisations of the parameters of the theory plus nonmultiplicative field redefinitions.

The paper is organised as follows. The model and the background field method are introduced in section 2. Section 3 is devoted to the computation of the full divergent part

of the one-loop effective action: first, a method is outlined which allows to obtain the full result by calculating a minimum number of diagrams, whose divergent parts are then computed in dimensional regularisation; following this the full gauge invariant expression is finally reconstructed. The renormalisability of the theory, both for arbitrary finite and large N , is studied in section 4, and then conclusions are drawn in section 5. Two appendices are included, the first one with some Lie and Dirac algebra identities, and the second one displaying the Feynman rules employed in the computation.

2 The model and the background field method

The action of the model, in terms of noncommutative fields, is the following,

$$S = \int d^4x - \frac{1}{2g^2} \text{Tr} F_{\mu\nu} \star F^{\mu\nu} + \frac{i}{g^2} \text{Tr} \bar{\Lambda} \not{D} \star \Lambda, \\ F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu - i[A_\mu, A_\nu]_\star, D_{\star, \mu} = \partial_\mu - i[A_\mu,]_\star, \quad (2.1)$$

where the fields take values in the enveloping algebra of $U(N)$, $A_\mu = A_\mu^A T^A$, $\Lambda = \Lambda^A T^A$ and Λ is a Majorana spinor (see appendix A for conventions). The $U(N)$ fields will be taken in the fundamental representation. The noncommutative product \star is the usual Moyal product,

$$a \star b = a \exp \left[\frac{i\hbar}{2} \theta^{\mu\nu} \overleftarrow{\partial}_\mu \overrightarrow{\partial}_\nu \right] b,$$

with \hbar setting the noncommutative scale. The model has $\mathcal{N} = 1$ supersymmetry in terms of the noncommutative fields; it can be formulated in terms of a noncommutative vector superfield in the Wess-Zumino gauge.

The noncommutative fields are defined in terms of $U(N)$ Lie algebra valued ordinary fields, which we denote by $\mathbf{a}_\mu, \mathfrak{l}$, by means of the following Seiberg-Witten maps,

$$A_\mu = \mathbf{a}_\mu - \frac{\hbar}{4} \theta^{\alpha\beta} \{ \mathbf{a}_\alpha, \partial_\beta \mathbf{a}_\mu + \mathfrak{f}_{\beta\mu} \} + \hbar \mathfrak{S}_\mu + O(\hbar^2), \\ \Lambda = \mathfrak{l} - \frac{\hbar}{4} \theta^{\alpha\beta} \{ \mathbf{a}_\alpha, 2D_\beta \mathfrak{l} + i[\mathbf{a}_\beta, \mathfrak{l}] \} + \hbar \mathfrak{L} + O(\hbar^2), \quad (2.2)$$

where $D_\mu = \partial_\mu - i[\mathbf{a}_\mu,]$, $\mathfrak{f}_{\mu\nu} = \partial_\mu \mathbf{a}_\nu - \partial_\nu \mathbf{a}_\mu - i[\mathbf{a}_\mu, \mathbf{a}_\nu]$, and $\mathfrak{S}_\mu, \mathfrak{L}$ represent the ambiguities in the map at order \hbar , given by sums of terms which involve a contraction with $\theta^{\mu\nu}$, have the appropriate mass dimensions and transform in the adjoint representation of the gauge group; they can be argued to be equivalent to field redefinitions, as will be seen in section 4.

We will work with the following decomposition of the $U(N)$ fields in the fundamental representation into their $SU(N)$ and $U(1)$ parts:

$$\mathbf{a}_\mu = a_\mu^a T^a + b_\mu \frac{\mathbb{1}}{\sqrt{2N}}, \quad \mathfrak{f}_{\mu\nu} = f_{\mu\nu}^a T^a + g_{\mu\nu} \frac{\mathbb{1}}{\sqrt{2N}}, \\ \mathfrak{l} = \lambda^a T^a + u \frac{\mathbb{1}}{\sqrt{2N}}. \quad (2.3)$$

This will allow us to study the properties of both the U(N) theory and the SU(N) theory that results from suppressing the U(1) degrees of freedom b_μ, u .

We will argue in the next section that, for the purpose of checking renormalisability, it suffices to compute the divergent part of the effective action ignoring at tree-level the ambiguities $\mathfrak{S}_\mu, \mathfrak{L}$ of the Seiberg-Witten maps in eq. (2.2); the ambiguities, however, have to be taken into account when considering the allowed counterterms. The action in terms of ordinary fields, after expanding (2.1) with eqs. (2.2) with $\mathfrak{S}_\mu = \mathfrak{L} = 0$, turns out to be the following

$$\begin{aligned}
 S &= S^{(0)} + hS^{(1)} + O(h^2), \\
 S^{(0)} &= -\frac{1}{2g^2} \int d^4x \operatorname{Tr} f_{\mu\nu} f^{\mu\nu} + \frac{i}{g^2} \int d^4x \operatorname{Tr} \bar{l} \not{D} l, \\
 S^{(1)} &= \frac{1}{4g^2} \int d^4x \operatorname{Tr} \theta^{\alpha\beta} f_{\mu\nu} f^{\mu\nu} f^{\alpha\beta} - \frac{1}{g^2} \int d^4x \operatorname{Tr} \theta^{\alpha\beta} f_{\alpha\mu} f_{\beta\nu} f^{\alpha\beta} - \frac{i}{4} \int d^4x \operatorname{Tr} \theta^{\alpha\beta} \bar{l} \gamma^\mu \{D_\mu l, f_{\alpha\beta}\} \\
 &\quad - \frac{i}{2} \int d^4x \operatorname{Tr} \theta^{\alpha\beta} \bar{l} \gamma^\mu \{D_\beta l, f_{\mu\alpha}\}.
 \end{aligned} \tag{2.4}$$

In the previous action, all the noncommutative terms involve traces of the type $\operatorname{Tr} T^A \{T^B, T^C\} = \frac{1}{2} d^{ABC}$ (see appendix A). For $N < 3$, the SU(N) part of the Lie algebra, for arbitrary representations, has $d^{abc} = 0$, which means that the SU(N) theory obtained by eliminating the U(1) degrees of freedom is, to order h , equivalent to its commutative limit. Therefore, when studying the SU(N) theory we will only consider $N \geq 3$. As shown in ref. [3] (see also [30]) the fields in the action in eq.(2.4) carry a nonlinear realisation of $\mathcal{N} = 1$ supersymmetry which define supersymmetry transformations that leave that action invariant.

In the enveloping algebra approach, quantisation is performed on the ordinary fields. In order to compute the effective action with the background field method [31], we split the gauge field \mathfrak{a}_μ in a background part b_μ and a quantum part q_μ ,

$$\mathfrak{a}_\mu = b_\mu + q_\mu. \tag{2.5}$$

A gauge transformation of \mathfrak{a}_μ , $\delta\mathfrak{a}_\mu = D_\mu c$, can be generated by two types of transformations of the fields b, q :

$$\text{Quantum gauge transformations: } \delta q_\mu = D[q]_\mu c, \delta b_\mu = -i[b_\mu, c], \quad D[q]_\mu = \partial_\mu - i[q_\mu,], \tag{2.6}$$

$$\text{Background gauge transformations: } \delta q_\mu = -i[q_\mu, c], \delta b_\mu = D[b]_\mu c, \quad D[b]_\mu = \partial_\mu - i[b_\mu,]. \tag{2.7}$$

In order to quantise q with the path integral formalism, a gauge fixing procedure is needed for the transformations in eq. (2.6). The background field method relies in a clever choice of the gauge-fixing function which is covariant under the transformations (2.7). With the gauge-fixing choice $G = D_\mu^{[b]} q^\mu = 0$, the gauge-fixing and ghost action are the following

$$S_{gf} = -\frac{1}{2\alpha} \int d^4x (D_\mu^{[b]} q^\mu)^2, \quad S_{gh} = \int d^4x \bar{c} D_\mu^{[b]} D^{[b+q]\mu} c. \tag{2.8}$$

Quantising the fields $q_\mu, \mathfrak{l}, \bar{\mathfrak{l}}$, the generating functional of the background Green functions is given by

$$\tilde{Z}[\tilde{J}, \tilde{\sigma}, \tilde{\bar{\sigma}}; b] = \int [dq][d\mathfrak{l}][d\bar{\mathfrak{l}}] \exp[i(S[b + q, \mathfrak{l}, \bar{\mathfrak{l}}] + S_{gf}[q; b] + S_{gh}[c, \bar{c}, q; b] + \tilde{J}_\mu q^\mu + \tilde{\sigma}\mathfrak{l} + \tilde{\bar{\sigma}}\bar{\mathfrak{l}})], \quad (2.9)$$

where $\tilde{J}, \tilde{\sigma}, \tilde{\bar{\sigma}}$ are sources for the gauge field and Majorana fermions. Note the use of “ $\tilde{}$ ” to distinguish the background currents and functional generator \tilde{Z} from the ones defining the true Green functions of the theory, when the splitting of eq. (2.5) is not used and functional integration is performed over \mathbf{a} . The generator of connected background Green functions is given by

$$\tilde{W}[\tilde{J}, \tilde{\sigma}, \tilde{\bar{\sigma}}; b] = -i \ln \tilde{Z}[\tilde{J}, \tilde{\sigma}, \tilde{\bar{\sigma}}; b].$$

Defining the background classical fields as

$$\tilde{q} = \frac{\delta \tilde{W}}{\delta \tilde{J}}, \quad \tilde{\mathfrak{l}} = \frac{\delta \tilde{W}}{\delta \tilde{\sigma}}, \quad \tilde{\bar{\mathfrak{l}}} = -\frac{\delta \tilde{W}}{\delta \tilde{\bar{\sigma}}},$$

then by performing a Legendre transformation we get the functional $\tilde{\Gamma}$ which generates the 1PI connected background Green functions:

$$\tilde{\Gamma}[\tilde{q}, \tilde{\mathfrak{l}}, \tilde{\bar{\mathfrak{l}}}; b] = \tilde{W}[J, \tilde{\sigma}, \tilde{\bar{\sigma}}; b] - \int d^4x \tilde{J}_\mu \tilde{q}^\mu - \int d^4x \tilde{\sigma} \tilde{\mathfrak{l}} - \int d^4x \tilde{\bar{\sigma}} \tilde{\bar{\mathfrak{l}}}. \quad (2.10)$$

In a similar fashion, without using the splitting of eq. (2.5), one can define the true Green function generators $Z[J, \sigma, \bar{\sigma}]$ and $W[J, \sigma, \bar{\sigma}]$ as well as the true classical fields $\hat{\mathbf{a}}, \hat{\mathfrak{l}}, \hat{\bar{\mathfrak{l}}}$. Standard formal manipulations show that the effective action of the theory $\Gamma[\hat{\mathbf{a}}, \hat{\mathfrak{l}}, \hat{\bar{\mathfrak{l}}}]$ is related to $\tilde{\Gamma}[\tilde{q}, \tilde{\mathfrak{l}}, \tilde{\bar{\mathfrak{l}}}; b]$ of eq. (2.10) by the following identity [31]:

$$\Gamma[\hat{\mathbf{a}}, \hat{\mathfrak{l}}, \hat{\bar{\mathfrak{l}}}] = \tilde{\Gamma}[0, \tilde{\mathfrak{l}}, \tilde{\bar{\mathfrak{l}}}; b]_{b=\hat{\mathbf{a}}, \tilde{\mathfrak{l}}=\hat{\mathfrak{l}}, \tilde{\bar{\mathfrak{l}}}=\hat{\bar{\mathfrak{l}}}}, \quad (2.11)$$

where Γ is computed with an unusual gauge-fixing. From the r.h.s. of eq. (2.11) it is clear that the effective action is obtained by calculating the background effective action for the Majorana fields after integrating out the quantum fields q , with the background fields b_μ taken as external sources. We thus can write

$$\Gamma[\hat{\mathbf{a}}, \hat{\mathfrak{l}}, \hat{\bar{\mathfrak{l}}}] = \int d^4x \sum_k \frac{-i}{2^k (k!)^2} \tilde{\Gamma}[\hat{\mathbf{a}}]_{\substack{i_1, \dots, i_k, \\ A_1, \dots, A_k}}^{(k)} \prod_{l=1}^k \hat{\mathfrak{l}}_{i_l}^{A_l} \prod_{p=1}^k \hat{\bar{\mathfrak{l}}}_{j_p}^{B_p},$$

where the factor $(k!)^2$ takes into account the permutations of the i 's and j 's, while the factor 2^k comes from the fact that, since the Majorana fermions are self-conjugate, it is always possible to interchange one \mathfrak{l} with an $\bar{\mathfrak{l}}$. $\tilde{\Gamma}[\hat{\mathbf{a}}]^{(k)}$ is nothing but the sum of background 1PI diagrams with k fermionic legs, k anti-fermionic legs and no quantum gauge field legs, and with the background field b renamed as $\hat{\mathbf{a}}$. Expanding $\tilde{\Gamma}[\hat{\mathbf{a}}]^{(k)}$ in the number of background gauge fields, one gets

$$\Gamma[\hat{\mathbf{a}}, \hat{\mathfrak{l}}, \hat{\bar{\mathfrak{l}}}] = \int d^4x \sum_k \sum_n \frac{-i}{2^k (k!)^2} \tilde{\Gamma}^{(n,k)}_{\substack{i_1, \dots, i_k, \\ A_1, \dots, A_k}} \prod_{l=1}^k \hat{\mathfrak{l}}_{i_l}^{A_l} \prod_{p=1}^k \hat{\bar{\mathfrak{l}}}_{j_p}^{B_p} \prod_{m=1}^n \hat{\mathbf{a}}_{\mu_m}^{C_m}. \quad (2.12)$$

In the previous formula $\tilde{\Gamma}^{(n,k)}$ is equivalent to a background 1PI diagram with n background gauge field legs, k fermionic legs and k anti-fermionic legs. Note that our definitions do not involve any symmetrisation over the background gauge fields. Symmetrising over them we can make contact with the usual expansion of the effective action in terms of 1PI Green functions:

$$\Gamma[\hat{a}, \hat{l}, \hat{\bar{l}}] = \int d^4x \sum_k \sum_n \frac{-i}{n!2^k(k!)^2} \Gamma_{\substack{i_1, \dots, i_k, \\ A_1, \dots, A_k}}^{(n,k)} \prod_{l=1}^k \hat{l}_{i_l}^{A_l} \prod_{p=1}^k \hat{l}_{j_p}^{B_p} \prod_{m=1}^n \hat{a}_{\mu_m}^{C_m},$$

where $\Gamma^{(n,k)}$, which is obtained from $\tilde{\Gamma}^{(n,k)}$ by summing over the permutations of the background gauge fields, is the 1PI Green function with n gauge fields and k fermion pairs.

The advantage of using background diagrams coming from the functional generator in eq. (2.9) is that $\tilde{\Gamma}[0, \tilde{l}, \tilde{\bar{l}}; b]$ is gauge invariant, so that the effective action $\Gamma[\hat{a}, \hat{l}, \hat{\bar{l}}]$ is indeed gauge invariant. As explained in the next section, this can be used to simplify the computation of the divergent part of the effective action.

3 Computation of the divergent part of the effective action

The aim of this section is to compute the divergent part of the effective action at first order in $\hbar\theta$, by calculating the background 1PI diagrams $\tilde{\Gamma}^{(n,k)}$ with no external quantum gauge fields of eq. (2.12) using the Feynman rules associated with the functional generator in eq. (2.9). These rules can be derived from the expressions for the action, gauge fixing and ghost terms given in eqs. (2.4), (2.8), keeping in mind the splitting (2.5).

Before plunging into the computation, we will justify a number of simplifications that do not imply a loss of generality on the final result concerning the regularisation and renormalisation of the theory.

- We shall carry out our computations in dimensional regularisation with $D = 4 - 2\epsilon$ –it is always advisable to keep an eye on dimensional reduction. That this regularisation does not preserve supersymmetry will have no bearing on our conclusions since our computations are one-loop and the inclusion of the ϵ -scalars of dimensional reduction to turn our dimensionally regularised theory into a theory regularised by dimensional reduction –and thus supersymmetric– will not modify the value of UV divergences that we will compute, but will add new ones which would be subtracted by introducing counterterms made out of “evanescent” operators and couplings –see ref. [32, 33] for further details.
- Choice of gauge $\alpha = 1$ in the gauge-fixing term in eq.(2.8). This choice of gauge simplifies the gauge field propagator. This brings up the question of whether, if problematic divergences appear for $\alpha = 1$ that make the theory nonrenormalisable, the consideration of an arbitrary α might help remove these divergences. The answer is negative whenever any of the problematic divergences appearing at $\alpha = 1$ do not go away on the mass shell. This is due to the results in ref. [34] (see also [35]) which establish that the background field effective action is independent of the gauge-fixing

term if the background fields are on shell. Thus, when the background fields are on shell any divergent contribution remaining will be independent of any gauge-fixing term that we chose.

- Setting to zero the tree-level ambiguities $\mathfrak{S}_\mu, \mathfrak{L}$ of the Seiberg-Witten map of eq. (2.2). This choice simplifies greatly the computation of the diagrams, though when studying renormalisability one can still contemplate infinite renormalisations of $\mathfrak{S}_\mu, \mathfrak{L}$, which tantamounts to consider the most general field redefinitions that cannot be reabsorbed by gauge transformations, as will be explained in section 4. Again, one may still object that considering arbitrary $\mathfrak{S}_\mu, \mathfrak{L}$ at tree level might be of use to cancel possible pathological divergences (i.e., that cannot be removed by field redefinitions or multiplicative renormalisation) appearing for $\mathfrak{S}_\mu^{\text{tree}} = \mathfrak{L}^{\text{tree}} = 0$. This possibility is precluded by the arguments presented in ref. [36], proven there for a specific model but expected to have general validity. In this reference the authors claim that, given a theory which is multiplicatively renormalisable, then by quantising the theory after performing a field redefinition, the divergences in terms of the new fields can be reabsorbed by the same multiplicative renormalisations of physical parameters as in the original case, plus infinite field redefinitions. In our case, we worry about possible divergences at order $h\theta$ for $\mathfrak{S}_\mu^{\text{tree}} = \mathfrak{L}^{\text{tree}} = 0$ which cannot be removed by infinite field redefinitions. The theory at order h^0 is known to be multiplicatively renormalisable, and considering arbitrary $\mathfrak{S}_\mu^{\text{tree}}, \mathfrak{L}^{\text{tree}}$ is equivalent to performing finite field redefinitions of order h on the ordinary fields $\mathfrak{a}_\mu, \mathfrak{l}, \bar{\mathfrak{l}}$. Thus, the additional divergences dependent on $\mathfrak{S}_\mu^{\text{tree}}, \mathfrak{L}^{\text{tree}}$ that might appear would be equivalent to infinite field redefinitions and therefore by assumption would not be useful to cancel the original problematic divergences at $\mathfrak{S}_\mu^{\text{tree}} = \mathfrak{L}^{\text{tree}} = 0$. It follows that the conclusions about the renormalisability of the theory obtained for $\mathfrak{S}_\mu^{\text{tree}} = \mathfrak{L}^{\text{tree}} = 0$ have a general validity.
- Computing a minimum number of diagrams. The use of the background field method guarantees that the result for the effective action will be gauge invariant. Furthermore, its divergent part computed in dimensional regularisation will be local. Thus, if one chooses a basis of all possible local gauge invariant terms up to order h , the divergent part of the effective action will be a linear combination of these terms. The coefficients in this linear combination can be determined by identifying its contributions with any given number and types of fields with the poles in the dimensional regularisation parameter ϵ of the corresponding 1PI Green functions with the same number and types of external fields. By appropriately choosing the basis, it can be guaranteed that the contributions to its elements with a minimum number of fields are also independent of each other, so that the unknown coefficients in the expansion of the divergent part of the effective action in terms of the basis can be determined from the diagrams with lowest number of fields.

We have thus argued that we can determine unambiguously the renormalisability of the theory by computing the effective action for $\alpha = 1, \mathfrak{S}_\mu^{\text{tree}} = \mathfrak{L}^{\text{tree}} = 0$. Under these assumptions, the Feynman rules relevant to our computations are those given in appendix B; they

use a compact notation for the Lie algebra indices, following ref. [37, 38], in which the U(N) field expansion in the Lie algebra generators in the fundamental representation is taken as

$$\mathbf{a}_\mu = \mathbf{a}_\mu^A T^A,$$

where $T^A = \{T^0, T^a\}$, with $T^0 = \frac{\mathbb{1}}{\sqrt{2N}}$ the U(1) generator and T^a denoting the SU(N) generators; more details are given in appendix A. This allows to compute simultaneously diagrams involving both SU(N) and U(1) fields, and the results for the SU(N) theory can also be easily obtained by setting the external “A” indices to SU(N) indices “a”, and by taking care to drop the contributions of U(1) indices in terms involving contractions of internal U(N) Lie algebra indices “A”.

Let us start by identifying the diagrams that need to be computed by constructing the appropriate basis of local gauge invariant terms whose integrals are independent. We use the decomposition in eq. (2.3). Local gauge invariant terms are then constructed from traced products of the field strengths and fermion fields and their covariant derivatives; we can classify them in three sectors: SU(N) sector -only including fields in the Lie algebra of SU(N)- U(1) sector, and mixed sector. A list follows:

- SU(N) sector:

$$\begin{aligned}
 t_1 &= \theta^{\alpha\beta} \text{Tr} f_{\alpha\beta} f_{\mu\nu} f^{\mu\nu}, & t_2 &= \theta^{\alpha\beta} \text{Tr} f_{\alpha\mu} f_{\beta\nu} f^{\mu\nu}, \\
 t_3 &= \theta^{\alpha\beta} \text{Tr} \bar{\lambda} \gamma_\alpha D^2 D_\beta \lambda, & t_4 &= \theta^{\alpha\beta} \text{Tr} \bar{\lambda} \gamma_{\alpha\beta} D^2 D_\mu \lambda, \\
 t_5 &= \theta^{\alpha\beta} \text{Tr} \bar{\lambda} \gamma^\mu \{f_{\mu\beta}, D_\alpha \lambda\}, & t_6 &= \theta^{\alpha\beta} \text{Tr} \bar{\lambda} \gamma^\mu \{f_{\alpha\beta}, D_\mu \lambda\}, \\
 t_7 &= \theta^{\alpha\beta} \text{Tr} \bar{\lambda} \gamma_\alpha \{f_{\beta\mu}, D^\mu \lambda\}, & t_8 &= \theta^{\alpha\beta} \text{Tr} \bar{\lambda} \gamma_{\alpha\beta} \{D^\nu f_{\mu\nu}, \lambda\}, \\
 t_9 &= \theta^{\alpha\beta} \text{Tr} \bar{\lambda} \gamma_\alpha^{\rho\sigma} \{D_\beta f_{\rho\sigma}, \lambda\}, & t_{10} &= \theta^{\alpha\beta} \text{Tr} \bar{\lambda} \gamma^\mu [D_\mu f_{\alpha\beta}, \lambda], \\
 t_{11} &= \theta^{\alpha\beta} \text{Tr} \bar{\lambda} \gamma_\alpha [D^\mu f_{\beta\mu}, \lambda], & t_{12} &= \theta^{\alpha\beta} \text{Tr} \bar{\lambda} \gamma_{\alpha\beta} [f_{\mu\nu}, D^\nu \lambda], \\
 t_{13} &= \theta^{\alpha\beta} \text{Tr} \bar{\lambda} \gamma_\alpha^{\rho\sigma} [f_{\rho\sigma}, D_\beta \lambda], & t_{14} &= \theta^{\alpha\beta} \text{Tr} \bar{\lambda} \gamma_\alpha^{\rho\sigma} [f_{\beta\sigma}, D_\rho \lambda], \\
 t_{15} &= \theta^{\alpha\beta} \text{Tr} \bar{\lambda}_i (\gamma_\alpha)_{ij} [\{\bar{\lambda}_k, (\gamma_\beta \lambda)_k\}, \lambda_j], & t_{16} &= \theta^{\alpha\beta} \text{Tr} \bar{\lambda}_i (\gamma^\mu)_{ij} [[\bar{\lambda}_k, (\gamma_{\mu\alpha\beta} \lambda)_k], \lambda_j].
 \end{aligned} \tag{3.1}$$

- U(1) sector:

$$\begin{aligned}
 u_1 &= \theta^{\alpha\beta} g_{\alpha\beta} g^{\mu\nu} g_{\mu\nu}, & u_2 &= \theta^{\alpha\beta} g_{\alpha\mu} g_{\beta\nu} g^{\mu\nu}, & u_3 &= \theta^{\alpha\beta} \bar{u} \gamma_\alpha \partial^2 \partial_\beta u, \\
 u_4 &= \theta^{\alpha\beta} \bar{u} \gamma_{\alpha\beta} \partial^2 \partial_\mu u, & u_5 &= \theta^{\alpha\beta} \bar{u} \gamma^\mu \partial_\alpha u g_{\mu\beta}, & u_6 &= \theta^{\alpha\beta} \bar{u} \gamma^\mu \partial_\mu u g_{\alpha\beta}, \\
 u_7 &= \theta^{\alpha\beta} \bar{u} \gamma_\alpha \partial^\mu u g_{\beta\mu}, & u_8 &= \theta^{\alpha\beta} \bar{u} \gamma_{\alpha\beta} u \partial^\nu g_{\mu\nu}, & u_9 &= \theta^{\alpha\beta} \bar{u} \gamma_\alpha^{\rho\sigma} u \partial_\beta g_{\rho\sigma}.
 \end{aligned} \tag{3.2}$$

- Mixed sector:

$$\begin{aligned}
 v_1 &= \theta^{\alpha\beta} \text{Tr} g_{\alpha\beta} f^{\mu\nu} f_{\mu\nu}, & v_2 &= \theta^{\alpha\beta} \text{Tr} g^{\mu\nu} f_{\alpha\mu} f_{\beta\nu}, & v_3 &= \theta^{\alpha\beta} \text{Tr} g_{\alpha\mu} f_{\beta\nu} f^{\mu\nu}, \\
 v_4 &= \theta^{\alpha\beta} \text{Tr} g_{\mu\nu} f_{\alpha\beta} f^{\mu\nu}, & v_5 &= \theta^{\alpha\beta} \text{Tr} \bar{u} \gamma^\mu f_{\mu\beta} D_\alpha \lambda, & v_6 &= \theta^{\alpha\beta} \text{Tr} \bar{u} \gamma^\mu f_{\alpha\beta} D_\mu \lambda, \\
 v_7 &= \theta^{\alpha\beta} \text{Tr} \bar{u} \gamma^\mu D_\mu f_{\alpha\beta} \lambda, & v_8 &= \theta^{\alpha\beta} \text{Tr} \bar{u} \gamma_\alpha f_{\beta\mu} D^\mu \lambda, & v_9 &= \theta^{\alpha\beta} \text{Tr} \bar{u} \gamma_\alpha D^\mu f_{\beta\mu} \lambda, \\
 v_{10} &= \theta^{\alpha\beta} \text{Tr} \bar{u} \gamma_{\alpha\beta} D^\nu f_{\mu\nu} \lambda, & v_{11} &= \theta^{\alpha\beta} \text{Tr} \bar{u} \gamma_{\alpha\beta} f_{\mu\nu} D^\nu \lambda, & v_{12} &= \theta^{\alpha\beta} \text{Tr} \bar{u} \gamma_\alpha^{\rho\sigma} D_\beta f_{\rho\sigma} \lambda,
 \end{aligned} \tag{3.3}$$

$$\begin{aligned}
v_{13} &= \theta^{\alpha\beta} \text{Tr} \bar{u} \gamma_\alpha^{\rho\sigma} f_{\rho\sigma} D_\beta \lambda, & v_{14} &= \theta^{\alpha\beta} \text{Tr} \bar{u} \gamma_\alpha^{\rho\sigma} f_{\beta\sigma} D_\rho \lambda, & v_{15} &= \theta^{\alpha\beta} \text{Tr} \bar{\lambda} \gamma^\mu D_\alpha \lambda g_{\mu\beta}, \\
v_{16} &= \theta^{\alpha\beta} \text{Tr} \bar{\lambda} \gamma^\mu D_\mu \lambda g_{\alpha\beta}, & v_{17} &= \theta^{\alpha\beta} \text{Tr} \bar{\lambda} \gamma_\alpha D^\mu \lambda g_{\beta\mu}, & v_{18} &= \theta^{\alpha\beta} \text{Tr} \bar{\lambda} \gamma_{\alpha\beta}{}^\mu \lambda \partial^\nu g_{\mu\nu}, \\
v_{19} &= \theta^{\alpha\beta} \text{Tr} \bar{\lambda} \gamma_\alpha^{\rho\sigma} \lambda \partial_\beta g_{\rho\sigma}.
\end{aligned}$$

In the formulae above, “Tr” denotes the trace over the SU(N) generators. The list of terms spans modulo total derivatives all the possible gauge invariant terms of order $h\theta^{\mu\nu}$ with the appropriate dimensions with zero or two Majorana fields. Again, the Majorana properties (A.3) and (A.4) have been used, so that any term with two Majorana fermions not present above can be expressed as a linear combination of the t_i, u_i and v_i , again modulo total derivatives. In the case of terms with four Majorana fermions, t_{15} and t_{16} do not span all the allowed contributions, but the missing ones will play no role in our calculations and we will safely ignore them.

The contributions to the previous list of terms with a minimum number of fields are independent of each other, which, as explained before, allows to fix the coefficients of the expansion of the divergent part of the effective action, Γ^{div} , in terms of the t_i, u_i, v_i by computing only the 1PI diagrams with the least possible number of fields. Let us identify the diagrams that need to be computed, using the notation in eq. (2.12) for the 1PI background Green functions.

At order h^0 , the possible gauge invariant terms are $\text{Tr} f_{\mu\nu} f^{\mu\nu}$ and $\text{Tr} \bar{\lambda} \not{D} \lambda$. Thus, using the notation of eq. (2.12) only the diagrams contributing to $\tilde{\Gamma}^{(2,0)}$ -with two external background gauge field legs- and $\tilde{\Gamma}^{(0,1)}$ -with two external quantum fermionic legs- need to be computed. At order h , we have, schematically, the following types of terms:

- Terms of the type $\text{Tr} \theta f f f, \text{Tr} \theta g f f, \theta g g g$, which are spanned by t_1, t_2, u_1, u_2 and $v_1 - v_4$ in eqs. (3.1), (3.2) and (3.3), whose contributions with three gauge fields are independent. Thus it suffices to compute diagrams with three external gauge fields, contributing to $\tilde{\Gamma}^{(3,0)}$.
- Terms of the type $\text{Tr} \theta \bar{\lambda} D^3 \lambda, \theta \bar{u} \partial^3 u$, which are spanned by t_3, t_4 and u_3, u_4 in eqs. (3.1) and (3.2). They involve at least two fermionic fields, so that their coefficients in the expansion of Γ^{div} can be fixed by computing $\tilde{\Gamma}^{(0,1)}$, which arises from diagrams with two fermionic legs.
- Terms of the type -neglecting ordering- $\text{Tr} \theta \bar{\lambda} D f \lambda, \text{Tr} \theta \bar{\lambda} f D \lambda, \theta \bar{u} \partial g u, \theta \bar{u} g \partial u, \text{Tr} \bar{u} D f \lambda, \text{Tr} \bar{u} f D \lambda, \text{Tr} \bar{\lambda} D \lambda g, \text{Tr} \bar{\lambda} \lambda \partial g$, which are spanned by $t_5 - t_{14}, u_5 - u_9, v_5 - v_{19}$ in eqs. (3.1), (3.2) and (3.3). Their contributions with one gauge field and two Majorana fields are again independent, so that it suffices to compute the diagrams contributing to $\tilde{\Gamma}^{(1,1)}$, i.e., with one background gauge field leg and two quantum fermionic legs.
- Terms of the type $\text{Tr} \theta \bar{\lambda} \lambda \bar{\lambda} \lambda$, such as t_{15}, t_{16} in eq. (3.1). Though t_{15}, t_{16} do not span all possibilities, it is clear that the computation of $\tilde{\Gamma}^{(0,2)}$ (diagrams with four external fermionic legs) will completely determine the corresponding contribution to the effective action Γ .




Figure 1. Diagrams contributing to $\tilde{\Gamma}^{(3,0)}$ at order h .

Summarising, at order h the only diagrams that have to be computed are those contributing to the 1PI Green functions $\tilde{\Gamma}^{(3,0)}$, $\tilde{\Gamma}^{(0,1)}$, $\tilde{\Gamma}^{(1,1)}$ and $\tilde{\Gamma}^{(0,2)}$. We proceed in the next sections, using dimensional regularisation at $D = 4 - 2\epsilon$ dimensions, with the Feynman rules displayed in appendix B. The calculations are quite involved and were done with the symbolic manipulation software `Mathematica`.

3.1 Commutative limit

Here we quote the known commutative result for the dimensionally regularised divergent part of the effective action:

$$\Gamma_{[U(N)]}^{\text{ord,div}} = \int d^D x - \frac{3g^2 N}{16\pi^2 \epsilon} \text{Tr} \left[-\frac{1}{2g^2} f_{\mu\nu} f^{\mu\nu} \right] + \int d^D x \frac{N}{16\pi^2 \epsilon} [i \text{Tr} \bar{\lambda} \not{D} \lambda]. \quad (3.4)$$

For simplicity, we suppressed the “” symbols with which we denoted the classical fields in section 2; we will keep doing so in the rest of the paper. Note that the divergent part only involves the $SU(N)$ fields a, λ , since the $U(1)$ sector is free in the commutative limit. In fact, since the $U(1)$ sector is free, in the $SU(N)$ case the result is identical,

$$\Gamma_{[SU(N)]}^{\text{ord,div}} = \Gamma_{[U(N)]}^{\text{ord,div}}. \quad (3.5)$$

3.2 Noncommutative contributions to $\tilde{\Gamma}^{(3,0)}$

The diagrams that contribute are shown in figure 1. Note that, though we did not provide in appendix B the Feynman rule for the vertex appearing in the first diagram, this diagram is directly zero since it involves an integral of the type

$$\int d^D l \frac{\prod_i l_{\mu_i}}{(l^2)^k}, \quad (3.6)$$

which vanishes in dimensional regularisation.

The results for the diagrams are too lengthy to be displayed here individually. We will quote the final expression for the contribution to the divergent part of the effective action




Figure 2. Diagrams contributing to $\tilde{\Gamma}^{(0,1)}$ at order h .

in position space:

$$i\tilde{\Gamma}_{[U(N)]}^{(3,0),\text{NC,div}}{}_{A_1, A_2, A_3}{}^{\mu_1, \mu_2, \mu_3} \mathbf{a}_{\mu_1}^{A_1} \mathbf{a}_{\mu_2}^{A_2} \mathbf{a}_{\mu_3}^{A_3} = \frac{3g^2 N h}{16\pi^2 \epsilon} \left[\frac{1}{4g^2} t_1 - \frac{1}{g^2} t_2 \right] \Big|_{aaa} \quad (3.7)$$

$$+ \frac{2g^2 N h}{16\pi^2 \epsilon} \left[\frac{1}{4g^2 \sqrt{2N}} (v_1 + 2v_4) - \frac{1}{g^2 \sqrt{2N}} (v_2 + 2v_3) \right] \Big|_{baa} + O(h^2),$$

where “ $|_{aaa}$ ” and “ $|_{baa}$ ” denote the contributions with lowest number of fields, i.e., three SU(N) gauge fields and one U(1) and two SU(N) gauge fields, respectively. Recall that the t_i, u_i, v_i are the gauge invariant terms defined in eqs. (3.1), (3.2) and (3.3). To get the SU(N) result, the external Lie algebra indices of the diagrams have to be set to SU(N) indices, and any U(1) contributions to internal contractions have to be eliminated. It turns out that all diagrams involve contractions of the type appearing in eq. (A.1) of appendix A, which, when setting the uncontracted indices to SU(N) indices, do not involve any contributions from internal U(1) indices. This is equivalent to saying that the U(1) fields do not run in the loops when the external fields are the a_μ^a . From this we conclude that the SU(N) result is obtained from eq. (3.7) by simply setting to zero the U(1) fields:

$$i\tilde{\Gamma}_{[SU(N)]}^{(3,0),\text{NC,div}}{}_{a_1, a_2, a_3}{}^{\mu_1, \mu_2, \mu_3} a_{\mu_1}^{a_1} a_{\mu_2}^{a_2} a_{\mu_3}^{a_3} = \frac{3g^2 N h}{16\pi^2 \epsilon} \left[\frac{1}{4g^2} t_1 - \frac{1}{g^2} t_2 \right] \Big|_{aaa} + O(h^2). \quad (3.8)$$

3.3 Noncommutative contributions to $\tilde{\Gamma}^{(0,1)}$

The diagrams contributing to the $\tilde{\Gamma}^{(0,1)}$ Green function at order θ are shown in figure 2. The first diagram is zero as it involves again an integral of the type shown in eq. (3.6). For external colour indices A, B, it is easily seen that the rest of the diagrams are zero since they are proportional to either $f^{ACD} d^{BCD} = 0$ or $f^{BCD} d^{ACD} = 0$. To get the SU(N) result one has to set the external indices to a, b and drop any U(1) contributions in the contractions of the internal indices. However, since $f^{bcd} d^{acd} = f^{bcd} d^{acd}$, no U(1) contributions must be eliminated, and the same argument as before applies. Therefore,

$$\tilde{\Gamma}_{[U(N)]}^{(0,1),\text{NC,div}} = \tilde{\Gamma}_{[SU(N)]}^{(0,1),\text{NC,div}} = O(h^2). \quad (3.9)$$

3.4 Noncommutative contributions to $\tilde{\Gamma}^{(1,1)}$

The diagrams contributing to the $\tilde{\Gamma}^{(1,1)}$ Green function at order θ are shown in figure 3. Again, we will write down the final result of the lengthy computation:

$$\frac{i}{2} \tilde{\Gamma}_{[U(N)]}^{(1,1),\text{NC,div}}{}_{A, B, C}{}^{i, j, \mu} \bar{t}_i^A t_j^B \mathbf{a}_\mu^C = - \frac{i N h}{16\pi^2 \epsilon} \left[\frac{1}{4} t_6 - \frac{1}{2} t_7 - \frac{1}{8} t_8 - \frac{1}{16} t_9 \right] \Big|_{a\lambda\lambda}$$




Figure 3. Diagrams contributing to $\tilde{\Gamma}^{(1,1)}$ at order h .

$$\begin{aligned}
 & + \frac{iNh}{16\pi^2\epsilon} \left(\frac{1}{\sqrt{2N}} \right) \left[v_5 - \frac{3}{2}v_6 + 2v_8 - \frac{1}{4}v_{10} - \frac{1}{2}v_{12} \right] \Big|_{a\bar{a}\lambda} \quad (3.10) \\
 & + \frac{iNh}{16\pi^2\epsilon} \left(\frac{1}{\sqrt{2N}} \right) \left[-v_{15} + \frac{1}{2}v_{16} + \frac{3}{4}v_{18} + \frac{3}{4}v_{19} \right] \Big|_{b\bar{b}\lambda} + O(h^2).
 \end{aligned}$$

To get the SU(N) result, using the same arguments as in the previous subsection it suffices to set the U(1) fields to zero:

$$\frac{i}{2} \tilde{\Gamma}_{[\text{SU}(N)]}^{(1,1),\text{NC,div}}{}_{a,b,c}{}^{i,j,\mu} \bar{\lambda}_i^a \lambda_j^b a_\mu^c = - \frac{iNh}{16\pi^2\epsilon} \left[\frac{1}{4}t_6 - \frac{1}{2}t_7 - \frac{1}{8}t_8 - \frac{1}{16}t_9 \right] \Big|_{a\bar{a}\lambda} + O(h^2). \quad (3.11)$$

3.5 Noncommutative contributions to $\tilde{\Gamma}^{(0,2)}$

The diagrams that contribute are shown in figure 4; it is easily seen that the box diagrams are finite since, though they would appear to be logarithmically divergent, one of the momenta in the noncommutative vertex is always external, as can be seen from the Feynman rule in appendix B. The final result is as follows:

$$\frac{i}{16} \tilde{\Gamma}_{[\text{U}(N)]}^{(0,2),\text{NC,div}}{}_{A,B,C,D}{}^{i,j,k,l} \bar{l}_i^A \bar{l}_k^C l_j^B l_l^D = - \frac{3iNh}{512\pi^2\epsilon} t_{16} + O(h^2). \quad (3.12)$$

Again, for external SU(N) fields no U(1) fields run in the loops, and the SU(N) result is identical.

$$\frac{i}{16} \tilde{\Gamma}_{[\text{SU}(N)]}^{(0,2),\text{NC,div}}{}_{a,b,c,d}{}^{i,j,k,l} \bar{\lambda}_i^a \bar{\lambda}_k^c \lambda_j^b \lambda_l^d = - \frac{3iNh}{512\pi^2\epsilon} t_{16} + O(h^2). \quad (3.13)$$




Figure 4. Diagrams contributing to $\tilde{\Gamma}^{(0,2)}$ at order h .

3.6 Final expression

From the previous discussions and the notation employed in the results of the 1PI Green functions in eqs. (3.7), (3.9), (3.10) and (3.12), which are expressed as the contributions with the lowest number of fields of linear combinations of the gauge invariant terms t_i, u_i, v_i of eqs. (3.1), (3.2) and (3.3), it is clear that the final result for the first-order noncommutative correction to the divergent part of the one-loop effective action is simply given by the integral of a sum of the t_i, u_i, v_i with the same coefficients as in eqs. (3.9), (3.7), (3.10) and (3.12):

$$\begin{aligned}
 \Gamma_{[\text{U}(N)]}^{\text{div,NC}} = & -h \int d^D x \left(\frac{3g^2 N}{16\pi^2 \epsilon} \left[\frac{1}{4g^2} t_1 - \frac{1}{g^2} t_2 \right] + \frac{2g^2 N}{16\pi^2 \epsilon} \left[\frac{1}{4g^2 \sqrt{2N}} (v_1 + 2v_4) - \frac{1}{g^2 \sqrt{2N}} (v_2 + 2v_3) \right] \right. \\
 & - \frac{iN}{16\pi^2 \epsilon} \left[\frac{1}{4} t_6 - \frac{1}{2} t_7 - \frac{1}{8} t_8 - \frac{1}{16} t_9 \right] + \frac{iN}{16\pi^2 \epsilon} \left(\frac{1}{\sqrt{2N}} \right) \left[v_5 - \frac{3}{2} v_6 + 2v_8 - \frac{1}{4} v_{10} - \frac{1}{2} v_{12} \right. \\
 & \left. \left. - v_{15} + \frac{1}{2} v_{16} + \frac{3}{4} v_{18} + \frac{3}{4} v_{19} \right] - \frac{3iN}{512\pi^2 \epsilon} t_{16} \right) + O(h^2). \tag{3.14}
 \end{aligned}$$

Similarly, the SU(N) result obtained from the expressions in eqs. (3.8), (3.9), (3.11) and (3.13) is

$$\begin{aligned}
 \Gamma_{[\text{SU}(N)]}^{\text{div,NC}} = & -h \int d^D x \left(\frac{3g^2 N}{16\pi^2 \epsilon} \left[\frac{1}{4g^2} t_1 - \frac{1}{g^2} t_2 \right] - \frac{iN}{16\pi^2 \epsilon} \left[\frac{1}{4} t_6 - \frac{1}{2} t_7 - \frac{1}{8} t_8 - \frac{1}{16} t_9 \right] - \frac{3iN}{512\pi^2 \epsilon} t_{16} \right) \\
 & + O(h^2). \tag{3.15}
 \end{aligned}$$

Equivalently, substituting the expressions in eqs. (3.1), (3.2) and (3.3), and adding the commutative contribution of eq. (3.4), we arrive to the following formula for the one-loop

divergent part of the effective action at first order in the noncommutative parameters

$$\begin{aligned}
 \Gamma_{[U(N)]}^{\text{div}} = & - \int d^D x \left(\frac{3g^2 N}{16\pi^2 \epsilon} \text{Tr} \left[-\frac{1}{2g^2} f_{\mu\nu} f^{\mu\nu} + \frac{h}{4g^2} \theta^{\mu\nu} f_{\alpha\beta} f_{\mu\nu} f^{\mu\nu} - \frac{h}{g^2} \theta^{\alpha\beta} f_{\alpha\mu} f_{\beta\mu} f^{\mu\nu} \right] \right. \\
 & - \frac{N}{16\pi^2 \epsilon} [i \text{Tr} \bar{\lambda} \not{D} \lambda] \\
 & + \frac{2g^2 N h}{16\pi^2 \epsilon} \text{Tr} \left[\frac{1}{4g^2 \sqrt{2N}} \theta^{\alpha\beta} (g_{\alpha\beta} f^{\mu\nu} f_{\mu\nu} + 2g_{\mu\nu} f_{\alpha\beta} f^{\mu\nu}) - \frac{1}{g^2 \sqrt{2N}} \theta^{\alpha\beta} (g^{\mu\nu} f_{\alpha\mu} f_{\beta\nu} + 2g_{\alpha\mu} f_{\beta\mu} f^{\mu\nu}) \right] \\
 & - \frac{N h}{16\pi^2 \epsilon} \text{Tr} \left[\frac{i}{4} \theta^{\alpha\beta} \bar{\lambda} \gamma^\mu \{f_{\alpha\beta}, D_\mu \lambda\} - \frac{i}{2} \theta^{\alpha\beta} \bar{\lambda} \gamma_\alpha \{f_{\beta\mu}, D^\mu \lambda\} - \frac{i}{8} \theta^{\alpha\beta} \bar{\lambda} \gamma_{\alpha\beta}{}^\mu \{D^\nu f_{\mu\nu}, \lambda\} \right. \\
 & - \left. \frac{i}{16} \theta^{\alpha\beta} \bar{\lambda} \gamma_\alpha{}^{\rho\sigma} \{D_\beta f_{\rho\sigma}, \lambda\} \right] + \frac{N h}{16\pi^2 \epsilon} \left(\frac{1}{\sqrt{2N}} \right) \text{Tr} \left[i \theta^{\alpha\beta} \bar{u} \gamma^\mu f_{\mu\beta} D_\alpha \lambda - \frac{3}{2} i \theta^{\alpha\beta} \bar{u} \gamma^\mu f_{\alpha\beta} D_\mu \lambda \right. \\
 & + \left. 2i \theta^{\alpha\beta} \bar{u} \gamma_\alpha f_{\beta\mu} D^\mu \lambda - \frac{i}{4} \theta^{\alpha\beta} \bar{u} \gamma_{\alpha\beta}{}^\mu D^\nu f_{\mu\nu} \lambda - \frac{i}{2} \theta^{\alpha\beta} \bar{u} \gamma_\alpha{}^{\rho\sigma} D_\beta f_{\rho\sigma} \lambda \right] \quad (3.16) \\
 & + \frac{N h}{16\pi^2 \epsilon} \left(\frac{1}{\sqrt{2N}} \right) \text{Tr} \left[-i \theta^{\alpha\beta} \bar{\lambda} \gamma^\mu D_\alpha \lambda g_{\mu\beta} + \frac{i}{2} \theta^{\alpha\beta} \bar{\lambda} \gamma^\mu D_\mu \lambda g_{\alpha\beta} + \frac{3i}{4} \theta^{\alpha\beta} \bar{\lambda} \gamma_{\alpha\beta}{}^\mu \lambda \partial^\nu g_{\mu\nu} \right. \\
 & \left. + \frac{3i}{4} \theta^{\alpha\beta} \bar{\lambda} \gamma_\alpha{}^{\rho\sigma} \lambda \partial_\beta g_{\rho\sigma} \right] - \frac{3i N h}{512\pi^2 \epsilon} \theta^{\alpha\beta} \text{Tr} \bar{\lambda}_i (\gamma^\mu)_{ij} [[\bar{\lambda}_k, (\gamma_{\mu\alpha\beta} \lambda)_k], \lambda_j] \Big) + O(h^2).
 \end{aligned}$$

The corresponding expression in the SU(N) case is

$$\begin{aligned}
 \Gamma_{[SU(N)]}^{\text{div}} = & - \int d^D x \left(\frac{3g^2 N}{16\pi^2 \epsilon} \text{Tr} \left[-\frac{1}{2g^2} f_{\mu\nu} f^{\mu\nu} + \frac{h}{4g^2} \theta^{\mu\nu} f_{\alpha\beta} f_{\mu\nu} f^{\mu\nu} - \frac{h}{g^2} \theta^{\alpha\beta} f_{\alpha\mu} f_{\beta\mu} f^{\mu\nu} \right] \right. \\
 & - \frac{N}{16\pi^2 \epsilon} [i \text{Tr} \bar{\lambda} \not{D} \lambda] \\
 & - \frac{N h}{16\pi^2 \epsilon} \text{Tr} \left[\frac{i}{4} \theta^{\alpha\beta} \bar{\lambda} \gamma^\mu \{f_{\alpha\beta}, D_\mu \lambda\} - \frac{i}{2} \theta^{\alpha\beta} \bar{\lambda} \gamma_\alpha \{f_{\beta\mu}, D^\mu \lambda\} - \frac{i}{8} \theta^{\alpha\beta} \bar{\lambda} \gamma_{\alpha\beta}{}^\mu \{D^\nu f_{\mu\nu}, \lambda\} \right. \\
 & \left. - \frac{i}{16} \theta^{\alpha\beta} \bar{\lambda} \gamma_\alpha{}^{\rho\sigma} \{D_\beta f_{\rho\sigma}, \lambda\} \right] - \frac{3i N h}{512\pi^2 \epsilon} \theta^{\alpha\beta} \text{Tr} \bar{\lambda}_i (\gamma^\mu)_{ij} [[\bar{\lambda}_k, (\gamma_{\mu\alpha\beta} \lambda)_k], \lambda_j] \Big) + O(h^2). \quad (3.17)
 \end{aligned}$$

It is worth noting that, for $N = 2$, all the terms with SU(N) fields whose traces yield factors $d^{abc} = 2\text{Tr} T^a \{T^b, T^c\}$ in eqs. (3.16) and eq. (3.17) vanish. This means that all terms involving only SU(N) fields vanish; in the U(2) case, we are only left with SU(2)-U(1) mixed terms, while for the SU(2) theory the noncommutative divergences disappear. This fact is independent of the representation considered since, for a representation R of SU(N), $\text{Tr}_R T^a \{T^b, T^c\} \propto \text{Tr}_F T^a \{T^b, T^c\}$.

In the U(1) case, it is also clear from eqs. (3.16) and eq. (3.17) that, as in the SU(2) theory, the divergent part of the effective action reduces to its commutative counterpart.

4 Analysing renormalisability

In this section we will analyse whether the divergences in the effective actions, given in eqs. (3.16) and (3.17), can be subtracted from appropriate multiplicative renormalisations of fields and parameters and infinite shifts on the Seiberg-Witten map ambiguities \mathfrak{S}, L

-see eq. (2.2). We will use the minimal subtraction scheme. The counterterms in the action that cancel the divergences of the effective action are trivially given by

$$\int d^D x \mathcal{L}^{ct} = - \int d^D x \Gamma^{\text{div}}.$$

Were the theory to be renormalisable, these counterterms would arise from multiplicative renormalisation and from ambiguities of the SW maps, which, as will be argued, are equivalent to field redefinitions. We define the multiplicative renormalisation as

$$a_\mu = Z_a^{1/2} a_\mu^R, \quad b_\mu = Z_b^{1/2} b_\mu^R, \quad \lambda = Z_\lambda^{1/2} \lambda^R, \quad u = Z_u^{1/2} u^R, \quad g = Z_g g^R, \quad h = Z_h h^R, \quad (4.1)$$

with $Z_i = 1 + \delta Z_i$. It is easily seen that gauge invariance forces

$$\delta Z_a = 0.$$

On the other hand, the SW map ambiguities at order h , \mathfrak{S}_μ and \mathfrak{L} , are given by terms transforming in the adjoint representation of the gauge group, with the appropriate mass dimensions and index structure, involving a contraction with $\theta^{\alpha\beta}$; under a $U(N)$ gauge transformation of the fields with gauge parameter $\mathfrak{c} = c^a T^a + \frac{1}{\sqrt{2N}} C$, they transform as

$$s\mathfrak{S}_\mu = i[c^a T^a, \mathfrak{S}_\mu], \quad s\mathfrak{L} = i[c^a T^a, \mathfrak{L}].$$

We restrict ourselves to ambiguities that respect the parity transformation properties of the fields; doing so is justified since considering additional types of field redefinitions would yield terms in the modified action involving an odd number of $\epsilon^{\mu\nu\rho\sigma}$ tensors, which do not appear in the divergent parts of the effective actions given in eqs. (3.16) and (3.17) and thus need not be considered when checking renormalisability. The most general solution we found satisfying the specified requirements is of the form

$$\mathfrak{S}_\mu = S_\mu + T_\mu \mathbb{1}, \quad \mathfrak{L} = L + M \mathbb{1}, \quad (4.2)$$

such that

$$\begin{aligned} S_\mu &= y_1 \theta^{\alpha\beta} D_\mu f_{\alpha\beta} + y_2 \theta_\mu^\alpha D^\nu f_{\nu\alpha} + y_3 \theta_\mu^\alpha \{\bar{\lambda}_i, (\gamma_\alpha \lambda)_i\} + i y_4 \theta^{\alpha\beta} [\bar{\lambda}_i, (\gamma_{\mu\alpha\beta} \lambda)_i] + i y_5 \theta_\mu^\alpha \bar{u} \gamma_\alpha \lambda \\ &\quad + y_6 \theta^{\rho\sigma} \bar{u} \gamma_{\mu\rho\sigma} \lambda, \quad y_k \in \mathbb{R}, \\ L &= k_1 \theta^{\alpha\beta} \{f_{\alpha\beta}, \lambda\} + k_2 \theta^{\alpha\beta} D^2 \lambda + k_3 \theta^{\alpha\beta} \gamma_\alpha^\mu [f_{\beta\mu}, \lambda] + k_4 \theta^{\alpha\beta} \gamma_\alpha^\mu \{f_{\beta\mu}, \lambda\} + k_5 \theta^{\alpha\beta} \gamma_\alpha^\mu \{D_\mu, D_\beta\} \lambda \\ &\quad + k_6 \tilde{\theta}^{\alpha\beta} \gamma_5 \{f_{\alpha\beta}, \lambda\} + k_7 \tilde{\theta}^{\alpha\beta} \gamma_5 [f_{\alpha\beta}, \lambda] + k_8 \theta^{\alpha\beta} g_{\alpha\beta} \lambda + k_9 \theta^{\alpha\beta} \gamma_\alpha^\mu g_{\beta\mu} \lambda + k_{10} \tilde{\theta}^{\alpha\beta} \gamma_5 g_{\alpha\beta} \lambda \\ &\quad + k_{11} \theta^{\alpha\beta} f_{\alpha\beta} u + k_{12} \theta^{\alpha\beta} \gamma_\alpha^\mu f_{\beta\mu} u + k_{13} \tilde{\theta}^{\alpha\beta} \gamma_5 f_{\alpha\beta} u, \quad k_i \in \mathbb{C}, \\ T_\mu &= z_1 \theta^{\alpha\beta} \partial_\mu g_{\alpha\beta} + z_2 \theta_\mu^\alpha \partial^\nu g_{\nu\alpha} + i z_3 \theta^{\alpha\beta} \bar{u} \gamma_{\mu\alpha\beta} u + i z_4 \text{Tr} \theta^{\alpha\beta} [\bar{\lambda}_i, (\gamma_{\mu\alpha\beta} \lambda)_i], \quad z_k \in \mathbb{R}, \\ M &= l_1 \theta^{\alpha\beta} g_{\alpha\beta} u + l_2 \theta^{\alpha\beta} \partial^2 u + l_3 \theta^{\alpha\beta} \gamma_\alpha^\mu g_{\beta\mu} u + l_4 \theta^{\alpha\beta} \gamma_\alpha^\mu \partial_\mu \partial_\beta u + l_5 \tilde{\theta}^{\alpha\beta} \gamma_5 g_{\alpha\beta} u + l_6 \text{Tr} \theta^{\alpha\beta} \{f_{\alpha\beta}, \lambda\} \\ &\quad + l_7 \text{Tr} \theta^{\alpha\beta} \gamma_\alpha^\mu \{f_{\beta\mu}, \lambda\} + l_8 \text{Tr} \tilde{\theta}^{\alpha\beta} \gamma_5 \{f_{\alpha\beta}, \lambda\}, \quad l_i \in \mathbb{C}, \end{aligned} \quad (4.3)$$

where $\tilde{\theta}^{\alpha\beta} = \frac{1}{2} \epsilon^{\alpha\beta\rho\sigma} \theta_{\rho\sigma}$. In the $U(N)$ case, since the enveloping algebra coincides with the Lie algebra, the previous ambiguities are equivalent to field redefinitions of b_μ, a_μ, λ, u . The ambiguities in the $SU(N)$ case are obtained by setting $b_\mu = u = 0$. Since in principle there

still remain contributions along the identity operator after setting $b_\mu = u = 0$ in eq. (4.3), it would seem that the $SU(N)$ ambiguities are not equivalent to field redefinitions, which would invalidate our arguments concerning the possibility of setting the ambiguities to zero at tree-level without losing generality when dealing with the renormalisability of the theory. However, in the $SU(N)$ case it is easily seen that the contributions to the ambiguities along the identity, coming from the terms of eq. (4.3) proportional to $y_4, k_1, k_4, k_6, z_4, l_6, l_7, l_8$, do not yield modifications of the action at order \hbar , so that these contributions can be ignored and the ambiguities can be thought as Lie algebra valued and thus equivalent to field redefinitions. Note that the field redefinitions proportional to y_4, k_1, k_4, k_6 not only yield contributions along the identity but also on the Lie Algebra, and thus still have to be taken into account.

4.1 Commutative renormalisation

The $U(N)$ and $SU(N)$ $-N > 1$ - theories at order $\hbar = 0$ are multiplicatively renormalisable; the divergences appearing in eq. (3.4) -see also eq. (3.5)- can be absorbed by the following values of the renormalisation constants in eq. (4.1):

$$\delta Z_a = 0, \quad \delta Z_g = -\frac{3g^2 N}{32\pi^2 \epsilon}, \quad \delta Z_b = -\frac{3g^2 N}{16\pi^2 \epsilon}, \quad \delta Z_\lambda = -\frac{g^2 N}{4\pi^2 \epsilon}, \quad \delta Z_u = -\frac{3g^2 N}{16\pi^2 \epsilon}. \quad (4.4)$$

4.2 Renormalisation of the noncommutative bosonic sector

Starting with the $U(N)$ case, $N > 1$, let us consider the order \hbar noncommutative divergences only involving gauge fields in eq. (3.16). A key issue is that the ambiguities in the SW map given in eq. (4.3), when introduced in the action by means of eqs. (4.2), (4.3), (2.2) and (2.1), do not generate any purely bosonic terms. Thus the purely bosonic divergences can only be renormalised, if at all, by means of multiplicative renormalisations. In terms of the basis of gauge invariant terms given in eqs. (3.1), (3.2) and (3.3), the tree-level noncommutative contribution to the bosonic part of the action is

$$S_{[U(N)]_{\text{bos}}}^{\text{tree,NC}} = \int d^D x \left(\frac{\hbar}{4g^2} t_1 - \frac{\hbar}{g^2} t_2 + \frac{\hbar}{8g^2 \sqrt{2N}} u_1 - \frac{\hbar}{2g^2 \sqrt{2N}} u_2 + \frac{\hbar}{4g^2 \sqrt{2N}} (v_1 + 2v_4) - \frac{\hbar}{g^2 \sqrt{2N}} (v_2 + 2v_3) \right) + O(\hbar^2), \quad (4.5)$$

so that the counterterm action originated by the multiplicative renormalisations of eq. (4.1) would be, keeping in mind $\delta Z_a = 0$ and suppressing the “ R ” superindices for simplicity:

$$S_{[U(N)]_{\text{bos}}}^{\text{ct,NC}} = \int d^D x \left((-2\delta Z_g + \delta Z_h) \left(\frac{\hbar}{4g^2} t_1 - \frac{\hbar}{g^2} t_2 \right) + \left(-2\delta Z_g + \delta Z_h + \frac{3}{2} \delta Z_b \right) \left(\frac{\hbar}{8g^2 \sqrt{2N}} u_1 - \frac{\hbar}{2g^2 \sqrt{2N}} u_2 \right) + \left(-2\delta Z_g + \delta Z_h + \frac{1}{2} \delta Z_b \right) \left(\frac{\hbar}{4g^2 \sqrt{2N}} (v_1 + 2v_4) - \frac{\hbar}{g^2 \sqrt{2N}} (v_2 + 2v_3) \right) \right) + O(\hbar^2),$$

which should be made equivalent with minus the bosonic part of the divergent part of the effective action in eq. (3.14)

$$\Gamma_{[\text{U}(N)]^{\text{bos}}}^{\text{div,NC}} = - \int d^D x \left(\frac{3g^2 N}{16\pi^2 \epsilon} \left[\frac{h}{4g^2} t_1 - \frac{h}{g^2} t_2 \right] + \frac{2g^2 N}{16\pi^2 \epsilon} \left[\frac{h}{4g^2 \sqrt{2N}} (v_1 + 2v_4) - \frac{h}{g^2 \sqrt{2N}} (v_2 + 2v_3) \right] \right) + O(h^2). \quad (4.6)$$

For $N \geq 3$, for which the terms t_1 and t_2 are nonzero, this forces the three following identities,

$$\begin{aligned} \frac{3g^2 N}{16\pi^2 \epsilon} &= -2\delta Z_g + \delta Z_h, \\ 0 &= -2\delta Z_g + \delta Z_h + \frac{3}{2}\delta Z_b, \\ -\frac{2g^2 N}{16\pi^2 \epsilon} &= -2\delta Z_g + \delta Z_h + \frac{1}{2}\delta Z_b. \end{aligned} \quad (4.7)$$

Using the commutative results in eq. (4.4), the first equation implies

$$\delta Z_h = 0,$$

as has been obtained for a number of other noncommutative theories, but then the second and third identities in eq. (4.7) are not satisfied. In the $N = 2$ case, only the last two identities are relevant, since $t_1 = t_2 = 0$; again, they are incompatible with the $O(h^0)$ results of eq. (4.4).

From this we arrive to the first conclusions of our paper: the $\text{U}(N)$ theory is not renormalisable, for $N > 1$. In principle we have derived this only for our choice of gauge-fixing; to extend the result for arbitrary gauge-fixing, we have to consider the on-shell divergences, which are independent of the gauge-fixing. The equations of motion are of the form

$$\begin{aligned} (D_\mu f^{\mu\nu}) &= \frac{1}{2} \{ \bar{\lambda}_i, \gamma_{ij}^\nu \lambda_j \} + O(h), \quad \partial_\mu g^{\mu\nu} = O(h), \quad \not{D}u = O(h), \\ (\not{D}\lambda)^a &= \frac{h}{2} \theta^{\alpha\beta} \text{Tr} T^a \gamma^\mu \{ D_\mu \lambda, f_{\alpha\beta} \} + h \theta^{\alpha\beta} \text{Tr} T^a \gamma^\mu \{ D_\beta \lambda, f_{\mu\alpha} \} + O(h^2), \end{aligned} \quad (4.8)$$

where the details of the $O(h)$ part in the equations in the first line of (4.8) will not be relevant to our purposes. Since the bosonic divergences of eq. (4.6) –see eqs. (3.1) and (3.3)– do not involve covariant derivatives of field strenghts, it is easy to see that the on-shell conditions of eq. (4.8) cannot be used to relate the bosonic divergences in eq. (4.6) among themselves or with the fermionic divergences. Thus the on-shell bosonic divergences have the same form as in eq. (4.6) and the same conclusions about nonrenormalisability apply.

However, in the large N limit, $N \rightarrow \infty$ while keeping the 't Hooft coupling $g^2 N$ finite, both the tree-level contributions in eq. (4.5) and the problematic divergences in eq. (4.6) associated with the u_i terms are subleading, so that the second and third identities in eq. (4.7) need not be considered, and therefore the gauge sector is renormalisable in this limit with $\delta Z_h = 0$, in keeping with the expectations raised by the quantum behaviour of the SW duals of the NC $\text{U}(N)$ theories in the enveloping algebra approach.

In the $SU(N)$ case, $N \geq 3$, we only have the divergences coming from the t_i terms, which are multiplicatively renormalisable with $\delta Z_h = 0$; thus, the $SU(N)$ gauge sector is renormalisable, as has been obtained already for other NC theories in the enveloping algebra approach with different matter content [20–25].

It remains to examine the renormalisability of the gaugino sector. Given the previous conclusions, it suffices to study only the large N limit of $U(N)$ or the $SU(N)$ theory for $N \geq 3$, since the $U(N)$ theory fails to be renormalisable for finite $N > 1$.

4.3 Renormalisation of the noncommutative gaugino sector

In the large N limit, given the decomposition in eq. (2.3), the tree-level interactions involving $U(1)$ fields are subleading with respect to those of $SU(N)$ fields, so that at leading order in N the $U(1)$ fields are free. This is also reflected at the quantum level, since the divergences involving $U(1)$ fields in the effective action in eq. (3.16) are subleading. Thus, for large N the $U(1)$ fields can be neglected, and the problem of renormalisability is the same as for the $SU(N)$ theory. The tree level part of the $SU(N)$ action involving gaugino fields and taking into account the SW map ambiguities of eq. (4.3) restricted to the $SU(N)$ case is given, for $N \geq 3$ and in the basis of gauge invariant terms of eq. (3.1), by

$$S_{[SU(N)]\text{gaugino}}^{\text{tree,NC}} = \frac{h}{g^2} \int d^D x \sum_{i=3}^{16} C_i t_i + O(h^2),$$

$$\begin{aligned} C_3 &= -4i\text{Re}k_5 - 4i\text{Re}k_2, & C_4 &= -2\text{Im}k_2, \\ C_5 &= \frac{i}{2} - 2i\text{Re}k_4, & C_6 &= -\frac{i}{4} + 2i\text{Re}k_1, \\ C_7 &= -2i\text{Re}k_4, & C_8 &= -2iy_4 + \text{Re}k_6 + i\text{Im}k_6, \\ C_9 &= -\frac{i}{2}\text{Re}k_4 + \text{Re}k_6 + i\text{Im}k_6, & C_{10} &= \frac{1}{2}(2y_1 + \text{Im}k_3 - \text{Re}k_5), \\ C_{11} &= \frac{1}{2}(4y_3 - 2y_2 + 2\text{Im}k_3 - 2\text{Re}k_5), & C_{12} &= 2\text{Re}k_2 + 2i\text{Im}k_2 + 2\text{Re}k_6 + 2\text{Re}k_7, \\ C_{13} &= -2\text{Re}k_5 + 2\text{Re}k_6 + 2\text{Re}k_7, & C_{14} &= 2\text{Im}k_3 - 2\text{Re}k_5, \\ C_{15} &= y_3, & C_{16} &= iy_4. \end{aligned}$$

The previous formulae follow from eq. (2.1), the SW map in eq. (2.2) for $\mathbf{a}_\mu = a_\mu$ and the ambiguities of eq. (4.3) for $T_\mu = M = b_\mu = u = 0$. The Dirac algebra identities in eq. (A.2) were extensively used. The counterterm Lagrangian generated by the multiplicative renormalisations of fields and parameters of eq. (4.1) and by infinite shifts of the ambiguity parameters $y_i = y_i^R + \delta y_i^R$, $k_i = k_i^R + \delta k_i^R$ is given, for $y_i^R = k_i^R = 0$ -recall that we computed the divergences in the effective action for zero values of the SW map ambiguities-, by

$$S_{[SU(N)]\text{gaugino}}^{\text{ct,NC}} = \frac{h}{g^2} \int d^D x \sum_{i=3}^{16} (\delta C_i) t_i + O(h^2), \quad (4.9)$$

$$\delta C_3 = -4i\delta\text{Re}k_5 - 4i\delta\text{Re}k_2, \quad \delta C_4 = -2\delta\text{Im}k_2,$$

$$\begin{aligned}
 \delta C_5 &= \frac{i}{2}(-2\delta Z_g + \delta Z_\lambda) - 2i\delta \text{Re}k_4, & \delta C_6 &= -\frac{i}{4}(-2\delta Z_g + \delta Z_\lambda) + 2i\delta \text{Re}k_1, \\
 \delta C_7 &= -2i\delta \text{Re}k_4, & \delta C_8 &= -2i\delta y_4 + \delta \text{Re}k_6 + i\delta \text{Im}k_6, \\
 \delta C_9 &= -\frac{i}{2}\delta \text{Re}k_4 + \delta \text{Re}k_6 + i\delta \text{Im}k_6, & \delta C_{10} &= \frac{1}{2}(2\delta y_1 + \delta \text{Im}k_3 - \delta \text{Re}k_5), \\
 \delta C_{11} &= \frac{1}{2}(4\delta y_3 - 2\delta y_2 + 2\delta \text{Im}k_3 - 2\delta \text{Re}k_5), & \delta C_{12} &= 2\delta \text{Re}k_2 + 2i\delta \text{Im}k_2 + 2\delta \text{Re}k_6 + 2\delta \text{Re}k_7, \\
 \delta C_{13} &= -2\delta \text{Re}k_5 + 2\delta \text{Re}k_6 + 2\delta \text{Re}k_7, & \delta C_{14} &= 2\delta \text{Im}k_3 - 2\delta \text{Re}k_5, \\
 \delta C_{15} &= \delta y_3, & \delta C_{16} &= i\delta y_4,
 \end{aligned}$$

where the “ R ” superindices have been suppressed for simplicity and the result $\delta Z_h = 0$ of the previous subsection was used.

For the theory to be renormalisable, the previous counterterm action has to be matched with minus the divergent part of the effective action in eqs. (3.15) and (3.17) involving gaugino fields. This contribution involving fermion fields, expressed in the basis of gauge invariant terms of eq. (3.1), is

$$\Gamma_{\text{gaugino}, [\text{SU}(N)]}^{\text{div, NC}} = \int d^D x \left(\frac{iNh}{16\pi^2\epsilon} \left[\frac{1}{4}t_6 - \frac{1}{2}t_7 - \frac{1}{8}t_8 - \frac{1}{16}t_9 \right] + \frac{3iNh}{512\pi^2\epsilon} t_{16} \right) + O(h^2). \quad (4.10)$$

Matching eqs. (4.9) and (4.10), we get the following system of equations:

$$\begin{aligned}
 t_3 : \delta \text{Re}k_5 + \delta \text{Re}k_2 &= 0, & t_4 : \delta \text{Im}k_2 &= 0, \\
 t_5 : \frac{i}{2}(-2\delta Z_g + \delta Z_\lambda) - 2i\delta \text{Re}k_4 &= 0, & t_6 : \frac{i}{4}(-2\delta Z_g + \delta Z_\lambda) - 2i\delta \text{Re}k_1 &= \frac{iNg^2}{64\pi^2\epsilon}, \\
 t_7 : -2i\delta \text{Re}k_4 = \frac{iNg^2}{32\pi^2\epsilon}, & & t_8 : -2i\delta y_4 + \delta \text{Re}k_6 + i\delta \text{Im}k_6 &= \frac{iNg^2}{128\pi^2\epsilon}, \\
 t_9 : -\frac{i}{2}\delta \text{Re}k_4 + \delta \text{Re}k_6 + i\delta \text{Im}k_6 = \frac{iNg^2}{256\pi^2\epsilon}, & t_{10} : 2\delta y_1 + \delta \text{Im}k_3 - \delta \text{Re}k_5 &= 0, \\
 t_{11} : 4\delta y_3 - 2\delta y_2 + 2\delta \text{Im}k_3 - 2\delta \text{Re}k_5 &= 0, & t_{12} : \delta \text{Re}k_2 + i\delta \text{Im}k_2 + \delta \text{Re}k_6 + \delta \text{Re}k_7 &= 0, \\
 t_{13} : -\delta \text{Re}k_5 + \delta \text{Re}k_6 + \delta \text{Re}k_7 &= 0, & t_{14} : \delta \text{Im}k_3 - \delta \text{Re}k_5 &= 0, \\
 t_{15} : \delta y_3 = 0, & & t_{16} : i\delta y_4 = -\frac{3iNg^2}{512\pi^2\epsilon}. & \quad (4.11)
 \end{aligned}$$

Projecting into real and imaginary parts, there are 17 real equations for 13 real variables,

$$\delta y_1, \delta y_2, \delta y_3, \delta y_4, \delta \text{Re}k_1, \delta \text{Re}k_2, \delta \text{Im}k_2, \delta \text{Im}k_3, \delta \text{Re}k_4, \delta \text{Re}k_5, \delta \text{Re}k_6, \delta \text{Im}k_6, \delta \text{Re}k_7.$$

Remarkably, the equations are not independent and there is a one-paramter family of solutions,

$$\begin{aligned}
 \delta y_1 = \delta y_2 = \delta y_3 = \delta \text{Im}k_2 = \delta \text{Re}k_6 &= 0, \\
 \delta y_4 &= -\frac{3g^2 N}{512\pi^2\epsilon}, \\
 \delta \text{Im}k_3 = \delta \text{Re}k_7 = -\delta \text{Re}k_2 = \delta \text{Re}k_5,
 \end{aligned}$$

$$\delta \text{Re}k_1 = \delta \text{Re}k_4 = -\frac{g^2 N}{64\pi^2 \epsilon},$$

$$\delta \text{Im}k_6 = -\frac{Ng^2}{256\pi^2 \epsilon}.$$

This shows that, at one-loop and order θ , the $U(N)$ theory in the large N limit and the $SU(N)$ theory are renormalisable.

5 Conclusions

In this paper we have calculated the $O(\theta)$ divergent part of the background field effective action for the classical dual under the Seiberg-Witten map of noncommutative $\mathcal{N} = 1$ $U(N)$ super Yang-Mills, as well as for the $SU(N)$ theory that results from suppressing the $U(1)$ degrees of freedom in the former $U(N)$ theory. Our results can be summarised as follows: the quantisation of the classical dual under the Seiberg-Witten map of $\mathcal{N} = 1$ $U(N)$, $N > 1$, super Yang-Mills yields an $U(N)$ supersymmetric ordinary quantum theory that is not renormalisable –and neither is its gauge sector– for finite values of N . In the large N limit, however, the $U(N)$ theory remarkably becomes renormalisable. On the other hand, the $SU(N)$ theory for arbitrary $N > 2$ also turns out to be renormalisable.

That both –i.e., the standard, quantised in terms of noncommutative fields, and the θ -expanded, defined by means of the Seiberg-Witten map– large N supersymmetric $U(N)$ theories are one-loop renormalisable and have the same running coupling constant hints at the fact that the classical duality between noncommutative theories established by the Seiberg-Witten map may survive at the quantum level: as was mentioned in the introduction, the noncommutative $U(N)$ super Yang-Mills theory, at one-loop and large N , is renormalisable and has a smooth $\theta^{\mu\nu} \rightarrow 0$ limit, and this behaviour is reproduced in the dual theory formulated by means of the Seiberg-Witten map. This is in agreement with previous studies regarding the survival at the quantum level of the Seiberg-Witten map duality, although they focused on UV finite theories such as noncommutative Chern-Simons [28].

On the other hand, the $SU(N)$ result represents the first case in the literature in which a noncommutative theory in the enveloping algebra approach involving fermion fields turns out to be one-loop renormalisable at order θ . This could be attributed to the consideration of Majorana fermions and a noncommutative covariant derivative involving a star product commutator, or perhaps, more likely, to the effects surviving in the large N limit of the supersymmetry present in the parent $U(N)$ theory. Though the role of supersymmetry in the $SU(N)$ case asks for further analysis, the fact that the fermionic divergences can be renormalised, in contrast to the cases previously studied in the literature –in which the bosonic sector was found to be renormalisable, but not so the fermionic sector– suggests that a symmetry relating fermions and bosons may actually be present, making the corresponding divergences not independent. Our result encourages the study of noncommutative models sharing features with the $SU(N)$ theories studied here, with either Majorana fermions or supersymmetry in terms of noncommutative fields; it raises the hope of constructing renormalisable noncommutative gauge theories in the enveloping algebra approach with matter

fields. Also, the effect on the ordinary fields of the supersymmetry of the noncommutative fields in the $SU(N)$ case is still not understood and needs further investigation.

Acknowledgments

This work has been financially supported in part by MICINN through grant FPA2008-04906, and by the National Science Foundation under Grant No. PHY05-51164. The work of C. Tamarit has also received financial support from MICINN and the Fulbright Program through grant 2008-0800.

A Lie algebra, Dirac algebra, Majorana spinors

Following the notation of ref. [38], we denote the Lie algebra generators of $U(N)$ in the fundamental representation as $T^A = \{T^0, T^a\}$, $a = 1, \dots, N^2 - 1$, with $T^0 = \frac{1}{\sqrt{2N}}$ and T^a the standard $SU(N)$ generators. The generators satisfy

$$[T^A, T^B] = i f^{ABC} T^C, \quad \{T^A, T^B\} = d^{ABC} T^C,$$

where f^{ABC} are totally antisymmetric, with f^{abc} having their usual $SU(N)$ values and $f^{0BC} = 0$, whereas d^{ABC} are totally symmetric, d^{abc} having their usual $SU(N)$ values and $d^{0BC} = \sqrt{2/N} \delta^{BC}$, $d^{00c} = 0$, $d^{000} = \sqrt{2/N}$. We will make use of the following identities:

$$\begin{aligned} f^{ACD} f^{BCD} &= N c_A \delta^{AB}, \\ f^{DAE} f^{EBF} f^{FCD} &= -\frac{N}{2} f^{ABC}, \\ f^{DAE} f^{EBF} d^{FCD} &= -\frac{N}{2} d^{ABC} c_A c_B d_C, \\ c_A &= 1 - \delta_{A,0}, & d_A &= 2 - c_A. \end{aligned} \tag{A.1}$$

Concerning Dirac γ matrices, satisfying $\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu}$, we use a basis of operators in the space of spinors constructed from antisymmetrised products of these matrices: $\{\gamma^\mu, \gamma^{\mu\nu}, \gamma^{\mu\nu\rho}, \gamma_5\}$, with the following definitions

$$\begin{aligned} \gamma^{\mu\nu} &= \frac{1}{2}(\gamma^\mu \gamma^\nu - \gamma^\nu \gamma^\mu), & \gamma^{\mu\nu\rho} &= \frac{1}{6}(\gamma^\mu \gamma^\nu \gamma^\rho + \gamma^\nu \gamma^\rho \gamma^\mu + \gamma^\rho \gamma^\mu \gamma^\nu - \gamma^\mu \gamma^\rho \gamma^\nu - \gamma^\nu \gamma^\mu \gamma^\rho - \gamma^\rho \gamma^\nu \gamma^\mu), \\ \gamma_5 &= -\frac{i}{4!} \epsilon_{\mu\nu\rho\sigma} \gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma. \end{aligned}$$

In order to express products of γ matrices in terms of the previous basis, the following identities can be used:

$$\begin{aligned} \gamma^\mu \gamma^\nu &= \eta^{\mu\nu} + \gamma^{\mu\nu}, \\ \gamma^\mu \gamma^\nu \gamma^\rho &= \eta^{\nu\rho} \gamma^\mu - \eta^{\mu\rho} \gamma^\nu + \eta^{\mu\nu} \gamma^\rho + \gamma^{\mu\nu\rho}, \\ \gamma^\mu \gamma^\nu \gamma^\lambda \gamma^\rho &= \eta^{\mu\nu} \eta^{\lambda\rho} + \eta^{\mu\rho} \eta^{\nu\lambda} - \eta^{\mu\lambda} \eta^{\nu\rho} + \eta^{\mu\nu} \gamma^{\lambda\rho} + \eta^{\mu\rho} \gamma^{\nu\lambda} - \eta^{\mu\lambda} \gamma^{\nu\rho} + \eta^{\nu\lambda} \gamma^{\mu\rho} - \eta^{\nu\rho} \gamma^{\mu\lambda} + \eta^{\lambda\rho} \gamma^{\mu\nu} \\ &\quad - i \epsilon^{\mu\nu\lambda\rho} \gamma_5. \end{aligned} \tag{A.2}$$

Majorana spinors are self-conjugate, satisfying

$$\lambda = C\bar{\lambda}^T, \quad \bar{\lambda} = -\lambda^T C^{-1} \quad (\text{A.3})$$

for a charge conjugation matrix C such that

$$C^\dagger = C^{-1}, \quad C^T = -C, \quad C\Gamma_i^T C^{-1} = \eta_i \Gamma_i, \quad \eta_i = \begin{cases} +1, & \Gamma_i = \mathbb{I}, \gamma_5, \gamma^{\mu\nu\rho} \\ -1, & \Gamma_i = \gamma^\mu, \gamma^{\mu\nu}. \end{cases} \quad (\text{A.4})$$

B Feynman rules for $\alpha = 1$, $\mathfrak{S}_\mu^{\text{tree}} = \mathfrak{L}^{\text{tree}} = 0$

The background field legs are denoted by an encircled "b". We define the Feynman rules without symmetrising over these background field legs, which is consistent with the definition of the expansion of the effective action in terms of diagrams provided in eq. (2.12). Since we are doing one-loop calculations, only vertices with two quantum gauge fields contribute; vertices with one quantum gauge field are ignored since they do not contribute to 1PI diagrams. Since we are dealing with self-conjugate Majorana fermions, the vertices with Majorana fermions have to be symmetrised with respect to the conjugation of the interaction in each fermion pair, using (A.4) [39]. The Feynman rules used in our computations are then the following:

$$\begin{aligned}
 & \mu, A \text{ wavy } \xrightarrow{p} \nu, B \leftrightarrow \frac{-ig^2 \delta^{AB} \eta^{\mu\nu}}{p^2 + i\epsilon} \quad i, A \xrightarrow{p} j, B \leftrightarrow \frac{ig^2 (\not{p})_{ij} \delta^{AB}}{p^2 + i\epsilon} \\
 & A \cdots \xrightarrow{p} \cdots B \leftrightarrow \frac{i\delta^{AB}}{p^2 + i\epsilon} \\
 & \begin{array}{c} \text{b} \\ \circ \\ \mu, A \end{array} \xrightarrow{k_1} \begin{array}{c} \nu, B \\ \text{wavy} \\ k_2 \end{array} \xrightarrow{k_3} \begin{array}{c} \rho, C \\ \text{wavy} \\ k_3 \end{array} \leftrightarrow \frac{1}{g^2} f^{ABC} [\eta^{\mu\rho} (k_1 - k_3 - k_2)^\nu + \eta^{\nu\rho} (k_3 - k_2)^\mu \\
 & \quad + \eta^{\mu\nu} (k_2 - k_1 + k_3)^\rho] \\
 & \begin{array}{c} \mu, C \\ \text{b} \\ \circ \end{array} \xrightarrow{k} \begin{array}{c} i, A \\ \text{wavy} \\ q \end{array} \xrightarrow{p} \begin{array}{c} j, B \\ \text{wavy} \\ p \end{array} \leftrightarrow \frac{1}{g^2} (\gamma^\mu)_{ij} f^{ABC} \\
 & \begin{array}{c} \mu, A \\ \text{b} \\ \circ \end{array} \xrightarrow{k_1} \begin{array}{c} \nu, B \\ \text{wavy} \\ k_2 \end{array} \xrightarrow{k_3} \begin{array}{c} \lambda, C \\ \text{b} \\ \circ \end{array} \xrightarrow{k_4} \begin{array}{c} \rho, D \\ \text{wavy} \\ k_4 \end{array} \leftrightarrow -\frac{i}{2g^2} [f^{ABF} f^{FCD} (\eta^{\mu\lambda} \eta^{\nu\rho} - \eta^{\mu\rho} \eta^{\nu\lambda} + \eta^{\mu\nu} \eta^{\lambda\rho}) \\
 & \quad f^{ADF} f^{FBC} (\eta^{\mu\nu} \eta^{\lambda\rho} - \eta^{\mu\lambda} \eta^{\nu\rho} - \eta^{\mu\rho} \eta^{\nu\lambda})] \\
 & \quad f^{ACF} f^{FBD} (\eta^{\mu\nu} \eta^{\lambda\rho} - \eta^{\mu\rho} \eta^{\nu\lambda})]
 \end{aligned}$$

$$\leftrightarrow \frac{1}{4g^2} \theta_{\alpha\beta} d^{ABC} [k_1^\alpha k_2 \cdot k_3 \eta^{\mu\beta} \eta^{\nu\rho} - k_1^\alpha k_2^\rho k_3^\nu \eta^{\mu\beta} - 2(k_1^\alpha k_2^\beta k_3^\mu \eta^{\nu\rho} - k_1^\alpha k_2^\rho k_3^\mu \eta^{\nu\beta} - k_1 \cdot k_3 k_2^\beta \eta^{\nu\rho} \eta^{\mu\alpha} + k_1 \cdot k_3 k_2^\rho \eta^{\mu\alpha} \eta^{\nu\beta})] + (\text{permutations of all legs}) \quad (\text{B.1})$$

$$\leftrightarrow \frac{1}{4g^2} \theta_{\alpha\beta} d^{ABC} (\gamma_\rho)_{ij} [-\eta^{\mu\alpha} q^\rho p^\beta + 2\eta^{\rho\mu} q_\alpha p_\beta + \eta^{\mu\alpha} p_\rho q_\beta + \eta^{\mu\alpha} p^\rho q^\beta - \eta^{\mu\alpha} q_\rho p_\beta]$$

$$\leftrightarrow \frac{-i}{16g^2} \theta_{\alpha\beta} f^{ABF} d^{CDE} [k_3 \cdot k_4 \eta^{\mu\alpha} \eta^{\nu\beta} \eta^{\lambda\rho} - k_3^\rho k_4^\lambda \eta^{\mu\alpha} \eta^{\nu\beta} + 4k_3^\alpha k_4^\mu \eta^{\nu\rho} \eta^{\lambda\beta} - 4(k_3^\beta k_4^\nu \eta^{\mu\alpha} \eta^{\lambda\rho} - k_3^\beta k_4^\lambda \eta^{\mu\alpha} \eta^{\nu\rho} - k_3^\rho k_4^\nu \eta^{\mu\alpha} \eta^{\lambda\beta} + k_3 \cdot k_4 \eta^{\mu\alpha} \eta^{\nu\rho} \eta^{\lambda\beta}) - 2(k_3^\alpha k_4^\beta \eta^{\mu\lambda} \eta^{\nu\rho} - k_3^\alpha k_4^\nu \eta^{\mu\lambda} \eta^{\rho\beta} - k_3^\mu k_4^\beta \eta^{\nu\rho} \eta^{\lambda\alpha} + k_3^\mu k_4^\nu \eta^{\lambda\alpha} \eta^{\rho\beta})] + (\text{permutations of all legs})$$

$$\leftrightarrow \frac{-i}{2g^2} \theta_{\alpha\beta} (\gamma_\rho)_{ij} \left[\frac{1}{2} (d^{ACE} f^{BDE} - d^{BCE} f^{ADE}) [k_1^\alpha (\eta^{\mu\beta} \eta^{\rho\nu} - \eta^{\rho\mu} \eta^{\nu\beta}) + k_1^\rho \eta^{\mu\alpha} \eta^{\nu\beta}] + \frac{1}{2} (d^{ADE} f^{BCE} - d^{BDE} f^{ACE}) [k_2^\alpha (\eta^{\nu\beta} \eta^{\rho\mu} - \eta^{\rho\nu} \eta^{\mu\beta}) + k_2^\rho \eta^{\nu\alpha} \eta^{\mu\beta}] + \frac{1}{2} d^{ABE} f^{CDE} [(p+q)^\rho \eta^{\mu\alpha} \eta^{\nu\beta} + (p+q)^\alpha (\eta^{\mu\beta} \eta^{\rho\nu} - \eta^{\nu\beta} \eta^{\rho\mu})] \right]$$

References

- [1] N. Seiberg and E. Witten, *String theory and noncommutative geometry*, *JHEP* **09** (1999) 032 [[hep-th/9908142](#)] [[SPIRES](#)].
- [2] C.-S. Chu and F. Zamora, *Manifest supersymmetry in non-commutative geometry*, *JHEP* **02** (2000) 022 [[hep-th/9912153](#)] [[SPIRES](#)].
- [3] C.P. Martin and C. Tamarit, *The Seiberg-Witten map and supersymmetry*, *JHEP* **11** (2008) 087 [[arXiv:0809.2684](#)] [[SPIRES](#)].
- [4] J. Madore, S. Schraml, P. Schupp and J. Wess, *Gauge theory on noncommutative spaces*, *Eur. Phys. J. C* **16** (2000) 161 [[hep-th/0001203](#)] [[SPIRES](#)].
- [5] B. Jurčo, S. Schraml, P. Schupp and J. Wess, *Enveloping algebra valued gauge transformations for non-Abelian gauge groups on non-commutative spaces*, *Eur. Phys. J. C* **17** (2000) 521 [[hep-th/0006246](#)] [[SPIRES](#)].

- [6] B. Jurčo, L. Möller, S. Schraml, P. Schupp and J. Wess, *Construction of non-Abelian gauge theories on noncommutative spaces*, *Eur. Phys. J. C* **21** (2001) 383 [[hep-th/0104153](#)] [[SPIRES](#)].
- [7] S. Minwalla, M. Van Raamsdonk and N. Seiberg, *Noncommutative perturbative dynamics*, *JHEP* **02** (2000) 020 [[hep-th/9912072](#)] [[SPIRES](#)].
- [8] M. Nishimura, *A note on supersymmetry in noncommutative field theories*, *Phys. Lett. B* **570** (2003) 105 [[hep-th/0306095](#)] [[SPIRES](#)].
- [9] A. Matusis, L. Susskind and N. Toumbas, *The IR/UV connection in the non-commutative gauge theories*, *JHEP* **12** (2000) 002 [[hep-th/0002075](#)] [[SPIRES](#)].
- [10] F.R. Ruiz, *Gauge-fixing independence of IR divergences in non-commutative U(1), perturbative tachyonic instabilities and supersymmetry*, *Phys. Lett. B* **502** (2001) 274 [[hep-th/0012171](#)] [[SPIRES](#)].
- [11] D. Zanon, *Noncommutative $N = 1, 2$ super U(N) Yang-Mills: UV/IR mixing and effective action results at one loop*, *Phys. Lett. B* **502** (2001) 265 [[hep-th/0012009](#)] [[SPIRES](#)].
- [12] H.O. Girotti, M. Gomes, V.O. Rivelles and A.J. da Silva, *A consistent noncommutative field theory: the Wess-Zumino model*, *Nucl. Phys. B* **587** (2000) 299 [[hep-th/0005272](#)] [[SPIRES](#)].
- [13] I. Jack and D.R.T. Jones, *Ultra-violet finite noncommutative theories*, *Phys. Lett. B* **514** (2001) 401 [[hep-th/0105221](#)] [[SPIRES](#)].
- [14] A.F. Ferrari et al., *Towards a consistent noncommutative supersymmetric Yang-Mills theory: superfield covariant analysis*, *Phys. Rev. D* **70** (2004) 085012 [[hep-th/0407040](#)] [[SPIRES](#)].
- [15] M. Arai, S. Saxell and A. Tureanu, *A noncommutative version of the minimal supersymmetric standard model*, *Eur. Phys. J. C* **51** (2007) 217 [[hep-th/0609198](#)] [[SPIRES](#)].
- [16] J.M. Gracia-Bondia and C.P. Martin, *Chiral gauge anomalies on noncommutative R^4* , *Phys. Lett. B* **479** (2000) 321 [[hep-th/0002171](#)] [[SPIRES](#)].
- [17] L. Bonora, M. Schnabl and A. Tomasiello, *A note on consistent anomalies in noncommutative YM theories*, *Phys. Lett. B* **485** (2000) 311 [[hep-th/0002210](#)] [[SPIRES](#)].
- [18] F. Brandt, C.P. Martin and F.R. Ruiz, *Anomaly freedom in Seiberg-Witten noncommutative gauge theories*, *JHEP* **07** (2003) 068 [[hep-th/0307292](#)] [[SPIRES](#)].
- [19] A. Bichl et al., *Renormalization of the noncommutative photon self-energy to all orders via Seiberg-Witten map*, *JHEP* **06** (2001) 013 [[hep-th/0104097](#)] [[SPIRES](#)].
- [20] R. Wulkenhaar, *Non-renormalizability of θ -expanded noncommutative QED*, *JHEP* **03** (2002) 024 [[hep-th/0112248](#)] [[SPIRES](#)].
- [21] M. Burić and V. Radovanović, *The one-loop effective action for quantum electrodynamics on noncommutative space*, *JHEP* **10** (2002) 074 [[hep-th/0208204](#)] [[SPIRES](#)].
- [22] M. Burić and V. Radovanović, *Non-renormalizability of noncommutative SU(2) gauge theory*, *JHEP* **02** (2004) 040 [[hep-th/0401103](#)] [[SPIRES](#)].
- [23] M. Burić, D. Latas and V. Radovanović, *Renormalizability of noncommutative SU(N) gauge theory*, *JHEP* **02** (2006) 046 [[hep-th/0510133](#)] [[SPIRES](#)].
- [24] M. Burić, V. Radovanović and J. Trampetic, *The one-loop renormalization of the gauge sector in the noncommutative standard model*, *JHEP* **03** (2007) 030 [[hep-th/0609073](#)] [[SPIRES](#)].

- [25] C.P. Martin, D. Sánchez-Ruiz and C. Tamarit, *The noncommutative U(1) Higgs-Kibble model in the enveloping-algebra formalism and its renormalizability*, *JHEP* **02** (2007) 065 [[hep-th/0612188](#)] [[SPIRES](#)].
- [26] C.P. Martin and C. Tamarit, *Renormalisability of the matter determinants in noncommutative gauge theory in the enveloping-algebra formalism*, *Phys. Lett. B* **658** (2008) 170 [[arXiv:0706.4052](#)] [[SPIRES](#)].
- [27] M. Burić, D. Latas, V. Radovanović and J. Trampetic, *The absence of the 4ψ divergence in noncommutative chiral models*, *Phys. Rev. D* **77** (2008) 045031 [[arXiv:0711.0887](#)] [[SPIRES](#)].
- [28] K. Kaminsky, Y. Okawa and H. Ooguri, *Quantum aspects of Seiberg-Witten map in noncommutative Chern-Simons theory*, *Nucl. Phys. B* **663** (2003) 33 [[hep-th/0301133](#)] [[SPIRES](#)].
- [29] P. Schupp and J. You, *UV/IR mixing in noncommutative QED defined by Seiberg-Witten map*, *JHEP* **08** (2008) 107 [[arXiv:0807.4886](#)] [[SPIRES](#)].
- [30] O.F. Dayi, K. Ulker and B. Yapiskan, *Duals of noncommutative supersymmetric U(1) gauge theory*, *JHEP* **10** (2003) 010 [[hep-th/0309073](#)] [[SPIRES](#)].
- [31] L.F. Abbott, *The background field method beyond one loop*, *Nucl. Phys. B* **185** (1981) 189 [[SPIRES](#)].
- [32] D.M. Capper, D.R.T. Jones and P. van Nieuwenhuizen, *Regularization by dimensional reduction of supersymmetric and nonsupersymmetric gauge theories*, *Nucl. Phys. B* **167** (1980) 479 [[SPIRES](#)].
- [33] I. Jack and D.R.T. Jones, *Regularisation of supersymmetric theories*, [hep-ph/9707278](#) [[SPIRES](#)].
- [34] E.S. Fradkin and A.A. Tseytlin, *On the new definition of off-shell effective action*, *Nucl. Phys. B* **234** (1984) 509 [[SPIRES](#)].
- [35] A. Rebhan, *The Wilkowsky-de Witt effective action and its application to Yang-Mills theories*, *Nucl. Phys. B* **288** (1987) 832 [[SPIRES](#)].
- [36] G. Bonneau and F. Delduc, *Nonlinear renormalization and the equivalence theorem*, *Nucl. Phys. B* **266** (1986) 536 [[SPIRES](#)].
- [37] J.A. de Azcarraga, A.J. Macfarlane, A.J. Mountain and J.C. Perez Bueno, *Invariant tensors for simple groups*, *Nucl. Phys. B* **510** (1998) 657 [[physics/9706006](#)] [[SPIRES](#)].
- [38] L. Bonora and M. Salizzoni, *Renormalization of noncommutative U(N) gauge theories*, *Phys. Lett. B* **504** (2001) 80 [[hep-th/0011088](#)] [[SPIRES](#)].
- [39] A. Denner, H. Eck, O. Hahn and J. Kublbeck, *Compact Feynman rules for Majorana fermions*, *Phys. Lett. B* **291** (1992) 278 [[SPIRES](#)].