
This article was downloaded by: On: *16 January 2011* Access details: *Access Details: Free Access* Publisher *Taylor & Francis* Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



To cite this Article (2000) 'Subject Index to Volume 21', Journal of Immunoassay and Immunochemistry, 21: 4, 429 – 431 To link to this Article: DOI: 10.1080/01971520009349548 URL: http://dx.doi.org/10.1080/01971520009349548

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: http://www.informaworld.com/terms-and-conditions-of-access.pdf

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

SUBJECT INDEX TO VOLUME 21

A

 β -Adrenergic agonist ractopamine, development of an immunoassay for, 1-23 Agglutination, and precipitation, 143-164 Antibodies anti-GM1 ganglioside, latex agglutination assay for detection in patients with neuropathy, 377-386 recognition of proteins and peptides by, 85-108 Antibody affinity, determination of, 211-234 Antibody receptors, in immunosensors, electroplated conducting polymers as, 51–64 Antigen-antibody bond, nature of, 109-142 Anti-GM1 ganglioside antibodies, detection in patients with neuropathy by a latex agglutination assay, 377-386

С

Cell-capture enzyme immunoassay, a Giemsa assay for determination of optimal conditions for immobilization of cells in, 297–314
CGP 60536, direct micro-radioimmunoassay of, 65–84
Conducting polymers, electroplated, as antibody receptors in immunosensors, 51–64
Cortisol, one-step ELISA for direct estimation in the serum, 39–50

D

Dot-blot immunogold assay, for identification of a proliferative antigen in the initial cells of a wheat stem meristem, 401–410 Dot-blotting, methods and applications, 273–296

В

Blocking agents, comparison for an ELISA for lipopolysaccharide, 341-354 Electroelution, of individual protein bands from SDS polyacrylamide gels for direct study in cellular assays, 355–375

Ε

Electroplated conducting polymers, as antibody receptors in immunosensors, 51-64 ELISA. See Enzyme-linked immunosorbent assay ELISPOT. See Enzyme-linked immunospot Enzyme immunoassay, cell capture, a simple Giemsa assay for determination of optimal conditions for the immobilization of cells in, 297-314 Enzyme immunomicroscopy, 235-253 Enzyme-linked immunosorbent assay (ELISA), 165-209 for lipopolysaccharide, comparison of blocking agents for, 341-354 one step, for direct estimation of serum cortisol, 39-50 peptide-based sandwich, for human tissue prokallikrein, 411-426 sandwich, for osteoclast differentiation factor in human osteosarcoma cell line, 327-340 Enzyme-linked immunospot (ELISPOT) assay for detection of interleukin-12-secreting cells in normal mice, 25-37 methods and applications, 273-296

F

Flow cytometry, multiparameter, 255–272

G

Giemsa assay, for determination of optimal conditions for the immobilization of cells in a cell-capture enzyme immunoassay, 297–314

I

Immunoblotting, methods and applications, 273–296
Immunofluorescence methods, 235–253
Immunogold assay, dot-blot, for identification of a proliferative antigen in the initial cells of a wheat stem meristem, 401–410
Immunosensors, electroplated conducting polymers as antibody receptors in, 51–64
Inflammatory activity, as an indicator of water quality, 387–399
Interleukin-12–secreting cells, detection with ELISPOT assay, 25–37

K

Kallikrein, mature, development of a peptide-based sandwich ELISA for tissue prokallikrein without cross-reactivity to, 411–426

L

Latex agglutination for detection of anti-GM1 ganglioside antibodies in patients with neuropathy, 377–386 ultrasound enhanced, for rotavirus detection, 315–325 See also Agglutination

SUBJECT INDEX TO VOLUME 21

Ligand-receptor bonds, nature of, 109-142 Lipopolysaccharide, comparison of blocking agents for an ELISA for, 341-354

М

Micro-radioimmunoassay, direct, of the renin inhibitor CGP 60536, 65–84 Multiparameter flow cytometry,

255-272

0

Osteoclast differentiation factor, in human osteosarcoma cell line, 327–340

Р

Peptides, antibody recognition of, 85-108 Precipitation, and agglutination, 143-164 Prokallikrein, human, development of a peptide-based sandwich ELISA for, 411-426 Proliferative antigen, in the initial cells of a wheat stem meristem, a dot-blot immunogold assay for identification of, 401-410 Protein bands, electroelution from SDS polyacrylamide gels for direct study in cellular assays, 355-375 Proteins, antibody recognition of, 85 - 108

R

Ractopamine, development of an immunoassay for, 1–23 Renin inhibitor CGP 60536, direct micro-radioimmunoassay of, 65–84

Rotavirus, detection using ultrasoundenhanced latex agglutination and turbidimetry, 315–325

S

SDS polyacrylamide gels, electroelution of individual protein bands for direct study in cellular assays, 355–375

Stem meristem, wheat, a dot-blot immunogold assay for identification of a proliferative antigen in the initial cells of, 401–410

Т

Turbidimetry, for rotavirus detection, 315–325

U

Ultrasound-enhanced latex agglutination, for rotavirus detection, 315-325

W

Water quality, inflammatory activity as an indicator of, 387–399 Whole blood culture, human, to monitor inflammatory activity as an indicator of water quality, 387–399