



Short Communication

The Influence of the Guest Polarity on the Clathrate Structure Formed by the Werner Complex

A.YU. MANAKOV*

Institute of Inorganic Chemistry, Siberian Branch of Russian Academy of Sciences, Lavrentyeva 3, 630090, Novosibirsk, Russia.

J. LIPKOWSKI

Institute of Physical Chemistry, Polish Academy of Sciences, Kasprzaka 44/52, 01 224 Warsaw, Poland.

(Received: 20 July 1998; in final form: 31 July 1998)

The formation of several types of Werner clathrates has been reported for systems of $[\text{Ni}(4\text{-MePy})_4(\text{NCS})_2]$ and substituted benzene derivatives (4-MePy—4-methylpyridine) [1]. Tetrahedral clathrates (so-called β -phases with a 1 : 1 host : guest stoichiometry) are the most general ones, while formation of trigonal clathrates (1 : 0.67 host : guest ratio) was reported only for the *o*-, *m*-, and *p*-nitrotoluene (powder diffraction data), and *m*-bromonitrobenzene guest molecules [1, 2]. This fact may be easily explained in the case of the *o*- and *m*-isomers of the guest because formation of the β -clathrates with *ortho*- and *meta*-substituted benzene derivatives is extremely unfavorable (in these clathrates the channel width is most favorable for location of the *para*-substituted benzene derivatives) [1]. As discussed in [3], the volume of the NO_2 group in *p*-nitrotoluene allows inclusion of this molecule into the β -clathrates, and formation of this type of clathrate could be expected for this guest. To clarify this problem further we have studied the single crystal structure of the $[\text{Ni}(4\text{-MePy})_4(\text{NCS})_2]$ host with the *p*-nitrotoluene guest molecule. The product, however, appears to be a typical trigonal Werner clathrate ($R\bar{3}$ space group) [4, 5]; the parameters of the unit cell ($a = 27.650 \text{ \AA}$, $c = 11.165 \text{ \AA}$) are in good agreement with parameters determined in [2]. Formation of the metastable octahedral crystals (β -phase, as determined by X-ray diffraction) was observed during the initial stages of crystallization. In our opinion, the most probable explanation of the formation of the trigonal clathrate with the *p*-nitrotoluene guest molecule is the energetically favorable coupling of the dipoles of the guest molecules in this type of clathrate (Figure 1a). In the case of β -clathrates,

* Author for correspondence.




Figure 1. (a) Experimentally determined arrangement of the *p*-nitrotoluene guest molecules in the cavity of the trigonal clathrate. Dipoles of the *p*-nitrotoluene molecules are coupled. (b) Schematic representation of the arrangement of the polar *para*-substituted benzene guest molecules in the channels of the β -[Ni(4-MePy)₄(NCS)₂] clathrate phase. Accommodation of this type of guest molecules about the $\bar{4}$ symmetry positions causes unfavorable orientation of the guest molecule dipoles.

Table I. Compositions of the β -[Ni(4-MePy)₄(NCS)₂] \cdot *x*(toluene) \cdot *y*(4-MePy) (*x* + *y* = 1) clathrates and mother solutions in equilibrium at room temperature

β -[Ni(4-MePy) ₄ (NCS) ₂] \cdot <i>x</i> (toluene) \cdot <i>y</i> (4-MePy)		Mother solution (mol. %)		
<i>x</i> (by difference)	<i>y</i>	Toluene (by difference)	4-MePy	Host
0.63	0.37 ± 0.01	43.6	56.3 ± 0.1	0.1
0.50	0.50 ± 0.01	21.5	78.3 ± 0.2	0.2

dipoles of the guest molecules are situated in less energetically favorable positions (Figure 1b). This conclusion may be confirmed too by our data concerning the compositions of the β -[Ni(4-MePy)₄(NCS)₂] clathrates with mixed toluene + 4-methylpyridine guest and their mother solutions (see Table I, clathrate phase enriches with the non-polar guest component). The data presented in this communication show that the polarity of the guest molecule can modify the processes of clathrate formation in systems containing the [Ni(4-MePy)₄(NCS)₂] host molecule. This fact should be taken into account when considering the separation processes with the β -[Ni(4-MePy)₄(NCS)₂] sorbent.

References

1. J. Lipkowski: in J. L. Atwood, J. E. D. Davies and D. D. MacNicol, *Inclusion Compounds*, Vol. 1, pp. 59–103. Academic Press, London (1984).
2. A. Lewartowska, S. Brzozowski, and W. Kemula: *J. Mol. Struct.* **75**, 113 (1981).
3. A. Yu. Manakov J. Lipkowski, K. Suwinska and M. Kitamura: *J. Incl. Phenom.* **26**, 1 (1996).
4. N. V. Pervuchina, N. V. Podberezskaya, I. V. Davydova, N. V. Kislych, and Yu. A. Dyadin: *J. Incl. Phenom.* **13**, 9 (1992).
5. J. Lipkowski and D. V. Soldatov: *J. Incl. Phenom.* **18**, 311 (1994).