

# Silver(I) and mercury(II) complexes of *meta*- and *para*-xylyl linked bis(imidazol-2-ylidenes)

Murray V. Baker · David H. Brown ·  
Rosenani A. Haque · Brian W. Skelton ·  
Allan H. White

Received: 26 June 2009 / Accepted: 2 July 2009 / Published online: 28 July 2009  
© Springer Science+Business Media B.V. 2009

**Abstract** Mononuclear silver and mercury complexes bearing bis-*N*-heterocyclic carbene (NHC) ligands with linear coordination modes have been prepared and structurally characterised. The complexes form metallocyclic structures that display rigid solution behaviour. A larger metallocycle of the form  $[L_2Ag_2]^{2+}$  [where L = *para*-bis(*N*-methylimidazolylidene)xylylene] has been isolated from the reaction of *para*-xylylene-bis(*N*-methylimidazolium) chloride and Ag<sub>2</sub>O. Reaction of silver- and mercury-NHC complexes with Pd(NCCH<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub> affords palladium-NHC complexes via NHC-transfer reactions, the mercury case being only the second example of a NHC-transfer reaction using a mercury-NHC complex.

**Keywords** Silver · Mercury · *N*-Heterocyclic carbenes · Carbene transfer

---

Dedicated to Jack Harrowfield on his retirement. An inimitable educator, scientist, colleague, friend, writer of mischievous anagrams ... and still a good bloke.

---

M. V. Baker (✉) · D. H. Brown · R. A. Haque ·  
B. W. Skelton · A. H. White  
Chemistry M313, School of Biomedical, Biomolecular and  
Chemical Sciences, The University of Western Australia, 35  
Stirling Hwy, Crawley, WA 6009, Australia  
e-mail: murray.baker@uwa.edu.au

D. H. Brown  
Department of Applied Chemistry, Nanochemistry Research  
Institute, Curtin University of Technology, Kent St, Bentley,  
WA 6102, Australia

R. A. Haque  
School of Chemical Sciences, Universiti Sains Malaysia, 11800  
Minden, Penang, Malaysia


## Introduction

*N*-Heterocyclic carbene (NHC) complexes of silver have become common-place in the organometallic literature. The interest in Ag-NHC complexes is largely due to their ease of synthesis (from azolium salts and Ag<sub>2</sub>O) and their ability to serve as useful precursors to other NHC-metal complexes by NHC-transfer reactions [1]. In recent years NHC-silver complexes have also been studied for possible antimicrobial and anti-cancer properties [2, 3].


Wanzlick and Schönherer reported the synthesis of a mercury NHC complex via the reaction of an imidazolium salt with mercury(II) acetate in 1968 [4]. Despite the fact that this complex was one of the earliest examples of an NHC-metal complex, NHC-mercury complexes have received little attention compared to NHC-silver complexes, and applications of mercury carbene complexes have not been widely explored. We recently reported the use of NHC-mercury complexes for redox-transmetallation chemistry, where the reaction of a NHC-mercury(II) complex with a palladium(0) source afforded an NHC-palladium(II) complex [5]. Redox-transmetallation using mercury complexes is a well established route in organometallic synthesis outside of NHC chemistry [6].

We, and others, have an interest in xylyl-linked NHCs and their metal complexes, particularly systems where the NHCs form part of a cyclophane structure [5, 7–18]. Cyclophane NHCs (comprising two NHC groups linked by two xylyl groups) provide a fascinating range of binding modes for metals, ranging from *cis*-chelating modes (e.g., 1) and *trans*-spanning chelating modes (e.g., 2, 3) through to dinuclear modes involving two mono-NHC-bound metals (e.g., 4) and dinuclear modes involving two bis(NHC)-bound metals (e.g., 5, 6). Metal complexes of cyclophane NHC ligands display interesting properties, including high


stability compared to other complexes of unidentate and chelating NHCs and interesting catalytic properties (Pd-NHC complexes) [19] and anti-mitochondrial and luminescent properties (Au-NHC complexes) [20]. Compared to the cyclophane bis(NHC)s, bis(NHC) ligands containing only one xylyl linker (non-cyclophane structures) provide additional conformational flexibility that can permit additional metal-coordination modes (e.g., **7**) [21].



Silver complexes derived from non-cyclic xylyl-linked bis(NHC) ligands have been reported (e.g., **8**, **9**) [22–24]. Lutidinediyl analogues of the *meta*-xylyl linked structures have also been reported (**10**) [21, 25, 26]. In general metal complexes with *para*-xylyl linked bis(NHC) ligands are rare, though some interesting structures have been characterised in the solid-state (e.g., **11**) [27–29].



Here we report the synthesis and characterisation of silver and mercury complexes with NHC ligands **12** and **13**, derived from *meta*- and *para*-xylyl linked bis-imidazolium cations. We also report the transmetalation (carbene-transfer) reactions of silver and mercury complexes to form a palladium complex.



## Experimental section

### General experimental

Nuclear magnetic resonance spectra were recorded using Bruker Avance 500 (500.13 MHz for  $^1\text{H}$  and 125.77 MHz for  $^{13}\text{C}$ ) and Bruker ARX300 (300.14 MHz for  $^1\text{H}$  and 75.48 MHz for  $^{13}\text{C}$ ) spectrometers at ambient temperature.  $^1\text{H}$  and  $^{13}\text{C}$  chemical shifts were referenced to solvent resonances. Mass spectra were obtained by Dr A. Reeder using a VG Autospec Mass Spectrometer via electrospray ionisation (low resolution). Microanalyses were performed by the Microanalytical Laboratory at the Research School of Chemistry, Australian National University, Canberra.

### Synthesis

$\text{PdCl}_2(\text{NCCCH}_3)_2$  was prepared by the method of Wimmer et al. [30]. 1,3-Di(bromomethyl)-2,4,6-trimethylbenzene was prepared by the method of van der Made and van der Made [31].

#### *Synthesis of the para-xylyl bis(imidazolium) salts $13\text{H}_2\cdot 2\text{Cl}$ and $13\text{H}_2\cdot 2\text{PF}_6$*

*N*-Methylimidazole (3.0 g, 17.1 mmol) was added to a stirred solution of  $\alpha,\alpha'$ -dichloro-*p*-xylene (9.0 g, 51.4 mmol) in dioxane (150 mL). The mixture was heated at 100 °C for 24 h under nitrogen. The product precipitated from solution as a light yellow solid, which was collected and washed with diethyl ether. Recrystallisation from a mixture of ethanol and diethyl ether yielded the product as yellow crystals (4.7 g). Yield 81%.  $^1\text{H}$  NMR (300.1 MHz,  $d_6$ -DMSO):  $\delta$  3.86 (6H, s, 2  $\times$  CH<sub>3</sub>), 5.47 (4H, br s,  $W_{\text{H}/2}$  = 2 Hz, 2  $\times$  CH<sub>2</sub>), 7.50 (4H, br s,  $W_{\text{H}/2}$  2 Hz, 4  $\times$  Ar CH), 7.74 (2H, d,  $^3J_{\text{H,H}}$  = 2 Hz, 2  $\times$  imidazolium H5), 7.86 (2H, d,  $^3J_{\text{H,H}}$  = 2 Hz, 2  $\times$  imidazolium H4) and 9.49 (2H, s, 2  $\times$  imidazolium H2);  $^{13}\text{C}$  NMR (75.5 MHz,  $d_6$ -DMSO):  $\delta$  35.9 (NCH<sub>3</sub>), 51.3 (CH<sub>2</sub>), 122.4 (imidazolium C5), 124.1 (imidazolium C4), 129.1 (Ar CH), 135.6 (imidazolium C2) and 136.9 (Ar C); Anal. Calc. for  $\text{C}_{16}\text{H}_{20}\text{N}_4\text{Cl}_2\cdot 3\text{H}_2\text{O}$ : C, 48.86; H, 6.36; N, 14.24. Found: C, 48.74; H, 6.36; N, 14.07%.

$13\text{H}_2\cdot 2\text{Cl}$  was converted into its hexafluorophosphate salt ( $13\text{H}_2\cdot 2\text{PF}_6$ ) by a salt metathesis reaction using  $13\text{H}_2\cdot 2\text{Cl}$  and  $\text{KPF}_6$  in methanol and was obtained as white crystalline solid, in 71% yield after recrystallisation from hot water. Crystals suitable for X-ray diffraction studies were grown from slow evaporation of a concentrated solution of the imidazolium salt in acetonitrile at 4 °C.  $^1\text{H}$  NMR (500.1 MHz,  $d_6$ -DMSO):  $\delta$  3.84 (6H, s, 2  $\times$  CH<sub>3</sub>), 5.41 (4H, br s,  $W_{\text{H}/2}$  2 Hz, 4  $\times$  CH<sub>2</sub>), 7.45 (4H, br s,  $W_{\text{H}/2}$  2 Hz, 4  $\times$  Ar CH), 7.70 (2H, d,  $^3J_{\text{H,H}}$  = 2 Hz, 2  $\times$  imidazolium

H5), 7.75 (2H, d,  $^3J_{\text{H,H}}$  = 2 Hz, 2  $\times$  imidazolium H4) and 9.23 (2H, s, 2  $\times$  imidazolium H2);  $^{13}\text{C}$  NMR (125.8 MHz,  $d_6$ -DMSO):  $\delta$  35.9 (CH<sub>3</sub>), 51.3 (CH<sub>2</sub>), 122.2 (imidazolium C5), 124.0 (imidazolium C4), 128.9 (Ar CH), 136.7 (imidazolium C2) and 135.4 (Ar C); Anal. Calc. for  $\text{C}_{16}\text{H}_{20}\text{N}_4\text{P}_2\text{F}_{12}$ : C, 34.42; H, 3.61; N, 10.04. Found: C, 34.29; H, 3.53; N, 9.92%. The tetrafluoroborate salt  $13\text{H}_2\cdot 2\text{BF}_4$  was obtained by a similar procedure.

#### *Synthesis of the meta-mesitylene bis(imidazolium) salt $12\text{H}_2\cdot 2\text{PF}_6$*

A solution of 1,3-di(bromomethyl)-2,4,6-trimethylbenzene (1 g, 3.3 mmol) and 1-methylimidazole (0.84 g, 10 mmol) in 1,4-dioxane (13 mL) was heated at 100 °C, under nitrogen, for 24 h. The mixture was allowed to cool to room temperature and the precipitate was collected, washed with diethyl ether (5 mL) and dried over  $\text{P}_2\text{O}_5$  to afford  $12\text{H}_2\cdot 2\text{Br}$  as a white powder (1.4 g, 90%). The bromide salt  $12\text{H}_2\cdot 2\text{Br}$  was converted to its hexafluorophosphate salt by a metathesis reaction using  $12\text{H}_2\cdot 2\text{Br}$  and  $\text{KPF}_6$  in methanol and was obtained as white crystalline solid, in 85% yield after recrystallisation from hot water. Crystals suitable for X-ray diffraction studies were grown from slow evaporation of a concentrated solution of the imidazolium salt in acetonitrile at 4 °C.  $^1\text{H}$  NMR (500.1 MHz,  $d_6$ -DMSO):  $\delta$  2.21 (3H, s, 1  $\times$  C2-CH<sub>3</sub>), 2.30 (6H, s, 1  $\times$  C4-CH<sub>3</sub>/1  $\times$  C6-CH<sub>3</sub>), 3.79 (6H, s, 2  $\times$  N-CH<sub>3</sub>), 5.43 (4H, s, 2  $\times$  CH<sub>2</sub>), 7.18 (1H, s, 1  $\times$  Ar H5), 7.60 (2H, d,  $^3J_{\text{H,H}}$  = 2 Hz, 2  $\times$  imidazolium H5), 7.71 (2H, d,  $^3J_{\text{H,H}}$  = 2 Hz, 2  $\times$  imidazolium H4) and 8.91 (2H, s, 2  $\times$  imidazolium H2);  $^{13}\text{C}$  NMR (125.8 MHz,  $d_6$ -DMSO):  $\delta$  15.6 (C2-CH<sub>3</sub>), 19.1 (C4-CH<sub>3</sub>/C6-CH<sub>3</sub>), 35.9 (NCH<sub>3</sub>), 47.3 (CH<sub>2</sub>), 122.2 (imidazolium C5'), 123.7 (imidazolium C4'), 131.2 (Ar CH), 136.0 (imidazolium C2'), 128.4 (Ar C), 139.0 (Ar C) and 139.7 (Ar C); Anal. Calc. for  $\text{C}_{16}\text{H}_{20}\text{N}_4\text{P}_2\text{F}_{12}$ : C, 38.01; H, 4.37; N, 9.33. Found: C, 37.87; H, 4.09; N, 9.50%.

#### *Synthesis of the dinuclear silver complex $17\cdot 2\text{Cl}$*

$\text{Ag}_2\text{O}$  (85 mg, 367 mmol) was added to a solution of  $13\text{H}_2\cdot 2\text{Cl}$  (105 mg, 310 mmol) in methanol (15 mL). The mixture was heated at 60 °C for 2 h in darkness. A clear solution with some black suspension was obtained. The mixture was filtered and the filtrate was evaporated to dryness to give a white powder. Recrystallisation of the powder from minimum amount of hot methanol yielded white crystals (90 mg). Yield 52%.  $^1\text{H}$  NMR (500.1 MHz,  $d_6$ -DMSO):  $\delta$  3.83 (12H, s, 4  $\times$  NCH<sub>3</sub>), 5.29 (8H, br s,  $W_{\text{H}/2}$  2 Hz, 4  $\times$  CH<sub>2</sub>), 7.14 (8H, br s,  $W_{\text{H}/2}$  2 Hz, 8  $\times$  ArH), 7.46 (4H, d,  $^3J_{\text{H,H}}$  = 2 Hz, 4  $\times$  imidazolium H5) and 7.53 (4H, d,  $^3J_{\text{H,H}}$  = 2 Hz, 4  $\times$  imidazolium H4);  $^{13}\text{C}$  NMR (125.8 MHz,  $d_6$ -DMSO):  $\delta$  38.2 (NCH<sub>3</sub>), 53.5 (CH<sub>2</sub>),

122.2 (imidazolium C5'), 123.5 (imidazolium C4'), 127.7 (Ar CH), 137.2 (Ar C) and 180.1 (br C-Ag). Anal. Calc. for  $C_{32}H_{36}Ag_2N_8Cl_2 \cdot 1.5CH_3OH \cdot 2H_2O$ : C, 44.54; H, 5.13; N, 12.40. Found: C, 44.70; H, 4.79; N, 12.19%.

#### Synthesis of the dinuclear silver complex **17**·2PF<sub>6</sub>

Ag<sub>2</sub>O (23 mg, 100 mmol) was added to a solution of **13**H<sub>2</sub>·2PF<sub>6</sub> (50 mg, 90 mmol) in acetonitrile (10 mL). The mixture was heated at 60 °C for 2 h in darkness. The mixture was filtered and the filtrate was concentrated in vacuo. The white residue was washed with dichloromethane (2 × 2 mL) and was then recrystallised from a mixture of acetonitrile and diethyl ether to afford a white powder (36 mg). Yield 77%. <sup>1</sup>H NMR (500.1 MHz, *d*<sub>6</sub>-DMSO): δ 3.83 (12H, s, 4 × NCH<sub>3</sub>), 5.29 (8H, br s, *W*<sub>h/2</sub> 2 Hz, 4 × CH<sub>2</sub>), 7.13 (8H, br s, *W*<sub>h/2</sub> 2 Hz, 8 × ArH), 7.46 (4H, d, <sup>3</sup>*J*<sub>H,H</sub> = 2 Hz, 4 × imidazolium H4) and 7.53 (4H, d, <sup>3</sup>*J*<sub>H,H</sub> = 2 Hz, 4 × imidazolium H5); <sup>13</sup>C NMR (125.8 MHz, *d*<sub>6</sub>-DMSO): δ 38.1 (NCH<sub>3</sub>), 53.5 (CH<sub>2</sub>), 122.2 (imidazolium C5'), 123.5 (imidazolium C4'), 127.6 (Ar CH), 137.1 (Ar C) and 180.2 (d, <sup>1</sup>*J*<sub>C-109Ag</sub> = 210 Hz, <sup>1</sup>*J*<sub>C-107Ag</sub> = 180 Hz, Ag-C); ES-MS *m/z* 891 (55% relative intensity), 892 (30%), 893 (100%), 894 (40%), 895 (55%), 896 (15%) [M-PF<sub>6</sub>]<sup>+</sup> ≡ [**17**-PF<sub>6</sub>]<sup>+</sup>; Anal. Calc. for  $C_{32}H_{36}N_8Ag_2P_2F_{12}$ : C, 37.02; H, 3.49; N, 10.79. Found: C, 36.77; H, 3.32; N, 10.71%.

#### Synthesis of the silver complex **15**·PF<sub>6</sub>

Ag<sub>2</sub>O (30 mg, 129 mmol) was added to a solution of **12**H<sub>2</sub>·2Br (25 mg, 104 mmol) in methanol (20 mL). The mixture was heated at 50 °C for 2 h in darkness. The mixture was filtered and the filtrate was concentrated in vacuo to afford **15**-Br as a white solid. The white solid residue was dissolved in water (10 mL) and the resulting solution was filtered into an aqueous solution of KPF<sub>6</sub> (38 mg, 206 mmol, 5 mL). A grey precipitate formed, which was collected and dried under reduced pressure (44 mg). Yield 76%. Analytically pure samples were obtained by recrystallisation of the complex from acetonitrile. <sup>1</sup>H NMR (500.1 MHz, *d*<sub>6</sub>-acetone): δ 2.17 (6H, s, 2 × C4/C6-CH<sub>3</sub>), 2.73 (3H, s, 1 × C2-CH<sub>3</sub>), 3.85 (6H, s, 2 × NCH<sub>3</sub>), 5.48 (2H, A part of AB multiplet, <sup>2</sup>*J*<sub>H,H</sub> = 14 Hz, 2 × benzylic CHH), 5.57 (2H, B part of AB multiplet, <sup>2</sup>*J*<sub>H,H</sub> = 14 Hz, 2 × benzylic CHH), 7.12 (1H, s, 1 × Ar H5), 7.42 (2H, d, <sup>3</sup>*J*<sub>H,H</sub> = 2 Hz, 2 × imidazolium H5) and 7.64 (2H, d, <sup>3</sup>*J*<sub>H,H</sub> = 2 Hz, 2 × imidazolium H4); <sup>13</sup>C NMR (125.8 MHz, *d*<sub>6</sub>-DMSO): δ 17.0 (C4/C6-CH<sub>3</sub>), 19.1 (C2-CH<sub>3</sub>), 39.4 (NCH<sub>3</sub>), 47.9 (CH<sub>2</sub>), 122.1 (imidazolium-C5'), 123.7 (imidazolium-C4'), 131.7 (Ar CH), 131.5 (Ar C), 136.4 (Ar C), 138.0 (Ar C) and 178.7 (d, Ag-C, <sup>1</sup>*J*<sub>C-109Ag</sub> = 212 Hz, <sup>1</sup>*J*<sub>C-107Ag</sub> = 188 Hz); Anal. Calc. for  $C_{19}H_{24}N_4AgPF_6 \cdot 0.5H_2O$ : C,

40.02; H, 4.42; N, 9.82. Found: C, 40.25; H, 4.36; N, 9.90%.

#### Synthesis of the mercury complex **16**·2PF<sub>6</sub>

Hg(OAc)<sub>2</sub> (35 mg, 109 mmol) was added to a solution of **12**H<sub>2</sub>·2PF<sub>6</sub> (50 mg, 100 mmol) in acetonitrile (40 mL). The mixture was heated at reflux for 3 days. A clear colourless solution resulted. The solution was concentrated in vacuo. The solid residue was washed with water (2 × 5 mL) and recrystallised from acetonitrile to afford colourless crystals (45 mg). Yield 55%. Analytically pure samples and crystals suitable for X-ray diffraction studies were obtained by the slow evaporation of a solution of the salt in acetonitrile/water at room temperature; <sup>1</sup>H NMR (500.1 MHz, *d*<sub>3</sub>-acetonitrile): δ 2.16 (6H, s, 2 × C4/C6-CH<sub>3</sub>), 2.53 (3H, s, 1 × C2-CH<sub>3</sub>), 3.79 (6H, s, 2 × NCH<sub>3</sub>), 5.58 (2H, A part of AB pattern, <sup>2</sup>*J*<sub>H,H</sub> = 15 Hz, 2 × benzylic CHH), 5.62 (2H, B part of AB pattern, <sup>2</sup>*J*<sub>H,H</sub> = 15 Hz, 2 × benzylic CHH), 7.34 (1H, s, 1 × Ar H5), 7.42 (2H, m, <sup>3</sup>*J*<sub>H,H</sub> = 2 Hz, 2 × imidazolium H5) and 7.68 (2H, d, <sup>3</sup>*J*<sub>H,H</sub> = 2 Hz, 2 × imidazolium H4); <sup>13</sup>C NMR (125.8 MHz, *d*<sub>6</sub>-acetonitrile): δ 16.7 (C4/C6-CH<sub>3</sub>), 17.0 (C2-CH<sub>3</sub>), 39.9 (NCH<sub>3</sub>), 49.6 (CH<sub>2</sub>), 126.4 (imidazolium-C5'), 126.6 (imidazolium-C4'), 132.0 (Ar CH), 135.4 (Ar C), 137.4 (Ar C), 142.4 (Ar C) and 173.3 (C-Hg); Anal. Calc. for  $C_{19}H_{24}N_4HgP_2F_{12}$ : C, 28.56; H, 3.03; N, 7.01. Found: C, 28.74; H, 3.09; N, 7.13%.

#### Synthesis of a palladium complex via NHC-transfer reactions

Method A (via a silver complex): Ag<sub>2</sub>O (28 mg, 120 mmol) was added to a solution of **13**H<sub>2</sub>·2Cl (32 mg, 94 mmol) in methanol (10 mL). The mixture was heated at 60 °C for 2 h in darkness. A clear solution with some black suspension was obtained. The mixture was filtered and the filtrate was evaporated to dryness to give a white powder. The white powder was suspended in acetonitrile (10 mL) and PdCl<sub>2</sub>(NCCH<sub>3</sub>)<sub>2</sub> (49 mg, 188 mmol) was added. The resulting mixture was stirred at 60 °C for 24 h, which resulted in a yellow solution with a dark precipitate. The mixture was filtered and the filtrate was dried in vacuo to leave a yellow solid. The solid was washed with a minimum amount of dichloromethane and then recrystallised from acetonitrile, collected and dried in vacuo to afford a yellow powder (38 mg). Yield: 59%. Crystals suitable for X-ray diffraction studies were grown from slow evaporation of a concentrated solution of the complex in acetonitrile at 4 °C.

Method B (via a mercury complex): Hg(OAc)<sub>2</sub> (40 mg, 125 mmol) was added to a solution of **13**H<sub>2</sub>·2Cl (40 mg, 118 mmol) in methanol (50 mL). The mixture was heated at reflux for 24 h resulting in a clear colourless solution.

The solution was concentrated in vacuo to afford a white solid. The white solid was washed with water (2 × 6 mL), diethyl ether (2 × 6 mL) and dried under vacuum (64 mg). [<sup>1</sup>H NMR (300.1 MHz, *d*<sub>6</sub>-DMSO): δ 3.89 (12H, s, 4 × NCH<sub>3</sub>), 5.48 (8H, br s, 4 × CH<sub>2</sub>), 7.46 (8H, s, 8 × ArH), 7.68 (4H, d, <sup>3</sup>*J*<sub>H,H</sub> = 2 Hz, 4 × imidazolium H5) and 7.73 (4H, d, <sup>3</sup>*J*<sub>H,H</sub> = 2 Hz, 4 × imidazolium H4)]. A portion of the solid (30 mg, 38 mmol) was suspended in acetonitrile (10 mL). A solution of PdCl<sub>2</sub>(NCCCH<sub>3</sub>)<sub>2</sub> (20 mg, 77 mmol) in acetonitrile (10 mL) was added and the mixture was heated at 60 °C. After 30 min, a precipitate began to form. The reaction was continued until all the original white solid dissolved (18 h). A yellow solution with grey precipitate resulted. The solution was filtered and the filtrate was concentrated in vacuo to give a yellow solid, which was then washed with water (2 × 5 mL) and dried (23 mg). Yield 91%.

<sup>1</sup>H NMR (500.1 MHz, *d*<sub>3</sub>-acetonitrile): δ 4.01 (6H, s, 2 × CH<sub>3</sub>), 5.67 (4H, br s, 2 × CH<sub>2</sub>), 6.94 (2H, d, <sup>3</sup>*J*<sub>H,H</sub> = 2 Hz, 2 × imidazolium H5), 7.07 (2H, d, <sup>3</sup>*J*<sub>H,H</sub> = 2 Hz, 2 × imidazolium H4) and 7.50 (4H, s, 4 × ArH); <sup>13</sup>C NMR (125.8 MHz, *d*<sub>3</sub>-acetonitrile): δ 39.0 (N-CH<sub>3</sub>), 55.0 (CH<sub>2</sub>), 123.6 (imidazolium-C5'), 125.9 (imidazolium-C4'), 130.6 (Ar CH), 137.9 (Ar C) and 147.3 (C-Pd); Anal. Calc. for C<sub>16</sub>H<sub>18</sub>N<sub>4</sub>Pd<sub>2</sub>Cl<sub>4</sub>: C, 30.95; H, 2.92; N, 9.02. Found: C, 31.01; H, 3.22; N, 8.85%.

#### Structure determinations

Full spheres of CCD area-detector diffractometer data were measured (monochromatic Mo *K*α radiation, λ = 0.71073 Å, ω-scans) yielding *N*<sub>(total)</sub> reflections, these merging to *N* unique (*R*<sub>int</sub> cited) after 'empirical'/multiscan 'absorption correction', these being used in the full matrix least squares refinements on *F*<sup>2</sup> (anisotropic displacement parameter refinement for the non-hydrogen atoms, hydrogen atom treatment following a riding model; reflection weights: (σ<sup>2</sup>(*F*<sub>o</sub><sup>2</sup>) + (*aP*)<sup>2</sup>)<sup>-1</sup> (*P* = (*F*<sub>o</sub><sup>2</sup> + 2*F*<sub>c</sub><sup>2</sup>)/3)); *N*<sub>o</sub> with *F*<sub>o</sub> > 4σ(*F*<sub>o</sub>) were considered 'observed'. Pertinent details are given below in the Figures, the latter showing 50% probability amplitude displacement ellipsoids for the non-hydrogen atoms, hydrogen atoms having arbitrary radii of 0.1 Å. Neutral atom complex scattering factors were employed within the SHELXL 97 program [32]. Full.cif depositions (excluding structure factor amplitudes) reside with the Cambridge Crystallographic Data Centre, CCDC 736772-736777, 737102.

#### Crystal/refinement data

**13H<sub>2</sub>·2PF<sub>6</sub>** ≡ C<sub>16</sub>H<sub>20</sub>F<sub>12</sub>N<sub>4</sub>P<sub>2</sub>, *M* = 558.3. Monoclinic, space group *P*2<sub>1</sub>/*c* (*C*<sub>2h</sub><sup>5</sup>, No. 14), *a* = 7.1194(8), *b* = 12.1150(10), *c* = 13.055(2) Å, β = 105.019(2)°,

*V* = 1088 Å<sup>3</sup> (*T* ca. 153 K). *D*<sub>c</sub> (*Z* = 2) = 1.70<sub>5</sub> g cm<sup>-3</sup>. μ<sub>Mo</sub> = 0.32 mm<sup>-1</sup>; specimen: 0.43 × 0.31 × 0.25 mm; *T*<sub>min/max</sub> = 0.82. 2θ<sub>max</sub> = 66°; *N*<sub>t</sub> = 14578, *N* = 3980 (*R*<sub>int</sub> = 0.022), *N*<sub>o</sub> = 3382. *R*<sub>1</sub> = 0.041, *wR*<sub>2</sub> = 0.123 (*a* = 0.067, *b* = 0.326), *S* = 1.05. |Δρ<sub>max</sub>| = 0.45 e Å<sup>-3</sup>.

**13H<sub>2</sub>·2BF<sub>4</sub>** ≡ C<sub>16</sub>H<sub>20</sub>B<sub>2</sub>F<sub>8</sub>N<sub>4</sub>, *M* = 442.0. Monoclinic, space group *P*2<sub>1</sub>/*c*, *a* = 4.9639(8), *b* = 12.913(2), *c* = 15.401(3) Å, β = 92.932(3)°, *V* = 985.9 Å<sup>3</sup> (*T* ca. 153 K). *D*<sub>c</sub> (*Z* = 2) = 1.48<sub>9</sub> g cm<sup>-3</sup>. μ<sub>Mo</sub> = 0.14 mm<sup>-1</sup>; specimen: 0.48 × 0.15 × 0.10 mm; *T*<sub>min/max</sub> = 0.80. 2θ<sub>max</sub> = 50°; *N*<sub>t</sub> = 9070, *N* = 1730 (*R*<sub>int</sub> = 0.036), *N*<sub>o</sub> = 1454. *R*<sub>1</sub> = 0.058, *wR*<sub>2</sub> = 0.148 (*a* = 0.068, *b* = 1.03), *S* = 1.05. |Δρ<sub>max</sub>| = 0.46 e Å<sup>-3</sup>.

**17·2NO<sub>3</sub>·2CH<sub>3</sub>OH** ≡ C<sub>34</sub>H<sub>44</sub>Ag<sub>2</sub>N<sub>10</sub>O<sub>8</sub>, *M* = 936.5. Triclinic, space group *P* $\bar{1}$ , *a* = 9.363(2), *b* = 9.460(2), *c* = 21.779(4) Å, α = 94.308(4), β = 91.836(3), γ = 93.606(3)°, *V* = 1919 Å<sup>3</sup> (*T* ca. 170 K). *D*<sub>c</sub> (*Z* = 2) = 1.62<sub>1</sub> g cm<sup>-3</sup>. μ<sub>Mo</sub> = 1.08 mm<sup>-1</sup>; specimen: 0.49 × 0.15 × 0.12 mm; *T*<sub>min/max</sub> = 0.85. 2θ<sub>max</sub> = 58°; *N*<sub>t</sub> = 15090, *N* = 8356 (*R*<sub>int</sub> = 0.047), *N*<sub>o</sub> = 6607. *R*<sub>1</sub> = 0.050, *wR*<sub>2</sub> = 0.116 (*a* = 0.040, *b* = 5.58); *S* = 1.08. |Δρ<sub>max</sub>| = 1.17 e Å<sup>-3</sup>.

**18·2CH<sub>3</sub>CN** ≡ C<sub>24</sub>H<sub>30</sub>Cl<sub>4</sub>N<sub>8</sub>Pd<sub>2</sub>, *M* = 785.2. Triclinic, space group *P* $\bar{1}$ (*C*<sub>i</sub><sup>1</sup>, No. 2), *a* = 7.2833(6), *b* = 7.8832(7), *c* = 14.8010(10) Å, α = 80.196(2), β = 81.669(2), γ = 65.593(2)°, *V* = 759.9 Å<sup>3</sup> (*T* ca. 170 K). *D*<sub>c</sub> (*Z* = 1) = 1.56<sub>4</sub> g cm<sup>-3</sup>. μ<sub>Mo</sub> = 1.56 mm<sup>-1</sup>; specimen: 0.48 × 0.20 × 0.08 mm; *T*<sub>min/max</sub> = 0.81. 2θ<sub>max</sub> = 67°; *N*<sub>t</sub> = 10432, *N* = 5365 (*R*<sub>int</sub> = 0.026), *N*<sub>o</sub> = 4826. *R*<sub>1</sub> = 0.029, *wR*<sub>2</sub> = 0.074 (*a* = 0.021, *b* = 0.71), *S* = 1.10. |Δρ<sub>max</sub>| = 0.86 e Å<sup>-3</sup>.

**12H<sub>2</sub>·2PF<sub>6</sub>** ≡ C<sub>19</sub>H<sub>26</sub>F<sub>12</sub>N<sub>4</sub>P<sub>2</sub>, *M* = 600.4. Monoclinic, space group *P*2<sub>1</sub>/*n* (*C*<sub>2h</sub><sup>5</sup>, No. 14; variant), *a* = 8.1345(5), *b* = 26.145(2), *c* = 11.9508(5) Å, β = 108.052(5)°, *V* = 2417 Å<sup>3</sup> (*T* ca. 100 K). *D*<sub>c</sub> (*Z* = 4) = 1.65<sub>0</sub> g cm<sup>-3</sup>. μ<sub>Mo</sub> = 0.29 mm<sup>-1</sup>; specimen: 0.32 × 0.20 × 0.07 mm; *T*<sub>min/max</sub> = 0.81. 2θ<sub>max</sub> = 68°; *N*<sub>t</sub> = 30072, *N* = 9496 (*R*<sub>int</sub> = 0.033), *N*<sub>o</sub> = 4980. *R*<sub>1</sub> = 0.041, *wR*<sub>2</sub> = 0.106 (*a* = 0.058), *S* = 0.87. |Δρ<sub>max</sub>| = 0.45 e Å<sup>-3</sup>.

**15·HCO<sub>3</sub>·H<sub>2</sub>O** ≡ C<sub>20</sub>H<sub>27</sub>AgN<sub>4</sub>O<sub>4</sub>, *M* = 495.3. Triclinic, space group *P* $\bar{1}$ , *a* = 8.3300(10), *b* = 10.8710(10), *c* = 11.9050(10) Å, α = 72.493(3), β = 82.538(3), γ = 77.284 (3)°, *V* = 1001 Å<sup>3</sup> (*T* ca. 150 K). *D*<sub>c</sub> (*Z* = 2) = 1.64<sub>4</sub> g cm<sup>-3</sup>. μ<sub>Mo</sub> = 1.04 mm<sup>-1</sup>; specimen: 0.22 × 0.15 × 0.10 mm; *T*<sub>min/max</sub> = 0.74. 2θ<sub>max</sub> = 75°; *N*<sub>t</sub> = 19861, *N* = 10153 (*R*<sub>int</sub> = 0.025), *N*<sub>o</sub> = 8618. *R*<sub>1</sub> = 0.039, *wR*<sub>2</sub> = 0.13 (*b* = 3.7), *S* = 1.28. |Δρ<sub>max</sub>| = 1.93 e Å<sup>-3</sup>.

*Variata*. Hydrogen atoms, located in associated with residues modelled as bicarbonate and water, were refined in (*x*, *y*, *z*, *U*<sub>iso</sub>).

**16·2PF<sub>6</sub>** ≡ C<sub>19</sub>H<sub>24</sub>F<sub>12</sub>HgN<sub>4</sub>P<sub>2</sub>, *M* = 799.0. Monoclinic, space group *P*2<sub>1</sub> (*C*<sub>2</sub><sup>2</sup>, No. 4), *a* = 8.5280(6), *b* = 12.5400(9), *c* = 11.7590(8) Å, β = 96.925(2)°, *V* = 1248 Å<sup>3</sup> (*T* ca. 153 K). *D*<sub>c</sub> (*Z* = 2) = 2.12<sub>6</sub> g cm<sup>-3</sup>. μ<sub>Mo</sub> = 6.4 mm<sup>-1</sup>;

specimen:  $0.38 \times 0.37 \times 0.33$  mm;  $T_{\min/\max} = 0.58$ .  $2\theta_{\max} = 75^\circ$ ;  $N_t = 24232$ ,  $N = 12340$  ( $R_{\text{int}} = 0.038$ ),  $N_o = 10236$ .  $R1 = 0.036$ ,  $wR2 = 0.077$  ( $a = 0.035$ ),  $S = 1.05$ .  $|\Delta\rho_{\max}| = 3.25 \text{ e } \text{Å}^{-3}$ .  $x_{\text{abs}} = -0.018(4)$ .

## Results and discussion

### Synthesis, reactivity and solution behaviour





Imidazolium salts **12H<sub>2</sub>·2Br**, **12H<sub>2</sub>·2PF<sub>6</sub>**, **13H<sub>2</sub>·2Cl**, and **13H<sub>2</sub>·2PF<sub>6</sub>**

The *meta*- and *para*-xylyl linked bis(imidazolium) salts **12H<sub>2</sub>·2Br** and **13H<sub>2</sub>·2Cl** were prepared by procedures based on the method used by Dias and Jin for the synthesis of a tri-imidazolium analogue of **12H<sub>2</sub><sup>2+</sup>** [33]. The reaction of  $\alpha,\alpha'$ -dichloro-*p*-xylene with two equivalents of *N*-methylimidazole in refluxing dioxane for 24 h afforded **13H<sub>2</sub>·2Cl**, after recrystallisation from ethanol-diethyl ether. In a similar fashion, **12H<sub>2</sub>·2Br** was prepared by the reaction of 1,3-di(bromomethyl)-2,4,6-trimethylbenzene with two equivalents of *N*-methylimidazole in refluxing dioxane. The bis-imidazolium salts **12H<sub>2</sub>·2Br** and **13H<sub>2</sub>·2Cl** are very soluble in polar solvents such as ethanol, methanol, DMSO, DMF and water, but are insoluble in less polar solvents such as dichloromethane, acetone, acetonitrile and diethyl ether. The halide salts **12H<sub>2</sub>·2Br** and **13H<sub>2</sub>·2Cl** were converted to their hexafluorophosphate counterparts by salt metathesis in methanol using KPF<sub>6</sub>. The <sup>1</sup>H and <sup>13</sup>C NMR spectra for solutions of the imidazolium cations **12H<sub>2</sub><sup>2+</sup>** and **13H<sub>2</sub><sup>2+</sup>**, as the halide and hexafluorophosphate salts, display the expected signals.

### Complexes derived from the meta-linked di(imidazolium) cation **12H<sub>2</sub>**

The reaction of **12H<sub>2</sub>·2Br** with Ag<sub>2</sub>O in methanol afforded complex **15·Br** (Scheme 1). The imidazolium salt **12H<sub>2</sub>·2Br** reacts relatively quickly under these conditions to form the silver complex **15**. This reactivity is substantial higher than that of the cyclophane analogue **14** [11], a result that may be a consequence of higher conformational mobility and/or accessibility of the imidazolium C2 positions in **12H<sub>2</sub><sup>2+</sup>** vs. **14**. Salt metathesis by the addition of an aqueous solution of **15·Br** to an aqueous solution of KPF<sub>6</sub> afforded **15·PF<sub>6</sub>** as a grey precipitate, which was isolated in an overall yield of 76%. The bromide salt **15·Br** is freely soluble in highly polar solvents such as DMF, DMSO and water, while **15·PF<sub>6</sub>** is poorly soluble in water but dissolves well in acetonitrile and acetone at room temperature.

<sup>13</sup>C NMR spectra of solutions of **15·PF<sub>6</sub>** in *d*<sub>6</sub>-DMSO display two doublets centred at ca.  $\delta$  179, due to the



**Scheme 1** Synthesis of **15·PF<sub>6</sub>**

carbene carbons bound to the silver centre. The presence of <sup>13</sup>C-<sup>109</sup>Ag and <sup>13</sup>C-<sup>107</sup>Ag coupling of ca. 200 Hz in these signals is consistent with the absence of NHC-exchange processes at the Ag centre [1, 11, 34]. The <sup>1</sup>H NMR spectra of solutions of **15·PF<sub>6</sub>** in *d*<sub>6</sub>-acetone display an AX pattern due to the benzylic methylene protons, indicating that **15** is conformationally rigid on the NMR time-scale at room temperature (i.e., that the mesitylene ring does not “flip”, e.g., by a process involving the 2-methyl group passing through the metallocyclic ring). In *d*<sub>6</sub>-DMSO solutions prepared from **15·PF<sub>6</sub>**, **15** displayed excellent thermal stability, no decomposition being observed by <sup>1</sup>H NMR when the solution was heated at 100 °C for 2 days.


The mononuclear mercury complex **16** was obtained by the reaction of **12H<sub>2</sub>·2PF<sub>6</sub>** with Hg(OAc)<sub>2</sub> in acetonitrile at reflux for 3 days (Scheme 2). The salt **16·2PF<sub>6</sub>** was obtained as colourless crystals in 55% yield after recrystallisation

from a mixture of acetonitrile and water. The NMR spectra for solutions of  $16 \cdot 2PF_6$  were similar to those of the silver analogue  $15 \cdot PF_6$ . The  $^1H$  NMR spectra of solutions of  $16 \cdot 2PF_6$  in  $d_3$ -acetonitrile display an AB pattern due to the benzylic methylene protons, indicating some conformational rigidity in the structure. In the  $^{13}C$  NMR spectra, for the same solution, a signal at ca.  $\delta$  173 is attributed to the carbene carbon bound to mercury, which is consistent with the literature [5].


#### Complexes derived from the *para*-linked di(imidazolium) cation $13H_2$

Reaction of  $13H_2 \cdot 2Cl$  with an excess of  $Ag_2O$  in methanol (Scheme 3) afforded the salt  $17 \cdot 2Cl$  in 52% yield after recrystallisation from hot methanol. The reaction also proceeds well in DMF or DMSO. The salt  $17 \cdot 2Cl$  is soluble in DMF, DMSO and ethanol, and sparingly soluble in methanol. The hexafluorophosphate salt  $17 \cdot 2PF_6$  was prepared similarly, by reacting  $17 \cdot 2PF_6$  with  $Ag_2O$  in acetonitrile, and was isolated in 77% yield. Removal of the associated  $AgPF_6$  by-product proved problematic. One procedure that can be used to isolate a hexafluorophosphate salt from a reaction mixture is the selective precipitation of the salt by addition of water ( $AgPF_6$  remains dissolved). This method could not be used here because on addition of water to the reaction mixture the material darkened immediately. However, careful washing of the crude reaction product with dichloromethane, followed by recrystallisation from acetonitrile/diethyl ether, afforded pure  $17 \cdot 2PF_6$ .

The results of an electrospray ionisation mass spectrometry study were consistent with the dimeric structure  $[L_2Ag_2]^{2+}$  for the cation **17**. The study showed a molecular ion cluster with prominent peaks near 891 (55% relative intensity), 892 (30%), 893 (100%), 894 (40%), 895 (55%), and 896 (15%) amu, as expected for the  $[17 \cdot PF_6]^+$  ion pair (i.e.,  $\{[L_2Ag_2]PF_6\}^+$ ). The  $^1H$  NMR spectra for solutions of  $17 \cdot 2PF_6$  in  $d_6$ -DMSO indicate that the cation **17** exhibits some conformational flexibility in solution at ambient temperature. For example, the  $^1H$  NMR signal attributed to the benzylic protons appears as a singlet, which is



**Scheme 2** Synthesis of  $16 \cdot 2PF_6$



**Scheme 3** Synthesis of  $17 \cdot 2Cl$


consistent with a situation where the two protons on each benzylic carbon have their environments rendered chemically equivalent by conformational changes that are rapid on the NMR timescale. Similar to the situation for  $15 \cdot PF_6$ , the  $^{13}C$  NMR spectra for solutions of  $17 \cdot 2PF_6$  in  $d_6$ -DMSO displayed two doublets centred near  $\delta$  180, attributed to the carbene carbons bound to  $^{107}Ag$  and  $^{109}Ag$  centres. For a solution of  $17 \cdot 2Cl$  in  $d_6$ -DMSO, however, the  $^{13}C$  NMR spectrum displayed only one broad signal at ca.  $\delta$  180. The lack of any observed  $^{13}C$ - $^{107/109}Ag$  coupling is common for silver complexes with halide counter anions, and has been attributed to halide-mediated Ag-NHC exchange processes [1, 35].

It is interesting that the *para*-linked di(imidazolium) salt  $13H_2 \cdot 2Cl$  reacted with  $Ag_2O$  to form a dimeric structure  $[L_2Ag_2]^{2+}$  (**17**), whereas **11** was isolated from the reaction of  $Ag_2O$  with a *para*-linked di(benzimidazolium) salt [29]. Two factors that might be responsible for the different courses of these reactions are the reaction solvents employed and the presence or absence of methyl substituents on the *para*-phenylene linker. In the reaction leading to **17**, the solvent was methanol, from which  $AgCl$  precipitates, leaving only one  $Ag^+$  per bis(NHC) ligand, whereas for the reaction leading to **11**, the solvent was dichloromethane, from which silver halides are less likely to precipitate, leaving excess silver and bromide in solution to form the  $Ag_2Br_2$  core in **11**. Alternatively, the permethylation of the *para*-phenylene linker in **11** may serve to disfavour formation of a product of form  $[L_2Ag_2]^{2+}$ , with the methyl groups providing the extra steric encumbrance that destabilise the  $L_2Ag_2$  macrocycle.

Similar to those of **15**.PF<sub>6</sub>, solutions of **17**.2PF<sub>6</sub> in *d*<sub>6</sub>-DMSO displayed no decomposition (as observed by <sup>1</sup>H NMR) when heated at 100 °C for 2 days.

Silver-NHC complexes provide a synthetic pathway to other NHC-metal complexes via transmetalation reactions [1]. Since examples of palladium complexes of *para*-xylyl-linked bis(NHC) ligands are rare [27], we have explored the use of **17**.2Cl in transmetalation reactions. The silver complex **17**.2Cl, prepared by the reaction of **13**H<sub>2</sub>.2Cl with Ag<sub>2</sub>O in methanol, was treated with two equivalents of PdCl<sub>2</sub>(NCCH<sub>3</sub>)<sub>2</sub> in acetonitrile at 60 °C. A precipitate, presumably AgCl, formed immediately. Filtration of the mixture, followed by drying of the filtrate under vacuum afforded a yellow powder. Results of elemental analysis of the bulk material were consistent with structures involving bridging chlorides, such as **19** (intermolecular chloride bridges) or **20** (intramolecular chloride bridges). However, the <sup>1</sup>H and <sup>13</sup>C NMR spectra of solutions of the powder in *d*<sub>3</sub>-acetonitrile were consistent with formation of a solvated complex of structure **18**. Disruption of the chloride bridging to afford a solvated complex **18** would occur on dissolution of either **19** or **20** in acetonitrile. While the reaction of **17**.2Cl with PdCl<sub>2</sub>(NCCH<sub>3</sub>)<sub>2</sub> was conducted in acetonitrile and thus might be expected to yield the solvated complex **18**, the vacuum-drying step presumably resulted in the isolation of a non-solvated product. It has been shown previously that the acetonitrile ligand in complexes of the type *trans*-(NHC)PdX<sub>2</sub>(NCCH<sub>3</sub>) is extremely labile, such that even washing the complex with diethyl ether can result in loss of the coordinated acetonitrile and formation of non-solvated bridging-halide complexes [36, 37]. After the yellow powder was carefully recrystallised by very slow evaporation of an acetonitrile solution and isolated without any drying step, an X-ray study indicated that the crystals contained the solvated structure **18** (see below).

We recently reported the first example of redox-transmetalation using a mercury-NHC complex [5]. In view of the apparently labile nature of mercury-NHC complexes, we have explored their use in non-redox transmetalation (“carbene-transfer”) reactions. The reaction of **13**H<sub>2</sub>.2Cl with Hg(OAc)<sub>2</sub> in methanol at reflux for 24 h afforded the mercury analogue of **17**. The Hg complex was identified on the basis of the similarity of its <sup>1</sup>H NMR spectrum with that of the Ag analog **17**. The mercury complex was treated with two equivalents of PdCl<sub>2</sub>(NCCH<sub>3</sub>)<sub>2</sub> in acetonitrile and heated at 60 °C. After 18 h the reaction mixture consisted of a yellow solution and a grey precipitate (presumably HgCl<sub>2</sub> with traces of palladium black). The mixture was filtered and the yellow solution was evaporated to dryness under vacuum to leave a yellow powder. Solutions of this yellow powder in *d*<sub>3</sub>-acetonitrile gave <sup>1</sup>H and <sup>13</sup>C NMR spectra identical to those of the solvated complex **18** isolated from the silver transfer reaction.




We have found that both the silver and the mercury transmetalation experiments can be performed without the need to use inert-atmosphere conditions. The mercury transmetalation route proceeded as well as, if not better than, the silver transmetalation. The mercury complexes made from Hg(OAc)<sub>2</sub> proved easy to purify, and any excess Hg(OAc)<sub>2</sub> or acetic acid was easily removed by washing with water. Silver complexes, especially those made from the bis-imidazolium systems in this study darken easily if exposed to light and therefore need to be protected during synthesis and isolation. Mercury complexes do not need special experimental conditions and perhaps, in this respect, have an advantage over silver complexes.

#### Structure determinations

Cation **13**H<sub>2</sub><sup>2+</sup> has been defined in both bis-hexafluorophosphate and -tetrafluoroborate salts. In both, one half of the formula unit comprises the asymmetric unit, i.e., one anion, devoid of crystallographic symmetry, and one half of the cation, the latter disposed about a crystallographic centre of symmetry in both cases. Both structures are well-ordered. The imidazole ring is twisted out of the plane of the central *p*-xylyl ring in each case (C<sub>3</sub>N<sub>2</sub>/C<sub>6</sub> interplanar dihedral angles 77.75(5), 86.7(3)°, respectively) (Fig. 1). In both salts there are cation⋯anion H⋯F contacts; in the BF<sub>4</sub> salt these are all ≥2.4 Å, but, in the PF<sub>6</sub> salt, contacts to the imidazole hydrogen atoms *ortho* to the xylyl attachment are 2.2<sub>4</sub>, 2.3<sub>9</sub> Å.






**Fig. 1** (Centrosymmetric) Cation  $13H_2^{2+}$  in its hexafluorophosphate salt; the conformation in the tetrafluoroborate is similar

Although silver(I) complexes of **13** and **12** were obtained in bulk as chloride and hexafluorophosphate, and hexafluorophosphate respectively, as described above, the materials obtained were inappropriate for single crystal X-ray studies; smaller quantities of other salts suitable for the X-ray work were obtained adventitiously in the course of variations on the described syntheses. With  $13H_2$ , a nitrate complex was obtained from methanol solution in ambience.

$17 \cdot 2NO_3 \cdot 2MeOH$  is an elegant structure, in which, although there is only one formula unit in the asymmetric unit of the structure, that is comprised of halves of independent dimers disposed pseudo-symmetrically about inversion centres in a triclinic cell (Fig. 2a). The two independent cations are very similar, each comprising a pair of **17** units with their pendant groups oriented quasi-trans about a quasi-inversion centre at the centre of the xylyl ring except that the imidazole rings are oriented so that the carbene

**Fig. 2** **a** Unit cell contents of  $17 \cdot 2NO_3 \cdot 2CH_3OH$  projected down *a*. **b** Projections of cation 1 (cation 2 is similar): (i) normal to, and (ii) through, the  $Ag_2(xylyl\ centroid)_2$  plane




groups are directed to the same 'side' of the ligand. The pair of ligands may then be bridged by a pair of silver atoms to form a 22-membered (centrosymmetric) metallocycle (Fig. 2b(i)), the ring in projection through the xylyl centroids (or nearly so) having a 'Z' aspect (Fig. 2b(ii)). Ag-C are 2.116, 2.088; 2.107, 2.082(4) Å (units 1;2), with C-Ag-C 172.8; 175.4(2)° comparable to other silver-carbene distances, below and elsewhere [35].  $C_3N_2/C_6$  interplanar dihedral angles are 73.8, 75.2; 81.0, 77.0(2)°. Within the nitrate ions, N-O are 1.212, 1.243(6), 1.263(5); 1.247(6), 1.247, 1.250(5) Å, with opposed O-N-O angles 121.3(5), 121.4(4), 117.3; 120.8, 119.3, 119.9(5)°. The methanol C-O distances are 1.379(7); 1.393(8) Å, the hydroxyl groups being hydrogen-bonded to the nitrate ions (H, O...O 1.96(6), 2.800(7); 2.11(7), 2.864(7) Å).

As a ligand, **13** also forms a bis(carbene) complex with acetonitrile solvated palladium(II) chloride, shown to be **18**·2CH<sub>3</sub>CN, the neutral complex molecule also a centrosymmetric species, with one half of the array comprising the asymmetric unit of the structure (Fig. 3); the  $C_3N_2/C_6$  interplanar dihedral angle is 81.4(1)°. The pair of *trans*-Pd-Cl distances in the quasi-square-planar palladium environment are 2.2964, 2.2998(6) Å, Cl-Pd-Cl 178.46(2)°. Pd-N(CH<sub>3</sub>CN) is 2.081(2) and Pd-C(carbene) 1.950(2) Å, N-Pd-C being 177.89(8)°, N-Pd-Cl are 90.50, 91.04(6) and C-Pd-Cl 88.92, 89.55(6)°. The Cl<sub>2</sub>PdNC array is essentially planar ( $\chi^2$  293), the dihedral angle to the associated  $C_3N_2$  plane being 66.99(8)°. Distances of the palladium atom to methyl and methylene hydrogen atoms to either side are 2.9 Å. The solvating acetonitrile molecules have an association which presents the aspect of a proto-inclusion complex (Fig. 3), lying above and below the central *p*-xylyl ring and quasi-parallel to it and to the imidazole ring to one side. Their closest contact, however, is from the nitrogen atom to methylene and imidazole hydrogen atoms of an adjacent complex (N...H, 2.6 Å (×2)).


Cation **12H<sub>2</sub><sup>2+</sup>** has also been isolated as a bis(hexafluorophosphate) salt. Here, the imidazole pendants lie to the same side of the central ring, both inclined at similar angles ( $C_3N_2/C_6$  interplanar dihedrals: 84.82, 79.58(5)°), so that the cation has quasi-m symmetry, (to which the imidazole methyl substituents do not conform), with a pair of ring hydrogen atoms, ortho to the pendants, directed 'inwards'/towards each other, so as to 'chelate' one of the fluorine atoms of one of the anions (H...F 2.2<sub>5</sub>, 2.3<sub>x</sub> Å), while one of the other ortho hydrogen atom contacts one of the fluorine atoms of the other anion (H...F 2.2<sub>8</sub> Å) (Fig. 4). The 'chelation' mode is consummated, with replacement of the pair of 'chelating' hydrogen atoms by a single metal atom in a bis(carbene) complex, in two crystallographically characterized derivatives of silver(I) and mercury(II):

**15·HCO<sub>3</sub>·H<sub>2</sub>O**. Here all hydroxylic hydrogen atoms are confirmed by refinement in (*x*, *y*, *z*, *U*<sub>iso</sub>) using good quality



**Fig. 3** Projections of **18**·2CH<sub>3</sub>CN (a) through and (b) normal to, the central *p*-xylyl ring


data, unambiguously establishing the nature of the complex, deposited from a solution of the bromide on standing in ambience. The pair of imidazole rings now lie quasi-normal to the central ring ( $C_3N_2/C_6$  interplanar dihedrals: 89.7, 88.9(1);  $C_3N_2/C_3N_2$ : 2.2(1)°), and almost mutually coplanar (Fig. 5a), Ag-C are 2.093, 2.094(3) Å, C-Ag-C 178.3(1)°, similar to values recorded for other bis(carbene)silver(I) arrays elsewhere [35] and above. The cation symmetry is quasi-m. The anions and solvent molecules form a separate one-dimensional array, disposed about crystallographic



**Fig. 4** The asymmetric unit of  $12\text{H}_2\cdot 2\text{PF}_6$ , showing the  $\text{H}\cdots\text{F}$  interactions, and the ‘chelation’ of one of the anions by the cation of quasi- $m$  symmetry


centres of symmetry along  $b$ :  $\cdots\text{OCO}\cdot\text{OH}(i)\text{HO}\cdot\text{OCO}$  ( $\text{HOH}$ ) $_2\text{OCO}\cdot\text{OH}\cdots$  (Fig. 5b). Within the bicarbonate anion,  $\text{C}\text{--}\text{O}(2,3)$  are 1.265, 1.206(4) Å with  $\text{O}(2)\text{--}\text{C}\text{--}\text{O}(3)$  126.6(3)°;  $\text{C}\text{--}\text{O}(1)(\text{H})$  is 1.356(4) Å, with  $\text{O}(1)\text{--}\text{C}\text{--}\text{O}(2,3)$  116.8, 116.6(3)°.  $\text{O}(3)\cdots\text{O}(4,4')$  are 2.809(5), 2.871(4) Å, with  $\text{H}\cdots\text{O}$  distances 1.98, 2.03(3) Å.  $\text{O}(1)\cdots\text{O}(2)$  are 2.610(3) with  $\text{H}(1)\cdots\text{O}(2)$  1.79(5) Å. A similar cation disposition is found in:

**Fig. 5 a** The cation **15** in  $15\cdot\text{HCO}_3\cdot\text{H}_2\text{O}$ . **b** The bicarbonate/water molecule polymeric strand




**16·2PF<sub>6</sub>** (Fig. 6). Here the  $\text{C}_3\text{N}_2/\text{C}_6$  interplanar dihedral angles are 80.0, 77.6(2)°, the  $\text{C}_3\text{N}_2/\text{C}_3\text{N}_2$  inclination here being greater (22.4(2)°), although  $\text{M}\text{--}\text{C}$  distances here are almost identical to those of the silver(I) complex ( $\text{Hg}\text{--}\text{C}$  2.078, 2.086(4) Å;  $\text{C}\text{--}\text{Hg}\text{--}\text{C}$  176.8(1)°); again  $\text{Hg}\text{--}\text{C}$  are similar to those recorded in other mercury/carbene arrays [5]. In this complex the two anions approach the mercury atom from either side of the  $\text{Hgim}_2$  ‘plane’,  $\text{Hg}\cdots\text{F}(11,12)$  being 3.085, 3.228(4), and  $\text{Hg}\cdots\text{F}(21,22)$  3.065, 3.040(4) Å (Fig. 6b).

The structure of the cation **15** contrasts with structures such as **21** and **22**, which are also based on bis(NHC)s with *meta*-xylyl-type linking motifs [16, 24, 26]. Different coordination modes may arise due to the presence of coordinating vs non-coordinating anions (**21** was formed in the presence of chloride), solvent effects, etc. In the present case, however, internal steric hindrance between the mesityl C2 methyl substituent and the imidazolyl units may disfavour dinuclear binding motifs such as exemplified by **22** [26] and **23** [11]. In the latter, for example, the mesityl groups are distinctly distorted from planarity, with the C2 methyl group pushed out of the arene plane away from the imidazolyl groups. The ligand **12** can be thought of as derived from the cyclophane ligand found in **23**, and the absence of the *ortho*-xylyl linker leaves **12** with



**Fig. 6** **a** The cation **16** in **16**·**2PF**<sub>6</sub>. **b** Projection showing the approaches of the anions

conformational freedom to adopt the interesting trans-spanning chelating coordinating mode rather than a sterically disfavoured dinuclear structure.



**Acknowledgements** We thank the Australian Research Council for a Discovery Grant (to M.V.B. and A.H.W.), the Universiti Sains Malaysia for a postgraduate research scholarship (to R.A.H.) and Curtin University of Technology for a Curtin Research and Teaching Fellowship (to D.H.B.).

## References

- Lin, I.J.B., Vasam, C.S.: Preparation and application of N-heterocyclic carbene complexes of Ag(I). *Coord. Chem. Rev.* **251**, 642–670 (2007)
- Kascatan-Nebioglu, A., Panzner, M.J., Tessier, C.A., Cannon, C.L., Youngs, W.J.: N-Heterocyclic carbene silver complexes: a new class of antibiotics. *Coord. Chem. Rev.* **251**, 884–895 (2007)
- Medvetz, D.A., Hindi, K.M., Panzner, M.J., Ditto, A.J., Yun, Y.H., Youngs, W.J.: Anticancer activity of Ag(I) N-heterocyclic carbene complexes derived from 4,5-dichloro-1H-imidazole. *Met. Based Drugs* (2008). doi:10.1155/2008/384010
- Wanzlick, H.-W., Schönherr, H.J.: Direct synthesis of a mercury salt-carbene complex. *Angew. Chem. Int. Ed. Engl.* **7**, 141–142 (1968)
- Baker, M.V., Brown, D.H., Haque, R.A., Simpson, P.V., Skelton, B.W., White, A.H., Williams, C.C.: Mercury complexes of N-heterocyclic carbenes derived from imidazolium-linked cyclophanes: synthesis, structure, and reactivity. *Organometallics* **28**, 3793–3803 (2009). doi:10.1021/om8011745
- Wardell, J.L.: Mercury, Chap. 17. In: Wilkinson, G., Stone, F.G.A., Abel, E.W. (eds.) *Comprehensive Organometallic Chemistry*, 1st edn. Pergamon Press, Oxford (1982)
- Baker, M.V., Brown, D.H.: Azolium cyclophanes. *Mini. Rev. Org. Chem.* **3**, 333–354 (2006)
- Baker, M.V., Bosnich, M.J., Williams, C.C., Skelton, B.W., White, A.H.: Imidazolium-linked cyclophanes. *Aust. J. Chem.* **52**, 823–825 (1999)
- Baker, M.V., Bosnich, M.J., Brown, D.H., Byrne, L.T., Hesler, V.J., Skelton, B.W., White, A.H., Williams, C.C.: Azolium-linked cyclophanes: a comprehensive examination of conformations by <sup>1</sup>H NMR spectroscopy and structural studies. *J. Org. Chem.* **69**, 7640–7652 (2004)
- Baker, M.V., Brown, D.H., Heath, C.H., Skelton, B.W., White, A.H., Williams, C.C.: Azolium-linked cyclophanes: effects of structure, solvent, and counteranions on solution conformation behavior. *J. Org. Chem.* **73**, 9340–9352 (2008)
- Baker, M.V., Brown, D.H., Haque, R.A., Skelton, B.W., White, A.H.: Dinuclear N-heterocyclic carbene complexes of silver(I), derived from imidazolium-linked cyclophanes. *Dalton Trans.*, 3756–3764 (2004)
- Baker, M.V., Brown, D.H., Hesler, V.J., Skelton, B.W., White, A.H.: Synthesis of a bis(N-heterocyclic carbene)palladium complex via oxidative addition of a C–C bond in a biimidazolium ion. *Organometallics* **26**, 250–252 (2007)
- Baker, M.V., Brown, D.H., Simpson, P.V., Skelton, B.W., White, A.H., Williams, C.C.: Palladium, rhodium and platinum complexes of *ortho*-xylyl-linked bis-N-heterocyclic carbenes: synthesis, structure and catalytic activity. *J. Organomet. Chem.* **691**, 5845–5855 (2006)
- Baker, M.V., Skelton, B.W., White, A.H., Williams, C.C.: Synthesis and characterization of a saddle-shaped nickel-carbene complex derived from an imidazolium-linked *meta*-cyclophane. *Organometallics* **21**, 2674–2678 (2002)
- Magill, A.M., McGuinness, D.S., Cavell, K.J., Britovsek, G.J.P., Gibson, V.C., White, A.J.P., Williams, D.J., White, A.H., Skelton, B.W.: Palladium(II) complexes containing mono-, bi- and tridentate carbene ligands. Synthesis, characterisation and

- application as catalysts in C-C coupling reactions. *J. Organomet. Chem.* **617–618**, 546–560 (2001)
16. Garrison, J.C., Simons, R.S., Talley, J.M., Wesdemiotis, C., Tessier, C.A., Youngs, W.J.: Synthesis and structural characterisation of an imidazolium-linked cyclophane and the silver complex of an N-heterocyclic carbene-linked cyclophane. *Organometallics* **20**, 1276–1278 (2001)
  17. Willans, C.E., Anderson, K.M., Junk, P.C., Barbour, L.J., Steed, J.W.: A small tris(imidazolium) cage forms an N-heterocyclic carbene complex with silver(I). *Chem. Commun.*, 3634–3636 (2007)
  18. Hahn, F.E., Radloff, C., Pape, T., Hepp, A.: Synthesis of silver(I) and gold(I) complexes with cyclic tetra- and hexacarbene ligands. *Chem. Eur. J.* **14**, 10900–10904 (2008)
  19. Baker, M.V., Skelton, B.W., White, A.H., Williams, C.C.: Palladium carbene complexes derived from imidazolium-linked *ortho*-cyclophanes. *J. Chem. Soc. Dalton Trans.*, 111–120 (2001)
  20. Barnard, P.J., Wedlock, L.E., Baker, M.V., Berners-Price, S.J., Joyce, D.A., Skelton, B.W., Steer, J.H.: Luminescence studies of the intracellular distribution of a dinuclear gold(I) N-heterocyclic carbene complex. *Angew. Chem. Int. Ed.* **45**, 5966–5970 (2006)
  21. Danopoulos, A.A., Tulloch, A.A.D., Winston, S., Eastham, G., Hursthouse, M.B.: Chelating and ‘pincer’ dicarbene complexes of palladium; synthesis and structural studies. *Dalton Trans.*, 1009–1015 (2003)
  22. Tulloch, A.A.D., Danopoulos, A.A., Winston, S., Kleinhenz, S., Eastham, G.: N-Functionalised heterocyclic carbene complexes of silver. *J. Chem. Soc. Dalton Trans.*, 4499–4506 (2000)
  23. Alcalde, E., Ceder, R.M., López, C., Mesquida, N., Muller, G., Rodríguez, S.: Coordination features of bis(N-heterocyclic carbene) and bis(oxazolines) with 1,3-alkylidene-2,4,6-trimethylbenzene spacers. Synthesis of the ligands and silver and palladium complexes. *Dalton Trans.*, 2696–2706 (2007)
  24. Chen, W., Wu, B., Matsumoto, K.: Synthesis and crystal structure of N-heterocyclic carbene complex of silver. *J. Organomet. Chem.* **654**, 233–236 (2002)
  25. Simons, R.S., Custer, P., Tessier, C.A., Youngs, W.J.: Formation of N-heterocyclic complexes of rhodium and palladium from a pincer silver(I) carbene complex. *Organometallics* **22**, 1979–1982 (2003)
  26. Nielsen, D.J., Cavell, K.J., Skelton, B.W., White, A.H.: A pyridine bridged dicarbene ligand and its silver(I) and palladium(II) complexes: synthesis, structure, and catalytic applications. *Inorg. Chim. Acta* **327**, 116–125 (2002)
  27. Schneider, S.K., Schwarz, J., Frey, G.D., Herdtweck, E., Herrmann, W.A.: Chiral, bridged bis(imidazolium-2-ylidene) complexes of palladium. *J. Organomet. Chem.* **692**, 4560–4568 (2007)
  28. Türkmen, H., Denizalti, S., Özdemir, I., Çetinkaya, E., Çetinkaya, B.: Synthesis and use of mono- or bisxylyl linked bis(benzimidazolium) bromides as carbene precursors for C-C bond formation reactions. *J. Organomet. Chem.* **693**, 425–434 (2008)
  29. Liu, Q.-X., Zhao, X.-J., Wu, X.-M., Yin, L.-N., Guo, J.-H., Wang, X.-G., Feng, J.-C.: Two new N-heterocyclic carbene silver(I) complexes with the  $\pi$ - $\pi$  stacking interactions. *Inorg. Chim. Acta* **361**, 2616–2622 (2008)
  30. Wimmer, F.L., Wimmer, S., Castan, P.: (2,2'-Bipyridine)dichloropalladium (II) and [*N'*-(2-diaminoethyl)-1,2-ethanediamine] chloropalladium(II) chloride. *Inorg. Synth.* **29** 185–187 (1992)
  31. van der Made, A.W., van der Made, R.H.: A convenient procedure for bromomethylation of aromatic compounds. Selective mono-, bis-, or tribromomethylation. *J. Org. Chem.* **58**, 1262–1263 (1993)
  32. Sheldrick, G.M.: A short history of SHELX. *Acta Crystallogr. Sect. A* **64**, 112–122 (2008)
  33. Dias, H.V.R., Jin, W.: A stable tridentate carbene ligand. *Tetrahedron Lett.* **35**, 1365–1366 (1994)
  34. Wang, H.M.J., Lin, I.J.B.: Facile synthesis of silver(I)-carbene complexes. Useful carbene transfer agents. *Organometallics* **17**, 972–975 (1998)
  35. Garrison, J.C., Youngs, W.J.: Ag(I) N-heterocyclic carbene complexes: synthesis, structure, and application. *Chem. Rev.* **105**, 3978–4008 (2005)
  36. Han, Y., Hong, Y.-T., Huynh, H.V.: Ag(I) and Pd(II) complexes of a 1,3-dibenzhydryl substituted benzannulated N-heterocyclic carbene: unexpected rearrangement, structures and catalytic studies. *J. Organomet. Chem.* **693**, 3159–3165 (2008)
  37. Huynh, H.V., Han, Y., Ho, J.H.H., Tan, G.K.: Palladium(II) complexes of a sterically bulky, benzannulated N-heterocyclic carbene with unusual intramolecular C-H...Pd and C<sub>carbene</sub>...Br interactions and their catalytic activities. *Organometallics* **25**, 3267–3274 (2006)