

A Rapid Synthesis of [ $^{82}\text{Br}$ ]-2-Deoxy-2-bromo-D-mannose and [ $^{82}\text{Br}$ ]-2-Deoxy-2-bromo-D-glucose using D-Glucal as Precursor

Yoshio Homma, Yuko Murase, Michiko Ishii and Kazue Sonehara

Kyoritsu College of Pharmacy

Shibakoen 1-5-30, Minato-ku, Tokyo 105, Japan

SUMMARY

A new rapid synthesis of [ $^{82}\text{Br}$ ]-2-deoxy-2-bromo-D-mannose and [ $^{82}\text{Br}$ ]-2-deoxy-2-bromo-D-glucose is described. Reaction of  $^{82}\text{Br}_2$  with D-glucal in  $\text{CHCl}_3$ , followed by treatment with 1 N HCl gave [ $^{82}\text{Br}$ ]-2-deoxy-2-bromo-D-mannose and [ $^{82}\text{Br}$ ]-2-deoxy-2-bromo-D-glucose in the yield of 61.1 % and 27.7 %, respectively. The synthesis time was 20 min. The radiochemical purity of [ $^{82}\text{Br}$ ]-2-deoxy-2-bromo-D-mannose and [ $^{82}\text{Br}$ ]-2-deoxy-2-bromo-D-glucose was greater than 95 %. The radiochemical yield of [ $^{82}\text{Br}$ ]-2-deoxy-2-bromo-D-mannose and [ $^{82}\text{Br}$ ]-2-deoxy-2-bromo-D-glucose was 14.3 % and 31.6 %, respectively. D-Glucal was shown to be a useful precursor to allow rapid production of usable amounts of glucose analogs such as [ $^{75}\text{Br}$ ]-2-deoxy-2-bromo-D-mannose and [ $^{75}\text{Br}$ ]-2-deoxy-2-bromo-D-glucose.

Key Words: glucose analog; bromination;  $^{75}\text{Br}$ ; FDG.

INTRODUCTION

[ $^{18}\text{F}$ ]-2-Deoxy-2-fluoro-D-glucose (2- $^{18}\text{F}$ FDG) (1-4), [ $^{18}\text{F}$ ]-3-deoxy-3-fluoro-D-glucose (5) and 1-[ $^{11}\text{C}$ ]-2-deoxy-D-glucose (6) are used

clinically to measure regional cerebral glucose metabolism by positron emission tomography (PET). In particular, 2-<sup>18</sup>F<sub>2</sub>FDG has been proposed as a tracer useful for in vivo PET measurement of glucose metabolism (7-10), and the advantages have been detailed in the literature (11-14).

Since Fischer, Bergmann, and Schotte reported the halogenation of 3,4,6-tri-O-acetyl-D-glucal (TAG) (15), most of the methods used for the synthesis of the halogenated hexose, including those for 2-<sup>18</sup>F<sub>2</sub>FDG, are based on the reaction of halogenation agents with TAG (11-14). The limitation of this radiopharmaceutical is that the reported production methods for the compound give relatively low chemical and radiochemical yields (1-4).

In previous reports, the present authors described the usefulness of D-glucal, which is a kind of the simplest unsaturated sugar derivatives, as a precursor for synthesis of glucose analogs and reported that the dihalogen addition products of D-glucal have reasonably greater brain uptakes than 2-deoxy-D-glucose and D-glucose (16,17). From the standpoint of blood-to-brain transport, these dihalogeno compounds are very interesting; however, the halogen at carbon atom 1 was found to be relatively unstable in vivo. Therefore, we have searched for methods which will improve the yield of glucose analogs, or analogs which will behave like 2-<sup>18</sup>F<sub>2</sub>FDG.

In this paper we report a rapid synthetic procedure for [<sup>82</sup>Br]-2-deoxy-2-bromo-D-mannose (2-<sup>82</sup>BrDM) and [<sup>82</sup>Br]-2-deoxy-2-bromo-D-glucose (2-<sup>82</sup>BrDG) which are obtained on direct addition of <sup>82</sup>Br<sub>2</sub> to D-glucal.

Advantages of this method are that a rapid and selective direct addition of <sup>82</sup>Br<sub>2</sub> to D-glucal, which had been synthesized and stored prior to the bromination reaction, leads to 2-<sup>82</sup>BrDM and 2-<sup>82</sup>BrDG in high yields.

Since our previous papers, the works of Ehrenkauffer et al. (18)

and Van Rijn et al. (19) have appeared, in which the usefulness of the experimental approach using D-glucal as a precursor for synthesis of glucose analogs has been described.

#### MATERIALS AND METHODS

The synthetic route adopted for the preparation of D-glucal and 2-<sup>82</sup>BrDM and 2-<sup>82</sup>BrDG is shown in Fig.1.

3,4,6-Tri-O-acetyl-D-glucal (1) (5.86 g) was dissolved in dry methanol (100 ml), and sodium (15 mg) was added. The solution was allowed to remain at room temperature for 24 hr, treated with carbon dioxide and evaporated under diminished pressure. The residue was extracted with hot ethyl acetate. The combined extracts were concentrated to afford D-glucal (2) m.p. 57-59 °C,  $[\alpha]_{19}^D -8.0$  (c 1.88 in water).

Bromination of D-glucal was performed quite readily by the direct addition of bromine to the unsaturated linkage of 2 as follows. A stirred solution of 0.03 g of 2 in 100 ml of CHCl<sub>3</sub> was cooled in an ice bath and protected from light. Bromine (<sup>82</sup>Br<sub>2</sub>) was added dropwise until the color caused by excess bromine was present. Then, nitrogen gas was bubbled into the solution to remove the excess bromine, and the solvent was evaporated *in vacuo* at 40-43 °C to give a mixture of [<sup>82</sup>Br]-2-deoxy-2-bromo-D-mannopyranosyl bromide (3) and [<sup>82</sup>Br]-2-deoxy-2-bromo-D-glucopyranosyl bromide(4). The specific activity of bromine (<sup>82</sup>Br<sub>2</sub>) was 67.3 μCi/mmol. It is noteworthy that the bromination process was completed within a few minutes in a 98 % yield. The mixture was refluxed with 1 N hydrochloric acid (2.5 ml) for 5 min. at 110 °C. Then the solution was cooled, neutralized with silver carbonate, filtered, and evaporated. Further purification was performed on a column consisting of charcoal, an ion-retardation resin and alumina. The reaction mixture was analysed on TLC.




Fig. 1. Synthesis of [<sup>82</sup>Br]-2-deoxy-2-bromo-D-mannose and [<sup>82</sup>Br]-2-deoxy-2-bromo-D-glucose.

#### RESULTS AND DISCUSSION

Examination of the hydrolysate by TLC [CHCl<sub>3</sub>-CH<sub>3</sub>OH (6:4)] indicates two major components with R<sub>f</sub> values of 0.86 and 0.77, whose configurations have been assigned to be 2-deoxy-2-bromo-D-mannose (**5**) and 2-deoxy-2-bromo-D-glucose (**6**), respectively, by the following reaction sequence. Thin layer radiochromatography (silica gel 20 cm glass plates, mobile phase chloroform-methanol 6:4, radiochromatogram scanner, Packard model 7201) also showed single spots with R<sub>f</sub> 0.86 and 0.77, respectively.

The R<sub>f</sub> values were identical to those of authentic 2-deoxy-2-bromo-D-mannose and 2-deoxy-2-bromo-D-glucose, which were obtained by hydrolyzing 3,4,6-tri-O-acetyl-2-deoxy-2-bromo-D-mannopyranosyl

bromide and 3,4,6-tri-O-acetyl-2-deoxy-2-bromo-D-glucopyranosyl bromide respectively, according to the method of Fischer et al. (15) and Lemieux et al. (20).

It is interesting to note that the addition of  $^{82}\text{Br}_2$  to D-glucal (2) in  $\text{CHCl}_3$  forms D-manno (5) and D-gluco (6) configurations in a 2.2 : 1 ratio, whereas the addition of  $\text{Br}_2$  to TAG (1) gives  $\alpha$ -D-manno and  $\alpha$ -D-gluco configurations in a 1 : 2 ratio (15). Experiments are being undertaken to further study the stereoselectivity of these reactions, and the results will be reported and discussed elsewhere.

The yield, which was determined by collecting and counting the peaks for 5 and 6 comparing the results with an equivalent sample of the reaction solution, was 61.1 % for 5 and 27.7 % for 6, respectively.

Under standard conditions (reaction time 5 min at room temperature, hydrolysis of the products with 1 N HCl at 110 °C for 5 min, followed by silica gel column purification) 4.2  $\mu\text{Ci}$  of 5 and 2.0  $\mu\text{Ci}$  of 6 were obtained with synthetic time of 20 - 25 min. The specific activity of the 2- $^{82}\text{BrDM}$  and 2- $^{82}\text{BrDG}$  was 33.6  $\mu\text{Ci}/\text{mmol}$ .

Advantages of this method are as follows:

- (1) The reported synthetic procedures of 2-FDG and 3-FDG require 1.5 - 2 h (1,5), whereas direct addition to D-glucal is completed within a few minutes and the overall sample preparation for clinical studies is about 35 min. Note that this method is especially advantageous for production of glucose analogs labelled with short half-lived radionuclides such as  $^{75}\text{Br}$ .
- (2) The precursor of the proposed method, D-glucal, is synthesized from TAG by way of several nonradioactive steps and can readily be available for years without any detectable chemical alteration provided the compound is kept in a refrigerator.
- (3) This method can readily be adapted for the synthesis of  $^{75}\text{Br}$ ,  $^{77}\text{Br}$  labelled D-glucose derivatives, usefulness of which as poten-

tial radiopharmaceutical for tracing D-glucose utilization in brain and heart had been described.

## REFERENCES

1. Ido T., Wan C-N., Casella V., et al. - J. Labelled Comp. Radiopharm. 14: 175 (1978).
2. Fowler J.S., MacGregor R.R., Wolf A.P., et al. - J. Nucl. Med. 22: 376 (1981).
3. Barrio J.R., MacDonald N.S., Robinson G.D., et al. - J. Nucl. Med. 22: 372 (1981).
4. Shaughnessy W.J., Gatley S.J., Hichwa R.D., et al. - Int. J. Appl. Radiat. Isot. 32: 23 (1981).
5. Tewson T.J., Welch M.J. and Raichle M.E. - J. Nucl. Med. 19: 1339 (1978).
6. MacGregor R.R., Fowler J.S., Wolf A.P., et al. - J. Nucl. Med. 22: 800 (1981).
7. Kuhl D.E., Phelps M.E., Kowell A.P., et al. - Ann. Neurol. 8: 47 (1980).
8. Kuhl D.E., Phelps M.E., Markham C., et al. - J. Cereb. Blood Flow Metab. 1: 459 (1981).
9. Kuhl D.E., Engel J. Jr., Phelps M.E., et al. - Ann. Neurol. 8: 348 (1980).
10. Widen L., Bergstron M. and Bromquist C. - J. Cereb. Blood Flow Metab. 1: 455 (1981).
11. Wolf A.P., Christman D.R., Fowler J.S., Lambrecht R.M., et al. - Radiopharmaceuticals and Labelled Compounds, Vol. 1, Copenhagen, IAEA 1973, p 345.
12. Lambrecht R.M. and Wolf A.P. - Radiopharmaceuticals and Labelled Compounds, Vol. 1, Vienna, IAEA, 1973, p 275.

13. Robinson G.D. - Radiopharmaceuticals, Subramanian G., Rhodes B.A., Cooper J.F. and Sodd V.J. eds., New York, Society of Nuclear Medicine, 1975, p 141.
14. Palmer A.J., Clark J.C. and Goulding R.W. - Int. J. Appl. Radiat. Isot. 28: 53 (1977).
15. Fischer E., Bergmann M. and Schotte H. - Ber. dt. chem. Ges. 53: 509 (1920).
16. Homma Y., Murase Y. and Ishii M. - J. Radioanal. Chem. 76: 283 (1983).
17. Homma Y., Murase Y. and Ishii M. - J. Labelled Comp. Radiopharm. 19: 1633 (1982).
18. Ehrenkaufner R.E., Potocki J.F. and Jewett D.M. - J. Nucl. Med. 25: 333 (1984).
19. Van Rijn C.J.S., Herscheid J.D.M., Visser G.W.M., et al. - Int. J. Appl. Radiat. Isot. 36: 111 (1985).
20. Lemieux R.U. and Fraser-Reid B. - Can. J. Chem. 42: 532 (1964).