

SYNTHESIS OF [³H]CI-980, ETHYL[5-AMINO-1,2-DIHYDRO-2(S)-METHYL-3-(3-³H)PHENYL)PYRIDO[3,4-*b*]PYRAZIN-7-YL]CARBAMATE ISETHIONATE SALT, A TUBULIN-BINDING, ANTIMITOTIC, BROAD-SPECTRUM ANTITUMOR AGENT

Helen T. Lee and Peter W. K. Woo
Parke-Davis Pharmaceutical Research,
Division of Warner-Lambert Company,
Ann Arbor, Michigan 48105, U.S.A.

SUMMARY

[³H]CI-980 (10b) was synthesized in an eight-step sequence with an overall yield of 2.4%. Reaction of *m*-bromophenyl lithium (2) with *N*-ethoxycarbonyl-L-alanine gave the chiral ketone 3. Reduction of 3 with sodium borohydride followed by alkaline *N*-deprotection and condensation with ethyl 6-amino-4-chloro-5-nitro-2-pyridine carbamate (6) gave 7. Chromium trioxide oxidation of 7 followed by reductive cyclization with iron-acetic acid gave the key bromo intermediate 9. Palladium catalyzed ³H-hydrogenolysis of 9 gave the free base form of [³H]CI-980 (10a), which was converted to the isethionate salt (10b) before use.


Key Words: [³H]CI-980, antitumor agent, catalytic ³H-hydrogenolysis

INTRODUCTION

As indicated in the previous communication, the biologically more active 2(S) enantiomeric component of ethyl[5-amino-1,2-dihydro-2-methyl-3-phenylpyrido[3,4-*b*]pyrazin-7-yl]carbamate (*R,S*-10), a broad spectrum antitumor agent,¹ is being developed clinically as CI-980. During the process, the labeled forms of CI-980 were needed for pharmacokinetics and drug metabolism studies. This paper presents the synthesis of [³H]CI-980.

RESULTS AND DISCUSSION

[³H]CI-980 was synthesized in eight steps (Scheme I). Lithiation of 1 with *n*-butyl lithium at -110°C, followed by reaction with *N*-ethoxycarbonyl-L-alanine²

Scheme I. Synthesis of [³H]CI-980

gave, after chromatography, 21% of pure 3. Reduction of 3 with sodium borohydride in methanol at room temperature, followed by alkaline hydrolysis with

potassium hydroxide in aqueous methanol (2 h, reflux), gave a crude product which was purified by trituration with ethanol; pure **5** (77%) was isolated in the mother liquor. Condensation of **5** with **6** in the presence of triethylamine in ethanol (24 h, reflux) gave, after chromatography, 30% of **7**. Oxidation of **7** with chromium trioxide in pyridine-dichloromethane (2 h, room temperature) gave 66% of **8**. Reductive cyclization of **8** with iron powder in acetic acid-ethanol (3.5 h, reflux) gave 75% of **9**.

Replacement of the bromine in **9** was readily accomplished in 1 h with either hydrogen or deuterium, using Pd/C as catalyst and THF as solvent. Tritiation was then carried out under similar conditions but in the presence of 15 equivalents of unlabeled CI-980 free base as carrier, as no product could be isolated in its absence. The product, after purification by silica gel chromatography, was characterized and stored as the free base **10a**. The free base was converted to [³H]CI-981 (**10b**), the isethionate salt, for pharmacokinetic and metabolic studies immediately before use, as the latter was found to be unstable.

EXPERIMENTAL

Tritiation was performed by Amersham Co. Radioactivity was determined with a Packard Tri-Carb 4530 liquid scintillation counter, using Beckman Ready-Gel as the counting medium. TLC plates, E. Merck silica gel 60 F-254, were scanned on a Berthold LB2832 automatic TLC linear analyzer. Column chromatography was performed using E. Merck silica gel, 230-400 mesh. HPLC was performed using a Spectra Physics SP8700 solvent delivery system, Kratos Spectroflow 773 variable wavelength UV detector, and Radiomatic Beta Flow I radioactivity flow detector. Unless otherwise specified, Alltech Econosil Columns, C18 4.7 mm x 20 cm were used. MS data were obtained by electron impact at 70 eV using a V G Analytical 7070E/HF mass spectrometer.

Ethyl [2-(3-Bromophenyl)-1(S)-methyl-2-oxoethyl]carbamate (**3**).

n-Butyl lithium (30 mL, 75 mmol, 2.5 M) was added dropwise to an ether solution (125 mL) of *m*-dibromobenzene (17.7 g, 75 mmol) at -110°C and stirred at this temperature for 0.5 h. *N*-(Ethoxycarbonyl)-L-alanine² (0.4 g, 24.8 mmol) in

ether (50 mL) was then added slowly. This reaction mixture was allowed to warm up from -110°C to -10°C . 1 M H_3PO_4 (100 mL) was added all at once, and the temperature was kept below 5°C . The residue from evaporation of the ether layer was purified by chromatography over silica gel (3:2, CHCl_3 :hexane) to give product **3** (1.6 g, 21%). ^1H NMR (CDCl_3): δ 1.05-1.5 (m, 6H, CH_3), 4.05 (q, 2H, CH_2), 5.0-5.36 (m, 1H, CH), 5.5 (b, 1H, NH), 7.25-8.0 (m, 4H, Ar). Elemental analysis: calc. for $\text{C}_{12}\text{H}_{14}\text{N}_1\text{O}_3\text{Br}_1$: C 48.02; H 4.70; N 4.67. Found: C 48.13; H 4.82; N 4.83.

α -(1-Aminoethyl)-3-bromobenzenemethanol (5).

Sodium borohydride (361 mg, 9.54 mmol) was added in two portions over 0.5 h to a stirred solution of **3** (1.9 g, 6.33 mmol) in CH_3OH (25 mL) at room temperature. After 0.5 h, acetic acid (2 mL) was added and the reaction was evaporated to dryness. The residue was distributed between CH_2Cl_2 and saturated NaHCO_3 . The aqueous layer was extracted with CH_2Cl_2 (2 x 20 mL), and the combined CH_2Cl_2 portions were evaporated to an oil. Aqueous MeOH (25 mL, 3:1, MeOH:H₂O) containing 0.9 g of KOH was added and the solution was refluxed for 2 h, cooled, and evaporated to dryness. An insoluble impurity was removed by recrystallization from EtOH. Evaporation of the mother liquor gave pure **5**, 1.12 g (77%).

Ethyl [6-Amino-4-[[2-(3-bromophenyl)-2-hydroxy-1-methylethyl]amino]-5-nitro-2-pyridinyl]carbamate (7).

A mixture of **6** (1.017 g, 4.45 mmol), **5** (1.12 g, 4.87 mmol), and Et_3N (0.65 mL) in EtOH (17 mL) was heated under reflux for 24 h, cooled, and evaporated to dryness. The residue was purified by silica gel column chromatography (1% EtOH in CH_2Cl_2) to give 670 mg of **7** (30%). ^1H NMR (CDCl_3): δ 1.18 (d, 3H, CH_3), 1.3 (t, 3H, CH_3), 3.8-4.0 (m, 1H, CH), 4.25 (q, 2H, CH_2), 5.05 (m, 1H, CH), 6.95-7.65 (m, 5H, Ar).

Ethyl [6-Amino-4-[[2-(3-bromophenyl)-1-methyl-2-oxoethyl]amino]-5-nitro-2-pyridinyl]carbamate (8).

A solution of 7 (640 mg, 1.41 mmol) in CH₂Cl₂ (25 mL) was added to a stirred solution of pyridine (1.77 mL in 25 mL CH₂Cl₂), followed by CrO₃ (1.078 g, 10.78 mmol). After stirring at room temperature for 2 h, the mixture was filtered through celite. The filtrate was evaporated, and the product (8) was obtained after purified by silica gel column chromatography (3% EtOH in CHCl₃) as yellow powder (420 mg, 66%). ¹H NMR (CDCl₃): δ 1.3 (t, 3H, CH₃), 1.6 (d, 3H, CH₃) 4.22 (q, 2H, CH₂), 5.16 (t, 1H, CH), 6.8 (s, H, NH), 7.1-7.98 (m, 4H, Ar), 8.05 (s, 1H, Ar). Elemental analysis: calc. for C₁₇H₁₈N₅O₅Br·0.15CHCH₃: C 43.81; H 3.89; N 14.90. Found: C 43.75; H 3.60; N 15.14. MS: m/z 452, 454 [(M+H)⁺; ⁷⁹Br, ⁸¹Br].

Ethyl [5-Amino-3-(3-bromophenyl)-1,2-dihydro-2(S)-methylpyrido[3,4-b]pyrazin-7-yl]carbamate (9).

Iron powder (39.6 mg, 0.71 nmol) was added to a solution of 8 (80 mg, 0.18 mmol) in 6.2 mL of glacial acetic acid and 6.2 mL of abs EtOH. The mixture was heated under reflux for 3.5 h. After addition of H₂O (1 mL) and neutralization with solid Na₂CO₃, the mixture was extracted with CH₂Cl₂ (3 x 20 mL), and the combined extracts were dried over MgSO₄ and concentrated under vacuum. Pure 9 was isolated by column chromatography on silica gel using Et₃N:EtOH:CH₂Cl₂ (3:3:94) as eluent. The yellow powder (9) weighed 70 mg (98%). MS: m/z 404, 406 [(M+H)⁺; ⁷⁹Br, ⁸¹Br].

[³H]CI-980 (10b)

The ²H analog of 10a was first prepared in order to optimize the reaction conditions and isolation procedures. Compound 9 (10 mg, 0.025 mmol) was subjected to ²H-hydrogenolysis in the presence of 5% Pd/C (10 mg) in 4 mL THF under deuterium for 1 h at 1 atm and room temperature. The mixture was filtered and washed with methanol, and the filtrate was neutralized with K₂CO₃ (0.1 g) and

evaporated to dryness. The pure ^2H analog (6 mg, 75%) was isolated by column chromatography using 2% EtOH in CHCl_3 as eluant. MS: m/z 327 $[(\text{M}+\text{H})^+]$; 326 for the reference ^1H analog).

Using similar procedure as for the ^2H analog above but in the presence of unlabeled CI-980 free base as carrier, ^3H -hydrogenolysis of **9** was carried out (8 mg of **9**, 100 mg of unlabeled **10a**, 20 mg Pd/C, 10 mL THF, 5 Ci of tritium gas), and the crude product was isolated as the free base (5 mL methanol rinse, 100 mg K_2CO_3). The crude product was coevaporated three times with ethanol to remove tritium gas, purified by chromatography, and further diluted with unlabeled **10a** to give the tritiated free base product (210 mg; 113 mCi 19.5% of theoretical). The purity of **10a**, according to TLC, was 98.5% (2:98 EtOH: CHCl_3 ; R_f 0.22), and according to HPLC (55% 0.05 M triethylamine adjust to pH 3.1 with formic acid, 45% acetonitrile, 1 mL/min, retention time 5.1 min), was 98.5% radiochemical and 100% chemical (270 nm). ^1H NMR (CDCl_3): δ 1.1-1.35 (m, 6H, CH_3), 4.1-4.35 (q, 2H, CH_2), 4.45 (b, 1H, NH), 4.8-4.9 (m, 1H, CH), 5.0 (b, 2H, NH_2), 7.35-7.05 (s, 1H, Ar), 7.75 (s, 1H, Ar), 7.9-8.05 (m, 2H, Ar). The specific activity was 175 mCi/mmol. Compound **10a** was treated with 1 eq of isethionic acid to give the salt **10b** immediately before pharmacokinetic and metabolic studies.

ACKNOWLEDGMENT

The authors thank Dr. C. C. Huang and Mr. David Lustgarten for helpful advice, discussions and assistance.

REFERENCES

1. Temple C. Jr. and Rener G.A. - J. Med. Chem. 32: 2089 (1989), and references therein.
2. Buckley T.F. III and Rapoport H. - J. Am. Chem. Soc. 103: 6157 (1981).