

Research Article

Synthesis of (*R*)- and (*S*)-[*O*-methyl-¹¹C]N-[2-[3-(2-cyano-phenoxy)-2-hydroxy-propylamino]-ethyl]-*N'*-(4-methoxy-phenyl)-urea as candidate high affinity β_1 -adrenoceptor PET radioligands

Stefan Wagner^{a,*}, Marilyn P. Law^a, Burkhard Riemann^a, Victor W. Pike^b, Hans-Jörg Breyholz^a, Carsten Höltke^a, Andreas Faust^a, Otmar Schober^a, Michael Schäfers^a and Klaus Kopka^a

^aDepartment of Nuclear Medicine, Albert-Schweitzer-Str. 33, University Hospital Münster, 48149 Münster, Germany

^bMolecular Imaging Branch, National Institute of Mental Health, National Institutes of Health, 10 Center Drive, Rm B3 C346A, MSC 1003, Bethesda, MD 20892-1003, USA

Summary

Molecular imaging and quantification of myocardial β_1 -adrenoceptor (AR) rather than total β -AR density is of great clinical interest since cardiac biopsy studies suggest that myocardial β_1 -AR density is reduced in patients with chronic heart failure whereas cardiac β_2 -AR density may vary. Positron emission tomography (PET), with appropriate radioligands, offers the possibility to assess β -AR density non-invasively in humans. However, no PET radioligand for the selective imaging of cardiac β_1 -ARs is clinically available. Here some derivatives of the well characterized β_1 -AR selective antagonist, ICI 89,406, namely the enantiomers of *N*-[2-[3-(2-cyano-phenoxy)-2-hydroxy-propylamino]-ethyl]-*N'*-(4-hydroxy-phenyl)-urea (**5a** and **5b**) were synthesized and evaluated *in vitro*. The (*R*)-isomer **5a** was more β_1 -selective but has lower affinity than its (*S*)-enantiomer **5b** (β_1 -AR selectivity: 6100 vs 1240; β_1 -affinity: $K_1 = 0.288$ nM vs $K_1 = 0.067$ nM). Etherification of the analogous desmethyl precursors, **5e** and **5f**, respectively, with [¹¹C]iodomethane gave ¹¹C-labelled versions of **5a** and **5b**, namely **5g** and **5h**, in $44 \pm 5\%$ radiochemical yield (decay-corrected) and $97.4 \pm 1.3\%$ radiochemical purity with specific radioactivities of 26.4 ± 9.4 GBq/ μ mol within 41.2 ± 3.4 min from the end of bombardment ($n = 14$). **5g** and **5h** are now

*Correspondence to: S. Wagner, Department of Nuclear Medicine, Albert-Schweitzer-Str. 33, University Hospital Münster, 48149 Münster, Germany. E-mail: stwagner@uni-muenster.de.

Contract/grant sponsor: DFG; contract/grant number: SFB 556

Contract/grant sponsor: DFG; contract/grant number: SCHA 758/5-1

Contract/grant sponsor: SFB; contract/grant number: 656

being evaluated as candidate radioligands for myocardial β_1 -ARs. Copyright © 2005 John Wiley & Sons, Ltd.

Key Words: 3-aryloxy-2-propanolamine; ICI 89,406; β_1 -adrenoceptor selective ligand; PET radioligand


Introduction

β -ARs belong to the rhodopsin/ β_2 -adrenergic receptor-like receptors, that represent a subfamily of the G protein-coupled-receptors (GPCRs).¹ They can be subdivided into at least three discrete subtypes, the β_1 -, β_2 -AR² and the atypical β_3 -AR.^{3,4} β_1 -AR agonists cause increased cardiac contractility and heart rate while β_2 -selective AR agonists mediate vasodepression and bronchodilation.⁵ The so-called atypical β_3 -ARs are involved in lipolysis.³

In heart disease, both the total β -AR density and the β_1/β_2 -AR ratio may change. For example, reduced myocardial β -AR density has been found in hypertension, heart failure, ischaemia and both hypertrophic (HCM) and dilated cardiomyopathies (DCM).^{6–11}

A selective reduction in β_1 -ARs without change in β_2 -AR density is often observed in the failing human heart.⁵ Therefore, a means for the visualization and quantification of the β_1 -AR density rather than total β -AR density in the human heart is of great interest in basic research and clinical application.¹² The molecular imaging techniques, single photon emission computed tomography (SPECT) and positron emission tomography (PET), with suitable radioligands may be ideal tools for assessing β -AR density non-invasively in humans.^{13,14} Some subtype- β_1 -AR selective radioligands have been developed for PET, for example (+/-)-[¹¹C]HX-CH 44,¹⁵ (*S*)-[¹¹C]bisoprolol,¹⁶ [¹¹C]CGP 20712A¹⁷ and its (*S*)-enantiomer, [¹¹C]CGP-26505.¹⁸ The clinical use of these radioligands, however, is limited because of high non-specific binding, rapid metabolism or tissue uptake that does not reflect binding to β -ARs.^{15–18}

ICI 89,406 (Scheme 1) is an example of a designed β_1 -selective AR antagonist. Although membrane studies show that the (*S*)-enantiomer is the more potent enantiomer even the racemate produces effective β_1 -AR blockade during exercise in patients with angina pectoris.^{19–21} Therefore, we chose this compound as a lead structure to develop new β_1 -selective AR radioligands. In our initial study, the β_1 -AR affinity and selectivity of a number of novel derivatives of ICI 89,406 were determined.²² One of these derivatives, racemic I-ICI-H and its radioiodinated analogue (Scheme 1), showed high affinity and selectivity for β_1 -ARs in myocardial membranes *in vitro* but high non-specific binding *in vivo*.²³ Another derivative, racemic I-ICI-COOH and its (*S*)-enantiomer, were also characterized by high β_1 -AR affinity and selectivity *in vitro*.²⁴ The ¹²⁵I/¹²³I-labelled counterparts (Scheme 1) were evaluated as



ICI 89,406	X: CN	Y: H	racemic
I-ICI-H	X: I, ^{125}I or ^{123}I	Y: H	racemic
I-ICI-COOH	X: I, ^{125}I or ^{123}I	Y: COOH	(<i>S</i>) or racemic


Scheme 1. Lead structure ICI 89,406 and its iodinated derivatives^{23,24}

potential SPECT ligands using Sprague Dawley rats but were unsuitable because of a high degree of deiodination.²⁴ It is apparent that high non-specific binding and rapid deiodination *in vivo* preclude the use of such iodinated ligands for SPECT or PET. Ligands labelled with the positron-emitter, carbon-11 ($t_{1/2} = 20.4$ min), may be less susceptible to problematic metabolism. The work presented here describes the synthesis and the *in vitro* pharmacology of the non-iodinated ICI 89,406 derivatives, (*R*)- and (*S*)-*N*-[2-[3-(2-cyano-phenoxy)-2-hydroxy-propylamino]-ethyl]-*N'*-(4-methoxy-phenyl)-urea (**5a** and **5b**, respectively) and their precursors (**5c**–**5f**). The *in vitro* studies showed that the (*R*)-isomer (**5a**) is more selective but has lower affinity for β_1 -ARs than the (*S*)-isomer (**5b**). Therefore, ^{11}C -labelled versions of each enantiomer (**5g** and **5h**) were synthesized as potential selective and high affinity β_1 -AR radioligands for PET (Scheme 2).

Results and discussion

The syntheses of the ureas **5a**–**f** were achieved through a five- or six-step sequence (Scheme 2). Treatment of the homochiral glycidyl-3-nitrobenzene sulphonates **2a** and **2b** with 2-cyano-phenol **1** under basic conditions yielded the corresponding oxiranes **3a** and **3b** in good yields (85–92%). Comparison of the measured optical rotations with literature values showed that the stereochemistry was retained; the absolute $[\alpha]_{\text{D}}$ values were within errors.²⁵ The ureas **5a**–**d** were prepared from the oxiranes **3a** and **3b** and the amines **4a** and **4b** via nucleophilic ring opening. Compounds **4a** and **4b** were synthesized through a three-step sequence as previously described (steps not shown in Scheme 2).²² The chemical yields ranged from 35 to 45%. Finally, **5c** and **5d** were deprotected by hydrogenolysis resulting in the phenol precursors **5e** and **5f** (51–56%).

The structure–activity relationships (SAR) between the ligands and β_1 -ARs were assessed in competition studies using [^{125}I]ICYP and mouse ventricular



Scheme 2. Synthesis of the chiral β_1 -AR ligands **5a–h**

membrane preparations. The high- and low-affinity IC_{50} values for the β_1 - and β_2 -ARs of the non-radioactive 3-aryloxy-2-propanolamine derivatives **5a–f** were calculated by non-linear regression analysis of membrane radioactivity. The high- and low-affinity inhibition constants (K_1 for the β_1 -ARs and K_2 for the β_2 -ARs) were obtained from the IC_{50} values by the method of

Table 1. Inhibition constants and calculated β_1 -AR selectivities of the ligands **5a–f** determined by a radioligand binding assay using mouse ventricular membrane preparations, plus calculated ligand logP and logD values of the ligands

Compound	K_1 (nM) ^a	K_2 (nM) ^a	β_1 -selectivity ^b	logP ^c	logD ^c
5a	0.288 ± 0.032	1760 ± 300	6100 ± 1540	1.22	-0.21
5b	0.067 ± 0.016	83 ± 33	1240 ± 280	1.22	-0.21
5c	20.7 ± 4.2	3900 ± 150	188 ± 45	2.87	1.55
5d	0.085 ± 0.019	119 ± 19	1400 ± 160	2.87	1.55
5e	0.269 ± 0.067	569 ± 277	2120 ± 450	0.83	-0.55
5f	0.328 ± 0.046	149 ± 27	453 ± 106	0.83	-0.55
ICI 89 406	0.28 ± 0.12	41 ± 3	149 ± 86	1.57	0.13

^aInhibition constants for non-selective β -AR ligand [¹²⁵I]ICYP binding at β_1 - and β_2 -ARs expressed in mouse ventricular membrane preparations given as mean ± SEM, $n = 3$.

^bThe ratios of the low- over the high-affinity inhibition constants (K_2/K_1) indicate the β_1 -AR selectivities of the non-radioactive β_1 -AR ligands, noted as mean ± SEM, $n = 3$.

^clogP values of the neutral form and logD values calculated by ACD/LogD Suite (logD = logP at physiological pH 7.4 with consideration of charged species).

Cheng–Prusoff²⁶ using the previously experimentally determined K_D value of [¹²⁵I]ICYP (32.3 ± 1.9 pM).²² The ratios of the low- to high-affinity inhibition constants (K_2/K_1) yield the β_1 -selectivities of the unlabelled compounds **5a–f** (Table 1). The calculated logP and logD values (ACD/LogD Suite) of compounds **5a–f** are also listed in Table 1 and indicate the change in lipophilicities ensuing chemical modifications of the lead compound, ICI 89,406. The logP and logD values of **5a** and **5b** are low and similar to the values for (*S*)-CGP 12177 (logP = 1.81),¹³ which when labelled with carbon-11 is an effective radioligand for imaging cell surface β -ARs in human myocardium with PET. All compounds **5a–f** possess high affinity to β_1 -ARs. With the exception of **5c** ($K_1 = 20.7$ nM) the ligands showed β_1 -AR affinities in the subnanomolar range ($K_1 = 0.067$ – 0.328 nM).

Usually the (*S*)-enantiomer of a β -blocker possesses an up to 100-fold higher affinity to β -ARs than the corresponding (*R*)-enantiomer.^{27,28} This feature was shown by the ligands **5a** and **5b**. The (*R*)-enantiomer **5a** has at least four times less β_1 -AR affinity than the (*S*)-isomer **5b** ($K_1 = 0.288$ nM vs $K_1 = 0.067$ nM). However, the β_1 -AR selectivity of **5a**, is five times greater than that of **5b** (6100 vs 1240). Furthermore, the β_1 -AR selectivities of the methoxy compounds **5a** and **5b** are at least an order of magnitude greater than that of the racemic lead compound, ICI 89,406 (149). A similar trend is seen in the benzyloxy-pair **5c** and **5d**. The (*R*)-enantiomer **5c** has about 240-times less β_1 -affinity than the (*S*)-isomer **5d** ($K_1 = 20.7$ nM vs $K_1 = 0.085$ nM). In contrast, the (*S*)-enantiomer **5d** exhibits higher β_1 -AR selectivity than its (*R*)-enantiomer **5c** (1400 vs 188). Contrary to expectations, both enantiomers of the hydroxy-pair **5e** and **5f** display nearly equal β_1 -AR affinities ($K_1 = 0.269$ and 0.328 nM). Similar to

compounds **5a** and **5b**, the (*R*)-enantiomer **5e** possesses a nearly five-fold higher β_1 -AR selectivity than its (*S*)-enantiomer **5f** (2120 vs 453).

For the compounds **5a–f**, with their specific substitution pattern of a 2-cyanophenoxy core and a 4-hydroxyphenyl or an 4-alkoxyphenyl unit, two SAR-conclusions can be drawn. Firstly, a more sterically demanding residue and/or a more lipophilic aromatic substituent at the oxygen of the phenyl-4-position (**5c** and **5d** R:Bn) leads to an increased difference between the β_1 -AR affinities of the (*R*)- and (*S*)-enantiomers compared to enantiomer-pairs with a smaller substituent at this position (**5a** and **5b** R:Me; **5e** and **5f** R:H). The (*R*)-enantiomer **5c** has 244-fold less β_1 -affinity than the (*S*)-enantiomer **5d**. The tendency is reduced in the methoxy-pair **5a** and **5b** (four-fold) and the hydroxy-pair **5e** and **5f** (0.8-fold). Secondly, an (*S*)-enantiomer with a benzyl-substituent at the phenolic oxygen is more β_1 -selective than its (*R*)-enantiomer, whereas smaller substituents shift the increased β_1 -AR selectivity to the (*R*)-enantiomer ((*R*)-enantiomer to (*S*)-enantiomer ratios of β_1 -AR selectivities: **5c** and **5d** R:Bn: 0.13; **5a** and **5b** R:Me: 4.9; **5e** and **5f** R:H:4.7).

As a result of these binding affinity determinations, the potent methoxy derivatives **5a** and **5b** were chosen for conversion into corresponding potential high affinity and selective β_1 -AR radioligands for PET, namely **5g** and **5h**, respectively. Thus, the desmethyl precursors **5e** and **5f** were treated with sodium hydride and [^{11}C]iodomethane in DMF at 40°C for 5 min to yield ^{11}C -labelled **5g** and **5h**, respectively (Scheme 2), which were then purified with HPLC. The activities of **5g** and **5h** ranged from 1.2 to 2.7 GBq (EOS), representing $44.1 \pm 5.0\%$ decay-corrected radiochemical yields at 41.2 ± 3.4 min from the end of radionuclide production. Radiochemical purities of $97.4 \pm 1.3\%$ and specific activities of 26.4 ± 9.4 GBq/ μmol (13.3–47.4 GBq/ μmol) were achieved ($n=14$), as determined by analytical radio-HPLC. The new radioligands **5g** and **5h** are now being evaluated in rodents with PET.

Experimental

General methods

All chemicals, reagents, and solvents for the synthesis of the compounds were analytical grade and purchased from commercial sources.

The melting points (uncorrected) were determined on a Stuart Scientific SMP3 capillary melting point apparatus. ^1H -NMR and ^{13}C -NMR spectra were recorded on a Bruker ARX 300 spectrometer and an AMX 400 spectrometer, respectively. Mass spectrometry was performed using a Varian MAT 212 (EI=70 eV) spectrometer and a Bruker MALDI-TOF-MS Reflex IV (matrix: DHB). $[\alpha]_{\text{D}}$ values were determined on a Perkin-Elmer 341 polarimeter. Elemental analysis was realised by a Vario EL III analyser.

Radiosyntheses were carried out using an automated PET tracer synthesizer (TRACERLab Fx_c; GE Functional Imaging GmbH). Separation of the radiosynthesized compounds, and determinations of the radiochemical yields were performed with radio-HPLC using a Syknm S1021 pump, a Knauer K-2001 UV-detector (wavelength 254 nm), a Raytest Ramona-90/92 γ -detector, a Nucleosil 100-10 C18 precolumn ($20 \times 8 \text{ mm}^2$) and a Nucleosil 100-10 C18 column ($250 \times 8 \text{ mm}^2$). Sample injection was carried out using a VICI injector block (type C6W incl. 1000 μl loop). The recorded data were processed by the TRACERLab C software (GE Functional Imaging GmbH). The radiochemical purities and the specific activities were acquired with a radio-HPLC system composed of a Syknm S1021 pump, a Knauer K-2501 UV-detector (wavelength 254 nm), a Crismatec Na(Tl) Scintibloc 51 SP51 γ -detector, a Nucleosil 100-3 C18 column ($200 \times 3 \text{ mm}^2$), a VICI injector block (type C1 incl. 20 μl loop) and the NINA version 4.8, Rev. 4 software (GE Functional Imaging GmbH).

Synthetic methods

Synthesis of oxiranes (**3a–b**)

General procedure. 2-Cyano-phenol **1** (2 eq.), (*R*)-glycidyl-3-nitrobenzene sulphonate **2a** or (*S*)-glycidyl-3-nitrobenzene sulphonate **2b** (1 eq.) and anhydrous K_2CO_3 (6 eq.) were refluxed in dry 2-butanone (3.8 ml/mmol sulphonate) for 6–7 h and then stirred at RT for 16–72 h under an argon atmosphere. The mixture was filtered and the filter cake washed with 2-butanone. Solvent from the combined filtrates was evaporated off. The residue was dissolved in water and CH_2Cl_2 . The pH of the aqueous layer was adjusted to be within the range 11–13 and the aqueous layer was extracted. The layers were separated. The aqueous layer was extracted with CH_2Cl_2 twice, the combined organic layers dried (Na_2SO_4) and the solvent evaporated off. The residue was recrystallized from diisopropyl ether– CHCl_3 ($\sim 3:2 \text{ v/v}$) at -30°C to provide **3a–b** as colourless solids.

(*R*)-2-(2-cyano-phenoxyethyl)-oxirane (**3a**). Yield: 85%. Mp.: $85\text{--}86^\circ\text{C}$. $^1\text{H-NMR}$ (300 MHz, CDCl_3): δ [ppm]: 7.58–7.50 (m, 2H, H_{Ar}), 7.09–7.01 (m, 2H, H_{Ar}), 4.37 (dd, $^2J=11.6 \text{ Hz}$, $^3J=3.2 \text{ Hz}$, 1H, 1 CH_2), 4.12 (dd, $^2J=11.6 \text{ Hz}$, $^3J=5.3 \text{ Hz}$, 1H, 1 CH_2), 3.40–3.37 (m, 1H, CH), 2.93 (dd, $^2J=4.8 \text{ Hz}$, $^3J=4.2 \text{ Hz}$, 1H, 1 CH_2), 2.84 (dd, $^2J=5.0 \text{ Hz}$, $^3J=2.4 \text{ Hz}$, 1H, 1 CH_2). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3): δ [ppm]: 160.01, 134.35, 133.87, 121.44, 116.26, 112.79, 102.50, 69.49, 49.85, 44.55. $[\alpha]_{\text{D}}^{20} = -15.8^\circ$ ($c=1.0$, EtOH); Lit.: $[\alpha]_{\text{D}}^{25} = -16.0^\circ$ ($c=1.0$, EtOH).²⁵ Analytically calculated for $\text{C}_{10}\text{H}_9\text{NO}_2$: C 68.56, H 5.12, N 8.00. Found: C 68.93, H 5.07, N 8.16.

(*S*)-2-(2-cyano-phenoxy-methyl)-oxirane (**3b**). Yield: 92%. Mp.: 86°C; Lit.: 88–89°C.²⁵ ¹H-NMR (300 MHz, CDCl₃): δ [ppm]: 7.58–7.49 (m, 2H, H_{Ar}), 7.10–7.01 (m, 2H, H_{Ar}), 4.37 (dd, ²J = 11.4 Hz, ³J = 3.0 Hz, 1H, 1CH₂), 4.12 (dd, ²J = 11.4 Hz, ³J = 5.4 Hz, 1H, 1CH₂), 3.41–3.37 (m, 1H, CH), 2.93 (t, ²J = 4.5 Hz, 1H, 1CH₂), 2.84 (dd, ²J = 5.0 Hz, ³J = 2.6 Hz, 1H, 1CH₂). ¹³C-NMR (75.5 MHz, CDCl₃): δ [ppm]: 159.99, 134.35, 133.84, 121.42, 116.26, 112.80, 102.45, 69.47, 49.85, 44.54. [α]_D²⁰ = +18.1° (c = 1.0, EtOH); Lit.: [α]_D²⁵ = +17.69° (c = 1.0, EtOH).²⁵ Analytically calculated for C₁₀H₉NO₂: C 68.56, H 5.12, N 8.00. Found: C 68.80, H 4.96, N 8.38.

Synthesis of the N-aryl-N'-[2-[3-aryloxy-2-hydroxy-propylamino]-ethyl]-urea derivatives (5a–d)

General procedure. **3a** or **3b** (1 eq.), *N*-(2-amino-ethyl)-*N'*-(4-methoxy-phenyl)-urea hydrochloride **4a** (1 eq.) or *N*-(2-amino-ethyl)-*N'*-(4-benzyloxy-phenyl)-urea hydrochloride **4b** (1 eq.), prepared as previously described,²² and 1 N NaOH (1.05 eq.) were heated in *n*-propanol (1.8–2.9 ml/mmol oxirane) and water (1.0–2.4 ml/mmol oxirane) up to 90°C for 2–3 h.

Purification procedure A (5a and 5c): The mixture was evaporated to dryness. Water was added and stirred. The product was filtered off and dried *in vacuo*. The crude product was refluxed in methanol, the hot suspension was filtered and the filtrate was left for crystallization at +4 to –30°C. Additional amounts of product were obtained by crystallization from the mother liquor.

Purification procedure B (5b and 5d): Water was added. The product was filtered off and washed with water–diethyl ether. The crude product was purified by silica gel chromatography (ethyl acetate–MeOH 4:1 v/v). The product fraction was evaporated and recrystallized from methanol.

After the purification procedures A or B, the urea derivatives **5a–d** were obtained as colourless solids.

(*R*)-*N*-[2-[3-(2-cyano-phenoxy)-2-hydroxy-propylamino]-ethyl]-*N'*-(4-methoxy-phenyl)-urea (**5a**). Yield: 35%. Mp.: 126–127°C. ¹H-NMR (400 MHz, DMSO-*d*₆): δ [ppm]: 8.29 (s, 1H, NH), 7.69 (dd, ³J = 7.7 Hz, ⁴J = 1.7 Hz, 1H, H_{Ar}), 7.61 (ddd, ³J = 8.7 Hz, ³J = 7.4 Hz, ⁴J = 1.4 Hz, 1H, H_{Ar}), 7.27–7.23 (m, 3H, H_{Ar}), 7.06 (dt, ³J = 7.5 Hz, ⁴J = 0.9 Hz, 1H, H_{Ar}), 6.78 (dd, ³J = 6.8 Hz, ⁴J = 2.3 Hz, 2H, H_{Ar}), 6.02 (t, ³J = 5.1 Hz, 1H, NH), 5.01 (s, br, 1H, OH), 4.16–3.92 (m, 3H, CH₂CH), 3.67 (s, 3H, CH₃), 3.14 (q, ³J = 5.9 Hz, 2H, CH₂), 2.75–2.60 (m, 4H, 2CH₂). ¹³C-NMR (100.6 MHz, DMSO-*d*₆): δ [ppm]: 160.52, 155.71, 154.03, 135.14, 133.89, 133.76, 121.13, 119.53, 116.58, 114.02, 113.35, 100.83, 71.79, 68.12, 55.30, 52.13, 49.53, 39.89. Analytically calculated for C₂₀H₂₄N₄O₄: C 62.49, H 6.29, N 14.57. Found: C 62.25, H 6.24, N 14.62.

(*S*)-*N*-[2-[3-(2-cyano-phenoxy)-2-hydroxy-propylamino]-ethyl]-*N'*-(4-methoxy-phenyl)-urea (**5b**). Yield: 45%. Mp.: 128–129°C. $^1\text{H-NMR}$ (400 MHz, DMSO- d_6): δ [ppm]: 8.30 (s, 1H, NH), 7.69 (d, $^3J=7.8$ Hz, 1H, H_{Ar}), 7.62 (t, $^3J=7.4$ Hz, 1H, H_{Ar}), 7.27–7.23 (m, 3H, H_{Ar}), 7.06 (t, $^3J=7.5$ Hz, 1H, H_{Ar}), 6.78 (d, $^3J=8.7$ Hz, 2H, H_{Ar}), 6.03 (t, $^3J=5.3$ Hz, 1H, NH), 5.02 (s, br, 1H, OH), 4.16–3.90 (m, 3H, CH_2CH), 3.68 (s, 3H, CH_3), 3.15 (q, $^3J=5.7$ Hz, 2H, CH_2), 2.76–2.61 (m, 4H, 2CH_2). $^{13}\text{C-NMR}$ (100.6 MHz, DMSO- d_6): δ [ppm]: 160.50, 155.68, 154.01, 135.13, 133.89, 133.77, 121.13, 119.48, 116.54, 114.01, 113.35, 100.82, 71.76, 68.05, 55.29, 52.09, 49.49, 39.91. Analytically calculated for $\text{C}_{20}\text{H}_{24}\text{N}_4\text{O}_4$: C 62.49, H 6.29, N 14.57. Found: C 62.11, H 6.12, N 14.17.

(*R*)-*N*-(4-benzyloxy-phenyl)-*N'*-[2-[3-(2-cyano-phenoxy)-2-hydroxy-propylamino]-ethyl]-urea (**5c**). Yield: 41%. Mp.: 117–118°C. $^1\text{H-NMR}$ (300 MHz, DMSO- d_6): δ [ppm]: 8.31 (s, 1H, NH), 7.68 (d, $^3J=7.5$ Hz, 1H, H_{Ar}), 7.63–7.58 (m, 1H, H_{Ar}), 7.43–7.23 (m, 8H, H_{Ar}), 7.06 (t, $^3J=7.4$ Hz, 1H, H_{Ar}), 6.87 (d, $^3J=9.0$ Hz, 2H, H_{Ar}), 6.04 (t, $^3J=5.6$ Hz, 1H, NH), 5.02 (s, 3H, OH and CH_2), 4.16–3.90 (m, 3H, CH_2CH), 3.15 (q, $^3J=5.1$ Hz, 2H, CH_2), 2.76–2.58 (m, 4H, 2CH_2), 1.85 (s, br, 1H, NH). $^{13}\text{C-NMR}$ (75.5 MHz, DMSO- d_6): δ [ppm]: 160.48, 155.63, 153.03, 137.52, 135.08, 134.13, 133.74, 128.46, 127.78, 127.69, 121.08, 119.39, 116.54, 115.06, 113.32, 100.81, 71.74, 69.59, 68.08, 52.12, 49.49. Analytically calculated for $\text{C}_{26}\text{H}_{28}\text{N}_4\text{O}_4$: C 67.81, H 6.13, N 12.17. Found: C 67.56, H 6.04, N 12.11.

(*S*)-*N*-(4-benzyloxy-phenyl)-*N'*-[2-[3-(2-cyano-phenoxy)-2-hydroxy-propylamino]-ethyl]-urea (**5d**). Yield: 36%. Mp.: 113–114°C. $^1\text{H-NMR}$ (300 MHz, DMSO- d_6): δ [ppm]: 8.31 (s, 1H, NH), 7.68 (d, $^3J=7.5$ Hz, 1H, H_{Ar}), 7.63–7.58 (m, 1H, H_{Ar}), 7.43–7.23 (m, 8H, H_{Ar}), 7.06 (t, $^3J=7.5$ Hz, 1H, H_{Ar}), 6.87 (d, $^3J=9.0$ Hz, 2H, H_{Ar}), 6.04 (t, $^3J=5.1$ Hz, 1H, NH), 5.02 (s, 3H, OH and CH_2), 4.16–3.90 (m, 3H, CH_2CH), 3.15 (q, $^3J=5.4$ Hz, 2H, CH_2), 2.76–2.61 (m, 4H, 2CH_2), 1.96 (s, br, 1H, NH). $^{13}\text{C-NMR}$ (75.5 MHz, DMSO- d_6): δ [ppm]: 160.50, 155.64, 153.02, 137.53, 135.10, 134.15, 133.75, 128.45, 127.78, 127.70, 121.09, 119.40, 116.55, 115.08, 113.32, 100.81, 71.74, 69.59, 68.09, 52.12, 49.49, 39.84. Analytically calculated for $\text{C}_{26}\text{H}_{28}\text{N}_4\text{O}_4$: C 67.81, H 6.13, N 12.17. Found: C 62.48, H 5.97, N 12.16.

Synthesis of the N-aryl-N'-[2-[3-aryloxy-2-hydroxy-propylamino]-ethyl]-urea derivatives (5e–f)

General procedure. **5c** or **5d** (1 eq.), methanol (46 ml/mmol) and Pd/C (10%, 47 mg/mmol) were stirred in a H_2 -atmosphere at RT for 15–16 h. Then the mixture was refluxed for 6–8.5 h. The hot reaction mixture was filtered and the

filtrate was evaporated to dryness. The residue was purified by silica gel chromatography (ethyl acetate–MeOH 4:1 v/v) to provide the urea derivative **5e** or **5f** (R_f : 0.15) as a colourless solid.

(*R*)-*N*-[2-[3-(2-cyano-phenoxy)-2-hydroxy-propylamino]-ethyl]-*N'*-(4-hydroxy-phenyl)-urea (**5e**). Yield: 51%. Mp.: 120–122°C. $^1\text{H-NMR}$ (300 MHz, DMSO- d_6): δ [ppm]: 8.87 (s, br, 1H, OH), 8.14 (s, 1H, NH), 7.69 (dd, $^3J=7.7$ Hz, $^4J=1.4$ Hz, 1H, H_{Ar}), 7.61 (ddd, $^3J=8.4$ Hz, $^3J=7.7$ Hz, $^4J=1.1$ Hz, 1H, H_{Ar}), 7.24 (d, $^3J=8.7$ Hz, 1H, H_{Ar}), 7.14–7.03 (m, 3H, H_{Ar}), 6.78 (dd, $^3J=6.8$ Hz, $^4J=2.1$ Hz, 2H, H_{Ar}), 5.98 (t, $^3J=5.4$ Hz, 1H, NH), 4.98 (s, br, 1H, OH), 4.16–3.88 (m, 3H, CH_2CH), 3.13 (q, $^3J=5.8$ Hz, 2H, CH_2), 2.75–2.59 (m, 4H, 2CH_2). $^{13}\text{C-NMR}$ (75.5 MHz, DMSO- d_6): δ [ppm]: 160.50, 155.77, 152.01, 135.11, 133.76, 132.27, 121.10, 119.93, 116.54, 115.19, 113.34, 100.81, 71.76, 68.07, 52.12, 49.55. Analytically calculated for $\text{C}_{19}\text{H}_{22}\text{N}_4\text{O}_4$: C 61.61, H 5.99, N 15.13. Found: C 61.24, H 5.98, N 14.92.

(*S*)-*N*-[2-[3-(2-cyano-phenoxy)-2-hydroxy-propylamino]-ethyl]-*N'*-(4-hydroxy-phenyl)-urea (**5f**). Yield: 56%. Mp.: 118–119°C. $^1\text{H-NMR}$ (300 MHz, DMSO- d_6): δ [ppm]: 8.87 (s, br, 1H, OH), 8.14 (s, 1H, NH), 7.69 (dd, $^3J=7.7$ Hz, $^4J=1.4$ Hz, 1H, H_{Ar}), 7.61 (ddd, $^3J=8.7$ Hz, $^3J=7.4$ Hz, $^4J=1.5$ Hz, 1H, H_{Ar}), 7.24 (d, $^3J=8.7$ Hz, 1H, H_{Ar}), 7.14–7.03 (m, 3H, H_{Ar}), 6.78 (dd, $^3J=6.8$ Hz, $^4J=2.0$ Hz, 2H, H_{Ar}), 5.97 (t, $^3J=5.4$ Hz, 1H, NH), 5.01 (s, br, 1H, OH), 4.14–3.89 (m, 3H, CH_2CH), 3.13 (q, $^3J=6.0$ Hz, 2H, CH_2), 2.75–2.60 (m, 4H, 2CH_2). $^{13}\text{C-NMR}$ (75.5 MHz, DMSO- d_6): δ [ppm]: 160.49, 155.81, 152.02, 135.12, 133.78, 132.27, 121.13, 119.96, 116.57, 115.18, 113.33, 100.82, 71.77, 68.08, 52.11, 49.56. Analytically calculated for $\text{C}_{19}\text{H}_{22}\text{N}_4\text{O}_4$: C 61.61, H 5.99, N 15.13. Found: C 61.14, H 5.94, N 14.88.

*Radiosynthesis of (R)- and (S)-[O-methyl- ^{11}C]N-[2-[3-(2-cyano-phenoxy)-2-hydroxy-propylamino]-ethyl]-*N'*-(4-methoxy-phenyl)-urea (**5g** and **5h**).* [^{11}C]Carbon dioxide was produced by the $^{14}\text{N}(p,\alpha)^{11}\text{C}$ nuclear reaction on research grade nitrogen containing 2.5% oxygen, using 11 MeV proton beams at currents of 40 μA from a CTI-RDS-111 cyclotron. The [^{11}C]carbon dioxide was trapped from the target gas in a stainless steel loop cooled with liquid nitrogen to -150°C . [^{11}C]Iodomethane was prepared from [^{11}C]carbon dioxide as previously described.²⁹ Phenol precursor **5e** or **5f** (1.0 mg; 2.70 μmol) was treated with NaH (60% oil suspension; 1.5 mg; 37.5 μmol) in DMF (200 μl) and [^{11}C]CH $_3$ I at 40°C for 5 min. After heating the reaction mixture to 50°C , water for injection (200 μl) was added and the crude mixture was loaded onto a semi-preparative HPLC-column (see Synthetic methods). Product **5g** or **5h** was eluted with 20 mM Na_2HPO_4 -buffer/EtOH 4/1 at a flow of 4 ml/min at

11.9–14.5 min. The product solution was filtered through a sterile filter REF F60 (0.2 μm , pvb Medizintechnik GmbH). The time of synthesis and purification was 41.2 ± 3.4 min from the EOB. The radiochemical yield averaged $44.1 \pm 5.0\%$ (decay-corrected) and the radiochemical purity, determined via radio-HPLC (see Synthetic methods, eluent: $\text{H}_2\text{O}/\text{MeCN}/\text{TFA}$ 700/300/1, flow: 0.3 ml/min, retention time: 11.1 min), was $97.4 \pm 1.3\%$ with a specific activity of 26.4 ± 9.4 GBq/ μmol (13.3–47.4 GBq/ μmol) at EOS ($n = 14$).

Ligand binding assay

Microsomes were prepared by homogenizing ventricles from DBA mice at 4°C for 90 s in buffer A (1 ml) containing 10 mM EDTA, 10 mM HEPES and 0.1 mM benzamidine (pH 7.4), using a Polytron PT 3000 (Kinematica, Lucerne, Switzerland). Homogenates were centrifuged at $45\,000 \times g_{\text{max}}$ for 15 min at 4°C . The pellets were resuspended again in buffer B (1 ml) containing 1 mM EDTA, 10 mM HEPES and 0.1 mM benzamidine (pH 7.4) and recentrifuged at $45\,000 \times g_{\text{max}}$ for 15 min at 4°C . The pellets were resuspended in buffer B (1 ml) and centrifuged at $10\,000 \times g_{\text{max}}$ for 10 min at 4°C . The supernatants were recentrifuged at $45\,000 \times g_{\text{max}}$ for 15 min at 4°C . The pellets, partially enriched membranes, were resuspended in buffer C (50 mM *Tris*·HCl, 5 mM MgCl_2 (pH 7.4)), and stored frozen at -80°C . For competition binding studies, the prepared membranes were resuspended in buffer D (10 mM *Tris*HCl, 154 mM NaCl, 0.1 mM ascorbic acid, pH 7.4). 15 μg of membranes were incubated with a constant concentration of [^{125}I]ICYP (80 pM) and with varying concentrations (1 pM–100 μM) of compounds **5a–f** (Scheme 2). Reactions were conducted at 37°C for 60 min. Reactions were stopped by filtering onto Whatman GF/B filters and washed with water for injection. The membrane bound radioactivity was determined in a γ -scintillation counter. Competition binding curves were analysed by non-linear regression analysis as previously described^{30–32} using the XMGRACE program (Linux software). The data of the ligands **5a–f** fitted a two-site model significantly better than a one-site model ($F = 2.15$, $p < 0.05$). The resulting high- and low-affinity IC_{50} values of all synthesized and non-radioactive 3-aryloxy-2-propranolamine derivatives **5a–f** were converted into the high- and low-affinity inhibition constants (K_1 for the β_1 -ARs and K_2 for the β_2 -ARs) by the method of Cheng–Prusoff²⁶ using the experimentally determined K_D value of [^{125}I]ICYP (32.3 ± 1.9 pM).²² The ratios of the low- to high-affinity inhibition constants (K_2/K_1) yield the β_1 -selectivities of the unlabelled derivatives **5a–f** (Table 1). These compounds show a significantly higher affinity to β_1 - than to β_2 -ARs. Additionally, the calculated logP and logD values (ACD/LogD Suite) for compounds **5a–f** are listed in Table 1 to indicate

the changes of the lipophilicities caused by the chemical modifications of the lead compound, ICI 89,406.

Conclusion

The syntheses, radiosyntheses and *in vitro* pharmacology of (*R*)- and (*S*)-*N*-[2-[3-(2-cyano-phenoxy)-2-hydroxy-propylamino]-ethyl]-*N'*-(4-methoxy-phenyl)-urea (**5a** and **5b**, respectively), and their ¹¹C-labelled versions (**5g** and **5h**, respectively), potential new high affinity β_1 -AR selective PET radioligands, are reported. The non-radioactive counterparts **5a** and **5b** of the radiolabelled target compounds **5g** and **5h** showed high affinity to β_1 -ARs in murine myocardial membranes and moderate lipophilicity. The radiosyntheses of **5g** and **5h** were achieved with good radiochemical and acceptable specific activities. The formulation of the preparations is suitable for preliminary *in vivo* studies using small animal PET.

Acknowledgements

This work was supported by the SFB 556, 656 and SCHA 758/5-1 of the Deutsche Forschungsgemeinschaft (DFG), Bonn, Germany. The authors gratefully thank Ms Monika Trub and Ms Sandra Schröer for technical assistance.

References

1. Gether U. *Endocr Rev* 2000; **21**: 90–113.
2. Lands AM, Arnold A, McAuliff JP, Luduena FP, Brown TG. *Nature* 1967; **214**: 597–598.
3. Arch JR, Ainsworth AT, Cawthorne MA, Piercy V, Sennitt MV, Thody VE, Wislon C, Wilson S. *Nature* 1984; **309**: 163–165.
4. Bond RA, Clarke DE. *Br J Pharmacol* 1988; **95**: 723–734.
5. Brodde OE. *Pharmacol Rev* 1991; **43**: 203–242.
6. Castellano M, Böhm M. *Hypertension* 1997; **29**: 715–722.
7. Khamssi M, Brodde OE. *J Cardiovasc Pharmacol* 1990; **16** (Suppl 5): S133–S137.
8. Brodde OE, Zerkowski HR, Doetsch N, Motomura S, Khamssi M, Michel MC. *J Am Coll Cardiol* 1989; **14**: 323–331.
9. Anthonio RL, Brodde OE, van Veldhuisen DJ, Scholtens E, Crijns HJ, van Gilst WH. *Int J Cardiol* 2000; **72**: 137–141.
10. Schäfers M, Dutka D, Rhodes CG, Lammertsma AA, Hermansen F, Schober O, Camici PG. *Circ Res* 1998; **82**: 57–62.
11. Yamada S, Ohkura T, Uchida S, Inabe K, Iwatani Y, Kimura R, Hoshino T, Kaburagi T. *Life Sci* 1996; **58**: 1737–1744.
12. Riemann B, Schäfers M, Law MP, Wichter T, Schober O. *Nuklearmedizin* 2003; **42**: 4–9.
13. Pike VW, Law MP, Osman S, Davenport RJ, Rimordi O, Giardinà D, Camici PG. *Pharm Acta Helv* 2000; **74**: 191–200.

14. van Waarde A, Visser TJ, Elsinga PJ, de Jong B, van der Mark TW, Kraan J, Elsinga PH, Pruijm J, Willemsen AT, Brodde OE, Visser GM, Paans AM, Vaalburg W. *J Nucl Med* 1997; **38**: 934–939.
15. Valette H, Dollé F, Guenther I, Demphel S, Rasetti C, Hinnen F, Fuseau C, Crouzel C. *Nucl Med Biol* 1999; **26**: 105–109.
16. Soloviev DV, Matarrese M, Moresco RM, Todde S, Buonasera TA, Sudati F, Simonelli P, Magni F, Colombo D, Carpinelli A, Kienle MG, Fazio F. *Neurochem Int* 2001; **38**: 169–180.
17. Elsinga PH, van Waarde A, Visser GM, Vaalburg W. *Nucl Med Biol* 1994; **21**: 211–217.
18. van Waarde A, Meeder JG, Blanksma PK, Bouwer J, Visser GM, Elsinga PH, Paans AM, Vaalburg W, Lie KI. *Int J Appl Instrum B* 1992; **19**: 711–718.
19. Imperial Chemical Industries Limited, London (UK), Patent CH 605666 1978; DE 2458908 1975; *Chem Abstr* 1976; **84**: 43599.
20. Majid PA, Schreuder JE, de Feyter PJ, Roos JP. *J Cardiovasc Pharmacol* 1980; **2**: 435–444.
21. Svendsen TL, Hartling O, Trap-Jensen J. *Eur J Clin Pharmacol* 1979; **15**: 223–228.
22. Kopka K, Wagner S, Riemann B, Law MP, Puke C, Luthra SK, Pike VW, Wichter T, Schmitz W, Schober O, Schäfers M. *Bioorg Med Chem* 2003; **11**: 3513–3527.
23. Riemann B, Law MP, Kopka K, Wagner S, Luthra SK, Pike VW, Neumann J, Kirchhefer U, Schmitz W, Schober O, Schäfers M. *Nuklearmedizin* 2003; **42**: 173–180.
24. Wagner S, Kopka K, Law MP, Riemann B, Pike VW, Schober O, Schäfers M. *Bioorg Med Chem* 2004; **12**: 4117–4132.
25. Nicola M, Depaoli A, Inglesi M. *Gazz Chim Ital* 1990; **120**: 393–396.
26. Cheng Y, Prusoff WH. *Biochem Pharmacol* 1973; **22**: 3099–3108.
27. Mutschler E. β -Adrenozeptor-antagonisten. In *Arzneimittelwirkungen*. Wissenschaftliche Verlagsgesellschaft mbH: Stuttgart, 1996; 291–292.
28. Mehvar R, Brocks DR. *J Pharm Pharmaceut Sci* 2001; **4**: 185–200.
29. Holschbach M, Schüller M. *Appl Radiat Isot* 1993; **44**: 779–780.
30. Nanoff C, Freissmuth M, Schütz W. *Naunyn Schmiedeberg's Arch Pharmacol* 1987; **336**: 519–525.
31. Engel G, Hoyer D, Berthold R, Wagner H. *Naunyn Schmiedeberg's Arch Pharmacol* 1981; **317**: 277–285.
32. DeLean A, Hancock AA, Lefkowitz RJ. *Mol Pharmacol* 1982; **21**: 5–16.