

Research Article

The synthesis of deuterium-labeled spermine, N¹-acetylspermine and N¹-acetylspermidine

VIJAY GAWANDI¹ and PAUL F. FITZPATRICK^{1,2,*}

¹Department of Biochemistry and Biophysics, 2128 Texas A&M University, College Station, TX 77843-2128, USA

²Department of Chemistry, 2128 Texas A&M University, College Station, TX 77843-2128, USA

Received 1 March 2007; Revised 30 March 2007; Accepted 2 April 2007

Abstract: The synthesis of deuterium-labeled spermine, N¹-acetylspermine and N¹-acetylspermidine is reported. 1,1,3,3,3-²H₄-N¹-Acetylspermine hydrochloride, 1,1,3,3,3-²H₄-N¹-acetylspermidine hydrochloride and 1,1,3,3,10,10,12,12-²H₈-spermine dihydrochloride were obtained in seven, four and three steps, respectively. All the syntheses were carried out by simple protection and deprotection steps from commonly used selective protecting reagents. These deuterium-labeled compounds can be used as mechanistic probes of polyamine oxidizing enzymes. Copyright © 2007 John Wiley & Sons, Ltd.

Keywords: deuterium labeling; spermine; N¹-acetylspermine; N¹-acetylspermidine; polyamine oxidase

Introduction

Polyamines such as putrescine, spermidine and spermine are naturally occurring, positively charged compounds found in virtually all living cells.^{1,2} These compounds bind DNA and have been implicated in a number of crucial cellular processes such as cell division, differentiation and membrane function.^{1–4} Inhibition of polyamine synthesis, or their depletion from cells, stops cell growth, suggesting that analogs of these compounds have potential for cancer treatment.^{4–6} In support of this, several polyamine analogues exhibit antitumor effects.^{7–10}

Spermine, N¹-acetylspermine and N¹-acetylspermidine are substrates for polyamine oxidase (PAO), whereas spermine is the substrate for spermine oxidase (SMO).^{11–14} PAO in the peroxisome oxidizes N¹-acetylspermine to spermidine and 3-acetamidopropanal and N¹-acetylspermidine to putrescine and 3-acetamidopropanal (Figure 1).¹¹ In contrast, the cytosolic enzyme SMO oxidizes spermine to spermidine and 3-aminopropanol but does not oxidize N¹-acetylspermine or N¹-acetylspermidine.^{12–14} Flavin adenine dinucleotide (FAD) is a noncovalently bound redox

cofactor in both PAO and SMO. During the oxidation of N¹-acetylspermine, N¹-acetylspermidine and spermine, FAD is reduced; it is subsequently oxidized by oxygen to generate hydrogen peroxide.¹⁵ Hydrogen peroxide and 3-aminopropanal (obtained as a side product of the enzymatic deacetylation¹⁶ of 3-acetamidopropanal) are cytotoxic.^{17–19} 3-Aminopropanal undergoes spontaneous conversion to acrolein which is extremely cytotoxic.¹⁶ It has been proposed that the hydrogen peroxide produced by SMO and PAO may play a role in the progression of apoptosis.^{20–24}

Relatively little mechanistic study of animal or human PAO or SMO has been reported. Syntheses of protiated analogues of spermine, N¹-acetylspermine and N¹-acetylspermidine have been reported by several researchers,^{25–27} but the syntheses of deuterated analogues have not. Syntheses of deuterated analogues have now been carried out using simple protection and deprotection steps. These deuterium-labeled compounds can be used as mechanistic probes of polyamine oxidizing enzymes.

Results and discussion

Although syntheses of a few polyamines, e.g. N¹-acetylspermine and N¹-acetylspermidine, have been reported,^{25–27} preparation of ²H₈-spermine, ²H₄-N¹-acetylspermine and ²H₄-N¹-acetylspermidine has not been described previously. Scheme 1 shows the

*Correspondence to: Paul F. Fitzpatrick, Department of Biochemistry and Biophysics, 2128 TAMU, College Station, TX 77843-2128, USA. E-mail: fitzpat@tamu.edu
Contract/grant sponsor: NIH; contract/grant number: R01 GM58698





Figure 1 The polyamine oxidase reaction.




Scheme 1

synthesis of 1,1,3,3-²H₄-N¹-acetylspermine. Commercially available 4-aminopropanol was treated with Boc anhydride to protect the amino group of 4-aminopropanol to yield N-Boc-3-aminopropanol **1**. This compound was then activated by treatment with methanesulfonyl chloride, a hydroxyl group-activating reagent, to give N-Boc-3-amino-1-methanesulfonylpropane **2**, which underwent nucleophilic substitution by the amino group of 4-aminobutanol to yield N-(N-Boc-3'-aminopropyl)-4-aminobutanol **3**. The secondary amino group of compound **3** was then protected to yield N-(N-Boc-3'-aminopropyl)-N-Boc-4-aminobutanol **4**. Activation of **4** by methanesulfonyl chloride yielded N-(N-Boc-3'-aminopropyl)-N-Boc-4-amino-1-methanesulfonylbutane **5**, which was coupled with the monoacetyl analogue of 1,1,3,3-²H₄-diaminopropane **6** to yield N¹-acetyl-1,1,3,3-²H₄-N⁹, N¹²-Boc₂-spermine **7**. The method described by Tabor *et al.*²⁷ was used


to prepare **6**. Crude 1,1,3,3-²H₄-N¹-acetylspermine hydrochloride was obtained by deprotection of **7** with 4 M hydrochloric acid in dioxane at 25°C. The final compound was then purified by reverse-phase column chromatography using 20% ethanol-water as the mobile phase.

Scheme 2 presents the synthesis of 1,1,3,3-²H₄-N¹-acetylspermidine. The procedure used for the synthesis of 1,1,3,3-²H₄-N¹-acetylspermidine is similar to that for 1,1,3,3-²H₄-N¹-acetylspermine.

Scheme 3 shows the synthetic route used to prepare 1,1,3,3,10,10,12,12-²H₈-spermine. 1,4-Dihydroxybutane was activated by treatment with methanesulfonyl chloride in dry pyridine to yield 1,4-dimethanesulfonylbutane **11**. Treatment of **11** with N¹-Boc-1,1,3,3-²H₄-1,3-diaminopropane **12** gave N¹, N¹²-Boc₂-²H₈-spermine **13**, which was deprotected with hydrochloric acid in dioxane to generate 1,1,3,3,10,10,12,12-²H₈-spermine.



Scheme 2



Scheme 3

Experimental

Materials

3-Aminopropanol, 4-aminobutanol and 1,4-dihydroxybutane were obtained from Aldrich Chemicals. 1,1,3,3- $^2\text{H}_4$ -1,3-Diaminopropane was obtained from C/D/N Isotopes, Inc. All other reagents were of the highest purity commercially available. ^1H NMR spectra were acquired on an Inova 300 MHz spectrometer.

Synthesis of 1,1,3,3- $^2\text{H}_4$ -N 1 -acetylspermine trihydrochloride

N-Boc-3-aminopropanol (1). To 3-aminopropanol (0.98 g, 13 mmol) in 5 mL of dry acetonitrile was added dropwise 1.1 equivalent (3.12 g, 14.3 mmol) of $(\text{Boc})_2\text{O}$ in 2.0 equivalent (1.4 mL, 26 mmol) of triethylamine. After 6 h at 25°C an additional 0.5 equivalent (1.42 g) of $(\text{Boc})_2\text{O}$ was added. The reaction was continued overnight until silica gel thin-layer chromatography in hexane–ethylacetate (8:2) showed one major spot. The solvent was evaporated *in vacuo*. The brown residue was treated with 1 M KHSO_4 (60 mL) and ether

(100 mL). The solution was extracted with ether (4×25 mL), and the combined organic layers were washed with 50 mL each of 1 M KHSO_4 and 1 M NaHCO_3 and twice with a saturated solution of NaCl . The yellowish extract was dried (MgSO_4) and the solvent removed under vacuum. The brown residue was chromatographed on silica using hexane–ethylacetate (8:2) to give 1.63 g (66%) N-Boc-3-aminopropanol **1**. ^1H NMR (CDCl_3): δ 7.38 (1H, s), 3.72 (2H, m), 1.71 (2H, m), 2.55 (2H, t), 1.32 (9H, s).

N-Boc-3-amino-1-methanesulfonyloxypropane (2). The Boc-protected aminopropanol (0.82 g, 4.7 mmol) was added to 1.2 equivalents (0.65 g, 5.64 mmol) of methanesulfonyl chloride in 2 mL anhydrous pyridine. After 2 h at 25°C , the reaction mixture was treated with 1 M NaHCO_3 (30 mL) and ether (70 mL). The solution was extracted with ether (3×25 mL), and the combined organic layers were washed with a NaCl solution (2×50 mL). A brownish crude product was obtained after evaporation of solvent. Flash silica gel column chromatography using hexane–ethyl acetate (8:2) as solvent yielded 0.79 g (62%) N-Boc-3-amino-1-methanesulfonyloxypropane **2**. ^1H NMR (CDCl_3): δ (ppm) 2.35 (s, 3H), 4.21 (t, 2H), 1.72 (m, 2H), 2.81 (t, 2H), 1.35 (s, 9H).

N-(N-Boc-3'-aminopropyl)-4-aminobutanol (3). Compound **2** (0.79 g, 3.1 mmol) was added to 1.0 equivalent (0.28 g) of 4-aminobutanol in 5 mL dry dichloromethane. After 5 h at 50°C, the reaction mixture was cooled, washed with saturated Na₂CO₃ and extracted with ether. Flash silica gel chromatography using hexane–ethyl acetate (9:1) as solvent gave 0.51 g (66%) N-(N-Boc-3'-aminopropyl)-4-aminobutanol **3**. ¹H NMR (CDCl₃): δ 7.40 (s, 1H), 2.55 (t, 2H), 1.75 (m, 2H), 2.60 (m, 2H), 2.55 (m, 2H), 1.81 (m, 2H), 1.72 (m, 2H), 3.75 (m, 2H), 1.35 (s, 9H).

N-(N-Boc-3'-aminopropyl)-N-Boc-4-aminobutanol (4). Compound **3** (0.51 g, 2.07 mmol) was added to 1.0 equivalent (0.45 g) of (Boc)₂O in 2.0 equivalent (0.6 mL, 4.14 mmol) of triethylamine in 10 mL of dry acetonitrile at 25°C; the reaction was continued overnight. Pure N-(N-Boc-3'-aminopropyl)-N-Boc-4-aminobutanol **4** (0.325 g, 86%) was obtained following the procedure described for **1**. ¹H NMR (CDCl₃): δ 2.60–2.66 (m, 6H), 1.78–1.82 (m, 6H), 3.72 (m, 2H), 1.40 (s, 18H).

N-(N-Boc-3'-aminopropyl)-N-Boc-4-amino-1-methanesulfonybutane (5). The reaction of **4** (0.33 g, 0.94 mmol) with 1.2 equivalent (0.13 g) methane sulfonyl chloride as described above for the preparation of **2** yielded 0.33 g (85%) N-(N-Boc-3'-aminopropyl)-N-Boc-4-amino-1-methanesulfony butane **5**. ¹H NMR (CDCl₃): δ 2.40 (s, 3H), 2.82–2.86 (m, 6H), 1.68–1.75 (m, 6H), 4.23 (m, 2H), 1.39 (s, 18H).

N-Acetyl-1,1,3,3-²H₄-1,3-diaminopropane (6). The synthesis of **6** was adapted from the procedure of Tabor *et al.*²⁷ 1,1,3,3-²H₄-1,3-Diaminopropane (0.6 g, 7.69 mmol) was added to 45 mL of cooled glacial acetic acid with magnetic stirring. The mixture was heated to 55–60°C. A mixture of 0.36 mL of acetic anhydride (0.5 equivalent) in 10 mL of glacial acetic acid was added dropwise with stirring during a 1 h period. The mixture was stored at 25°C overnight and then evaporated to dryness. The residue was dissolved in hot water, cooled and adjusted to acidic pH with 6N HCl. After evaporation to dryness *in vacuo*, the resultant solid was extracted twice with 50 mL 2-propanol, and the insoluble residue (unreacted 1,3-diaminopropane hydrochloride) was discarded. The combined filtrates were concentrated and cooled to –10°C. The crystals that formed were collected by filtration and recrystallized from 100 mL of hot 2-propanol. The yield of **6** was 0.48 g (73%). ¹H NMR (CDCl₃): δ 2.05 (s, 3H), 1.65 (s, 2H).

N¹-Acetyl-1,1,3,3-(²H₄)-N⁹,N¹²-Boc₂-spermine (7). Compound **5** (0.33 g, 0.84 mmol) was added to 1.3 equivalent (0.13 g, 1.09 mmol) of **6** in 5.0 mL of dry dichloromethane. After 5 h at 50°C, workup similar to that

described for **3** yielded 0.25 g (62%) of compound **7**. ¹H NMR (CDCl₃): δ 2.11 (s, 3H), 2.85–2.88 (m, 8H), 1.80–1.82 (m, 4H), 1.74–1.76 (m, 4H), 1.40 (s, 18H).

1,1,3,3-²H₄-N¹-acetylspermine trihydrochloride. Compound **7** (0.22 g, 0.5 mmol) was treated with 10 mL of 4 M HCl in dioxane (5 mL) with stirring at 25°C for 5 h. Most of the solvent was evaporated *in vacuo* and the sticky residue was then dissolved in distilled water (50 mL) and extracted with ether (3 x 20 mL). The aqueous layer was then passed through a C18 reverse-phase column. The compound was eluted with 20% ethanol–water. Fractions were collected and flushed with N₂ and lyophilized to afford 42 mg (24%) of 1,1,3,3-²H₄-N¹-acetylspermine trihydrochloride. ¹H NMR (D₂O): δ 1.98 (s, 3H), 3.11 (m, 8H), 2.06 (m, 4H), 1.83 (m, 4H). HRMS (*m* + *H*) theor.: 249.4121, found: 249.4130.

Synthesis of N¹-acetylspermidine dihydrochloride

N-Boc-4-aminobutanol (8). 4-Aminobutanol (1.02 g, 11.5 mmol) in 5 mL of dry acetonitrile was added to 1.1 equivalent (2.76 g) of (Boc)₂O in 2.0 equivalent (3.2 mL, 23 mmol) of triethylamine and kept stirring at 25°C overnight. Further workup and purification as described for **1** yielded 1.81 g (78%). ¹H NMR (CDCl₃): δ: 2.88 (s, 1H), 3.67 (m, 2H), 1.63 (m, 2H), 1.58 (m, 2H), 2.95 (m, 2H), 1.31 (s, 9H).

N-Boc-4-amino-1-methanesulfonybutane (9). Methane sulfonyl chloride (1.32 g, 11.5 mmol, 1.2 equivalent) in anhydrous pyridine was added dropwise to the Boc-protected aminobutanol **8** (1.81 g, 9.6 mmol). After 2 h at 25°C, workup and purification as described for **2** gave 1.9 g (69%). ¹H NMR (CDCl₃): δ 2.37 (s, 3H), 3.81 (m, 2H), 1.69 (m, 2H), 1.60 (m, 2H), 2.97 (t, 2H), 1.35 (s, 9H).

N¹-Acetyl-1,1,3,3-²H₄-N⁸-Boc-spermidine (10). Compound **6** (0.3 g, 1.0 equivalent) in 5.0 mL of dry dichloromethane was added to 2.5 mmol (0.67 g) **9**. After 5 h at 50°C, workup as described for **3** gave 0.25 g (34%) of compound **10**. ¹H NMR (CDCl₃): δ 2.105 (s, 3H), 2.88–2.92 (m, 8H), 1.78–1.82 (m, 4H), 1.33 (s, 9H).

1,1,3,3-²H₄-N¹-acetylspermidine dihydrochloride. Compound **10** (0.146 g, 0.5 mmol) was treated with 10 mL of 4 M HCl in dioxane (5 mL) with stirring at 25°C for 5 h. Pure 1,1,3,3-²H₄-N¹-acetylspermidine hydrochloride (50 mg, 38%) was obtained by following the procedure described for 1,1,3,3-²H₄-N¹-acetylspermine hydrochloride. ¹H NMR (D₂O): δ 1.99 (s, 3H), 3.150 (m, 4H), 2.12

(m, 2H), 1.73 (m, 4H); HRMS ($m + H$) theor.: 192.3224, found: 192.3214.

Synthesis of 1,1,3,3,10,10,12,12-²H₈-spermine tetrahydrochloride

1,4-Dimethanesulfonyl-butane (11). Methane sulfonyl chloride (3.15 g, 27.5 mmol, 2.5 equivalent) in anhydrous pyridine was added to 11 mmol (1.0 g) of 1,4-dihydroxybutane, and the mixture was stirred for 2 h at 25°C. Further steps as described for **2** gave 2.76 g (55%) of compound **11**. ¹H NMR (CDCl₃): δ 2.40 (s, 6H), 4.16 (t, 4H), 1.99 (m, 4H).

N¹-Boc-1,1,3,3-²H₄-1,3-diaminopropane (12). One equivalent (0.56 g) of (Boc)₂O in 2.0 equivalent (0.7 mL, 5.12 mmol) of triethylamine was added to 2.56 mmol (0.2 g) of 1,1,3,3-²H₄-1,3-diaminopropane in 5 mL of dry acetonitrile while stirring at 25°C overnight. Workup and purification as described for **1** gave 0.24 g (53%) of compound **12**. ¹H NMR (CDCl₃): δ 1.30 (s, 9H), 1.70 (s, 2H).

N¹,N¹²-Boc₂-1,1,3,3,10,10,12,12-²H₈-spermine (13). Compound **11** (0.172 g, 0.7 mmol) was treated with 2.0 equivalents (0.249 g, 1.4 mmol) of **12** in 4 mL dry dichloromethane at 50°C for 5 h. The crude product was purified by flash silica gel chromatography using hexane–ethylacetate (9:1) as solvent to afford 0.26 g (64%) of compound **13**. ¹H NMR (CDCl₃): δ 1.35 (s, 18H), 1.74 (s, 4H), 2.82 (m, 4H), 1.85 (m, 4H).

1,1,3,3,10,10,12,12-²H₈-spermine tetrahydrochloride. Compound **13** (0.254 g, 0.62 mmol) was further treated with 10 mL of 4 M hydrochloric acid in 5 mL of dioxane. The pure ²H₈-spermine tetrahydrochloride was obtained by the procedure described for 1,1,3,3-²H₄-N¹-acetylspermine hydrochloride. The yield of the final product was 60 mg (27%).

¹H NMR (D₂O): δ 2.61 (m, 4H), 1.62 (m, 4H), 1.51 (m, 4H); HRMS ($m + H$) theor.: 211.4113, found: 211.41125.

Acknowledgements

The authors thank Prof. Daniel Romo of the Department of Chemistry, Texas A&M University, for providing space in his laboratory for the syntheses. The authors thank Dr Satyamaheshwar Peddibhotla, Dr Gil Ma and Vikram Purohit from the Romo Group for their help and valuable suggestions.

REFERENCES

1. Tabor CW, Tabor H. *Microbiol Rev* 1985; **49**: 81–99.
2. Pegg AE. *Biochem J* 1986; **234**: 249–262.

3. Heby O. *Differentiation* 1981; **19**: 1–20.
4. Tabor CW, Tabor H. *Annu Rev Biochem* 1984; **53**: 749–790.
5. Wallace HM, Fraser AV, Hughes A. *Biochem J* 2003; **376**: 1–14.
6. Marton LJ, Pegg AE. *Annu Rev Pharmacol Toxicol* 1995; **35**: 55–91.
7. Pegg AE, Hu RH. *Cancer Lett* 1995; **95**: 247–252.
8. Sunkara PS, Baylin SB, Luk GD. Biological Significance and Basis for New Therapies. In *Inhibition of Polyamine Metabolism*. McCann PP, Pegg AE, Sjoerdsma A (eds). Academic Press: New York, 1987.
9. McCloskey DE, Woster PM, Casero Jr RA, Davidson NE. *Clin Cancer Res* 2000; **6**: 17–23.
10. Marton LJ, Pegg AE. *Annu Rev Pharmacol Toxicol* 1995; **35**: 55–91.
11. Wu T, Yankovskaya V, McIntire WS. *J Biol Chem* 2003; **278**: 20514–20525.
12. Wang Y, Devereux W, Woster PM, Stewart TM, Hacker A, Casero Jr RA. *Cancer Res* 2001; **61**: 5370–5373.
13. Vujcic S, Diegelman P, Bacchi CJ, Kramer DL, Porter CW. *Biochem J* 2002; **367**: 665–675.
14. Murray-Stewart T, Wang Y, Devereux W, Casero RA. *Biochem J* 2002; **368**: 673–677.
15. Seiler N. *Prog Brain Res* 1995; **106**: 333–344.
16. Houen G, Bock K, Jensen AL. *Acta Chem Scand* 1994; **48**: 52–60.
17. Ivanova S, Botchkina GI, Al-Abed Y, Meistrell M, Batliwalla F, Dubinsky JM, Iadecola C, Wang H, Gregersen PK, Eaton JW, Tracey KJ. *J Exp Med* 1998; **188**: 327–340.
18. Dogan A, Rao AM, Baskaya MK, Hatcher J, Temiz C, Rao VL, Dempsey RJ. *J Neurosurg* 1999; **90**: 1078–1082.
19. Dogan A, Rao AM, Hatcher J, Rao VL, Baskaya MK, Dempsey RJ. *J Neurosurg* 1999; **72**: 765–770.
20. Hu RH, Pegg AE. *Biochem J* 1997; **328**: 307–316.
21. Kramer DL, Vujcic S, Diegelman P, Alderfer J, Miller JT, Black JD, Bergeron RJ, Porter CW. *Cancer Res* 1999; **59**: 1278–1286.
22. Mank-Seymour AR, Murray TR, Berkey KA, Xiao L, Kern S, Casero RA. *Clin Cancer Res* 1998; **4**: 2003–2008.
23. Lindsay GS, Wallace HM. *Biochem J* 1999; **337**: 83–87.
24. Chopra S, Wallace HM. *Biochem Pharmacol* 1998; **55**: 1119–1123.
25. Mamos P, Karigiannis G, Athanassopoulos C, Bichta S, Kalpaxis D, Papaioannou, Sindona G. *Tetrahedron Lett* 1995; **36**: 5187–5190.
26. Almeida MLS, Grehn L, Ragnarsson U. *Acta Chem Scand* 1989; **43**: 990–994.
27. Tabor H, Tabor CW, Meis LD. *Methods Enzymol* 1971; **17**: 829–833.