

Research Article

Synthesis of radiolabelled photolabile fusidic acid analogues

SØREN CHRISTIAN SCHOU*, DITE RIBER and GUNNAR GRUE-SØRENSEN

Medicinal Chemistry Research, LEO Pharma, 55 Industriparken, DK-2750 Ballerup, Denmark


Received 27 July 2006; Revised 8 August 2007; Accepted 13 August 2007

Abstract: Two analogues of the antibiotic fusidic acid with photolabile groups, 4-azidophenyl and 4-benzoylphenyl, were successfully labelled with tritium via Pd/C catalysed tritiation of unsaturated precursors. Specific activities of 36 and 44 Ci/mmol were obtained. Copyright © 2007 John Wiley & Sons, Ltd.

Keywords: tritium; photolabile groups; fusidic acid analogues; Pd/C catalysed tritiation

Introduction

Fusidic acid is a unique antibiotic with a potent activity against *Staphylococcus aureus*. Fusidic acid inhibits the bacterial protein synthesis by interference with the elongation factor G (EF-G)/ribosome complex.¹ In an attempt to clarify the mechanism of action of fusidic acid, two radioactive, photolabile-labelled compounds ([24,25-³H₂]**5** and [24,25-³H₂]**8**) were prepared. Tritium was introduced via a Pd/C-catalysed tritiation.



Fusidic Acid

Results and discussion

Compounds **1** and **2** (24-*E/Z*; 1:3 mixture) were prepared according to a described method.¹ [24,25-³H₂]**3** was prepared by treating **1** and **2** with ³H₂ gas in MeOH for 70 min in the presence of Pd/C affording 600 mCi crude product. [24,25-³H₂]**3** was


transformed into the azide [24,25-³H₂]**4** by a procedure reported by Liu and Tor² by treating the amine with triflyl azide and triethylamine in the presence of CuSO₄ as catalyst. The protective group was removed by treating [24,25-³H₂]**4** with K₂CO₃ in MeOH at room temperature for 1.5 h. Final purification was achieved by prep-HPLC (RP-18, THF:H₂O:CH₃COOH (50:50:1)) affording [24,25-³H₂]**5** (37.8 mCi) with a specific activity of 36 Ci/mmol in 6.3% overall radiochemical yield. Because of an unexpected formation of 3-formyl-[24,25-³H₂]**5** when THF:H₂O:HCOOH (50:50:1) was used as eluent in the HPLC purification, a reduced yield was obtained. By replacing formic acid with acetic acid in the HPLC eluent no esterification of [24,25-³H₂]**5** took place (Scheme 1).

Compounds **6** and **7** (24-*E/Z*; 1:3 mixture), prepared as described in the literature,¹ were treated with ³H₂ gas in the presence of Pd/C in MeOH for 12 min affording 638 mCi of crude [24,25-³H₂]**8**. The crude product was filtered through silica gel to remove remains of catalyst followed by prep-HPLC purification (RP-18, THF:H₂O:CH₃COOH (37:63:0.5)) affording [24,25-³H₂]**8** (161 mCi) with a specific activity of 44 Ci/mmol in 25% overall radiochemical yield. Owing to difficulties in the HPLC separation of unreacted *E*-isomer (**6**) from [24,25-³H₂]**8**, not all [24,25-³H₂]**8** was isolated from the reaction mixture (Scheme 2).


Acknowledgements

We are indebted to Drs Stefan Nückel and Christian Meisterhans (catalytic tritiation) at RC TRITEC AG, Teufen, Switzerland, for the invaluable assistance.

*Correspondence to: Søren Christian Schou, Medicinal Chemistry Research, LEO Pharma, 55 Industriparken, DK-2750 Ballerup, Denmark. E-mail: sscdk@leo-pharma.com



Scheme 1



Scheme 2

REFERENCES

1. Riber D, Venkataramana M, Sanyal S, Duvold T.
J Med Chem 2006; **49**: 1503–1505.

2. Liu Q, Tor Y. *Org Lett* 2003; **5**: 2571–2572.