

# n-3 PUFA dietary supplementation inhibits proliferation and store-operated calcium influx in thymoma cells growing in Balb/c mice

Gabriella Calviello,\* Paola Palozza,\* Fiorella Di Nicuolo,\* Nicola Maggiano,<sup>†</sup> and Gianna M. Bartoli,<sup>1,\*</sup><sup>§</sup>

Institute of General Pathology\* and Institute of Pathology,<sup>†</sup> Catholic University, L. go F. Vito 1-00168, Rome, Italy, and Department of Biology,<sup>§</sup> Tor Vergata University, Rome, Italy

**Abstract** The antitumor effect of daily individual administration of eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) (2 g/kg body weight) in Balb/c mice bearing a transplantable thymoma was investigated. Mice received oleic acid (control group), EPA and DHA ethyl esters starting 10 days before tumor inoculation. Analysis of phospholipid composition of neoplastic cell revealed that EPA and DHA levels were significantly increased (63 and 22% increase) after EPA and DHA treatments, respectively. Conversely, decreased levels of arachidonic acid were found in both cases (19 and 24% decrease in EPA and DHA groups, respectively). EPA and DHA delayed the appearance of macroscopic ascites (100% of animal, from 7 to 28 days), prolonged animal survival (100% of animal, from 22 to 32 and 33 days, respectively) and reduced the percentage of proliferating tumor cells detected by immunostaining of proliferation cell nuclear antigen (PCNA) (80 and 85% decrease, respectively). Moreover, the regulatory effects of these dietary n-3 fatty acids on the influx of Ca<sup>2+</sup>, activated by depletion of intracellular stores with thapsigargin (Tg), were investigated. By using a Ca<sup>2+</sup>-free/Ca<sup>2+</sup>-reintroduction protocol and Fura-2 as fluorescent indicator of intracellular free Ca<sup>2+</sup> ([Ca<sup>2+</sup>]<sub>i</sub>), we observed that EPA and DHA treatments markedly decreased Tg-induced rise in [Ca<sup>2+</sup>]<sub>i</sub> (49 and 37% decrease, respectively). This effect was related to the inhibition of the store-operated Ca<sup>2+</sup> influx, as confirmed also by Mn<sup>2+</sup> influx experiments. The inhibitory action of EPA and DHA on the store-operated Ca<sup>2+</sup> influx could explain, at least in part, their antitumoral activity, as this Ca<sup>2+</sup> mobilization pathway appears to be involved in the cell signaling occurring in non-excitable cells to evoke many cellular processes, including cell proliferation.— Calviello, G., P. Palozza, F. Di Nicuolo, N. Maggiano, and G. M. Bartoli. N-3 PUFA dietary supplementation inhibits proliferation and store-operated calcium influx in thymoma cells growing in Balb/c mice. *J. Lipid Res.* 2000. 41: 182–188.

**Supplementary key words** DHA • EPA • fatty acids • tumor growth • phospholipids • signaling

There is increasing evidence for a protective effect of n-3 polyunsaturated fatty acids (PUFAs) against the de-

velopment and progression of several types of cancer (1, 2). We have previously reported the antiproliferative effect of a dietary supplementation with a mixture of EPA and DHA in the colonic mucosa of human subjects at high risk for colon cancer (3). Recently, we found that EPA dietary supplementation is able to inhibit the development of focal hepatic preneoplastic lesions during chemical carcinogenesis in rats (4). Moreover, we observed that individual administration of purified EPA and DHA inhibited the growth of Morris hepatocarcinoma 3924A transplanted in rats (5). In this study we found, in agreement with other authors (6–8), that the antitumoral action of EPA and DHA was related to the inhibition of cell proliferation and apoptosis.

The mechanism(s) underlying the antitumor effect of n-3 PUFAs have not been completely elucidated. The reduction in the synthesis of eicosanoids derived from arachidonic acid is considered to be involved in the growth inhibitory effect of n-3 PUFAs (9, 10). It has also been suggested that cancer cells are killed by lipid peroxidation products generated in larger amount in cells enriched with n-3 PUFAs (11). We recently suggested that the enhancement of  $\beta$ -oxidation activity by n-3 PUFAs and the consequent overproduction of H<sub>2</sub>O<sub>2</sub> may originate a mild oxidative stress able to inhibit neoplastic growth (4). Finally, different studies have reported that n-3 PUFAs can modulate the activity of components of intracellular signaling (12–14). It is well established that Ca<sup>2+</sup> is among the major intracellular factors involved in the signaling transduction pathways evoking cell growth and proliferation, and other key processes such as gene expression, contraction, secretion, and cellular metabolism (15, 16). Many of these processes require a sustained

Abbreviations: EPA, eicosapentaenoic acid; DHA, docosahexaenoic acid; PCNA, proliferating cell nuclear antigen; PUFAs, polyunsaturated fatty acids; SOCI, store-operated Ca<sup>2+</sup> influx; Tg, thapsigargin.

<sup>1</sup> To whom correspondence should be addressed.

increase in the concentration of intracellular free  $\text{Ca}^{2+}$  ( $[\text{Ca}^{2+}]_i$ ) that is mainly elicited by the influx of the cation into the cell from the extracellular compartment (16). Recently, it has been established that n-3 PUFAs suppress voltage-gated L-type  $\text{Ca}^{2+}$  currents in rat cardiac myocytes (17). In non-excitabile cells, the major pathway of  $\text{Ca}^{2+}$  entry is the so-called store-operated  $\text{Ca}^{2+}$  influx (SOCl), triggered by agonists (such as inositol 1,4,5-trisphosphate) that act by emptying the high affinity intracellular  $\text{Ca}^{2+}$  stores (18). In particular, SOCl has been involved in the cell signaling occurring in non-excitabile cells to evoke different cell processes including gene regulation and cell growth (19). Recently it has been observed that the addition of PUFAs to different strains of cells cultured in vitro modify the extent of SOCl (20). However, at the moment, the effect of a physiological membrane incorporation of PUFAs on this pathway of  $\text{Ca}^{2+}$  entry, obtained in vivo after a dietary treatment, has never been studied. In particular, the effect of n-3 PUFAs is almost completely unknown.

The purpose of the present study was to evaluate the effects of an oral administration of EPA and DHA on the growth of a highly malignant thymoma transplanted in the peritoneal cavity of Balb/c mice. We intended also to study the regulatory effect of these dietary fatty acids incorporated in cell phospholipids on the store-dependent pathway of  $\text{Ca}^{2+}$  influx.

## MATERIALS AND METHODS

### Animals and treatments

Male inbred Balb/c mice, aged 6 weeks (average body weight:  $23.4 \pm 1.7$  g), were housed in cages and fed a non-purified commercial diet (Altromin-Rieper, Bolzano, Italy). The composition of the diet was (% w/w) crude protein, 23; fat, 5.5; fiber, 5; mineral, 8; carbohydrates, 58.5; water, 12. The vitamin mixture added to the diet was 2.5 g/kg and the mineral mixture was 0.52 g/kg by weight. The diet PUFA content was the following: 7.5 g/kg n-6 PUFAs (linoleic acid, 18:2 n-6) and 1.02 g/kg n-3 PUFAs as 18:3, 18:4, 20:5, and 22:6. The animals were given free access to food and water.

Two sequential experiments were performed. In the first experiment 48 animals were randomly divided into three treatment groups (16 mice/group). Groups 1 and 2 received a daily dose of EPA or DHA ethyl esters (2 g/kg body weight), respectively, by gavage. To provide an equal caloric intake, the control group was supplemented with oleic acid ethyl esters (OA) at the same dosage. The dose of n-3 PUFAs was chosen to approximate the dose (1 g/kg body weight) that was recently found to inhibit the growth of a transplanted hepatocarcinoma in rats (5). This dose is also very similar to that used in our previous human study (3). Body surface area was used as the reference for conversion of the dosage among species (humans, rats, and mice) (21). Administration of n-3 PUFAs together with a normal dietary regimen ensured that the animals' essential fatty acid requirements were met. After EPA and DHA administration n-3/n-6 PUFA dietary ratio increased from 0.14 to 2.5. Purity of fatty acids (kindly provided by Hoffmann-La Roche, Basel, Switzerland) was verified by gas-liquid chromatography and calculated to be 90–92%. The animals received OA, EPA, or DHA for 10 days before inoculation of a highly malignant thymoma in ascites form and until the end of the experiment. Ten days represents the time required to

obtain the maximal incorporation of EPA and DHA (data not shown). All the animals of this experiment were used to study the appearance of tumor and survival.

In the second experiment the animals were divided into three groups (44 mice/group) and treated as reported above for the first experiment. Eight animals from each group were killed 4 days after tumor transplantation to evaluate tumor cell proliferation by immunostaining of proliferating cell nuclear antigen (PCNA). After 3 additional days, 20 animals from each group were killed and tumor thymocytes were harvested for cell counting and evaluation of fatty acid composition and  $\text{Ca}^{2+}$  influx. The remaining mice were used to study the appearance of tumor and survival. The animal-use protocol was approved by the Ministry of Health, Veterinary Service, Rome, Italy.

### Tumor cells

A highly malignant thymoma growing in ascitic form in the peritoneum of Balb/c mice was used. The neoplastic cell line derived from a strain of malignant spontaneous thymoma of Balb/c mice (originally propagated in the Institute of Pathology of the University of Perugia, Italy) (22). The ascites thymoma cells were maintained by weekly intraperitoneal transplantation in syngeneic Balb/c mice.

### Tumor cell inoculation and count

Tumor cells were aseptically inoculated ( $1 \times 10^7$  in 0.3 ml 0.9% NaCl) in the peritoneal cavity of Balb/c mice. Tumor cell count was performed after harvesting the cells from the peritoneum of mice 7 days after transplantation, because these cells grow exponentially for 5 days, reaching a plateau at 6–7 days. To ensure the complete collection of cells, the peritoneum was washed twice with 5 ml 0.9% NaCl. Cell suspension in ascitic fluid was diluted in Ringer-HEPES 50 mm, pH 7.4 (130 mm NaCl, 5 mm KCl, 1 mm  $\text{MgSO}_4$ , 1.3 mm  $\text{CaCl}_2$ , and 50 mm HEPES) to avoid agglutination, and tumor thymocytes were immediately counted using a Trypan Blue dye exclusion method using a Burkner chamber.

### Cell proliferation

The percentage of proliferating cells was determined by immunostaining of proliferating cell nuclear antigen (PCNA), a well-known cell cycle-related antigen detected by immunofluorescence in S-phase nuclei (23). Cells were harvested from peritoneum at the 4th day after transplantation, during the exponential phase of duplication. They were cytocentrifuged, fixed with acetone, and incubated for 1 h with the anti-PCNA antibody (PC 10, Boehringer Mannheim, Indianapolis, IN) at a dilution of 1:1,000. The cells were counterstained with hematoxylin after visualization of the reaction product with diaminobenzidine (DAB) and routine streptavidine–biotin–peroxidase treatment. The labeling index (LI%), defined as the percentage of positively staining nuclei, was determined from a count of 1000 cells at high power magnification (400 $\times$ ).

### Phospholipid fatty acid composition of tumor cells

Tumor cells were harvested from the peritoneum of mice at the 7th day after transplantation. The extraction of lipids from tumor cells was performed according to the method of Bligh and Dyer (24). Total phospholipids were separated from neutral lipids by thin-layer chromatography using toluene–diethyl ether–ethanol 35:3.5:1 (v/v/v) as the solvent system (25). After transesterification and extraction with n-hexane, fatty acid methyl esters were separated by gas-liquid chromatography using a Carlo Erba HRGC 5300 Mega-series chromatograph (5). Individual fatty acid methyl esters were identified by comparing their retention times to known standards (Sigma, St. Louis, MO). The areas under the peaks of each fatty acid were measured with a Carlo Erba Mega-

series integrator and expressed as percentages of the total area of all major fatty acid peaks.

### Intracellular free $\text{Ca}^{2+}$ measurements

$[\text{Ca}^{2+}]_i$  was evaluated by measuring the fluorescence of the  $\text{Ca}^{2+}$ -indicating dye, Fura-2 (26). The cells were loaded with 3  $\mu\text{M}$  Fura-2/AM (Fura-2/acetoxymethylester), an esterified derivative of Fura-2, as previously described (22). After loading, Fura-2/AM was hydrolyzed by intracellular esterases to Fura-2 that become entrapped in the cytosol and bind to free  $\text{Ca}^{2+}$ . Briefly, cells ( $2.5 \times 10^6/\text{ml}$ ) harvested at the 7th day from inoculum were washed twice and suspended in Ringer-HEPES 50 mM, pH 7.4. They were incubated in the dark at 34°C for 10 min and then supplemented with Fura-2/AM (Sigma, St. Louis, MO). After further 15 min at 34°C, to allow the incorporation of Fura-2/AM into the cells, the cell suspension was diluted 1:5 in the same buffer and incubated for an additional 10 min to allow complete hydrolysis of the Fura-2/AM. Cells were then washed and resuspended at the same initial cell concentration and kept at room temperature until used. Just before use, 1 ml of the stock suspension was rapidly centrifuged and resuspended at a final concentration of  $1.25 \times 10^6$  cells/ml in  $\text{Ca}^{2+}$ -free Ringer-HEPES 50 mM, pH 7.4, in which  $\text{CaCl}_2$  was omitted and 0.2 mM EGTA was added. The measurements were performed in a Perkin-Elmer LS3B fluorometer (excitation 340 nm, emission 510 nm) at 30°C, to minimize dye leakage from the cells. Calibration was performed according to the protocol described by Arslan et al. (27). A value of 184 nm was used as the  $K_D$  for the  $\text{Ca}^{2+}$ -Fura-2 complex at 30°C.

SOCI was investigated using thapsigargin (Tg), an endoplasmic reticulum  $\text{Ca}^{2+}$ -ATP-ase inhibitor that, leading to the depletion of intracellular  $\text{Ca}^{2+}$  stores, directly triggers this pathway of  $\text{Ca}^{2+}$  influx. SOCI was measured after cell pretreatment with 500 nM Tg in a  $\text{Ca}^{2+}$ -free medium and subsequent readmission of 1.5 mM  $\text{Ca}^{2+}$  (20).

$\text{Mn}^{2+}$  quenching of Fura-2 fluorescence was evaluated as surrogate of  $\text{Ca}^{2+}$  to trace  $\text{Ca}^{2+}$  influx mechanism (20). The measurements were performed at 360 nm, the excitation wavelength at which the indicator fluorescence is not influenced by changes in  $\text{Ca}^{2+}$  concentration (isosbestic point).

### Statistical analysis

The results were expressed as the means  $\pm$  SE. Comparisons among groups were made by one-way analysis of variance (ANOVA) using Minitab software (Minitab, State College, PA). When significant differences were found, post hoc comparisons of means were made using Fisher's test. Differences were considered significant at  $P < 0.05$ .

## RESULTS

Dietary supplementations with n-3 PUFAs at the dose used in this study did not modify daily food intake of mice. Weight gain, measured prior to tumor inoculation, was not significantly different (ANOVA,  $P < 0.001$ ) among the groups of animals (data not shown).

### Tumor growth

Figure 1 shows macroscopical ascites appearance and animal survival after transplantation. All the animals of the OA group exhibited maximal ascites growth within 7 days (panel A) and died within 21 days after tumor inoculation (panel B). Treatment with both EPA and DHA delayed ascites appearance and prolonged animal survival.

### Phospholipid fatty acids of neoplastic thymocytes

The content of the principal PUFAs in total phospholipids (PL) from OA-, EPA- and DHA-treated mice is shown in Fig. 2. The content of the PUFAs was significantly affected by the treatment with the two n-3 fatty acids with respect to the control group. EPA treatment significantly increased EPA (20:5n-3) and its metabolite docosapentaenoic acid (DPA, 22:5n-3) without modifying the level of DHA (22:6n-3). Conversely, DHA supplementation increased the amount of DHA itself (22:6n-3). Levels of linoleic acid were unchanged in both groups of animals and the content of arachidonic acid (20:4n-6) was significantly decreased in both groups.





Fig. 1. Development of tumor (A) and animal survival (B) after tumor cell transplantation in OA-, EPA-, and DHA- treated mice. Values are means of two experiments (16 mice for each experiment). Statistical comparisons were performed at the 7th (panel A) and 22nd day (panel B) after tumor transplantation, when 100% OA-treated animals had tumors or were dead, respectively. \* Significantly different from OA group ( $P \leq 0.001$ ; one-way ANOVA, followed by Fisher's test).



**Fig. 2.** Effect of EPA and DHA supplementation on the content of the principal PUFAs in total phospholipids of thymoma cells. Values are means  $\pm$  SEM for 10 mice from each group. \* Significantly different from OA-treated group ( $P < 0.005$ ; one-way ANOVA, followed by Fisher's test).


### Cell count and proliferation

**Figure 3** shows the effects of EPA or DHA on the number (A) and proliferation (B) of tumor cells growing in the peritoneal cavity of control and treated mice. Both n-3 PUFA treatments significantly reduced the number of neoplastic thymocytes (A) harvested 7 days after tumor inoculation. Cell proliferation (B) was evaluated by PCNA immunostaining at the 4th day after inoculum. PCNA-LI% of thymoma cells isolated from EPA and DHA was decreased 5- and 7-fold, respectively, as compared to cells from OA-treated mice.


### Store-dependent $\text{Ca}^{2+}$ influx of neoplastic thymocytes

**Figure 4** shows the modification in SOCI of tumor thymocytes enriched in vivo with EPA and DHA and deter-

mined by the  $\text{Ca}^{2+}$ -free/ $\text{Ca}^{2+}$ -reintroduction protocol after pretreatment with 500 nm Tg. Tg was used in supra-maximal concentration to ensure the complete release of  $\text{Ca}^{2+}$  stored in the endoplasmic reticulum. The  $[\text{Ca}^{2+}]_i$  increase, measured in the absence of extracellular  $\text{Ca}^{2+}$  and triggered by the Tg-induced depletion of intracellular stores, was small and not different among control and EPA- or DHA- enriched cells. After 6 min of Tg treatment, calcium readmission, at the concentration of 1.5 mM, resulted in a rapid  $[\text{Ca}^{2+}]_i$  rise, which is known to be dependent on activation of store-operated  $\text{Ca}^{2+}$  influx. The rise ended in a plateau phase and, as better shown by **Fig. 5**, both EPA and DHA significantly reduced  $\text{Ca}^{2+}$  increase at the same degree.



**Fig. 3.** Effect of EPA and DHA supplementation on cell number and cell proliferation. A: Cell number was evaluated at the 7th day after transplantation. B: Cell proliferation was evaluated at the 4th day as labeling index (LI) by PCNA immunostaining. LI, percentage of positively stained nuclei. Values are means  $\pm$  SEM for 8–10 mice from each group. \* Significantly different from OA-treated group ( $P < 0.05$ ; one-way ANOVA, followed by Fisher's test).




**Fig. 4.** Effect of EPA and DHA supplementation on  $\text{Ca}^{2+}$  influx induced by  $\text{Ca}^{2+}$  readmission in thymoma cells pretreated with thapsigargin in  $\text{Ca}^{2+}$ -free medium. The cells were loaded with Fura-2/AM and then resuspended in  $\text{Ca}^{2+}$ -free medium. Addition: thapsigargin (Tg, 500 nm) and  $\text{CaCl}_2$  (1.5 mM). Excitation, 340 nm; emission, 510 nm. The results are representative of at least 10 different experiments.

To exclude that the reduced increase in cytosolic  $\text{Ca}^{2+}$  concentration after Tg/ $\text{Ca}^{2+}$  addition was due to the enhancement of  $\text{Ca}^{2+}$  extrusion through the plasma membrane, the influx of  $\text{Mn}^{2+}$  was also analyzed (Fig. 6). This cation enters the cytosol through the same store-dependent pathway as  $\text{Ca}^{2+}$  and its flux is almost completely unidirectional towards the interior of the cell being poorly extruded from the cytosol (28). The figure shows the effect of the dietary supplementations with EPA or DHA on  $\text{Mn}^{2+}$  influx in Tg-pretreated neoplastic cells.  $\text{Mn}^{2+}$  influx has been measured as percentage of  $\text{Mn}^{2+}$  quenching of Fura-2 signal. As shown by the figure, both PUFAs markedly inhibited  $\text{Mn}^{2+}$  influx in Tg-pretreated neoplastic thymocytes.

## DISCUSSION


Different epidemiological and experimental studies demonstrate the antitumor action of n-3 PUFAs and support the hypothesis that they may have an anticancer effect also in humans (1-2, 29).



**Fig. 5.** Effect of on  $[\text{Ca}^{2+}]_i$  in thymoma cells pretreated with thapsigargin in  $\text{Ca}^{2+}$ -free medium, before (basal, 1 min) and after readmission of  $\text{Ca}^{2+}$  (12 min). Values are means  $\pm$  SEM for 10 mice from each group. \* Significantly different from OA-treated group ( $P < 0.005$ ; one-way ANOVA, followed by Fisher's test).

The present study investigated the effect of separate dietary administration of EPA and DHA on the growth of a highly malignant thymoma transplanted in mice. This tumor grows in the peritoneum as a cell suspension and is particularly suitable to study the effects of dietary manipulation on different cellular parameters.

Both EPA and DHA inhibited tumoral development and growth to the same extent, delaying macroscopical tumor appearance, and animal death. The antitumoral effect of EPA and DHA treatment was further proven by



**Fig. 6.** Effect of EPA and DHA supplementation on  $\text{Mn}^{2+}$  influx in thymoma cells pretreated with thapsigargin in  $\text{Ca}^{2+}$ -free medium. Cells ( $1.25 \times 10^6$ ) were pretreated with thapsigargin (Tg, 500 nm) 6 min before the addition of  $\text{Mn}^{2+}$  (0.1 mM  $\text{MnCl}_2$ ). The influx of  $\text{Mn}^{2+}$  was evaluated from the rate of  $\text{Mn}^{2+}$  quenching of Fura-2 fluorescence at the isosbestic point (excitation: 360 nm). The results are representative of at least 10 different experiments.

the reduced number of cells growing in the peritoneum and by the marked inhibition of thymoma cell proliferation, as demonstrated by PCNA immunostaining. We cannot exclude that EPA and DHA may act also by modulating the apoptotic process, as shown previously by Finstad et al. (8) in cell lines or by us in animal models (5, 30). However, in the present work, the number of apoptotic cells, evaluated in the peritoneum of Balb/c mice by the in situ nick end labeling method, was very low and it was not possible to appreciate the differences among the three treatment groups (data not shown). Presumably, the physiological phagocytosis occurring in the peritoneum in vivo did not allow the correct evaluation of the process.

Analysis of phospholipid fatty acid composition revealed increased levels of EPA and its metabolite docosapentaenoic acid (DPA, 22:5n-3) during EPA treatment and increased levels of DHA during DHA treatment. Conversely, the percentage of arachidonic acid was diminished in phospholipids from both groups. Such modifications confirmed those largely reported in *in vitro* and in *in vivo* models and in normal and tumor tissue and cells (5, 25, 31).

Among the mechanisms invoked to explain the n-3 PUFA inhibitory effect on cellular growth is their ability to modulate the activity of component of intracellular signaling (i.e., protein kinases, eicosanoids, reactive oxygen species, etc.) (4, 12-14). It is well known that  $Ca^{2+}$  is an essential component in the signaling pathways evoking key cellular processes including cell proliferation and growth (15, 16). For this reason we investigated  $Ca^{2+}$  metabolism in thymoma cells, whose proliferation was markedly inhibited by dietary treatment with EPA and DHA. In particular, we studied  $Ca^{2+}$  influx, which in non-excitable cells regulates different processes as exocytosis, contraction, enzyme control, gene regulation, cell proliferation, and apoptosis (19). The predominant  $Ca^{2+}$  entry pathway in non-excitable cells is SOCI, in which  $Ca^{2+}$  entry is regulated by the filling state of the agonist-sensitive intracellular  $Ca^{2+}$  stores (18). When the stores are full, they prevent extracellular  $Ca^{2+}$  entry, but after emptying,  $Ca^{2+}$  influx is promoted. Thymoma cells derived from both EPA- and DHA-treated mice showed a lower level of SOCI with respect to control cells. We studied SOCI after Tg-induced store depletion and subsequent  $Ca^{2+}$  addition. Moreover, the finding that n-3 PUFAs also inhibited  $Mn^{2+}$  influx in Tg-pretreated neoplastic cells strongly suggested that EPA and DHA treatments specifically inhibited the SOCI pathway. Our results are in good agreement with those recently obtained by Gamberucci, Fulceri, and Benedetti (20). They demonstrated the inhibition of the SOCI pathway by unsaturated fatty acids added *in vitro* to different strains of cells treated with Tg. This inhibition was related to the concentration and the degree of unsaturation of the fatty acids used. However, in the present work, thymoma cell phospholipids, in addition to an enrichment in n-3 PUFAs, showed also a decreased content of arachidonic acid after n-3 PUFA treatment. Concerning arachidonic acid, there is wide debate in the literature. Khordova and Astashkin (32) observed that arachidonic acid added to lymphocytes inhibited or induced store-dependent

$Ca^{2+}$  entry, depending on the concentration of the fatty acid. The activation of SOCI by arachidonic acid administration was also reported by Rzigalinski, Blackmore, and Rosenthal (33). Finally, it has also been recently shown (33, 34) that during the rise of  $Ca^{2+}$  associated to SOCI, phospholipase  $A_2$  ( $PLA_2$ ) is activated and arachidonic acid is released. These authors hypothesized that arachidonic acid may function as a second messenger involved in the coupling between store depletion and  $Ca^{2+}$  entry in SOCI. In agreement with these results, the decreased levels of arachidonic acid in thymoma phospholipids after treatments with EPA or DHA could explain the lower activity of SOCI, even though further work is needed to prove the existence of a  $PLA_2$ /arachidonic acid coupling signal between store emptying and  $Ca^{2+}$  influx also in thymoma cells.

Moreover, our results are in agreement with those obtained in muscle cells cultured *in vitro* by Hirafuji et al. (35). They observed that treatment with DHA for 48 h inhibited  $Ca^{2+}$  influx induced by angiotensin II. These authors hypothesized that the incorporation of DHA in membrane phospholipids is a necessary prerequisite for its inhibitory effect on angiotensin II-induced  $Ca^{2+}$  influx. Conversely, previous studies (36) had shown that EPA and DHA, added as free fatty acids in Jurkat cells cultured *in vitro*, induced a sustained increase of  $[Ca^{2+}]_i$  within a minute. However, this increase was related only to  $Ca^{2+}$  released from intracellular stores and not to  $Ca^{2+}$  influx through the plasma membrane.

The present data demonstrate for the first time that n-3 PUFAs incorporated into membrane phospholipids after a dietary treatment inhibited the store-dependent pathway of  $Ca^{2+}$  influx. This pathway, in addition to its more obvious role as a mechanism to refill empty stores, has been involved in cellular signalling (19), particularly in the regulation of cell proliferation (37, 38).

The results obtained suggest that the antitumoral effect of EPA and DHA is related to their inhibition of cell proliferation. This inhibition, in turn, could be explained, at least in part, by their ability to decrease the store-dependent  $Ca^{2+}$  entry. In agreement with our hypothesis, Kokoska, Smith and Miller (39) have recently found that SOCI is inhibited by indomethacin and other inhibitors of prostaglandin synthesis. They hypothesize that the known antiproliferative role of these nonsteroidal antiinflammatory drugs on gastric mucosa cells could be explained by their ability to inhibit the SOCI pathway in these cells.

In conclusion, the data herein presented demonstrate that dietary treatment with EPA or DHA inhibits the growth of highly malignant ascites thymoma and delays the animal's death. They suggest a possible use of these fatty acids as chemopreventive agents. The antitumoral effect of n-3 PUFAs can be explained by the inhibition of cell proliferation and the decrease in the activity of SOCI, a  $Ca^{2+}$  influx pathway that appears to be involved in the signaling that evokes many key cell processes. ■

The authors thank Hoffmann-La Roche Ltd, Basel, Switzerland for the generous provision of purified n-3 PUFA ethyl esters.

## REFERENCES

1. Reddy, B. S. 1992. Dietary fat and colon cancer: animal model studies. *Lipids*. **27**: 807–812.
2. Cave, W. T., Jr. 1994. Dietary n-3 ( $\omega$ -3) polyunsaturated fatty effects on animal tumorigenesis. *FASEB J.* **5**: 2160–2166.
3. Anti, M., G. Marra, F. Armelao, G. M. Bartoli, R. Ficarella, A. Percesepe, I. De Vitis, G. Maria, L. Sofo, G. L. Rapaccini, N. Gentiloni, E. Piccioni, and G. Miggiano. 1992. Effect of  $\omega$ -3 fatty acids on rectal mucosacell proliferation in subject at risk for colon cancer. *Gastroenterology*. **103**: 883–891.
4. Calviello, G., P. Palozza, P. Franceschelli, A. Frattucci, E. Piccioni, L. Tessitore, and G. M. Bartoli. Eicosapentaenoic acid inhibits the growth of liver preneoplastic lesions and alters membrane phospholipid composition and peroxisomal  $\beta$ -oxidation. *Nutr. Cancer*. **34**: 206–212.
5. Calviello, G., P. Palozza, E. Piccioni, N. Miggiano, A. Frattucci, P. Franceschelli, and G. M. Bartoli. 1998. Dietary supplementation with eicosapentaenoic and docosahexaenoic acid inhibits growth of Morris hepatocarcinoma 3924A in rats: effects on proliferation and apoptosis. *Int. J. Cancer*. **75**: 699–705.
6. Krokan, H., P. K. Rudra, S. Schonberg, W. Slettahjell, K. Solum, S. Mollerup, E. Eilertsen, L. Maele, and A. Haugen. 1993. Effect of n-3 fatty acids on the growth of human tumor cell lines in cell culture and in nude mice: *In Omega-3 Fatty Acids: Metabolism and Biological Effects*. C. A. Drevon, I. Baksas, and H. E. Krokan, editors. Birkhauser Verlag, Basel. 327–341.
7. Minami, M., and M. Noguchi. 1996. Effects of low-dose eicosapentaenoic acid, docosahexaenoic acid and dietary fat on the incidence, growth and cell kinetics of mammary carcinomas in rats. *Oncology*. **53**: 398–405.
8. Finstad, H. S., C. A. Drevon, M. A. Kulset, A. V. Synstad, and E. Knudsen. 1998. Cell proliferation, apoptosis and accumulation of lipid droplets in U937-1 cells incubated with eicosapentaenoic acid. *Biochem. J.* **336**: 451–459.
9. Karmali, R. A. 1987. Eicosanoids in neoplasia. *Prev. Med.* **16**: 493–502.
10. Rose, D. P., J. M. Connolly, J. Rayburn, and M. Coleman. 1995. Influence of diets containing eicosapentaenoic acid and docosahexaenoic acid on growth and metastasis of breast cancer cells in nude mice. *J. Natl. Cancer Inst.* **87**: 587–592.
11. Gonzales, M. J., R. A. Shemmel, L. Dugan, J. I. Gray, and C. W. Welsh. 1993. Dietary fish oil inhibits human breast carcinoma growth: a function of increased lipid peroxidation. *Lipids*. **28**: 827–832.
12. Rao, C. V., and B. S. Reddy. 1993. Modulating effect of amount and types of dietary fat ornithine decarboxylase, tyrosine protein kinase and prostaglandins production during colon carcinogenesis in male F344 rats. *Carcinogenesis*. **14**: 1327–1333.
13. Marignani, P. A., and R. J. Sebaldt. 1995. Formation of second messenger diradylglycerol in murine peritoneal macrophages is altered after in vivo (n-3) polyunsaturated fatty acid supplementation. *J. Nutr.* **125**: 3030–3040.
14. De Jonge, H. W., D. H. Dekkers, and J. M. Lamers. 1996. Polyunsaturated fatty acids and signalling via phospholipase C-beta and A2 in myocardium. *Mol. Cell Biochem.* **157**: 199–210.
15. Berridge, M. J. 1993. Inositol trisphosphate and calcium signaling. *Nature*. **361**: 315–325.
16. Berridge, M. J. 1995. Calcium signalling and cell proliferation. *Bioessays*. **17**: 491–500.
17. Xiao, Y. F., A. M. Gomez, J. P. Morgan, W. J. Lederee, and A. Leaf. 1997. Suppression of voltage-gated L type  $Ca^{2+}$  currents by polyunsaturated fatty acids in adult and neonatal rat cardiomyocytes. *Proc. Natl. Acad. Sci. USA*. **94**: 4182–4187.
18. Putney, J. W. 1990. Capacitative calcium entry revisited. *Cell Calcium*. **11**: 611–624.
19. Parekh, A. B., and R. Penner. 1997. Store depletion and  $Ca^{2+}$  influx. *Physiol. Rev.* **77**: 901–930.
20. Gamberucci, A., R. Fulceri, and A. Benedetti. 1997. Inhibition of store-dependent capacitative  $Ca^{2+}$  influx by unsaturated fatty acids. *Cell Calcium*. **21**: 375–385.
21. Freireich, E. J., E. A. Gehan, D. P. Rall, L. H. Schmidt, and H. E. Skipper. 1966. Quantitative comparison of toxicity of anticancer agents in mouse, rat, hamster, dog, monkey and man. *Cancer Chemother. Rep.* **50**: 219–243.
22. Calviello, G., P. Ricci, and G. M. Bartoli. 1993. *tert*-Butyl hydroperoxide induced  $[Ca^{2+}]$  increase in thymus and thymoma cells. *Biochem. Biophys. Res. Commun.* **197**: 859–868.
23. Sasano, H., S. Miyazaki, T. Nishihira, T. Sawai, and H. Nagura. 1992. The proliferative cell fraction in cytology specimens. *Am. J. Clin. Pathol.* **98**: 161–166.
24. Bligh, E. G., and W. J. Dyer. 1959. A rapid method of total lipid extraction and purification. *Can. J. Biochem. Physiol.* **37**: 911–917.
25. Calviello, G., P. Palozza, P. Franceschelli, and G. M. Bartoli. 1997. Low-dose eicosapentaenoic or docosahexaenoic acid administration modifies fatty acid composition and does not affect susceptibility to oxidative stress in rat erythrocytes and tissues. *Lipids*. **32**: 1075–1083.
26. Gryniewicz, G., M. Poenie, and R. Y. Tsien. 1985. A new generation of  $Ca^{2+}$  indicators with greatly improved fluorescence properties. *J. Biol. Chem.* **260**: 3440–3450.
27. Arslan P., F. Di Virgilio, M. Beltrame, R. Y. Tsien, and T. Pozzan. 1985. Cytosolic  $Ca^{2+}$  homeostasis in Ehrlich and Yoshida carcinomas. A new, membrane permeant chelator of heavy metals reveals that these ascites tumor cell lines have normal cytosolic free  $Ca^{2+}$ . *J. Biol. Chem.* **260**: 2719–2727.
28. Broad, L. M., D. A. Powis, and C. W. Taylor. 1996. Differentiation of BC3H1 smooth muscle cells changes the bivalent cation selectivity of the capacitative  $Ca^{2+}$  entry pathway. *Biochem. J.* **316**: 759–764.
29. Dyerberg, J., H. O. Bang, and N. Hjorne. 1975. Fatty acid composition of the plasma lipids in Greenland eskimos. *Am. J. Clin. Nutr.* **28**: 955–966.
30. Calviello, G., P. Palozza, N. Miggiano, E. Piccioni, P. Franceschelli, A. Frattucci, F. Di Nicuolo, and G. M. Bartoli. 1999. Cell proliferation, differentiation, and apoptosis are modified by n-3 polyunsaturated fatty acids in normal colonic mucosa. *Lipids*. **34**: 599–604.
31. Gronn, M., E. Christensen, T. A. Hage, and B. O. Christophersen. 1992. Effects of dietary purified eicosapentaenoic acid (20:5 n-3) and docosahexaenoic acid (22:6 n-3) on fatty acid desaturation and oxidation in isolated rat liver cells. *Biochim. Biophys. Acta*. **1125**: 35–43.
32. Khorodova, A. B., and E. I. Astashkin. 1994. A dual effect of arachidonic acid on  $Ca^{2+}$  transport systems in lymphocytes. *FEBS Lett.* **353**: 167–170.
33. Rzigalinski, B. A., P. F. Blackmore, and M. D. Rosenthal. 1996. Arachidonate mobilization is coupled to depletion of intracellular calcium stores and influx of extracellular calcium in differentiated U937 cells. *Biochim. Biophys. Acta*. **1299**: 342–352.
34. Gailly, P. 1998.  $Ca^{2+}$  entry in CHO cells, after  $Ca^{2+}$  stores depletion, is mediated by arachidonic acid. *Cell Calcium*. **24**: 293–304.
35. Hirafuji, M., T. Ebihara, F. Kawahara, and M. Minami. 1998. Effect of docosahexaenoic acid on smooth muscle cell functions. *Life Sci.* **62**: 1689–1693.
36. Chow, S. C., and M. Jondal. 1990. Polyunsaturated free fatty acids stimulated an increase in cytosolic  $Ca^{2+}$  by mobilizing the inositol 1,4,5-trisphosphate-sensitive  $Ca^{2+}$  pool in T cells through a mechanism independent of phosphoinositide turnover. *J. Biol. Chem.* **265**: 902–907.
37. Magni, M. G., J. Meldolesi, and A. Pandiella. 1991. Ionic events induced by epidermal growth factor. Evidence that hyperpolarization and stimulated cation influx play a role in the stimulation of cell growth. *J. Biol. Chem.* **266**: 6329–6335.
38. Charlesworth, A., and E. Rozengurt. 1994. Thapsigargin and *tert*-butylhydroquinone induced synergistic stimulation of DNA synthesis with phorbol ester and bombesin in Swiss 3T3 cells. *J. Biol. Chem.* **269**: 32528–32535.
39. Kokoska, E. R., G. S. Smith, and T. A. Miller. 1998. Store operated calcium influx in human gastric cells: role of endogenous prostaglandins. *Surgery*. **124**: 429–437.